

Monitoring of Pollen Micromorphology of *Acanthophyllum* C. A. Mey. (Caryophyllaceae) Genus from Central Anatolian in Turkey

Ali Savaş BÜLBÜL*

Kader VARLIK

Bartın University, Faculty of Science, Department of Molecular Biology and Genetics, 74100, Bartın, Turkey

*Corresponding Author:

E-mail: asavasbulbul@gmail.com

Received: October 04, 2016

Accepted: December 18, 2016

Abstract

Pollen morphology of *Acanthophyllum* C. A. Mey (Caryophyllaceae) species in Turkey were investigated by light and scanning electron microscopy in this study. There are 5 species in Turkey belonging this genus: *Acanthophyllum mucronatum* C. A. Meyer, *Acanthophyllum acerosum* Sosn., *Acanthophyllum microcephalum* Boiss., *Acanthophyllum oppositiflorum* Aytaç, *Acanthophyllum verticillatum* (Willd.) Hand.-Mazz. The pollen grains of *Acanthophyllum* species are radially symmetrical and isopolar, oblate-spheroidal and prolate-spheroidal, operculate, and pantoporate, their pores have conical spinules on operculum. Pore numbers and diameter, exine thickness, pollen ornamentation and operculum ornamentation vary between *Acanthophyllum* species.

Keywords: *Acanthophyllum* C.A.Mey., Light Microscope, Morphology, Pollen, Scanning Electron Microscope

INTRODUCTION

Caryophyllaceae is one of large angiosperm families. It contains 86 genera and almost 2200 species. These species are distributed mainly Mediterranean region [1-3]. There are 32 genera which includes over 470 species in Turkey [4]. Conventionally, there are 3 subfamilies among Caryophyllaceae: Alsinoideae Burnett, Caryophylloideae Arn., and Paronychioideae A.St. [1]. *Acanthophyllum* C. A. Mey. is a member of Caryophyllaceae family, subfamily Caryophylloideae [5]. Genus *Acanthophyllum* contains approximately 60 species and is distributed in the Irano-Turanian region [1]. In Turkey, there are 5 species representing the genus: *Acanthophyllum acerosum* Sosn., *Acanthophyllum microcephalum* Boiss., *Acanthophyllum mucronatum* C.A.Mey., *Acanthophyllum verticillatum* (Willd.) Hand.-Mazz., *Acanthophyllum oppositiflorum* Aytaç (endemic) [6, 7].

The Caryophyllaceae family has medicinal and ornamental properties which brings an importance [8]. This genus includes small pulvinate shrubs with spiny leaves that

grow in open habitats [9].

In this study, pollen morphology of 5 taxa, one of which is endemic belonging to Turkish *Acanthophyllum* C. A. Mey. were investigated by using light microscope (LM) and scanning electron microscope (SEM). The different and similar characteristics of the species were determined.

MATERIALS and METHODS

Specimens were collected in 2014 and 2016 from the localities given in "Table 1". The palynological investigations are based on the herbarium specimens obtained from Gazi University Faculty of Science Herbarium. Pollen slides were prepared according to Wodehouse (1935) from mature flowers. At least 50 pollen grains for each species were examined in light microscope and micrographs were taken by Leica DM 750 digital imaging system. Different morphological characters of pollen parts such as pollen diameter, pore diameter, distance between two pores and exine thickness were measured by means of light microscopy.

Table 1. List of investigated taxa and localities.

Collector Number	Species	GPS	Altitude	Collection Date	Habitat	Phytogeographic Region	Localities
MA 5693	<i>Acanthophyllum verticillatum</i> C.A.Mey.	39°09'29.6"N 39°29'01.7"E	970	24.7.2014	Step, Hillside	Irano-Turanian	B7Tunceli: Tunceli, Between Tunceli - Ovacık (Valley Munzur) 10. km
MA 5693	<i>Acanthophyllum oppositiflorum</i> Aytaç	38°53'41.2"N 39°14'52.9"E	875	13.8.2014	Step	Irano-Turanian Endemic	B7Tunceli: Pertek, Between Pertek - Çemişgezek 9. km, 2.5 km before Singeç Bridge
MA 6758	<i>Acanthophyllum mucronatum</i> C.A.Mey.	37° 47' 47.5"N 44° 05' 04.2"E	1825	31.7.2016	Naked Hillside	Irano-Turanian	C10Van: Başkale, Hakkari Provincial Border, 33 km to Hakkari from Başkale, Right-hand Hills
MA 6758	<i>Acanthophyllum microcephalum</i> Boiss.	37° 47' 47.5"N 44° 05' 04.2"E	1825	31.7.2016	Naked Hillside	Irano-Turanian	C10Van: Başkale, Hakkari Provincial Border, 33 km to Hakkari from Başkale, Right-hand Hills
MA 6758	<i>Acanthophyllum acerosum</i> Sosn.	37° 47' 47.5"N 44° 05' 04.2"E	1825	31.7.2016	Naked Hillside	Irano-Turanian	C10Van: Başkale, Hakkari Provincial Border, 33 km to Hakkari from Başkale, Right-hand Hills

For SEM, pollen grains were directly mounted on stubs and covered with gold and surface ornamentations of pollen grains were examined in detail with Jeol Tescan MAIA3 XMU model electron microscope in Bartın University Central Research Laboratory.

The pollen morphology terminology adopted by Fagri and Iversen (1989), Punt et al.(2007), Moore et al. (1991) was used[10-12]. Shape classification follows Erdtman (1969), based on P/E ratio in "Table 2-3". [13].

Table 2. Pollen morphology of species *Acanthophyllum* M: median, V: variation, S: standart deviation.

	Spheroidal	<i>A. acerosum</i>	<i>A. microcephalum</i>	<i>A. mucronatum</i>	<i>A. oppositiflorum</i>	<i>A. verticillatum</i>
Polar axes (µm)	M	35.03	30.11	30.32	28.95	33.9
	V	(28.54-45.82)	(28.87-34.16)	(26.24-33.87)	(25.81-30.65)	(26.78-42.17)
	S	± 2.95	± 2.11	± 1.65	± 1.48	± 2.16
Equatorial axes (µm)	M	35.08	29.28	30.38	27.59	34.17
	V	(27.74-46.82)	(26.39-31.44)	(27.97- 32.68)	(25.48-30.01)	(25.66-42.89)
	S	± 2.23	± 1.68	± 1.38	± 1.12	± 2.57
Exine thickness (µm)	M	1.96	2.14	1.79	1.88	1.86
	V	(1.43-2.29)	(1.92-2.37)	(1.40 -2.11)	(1.47-2.39)	(1.47-2.38)
	S	± 0.20	± 0.19	± 0.17	± 0.23	± 0.2
Intine thickness (µm)	M	0.51	0.57	0.51	0.52	0.52
	V	(0.38-0.66)	(0.51-0.64)	(0.41-0.64)	(0.44-0.64)	(0.41-0.69)
	S	± 0.07	± 0.05	± 0.07	± 0.05	± 0.06
Pore length (µm)	M	4.43	3.57	5.43	4.13	4.24
	V	(2.43-7.48)	(2.42-4.16)	(3.49-7.49)	(2.84-5.12)	(2.11-6.65)
	S	± 1.5	± 0.78	± 0.88	± 0.61	± 0.16
Pore width (µm)	M	4.46	3.34	5.55	4.12	4.52
	V	(2.74-8.75)	(3.14-3.69)	(3.97-7.76)	(3.09-5.32)	(2.76-7.53)
	S	± 1.64	± 0.24	± 0.83	± 0.54	± 1.33
Pore distance (µm)	M	7.19	4.13	8.12	7.65	6.94
	V	(4.38-11.10)	(3.07-4.2)	(5.97-10.39)	(4.96-10.29)	(3.71-9.17)
		± 1.91	± 0.17	± 1.09	± 1.06	± 1.32

Table 3. Pollen morphology of species *Acanthophyllum*

	<i>A. acerosum</i>	<i>A. microcephalum</i>	<i>A. mucronatum</i>	<i>A. oppositiflorum</i>	<i>A. verticillatum</i>
Pore number	7-13	7-11	6-9	6-13	6-12
Spinules on operculum	8-10	8-10	10-15	8-14	6-9
Spinules (100 µm ²)	55-60	45-50	55-58	90-95	50-55
Plg/plt	0.92	1.06	0.97	1	0.94
Operculum shape	Oblate-spheroidal	Prolate- spheroidal	Oblate-spheroidal	Spheroidal	Oblate- spheroidal
Pollen shape	Oblate- spheroidal	Prolate- spheroidal	Oblate-spheroidal	Prolate- spheroidal	Oblate- spheroidal
Pollen dimension(P/E)	0.99	1.02	0.99	1.01	0.99
Pollen class	Pantoporate	Pantoporate	Pantoporate	Pantoporate	Pantoporate
Pollen aperture type	Tectate	Tectate	Tectate	Tectate	Tectate
Exine structure	Spinulose-punctate	Spinulose-punctate	Spinulose-punctate	Spinulose-punctate	Spinulose-punctate
Pollen ornamentation	Scabrat-punctate	Scabrat-punctate	Scabrat-punctate	Scabrat-punctate	Scabrat-punctate

RESULTS

Acanthophyllum acerosum Sons.

Pollen grains were radially symmetrical and isopolar. Polar axes were 35.03 µm (28.54-45.82µm), equatorial axes were 35.08 µm (27.74-46.82µm). Pollen shape was P/E:0.99, oblate-spheroidal. The pollen grains were operculate and pantoporate, generally with 7-13 porate. Pores had polar length of 4.43(2.43-7.48) µm and equatorial length 4.46(2.74-8.75) µm. plg/plt: 0.92, oblate-spheroidal. There were 8-10 conical spinules on operculum. Distance between two pori was 7.19 (4.38-11.10)µm. The exine was tectate and 1.96 (1.43-2.29) µm in thickness. Intine thickness was 0.51(0.38-0.66) µm. Exine structure was spinulose-punctate. Polen ornamentation was scabrate-punctate. There were 55-60 spinules in 100 µm² area.

Acanthophyllum microcephalum Boiss

Pollen grains were radially symmetrical and isopolar. Polar axes was 30.11 (28.87-34.16) µm, equatorial axes was 29.28 (26.39-31.44) µm. Pollen shape was P/E:1.02, prolate-spheroidal. The pollen grains were operculate and pantoporate, generally 7-11 porate. Pores polar length of 3.57 (2.42-4.16)µm, equatorial length 29.28 (26.39-31.44) µm (plg/plt):1.06, prolate-spheroidal. There were 8-10 conical spinules on operculum. Distance between two pori was 4.13 (3.07-4.2)µm. The exine was tectate and 2.14(1.92-2.37) µm in thickness. Intine thickness was 0.57 (0.51-0.64) µm. Exine structure was spinulose-punctate. Polen ornamentation was scabrate-punctate. There were 45-50 spinules in 100 µm² area.

***Acanthophyllum mucronatum* C. A. Mey.**

Pollen grains were radially symmetrical and isopolar. Polar axes was 30.32 (26.24-33.87) μm , equatorial axes was 30.38 (27.97- 32.68) μm . Pollen shape was P/E:0.99, oblate-spheroidal. The pollen grains were operculate and pantoporate, generally 6-9 porate. Pores had polar length of 5.43 (3.49-7.49) μm and equatorial length 5.55 (3.97-7.76) μm (plg/plt):0.97, oblate-spheroidal. There were 10-15 conical spinules on operculum. Distance between two poriwass 8.12 (5.97-10.39) μm . The exine was tectate and 1.79 (1.40 -2.11) μm in thickness. Intine thickness was 0.51 (0.41-0.64) μm . Exine structure was spinulose-punctate. Pollen ornamentation was scabrate-punctate. There were 53-58 spinules in 100 μm^2 area.

***Acanthophyllum verticillatum* (Willd.) Hand.-Mazz.**

Pollen grains were radially symmetrical and isopolar. Polar axes was 33.9 (26.78-42.17) μm , equatorial axes was 34.17 (25.66-42.89) μm . Pollen shape was P/E:0.99, oblate-spheroidal. The pollen grains were operculate and pantoporate, generally 6-12 porate. Pores had polar length of 4.24 (2.11-6.65) μm and equatorial length 4.52 (2.76-7.53) μm . (plg/plt):0.94, oblate-spheroidal. There were 8-14 conical spinules on operculum. Distance between two poriwass 6.94 (3.71-9.17) μm . The exine was tectate and 1.86 (1.47-2.38) μm in thickness. Intine thickness was 0.52 (0.41-0.69) μm . Exine structure was spinulose-punctate. Pollen ornamentation was scabrate-punctate. There were 50-55 spinules in 100 μm^2 area.

***Acanthophyllum oppositiflorum* Aytaç**

Pollen grains were radially symmetrical and isopolar. Polar axes was 28.95 (25.81-30.65) μm , equatorial axes was 27.59 (25.48-30.01) μm . Pollen shape was P/E:1.01, prolate-spheroidal. The pollen grains were operculate and pantoporate, generally 6-13 porate. Pores had polar length of 4.13 (2.84-5.12) μm and equatorial length 4.12 (3.09-5.32) μm . (plg/plt):1, spheroidal. There were 6-9 conical spinules on operculum. Distance between two poriwass 7.65 (4.96-10.29) μm . The exine was tectate and 1.88 (1.47-2.39) μm in thickness. Intine thickness was 0.52 (0.44-0.64) μm . Exine structure was spinulose-punctate. Pollen ornamentation was scabrate-punctate. There were 90-95 spinules in 100 μm^2 area.

The pollen grains of *Acanthophyllum* were radially symmetrical and isopolar. Shape was oblate-spheroidal and prolate-spheroidal. Polar axes was 25.81-45.82 μm and the equatorial axes was 28.48-46.82 μm (Table 2, Figures 1-2). *A. acerosum*, *A. mucronatum* and *A. verticillatum* pollen shape was oblate-spheroidal while *A. microcephalum* and *A. oppositiflorum* pollen shape was prolate-spheroidal. *Acanthophyllum acerosum* had the biggest pollen grains with 35.03 μm polar axis and 35.08 μm equatorial axis lengths, while *Acanthophyllum oppositiflorum* had the smallest-sized pollen grains with 28.95 μm polar axes and 27.59 μm equatorial axis lengths.

The pollen grains were operculate and pantoporate, generally 6-13 porate. Pores length was 2.11-7.49 μm pore width of 2.74-8.75 μm . *A. mucronatum* had the largest pollen grains with 5.43 μm pore length and 5.55 μm pore widths while *A. microcephalum* had the smallest pollen pore 3.57 μm pore length and 3.34 μm pore width. *A. mucronatum* had the biggest pore distance of two pori 8.12 μm while *A. microcephalum* had the smallest pore distance of two pori 4.13 μm . Polar length/equatorial length (Plg/Plt) ratio was

between 0.92 μm and 1.06 μm . *A. acerosum*, *A. mucronatum* and *A. verticillatum* had oblate –spheroidal operculum shape while *A. microcephalum* prolate-spheroidal and *A. oppositiflorum* was spheroidal.

The exine structure was scabrate-punctate and 1.79-2.14 μm in thickness. Intine thickness ranged between 0.51-0.57 μm . There were spinules in the form of conical that vary from 6-15 in number over the operculum, according to the species.

DISCUSSION

According to Shamsabadi et al. (2013), palynological studies for 11 species of the *Acanthophyllum* sect. *Oligosperma* in northeastern Iran including *A. borsczowii* Litw., *A. speciosum* Rech. f. & Schiman-Czeika, *A. korshinskyi* Schischk., *A. pachystegium* Rech. f., *A. adenophorum* Freyn., *A. lilacinum* Schischk., *A. brevibracteatum* Lipsky., *A. diezianum* Hand.-Mzt., *A. laxiusculum* Schiman-Czeika, *A. squarrosom* Boiss., and *A. heratense* Schiman-Czeika [14]. Pollen grains were radially symmetrical and spheroidal and their type was pantoporate. The grains had median size (23-31 μm), which was different from 24.5 μm polar axes and 23.8 μm equatorial axis in *A. borsczowii* (smallest) to 30.7 μm polar axes and 30.3 μm equatorial axis in *A. adenophorum* (largest). While *Acanthophyllum* grains have median size (27-35 μm), this character is different from 28.95 μm polar axes and 27.59 μm equatorial axis in *Acanthophyllum oppositiflorum* (smallest) to 35.03 μm polar axes and 35.08 μm equatorial axis in *Acanthophyllum acerosum* (largest). Pollen ornamentation was scabrate-punctate and exine structure was spinulose-punctate. Our results are similar with M. Mahmoudi Shamsabad, H. Ejtehadi, J. Vaezi & F. Memariani results but our results include more knowledge. Also this is the first study to determine polar length/equatorial length (Plg/Plt) ratio is between 0.92 μm and 1.06 μm , number of spinules on operculum is between 6-15, *Acanthophyllum* species.

Except these studies that investigated pollen morphology of *A. borsczowii* Litw., *A. speciosum* Rech. f. & Schiman-Czeika, *A. korshinskyi* Schischk., *A. pachystegium* Rech. f., *A. adenophorum* Freyn., *A. lilacinum* Schischk., *A. brevibracteatum* Lipsky., *A. diezianum* Hand.-Mzt., *A. laxiusculum* Schiman-Czeika, *A. squarrosom* Boiss., and *A. heratense* Schiman-Czeika, there are no reported data about pollen morphology of other *Acanthophyllum* species.

Pollen morphology of some *Velezia* L. species has been studied by Poyraz, Ataslar (2010) and they reported the pollen grains of *V. tumicoidea*, *V. hispida*, *V. quadridentata*, *V. pseudorigida* and *V. rigida* has 11-14 porate while *Acanthophyllum* has 6-13 pores. *Velezia* L. has oblate-spheroidal shape with the polar axes of 27.92-39.64 μm and the equatorial axes of 28.21-39.80 μm . *Acanthophyllum* has oblate-spheroidal and prolate-spheroidal shape. Polar axes of 25.81-45.82 μm and the equatorial axes of 28.48-46.82 μm . *Acanthophyllum* has bigger pollen grains than *Velezia*. Pollen grains of *Velezia* species are oblate-spheroidal, operculate, polypantoporate with scabrate-microperforate or scabrate-foveolate ornamentation [15]. *Acanthophyllum* mainly differs from these taxa with exine structure is scabrate-punctate.

El Naggari (2004) reported the pollen grains of *Gypsophila pilosawere* monade, apolar, hexagonal-polyhedral or spheroidal, 18-19 x 18-19 μm . Aperture was pantoporate. Pores were circular, 3-4 μm and tectum




Figure 1. LM photos of pollen grains of *Acanthophyllum* species: *A. acerosum* A-B, *A. microcephalum* C-D, *A. mucronatum*. E-F, *A. verticillatum* G-H and *A. oppositiflorum* I-J.




Figure 2. SEM photos of pollen grains of *Acanthophyllum* species: A. *acerosum* A-B, *A. microcephalum* C-D, *A. mucronatum*. E-F, *A. verticillatum* G-H and *A. oppositiflorum* I-J.

was perforate [16]. *Acanthophyllum* mainly differs from these taxa with exine structure is scabrat-punctate and *Acanthophyllum* grains size is medium (27-35 µm). Also *Acanthophyllum* is spheroidal.

Yıldız, Minareci (2008) reported the pollen grains of *Silene urvillei* Schott. werespheroidal, numbers of pores were 15-24. The exine ornamentation was microechinate [17]. *Acanthophyllum* differs from these taxa with exine ornamentation is scabrat-punctate and numbers of pores are 6-13.

Yıldız, Dadandı (2009) reported the pollen grains of *Silene cirpicii* were spheroidal, ornamentation was microechinate, microperforate, number of pores per pollen was 23-28 µm, number of granules on pore 9-15. *Acanthophyllum* differs from these taxa with exine ornamentation is scabrat-punctate and numbers of pores are 6-13 [18].

Ataşlar, Erkara, Tokur (2009) were performed palynological studies for all 12 *Gypsophila* taxa which were polyporate and spheroidal. *Gypsophila* pollen grains ranged in size from 23.79 to 32.84 µm in non-acetolysed pollen and from 19.48 to 25.94 µm in acetolysed pollen. *Acanthophyllum* polar axes of 25.81-45.82 µm and the equatorial axes of 28.48-46.82 µm. The exine sculptures were clavate-microperforate ornamentation and granulate-microechinate - microperforate ornamentation [19]. The major difference of *Acanthophyllum* from this species is the exine ornamentation which is only scabret-punctate.

Pollen morphology of some *Dianthus* species has been studied by Vural (2008). According to that study, pollen grains of *D. Aytachii* were periporate, long axis 34.9 µm (32-37 µm), short axis 35.1 µm (33-38 µm). Pollen shape was oblate-spheroidal. Pore number was between 8-14, polar length/equatorial length (Plg/Plt):0.97, oblate-spheroidal, 5.5mm (4-6.5 mm) diameter, operculate with four to six echinae [20]. The major difference of *Acanthophyllum* from this species is the pantoporatelong axes of 25.81-45.82 µm and the short axes of 28.48-46.82 µm. Number of pores are 6-13. Polar length/equatorial length (Plg/Plt) ratio is between 0.92 µm and 1.06 µm. *A. acerosum* *A. mucronatum* and *A. verticillatum* have oblate -spheroidal operculum shape while *A. microcephalum* Prolate-spheroidal and *A. oppositiflorum* is spheroidal. The exine structure is scabrat-punctate.

Bozchaloyi and Keshavarzi (2014) determined some pollen features of *Stellaria* species using SEM. According to that study, pollen shapes in polar and equatorial views were spheroid. Average size of pollen grains in the *Stellaria* species and related genera varied from 22.5 µm (*S. alsinoides*) to 42µm (*Myosoton aquaticum*). Pore diameters varied between 4.45 µm and 7.63 µm. The average exine thickness was between 1.37 µm (*S. alsinoides*) and 2.61 µm (*S. holostea*) [21]. *Acanthophyllum* mainly differs from these taxa with long axes of 25.81-45.82 µm and the short axes of 28.48-46.82 µm. *Acanthophyllum* has 1.79-2.14 µm exine thickness.

Ataşlar (2003) reported the pollen grains of *Saponaria kotschy* Bois. hasspheroidae, 36 µm diameter, periporate, 12-pored. Operculum has 3 µm diameters with spinulate. The exine sculpture is granulatae microechinatae-microperforatae [22]. *Acanthophyllum* has 6-13 pores and pollen grains of *Acanthophyllum* long axes of 25.81-45.82 µm and the short axes of 28.48-46.82 µm.

Çinbilgel, et al. (2007) described the pollen grains of *Saponaria pamphylica* Boiss. & Heldr as pollen grains are spheroidae, with isopolar symmetry, size middle, average

42.795 µm diameter, periporate; numbers of pores are 11-12, average diameter 5.9 µm. Operculum is with 10-12 spinulate. The exine ornamentation is spinulatae (spinulosus) and microperforatae. [23] *Acanthophyllum* mainly differs from these taxa with pollen shape is oblate-spheroidal or prolate spheroidal.

In conclusion, according to the results of this study, pollen ornamentation, operculum ornamentation of pollen grains, pore numbers and diameter, exine and intine thickness are helpful characters in determining *Acanthophyllum* species.

Acknowledgments

I would like to thank to Z. Aytaç from the Herbarium of Gazi University (GAZI) and to I. Demir from Van (Turkey).

REFERENCES

- [1] Bittrich V. 1993. Caryophyllaceae, in Flowering Plants· Dicotyledons. Springer. 206-236.
- [2] Heywood VH. 1998. Flowering plants of the world, 3rd edition. Oxford University Press, Oxford.
- [3] Fior S et al. 2006. Molecular phylogeny of the Caryophyllaceae (Caryophyllales) inferred from chloroplast matK and nuclear rDNA ITS sequences. American journal of botany. 93(3):399-411.
- [4] Yıldız K. 2002. Seed morphology of Caryophyllaceae species from Turkey (north Anatolia). Pak J Bot. 34:161-171.
- [5] Shamsabad MM et al. 2013. Anatomical and pollen characters in *Acanthophyllum ca mey.* (caryophyllaceae) from northeast of Iran. Iranian Journal of Botany. 19(1):107-118.
- [6] Aytaç Z. 2001. A new species of *Acanthophyllum* (Caryophyllaceae) from central Anatolia, Turkey. Nordic Journal of Botany. 21(3):263-266.
- [7] Davis PH. 1965. Flora of Turkey and the east Aegean islands. Vol 10. Edinburgh University Press.
- [8] Guha Bakshi D. 1984. Flora of Murshidabad District, West Bengal, India.
- [9] Zohary M. 1973. Geobotanical foundations of the Middle East.
- [10] Faegri K. and Iversen J. 1989. Textbook of pollen analysis. 4th Edition. Wiley. New York.
- [11] Punt W et al. 2007. Glossary of pollen and spore terminology. Review of Palaeobotany and Palynology. 143(1):1-81.
- [12] Moore PD, Webb JA and Collison ME. 1991. Pollen analysis. Blackwell scientific publications.
- [13] Erdtman G. 1969. Handbook of palynology: Morphology, taxonomy, ecology. An introduction to the study of pollen grains and spores. Hafner.
- [14] Mahmoodi Shamsabadi M et al. 2013. Anatomical and pollen characters in *Acanthophyllum* CA Mey. (Caryophyllaceae) from northeast of Iran. Iranian Journal of Botany. 19.
- [15] Poyraz İE and Ataşlar E. 2010. Pollen and seed morphology of *Velezia* L. (Caryophyllaceae) genus in Turkey. Turkish Journal of Botany. 34(3):179-190.
- [16] El Naggat S. 2004. The seed coat and pollen morphology of *Gypsophila pilosa* (Caryophyllaceae). Flora Mediterranea. 14:109-114.
- [17] Yıldız K and Minareci E. 2008. Morphological, anatomical, palynological and cytological investigation on *Silene urvillei* Schott. (Caryophyllaceae). Journal of Applied Biological Sciences. 2(2):41-46.
- [18] Yıldız K and Dadandı MY. 2009. *Silene cirpicii*

(Caryophyllaceae), a new species from Turkey. in Annales Botanici Fennici. BioOne.

[19] Ataşlar E, Erkara İP and Tokur S. 2009. Pollen morphology of some *Gypsophila* L.(Caryophyllaceae) species and its taxonomic value. Turkish Journal of Botany.33(5):335-351.

[20] Vural C. 2008. A new species of *Dianthus* (Caryophyllaceae) from mount Erciyes, central Anatolia, Turkey. Botanical Journal of the Linnean Society. 158(1):55-61.

[21] Bozchaloyi SE and Keshavarz M. 2014. Micro-and macro-morphological study of *Stellaria* (Caryophyllaceae) and its closest relatives in Iran. Phytologia Balcanica: International Journal of Balkan Flora and Vegetation.20(2):179-197.

[22] Ataşlar E. 2004. Morphological and anatomical investigations on the *Saponaria kotschyi* Boiss. (Caryophyllaceae). Turkish Journal of Botany. 28(1-2):193-199.

[23] Çinbilgel İ, Karadeniz A and Gökceoğlu M. 2007. Morphological and anatomical study on endemic *Saponaria pamphylica* Boiss. & Heldr.(Caryophyllaceae). J. Appl. Biol. Sci. 1:19-25.