

# Micro-morphological Changes of Sandy Soils Through the Application of Compost Manure

M. M. Youssef<sup>1</sup>\* and Kh. M. Fathy<sup>2</sup>

<sup>1</sup>Soils & Water Use Dept., National Research Centre (NRC), Cairo, Egypt

<sup>2</sup>National Research Centre (NRC), El-Behouth ST., 12622, Cairo, Egypt

\*Corresponding Author  
e-mail: moniermorad@yahoo.com

Received: February 12, 2008  
Accepted: May 30, 2008

## Abstract

Compost material not only improves the structure of fine-textured soils but creates the structure coarse-textured soils as well. Compost was applied at the rates of 0 (Control) and 20 ton/ feddan /year to sandy soils (Typic Torripsamment) in order to investigate its potential for improving the soil properties. The experiment was conducted for two years in Wadi El-Natrun, El-Behaira Governorate, Egypt. The chemical and physical properties of the soils were affected directly by the compost application. Results from this study indicate that the soil structure and the cation exchange capacity were improved. Increasing in organic matter content and reduction in calcium carbonate content were due to the compost which is rich in humic acid. Thin sections of treated sandy samples were examined by polarized microscope. Macro and micro soil morphology were improved, especially the soil structure. Compared with the control treatment, the treated soil with compost had better aggregation in the surface layer due to change in fabric of ground mass and the altering changing of related distribution (C/F) from enaulic to chitonic and partly gefuric. The dominant voids (large vughs and chambers) were modified to simple packing voids. Organic materials can be seen as coatings on pore walls.

**Key words:** Compost, micromorphology, thin section.

## INTRODUCTION

Compost has been used extensively in reclamation of marginal and low quality soils. It has the unique ability to improve soil properties and the growing media physically (structurally), chemically (nutritionally) and biologically. Addition of compost to soil improves soil structure and lowers bulk density. Composting materials can increase macro aggregation and rhizospheric aggregate stability [1, 2]. The effects of compost additions on soil structure may be short-lived although they are generally positive on structural [3]. In the field [4] applied compost derived from crop residue alone and with other materials. They found that the improvements in soil physical properties (soil bulk density and water content) were obtained for compost treatments alone. Composts alone influenced soil C: N ratio, substantially increased soil organic carbon and soil microbial biomass carbon concentrations. Recently, [5] studied the effect of composted yard waste on the movement water in a sandy soil found that water and P retention in the soil were increased. [6] Found that after three years of compost application, organic matter content, water retention capacity, available water, and infiltration rate were significantly increased, whereas bulk density and aggregate instability index were decreased. The chemical properties of the soil were affected directly by the amendment compost. [7] Reported that in samples from the field trial, soil total C,N,P and Olsen P were increased markedly with increasing the compost application rate. Cation exchange capacity, exchangeable cation as well as total-extractable and EDTA-extractable metals (Cd, Cr, Cu, Ni, Pb and Zn) were also elevated. However, the total Cu reached to the allowable limit in bio-solids compost amended soil.

The physical properties of the amended soils were improved in all cases as far as the saturated and unsaturated hydraulic conductivity, water retention capacity, bulk density, total porosity, pore size distribution, soil resistance to penetration, aggregation and aggregate stability, were concerned, [8]. In this respect, [9], reported that the addition of organic materials of various origins to soil have been one of the most common rehabilitation practices to improve soil physical properties. [10] concluded that compost can help the formation of large number of water stable aggregates, through links between small particles, strong enough to stand the dispersing action of water. [11] Found that compost improved soil aggregation and increasing of water holding capacity. With respect to total porosity, [12] declared increasing the values of it as a result of applied compost. The value reached its maximum for banana compost to 50.91 % in the surface layer to the depth of 15cm. [13] reported that compost treatments increased micro pores of sandy soil which resulted in an increase in soil available water and decreased in large pore space. [14] Stressed on the decrease of soil bulk density with compost addition. Finally [15] reported that, both compost and manure improved soil pore system characteristics.

Therefore, the aim of this work is to study the effect of compost application to sandy soils through the examination of soil thin sections and investigation of some chemical characteristics.

## MATERIALS and METHODS

The compost was produced from manure, by Bio-Pianta Company, Egypt. Some properties of the used compost in this

study were presented. Field experiment was conducted between the years 2005 and 2007 in a farm at Wadi El-Natrun which is located 30° 18' 1" N and 30° 20' 5" E and lies some 100 km north west to Cairo, Egypt. In this experiment compost was applied at the rates of 0 (control), 20 ton/feddan/year to sandy soils (feddan = 4200m<sup>2</sup>). The compost incorporated by rotilling to the top 15 cm of the soil before seedling. Disturbed and undisturbed soil samples were taken from two different depths (0-15 and 15-30cm) of the treated and untreated soils after two years. Soil samples were analyzed for grain size distribution by dry sieving [16]. Soil pH and electrical conductivity (EC) were determined according to [17]. Some soil chemical properties analyzed according to [18]. Thin sections were examined using polarizing microscope at 25 times magnification to observe microstructure according to terminology developed by [19 and 20].

## RESULTS and DISCUSSION


Grain size distribution and chemical properties of the soil of the current experiment are presented in table 1. Various analyses showed that the soil is sandy, non-saline, low in carbonate content, exchange cation and poor in organic matter content. Accordingly the soil is classified at the subgroup level as Typic Torripsamment, [21].

The used compost in this experiment was rich in organic matter, (Table 2).

Moreover, this application of compost on sandy soil is resulted in some improvements in the soil treated with compost especially for the 0-15 cm layer. The pH values and calcium carbonate content slightly decreased, compared to the control. Similarly, [22] observed that the application of various organic materials decreased pH values since organic and inorganic acids formed when organic matter decomposed in the soil. Organic matter content and Cation exchange capacity of the

compost treated soil are markedly increased in the surface 30cm of the soil as it is compared with the control treatment. Similar results were reported by several authors, [14 and 23]. Due to the higher salt content of the used compost than the soil of the experiment, the total soluble salts of that soil is slightly increased in two years. However, the use of such compost for long period may cause a considerable increase of the soluble salt content of the treated soil. It is worth to mention that the morphological investigation of the soil samples after two years revealed that the soil colour became darker than that of the soil under 0 (control) compost applications. Most sandy soils always have a loose structure as a result of the soil aggregate absence. Compost is an important factor in the formation of soil aggregates and consequently their stability. Microscopic examination of thin sections revealed that the untreated soil contains groundmass which is dominated by coarse grains of mostly quartz with lots of air space and the related distribution (C/F) is enaulic with large vughs and chambers are dominated (figure1).

On the other hand, the compost treated soil had better aggregation in the surface layer. The compost fills the gaps between soil particles and the related distribution changed to chitonic and locally gefuric with packing voids (figure2).



**Figure1.** The groundmass is dominated by quartz grains and large chambers at 0- 15cm depth of sandy untreated soil. (X25).

**Table 1.** Some physical and chemical properties of the soil studied.

Location	Depth Cm	Grain size distribution				Texture	pH 1:2.5	Ec ds/m	CaCO <sub>3</sub> %	OM %	CEC Cmol kg <sup>-1</sup>
		*C. sand %	*F. sand %	Silt %	Clay %						
Wadi El-Natrun	0-15	50.1	40.6	4.2	5.1	Sand	8.35	1.23	3.24	0.51	5.32
	15-30	45.7	44.8	3.8	5.7	Sand	8.49	1.41	2.72	0.42	5.13

\*C. sand: Coarse sand. \*F. sand: Fine sand.

**Table 2.** Some chemical characteristics of compost manure.


pH	Ec ds/m	C/N ratio	Organic C %	N %	P %	K %
8.1	4.15	1:12	30.44	1.53	0.29	1.13

The effect of compost application on soil texture and some chemical properties after two years were presented in table 3.

**Table 3.** Some physical and chemical properties of the soil studied after two years of compost application.


Depth cm	Grain size distribution					pH 1:2.5	EC ds/m	CaCO <sub>3</sub> %	OM %	C.E.C Cmol Kg <sup>-1</sup>
	C. Sand %	F. Sand %	Silt %	Clay %	Texture					
0-15	43.4	38.7	6.4	11.5	L.S*	8.28	1.28	3.11	1.01	7.76
15-30	42.8	40.6	5.8	10.8	L.S*	8.41	1.47	2.62	0.95	7.23

\* L.S Loamy sand




**Figure 2.** The reduction of porosity in treated soil at depth 0-15cm. (X25).

The soil-binding properties of compost are due to its humus content. Humus is a stable residue resulting from a high degree of organic matter decomposition. The constituents of the humus act as soil “glue”, holding soil particles together, form soil aggregates, which lead to improve the soil’s ability to hold moisture (figure3).



**Figure 3.** Small and large of aggregates at 15-30cm depth of sandy treated soil. (X25).

The structure of sandy soil not only improves through the aggregate formation but also by coating the pore walls (figure 4). Therefore it will increase the water holding capacity and retain more nutrients in sandy soils. This finding is important in understanding the influence of compost application to coarse textured soils.



**Figure 4.** Pore walls are coated by the compost materials, at depth 15-30cm.(X25).

This finding is important in understanding the influence of compost application to coarse textured soils.

## CONCLUSION

Compost has a unique role in physical (soil structure), chemical (nutritionally), and biological (soil microbes). After two years organic matter content was increased and consequently humic acids content which helps in dissolving minerals in the soil resulting in more available nutrients to the plant. Compost can also improve the cation exchange capacity of soils, enabling them to retain nutrients longer.

The study of thin sections of the treated soil with compost clarified that the incorporating compost into sandy soils improves soil structure by changing type of voids and form of the related distribution (C/F concept) as well as by filling the gaps and coating the pore walls. That will increase the soil water holding capacity and adsorption capacity of nutrients, which are the main factors to be improved through reclamation of the sandy soils.

## REFERENCES

- [1] De Leon-Gonzalez, F., Hernandez-Serrano, M., Etchevers, J., Payan-Zelaya, F. and Ordaz-Chaparro, V. 2000. Short-term compost effect on macro aggregation in a sandy soil under low rainfall in the valley of Mexico. *Soil Tillage Res.* (56): 213–217.
- [2] Caravaca, F., Hernandez, T., Garcia, C. and Roldan, A. 2002. Improvement of rhizosphere aggregate stability of afforested semiarid plant species subjected to mycorrhizal inoculation and compost addition. *Geoderma* (108): 133–144.
- [3] Debosz, K., Petersen, S., Kure, L. and Ambus, P. 2002b. Evaluating effects of sewage sludge and household compost on soil physical, chemical and microbiological properties. *Appl. Soil Ecol.* (19): 237–248.
- [4] Lynch, D., Voroney, R. and Warman, P. 2005. Soil Physical Properties and Organic Matter Fractions Under Forages Receiving Composts, Manure or Fertilizer. *Compost Science & Utilization*. 13 (4): 252- 261.
- [5] Pandey, C. and Shukla, S. 2006. Effects of Composted Yard Waste On Water Movement in Sandy Soil. *Compost Science & Utilization*. 14(4): 252-259.
- [6] Tsadilas, C., Mitsios, I. and Golia, E. 2005. Influence of Biosolids Application on Some Soil Physical Properties. *Communications in Soil Science and Plant Analysis*. (36): 709 – 716.
- [7] Speir, T., Horswell, J., Schaik, A., McLaren, R. and Fietje, G. 2004. Composted biosolids enhance fertility of a sandy loam soil under dairy pasture. *Biology and Fertility of Soils*. 40(5): 349-358.
- [8] Aggelides, S. and Londra, P. 2000. Effects of compost produced from town wastes and sewage sludge on the properties of a loamy and a clay soil. *Bioresource Technology*. (71): 3253-259.
- [9] Celik, I., Ortas, I. and Kilic, S. 2004. Effects of compost, mycorrhiza manure and fertilizer on some physical properties of a Chromoxeret soil. *Soil and Tillage Research*. (78): 59-67.
- [10] Rizzi, L., Petruzzelli, G. and Vigna, G. 2004. Soil physical changes and plant availability of Zn and Pb in a treat ability test of phyto-stabilization. *Chemosphere*. (57): 1039-1046.

- [11] Shanjida, K. and Sarwar, K. 2002. Effect of water-hyacinth compost on physical, physico-chemical properties of soil and on rice yield. *Pakistan J. agronomy*. 1(2-3): 64-67.
- [12] Wanas, Sh. And Omran, W. 2006. Advantages of applying various compost types to different layers of sandy soil: 1-Hydro-physical properties. *J. App. Sci Rec.*, 2(12): 1298-1303.
- [13] Wanas, Sh. 2002. A compression study about the influence of inorganic fertilizers and organic composts on the structure and water characteristics of sandy soil cultivated with cowpea plants. *Egypt. J. App. Sci.*, 17(4): 362-375.
- [14] Tejada, M., Garcia, C., Gonezalez, J. and Hernanedz, L. 2006. Use of organic amendments as a strategy for saline soil remediation: Influence on the physical, chemical and biological properties of soil. *Soil Biology & biochemistry*.(38): 1413-1421.
- [15] Pagliai, M., Vignozzi, N. and Pellegrini, S. 2004. Soil structure and the effect of management practices. *Soil & Tillage Research*.(79):131-143.
- [16] Soil Conservation Service. 1984. Procedures for Collecting Soil Samples and Methods of Analysis for Soil Survey. USDA-SCS. Soil Surv. Invest. Rep.No.1. U.S. Govt. Print. Office, Washington, DC.
- [17] Smith, J. and Doran, J. 1996. Measurement and use of pH and electrical conductivity for soil quality analysis. p.169-185. In J.W. Doran and A.J. Jones (ed). *Methods of assessing soil quality*. SSSA Spec. Publ. SSSA, Madison. WI.
- [18] Black, C. A., Evans, D.D., White, J. I., Ensminger, L.E. and Clark, F. E. 1982. *Methods of Soil Analysis*. Amer. Soc. Agronomy. Inc. Publisher Madison, Wisconsin. U.S.A.
- [19] Stoops, G 1978. Provisional notes on micromopedology. International Training Centre for Post – Graduate Soil Scientists, Univ. of Gent, Belgium.
- [20] Bullock, P., Fedoroff, N., Jongerius, A. Stoops, G. and Tursina, T. 1984. *Handbook for soil thin section description*. Waine Research Publication (London).
- [21] Soil Survey Staff. 2003. *Key a to Soil Taxonomy by Soil Survey Staff Eighth Edition*, (USDA Agriculture Handbook, 436). Washington, D.C.
- [22] Guidi, G. and Hall, J.E., 1984. *Effects of Sewage Sludge on the Physical and Chemical Properties of Soils*. 3th International Symposium on Processing and Use of Sewage Sludge, Brighton, UK.
- [23] He, X.T., Traina, S.J., Logan, T.J. 1992. Chemical Properties of Municipal Solid Waste Composts. *Journal Environ. Qual.* 21 (3), 318-329.