

Yetişkin Türk Sigara İçicilerinde Sigara Bırakma Tetikleyicilerinde Cinsiyet ve Yaş Farklılıkları

Gender and Age Differences in the Triggers for Smoking Cessation in Adult Turkish Smokers

[Hülya Abalı](#) , [Fatma Tokgöz Akyıl](#) , [Nurdan Şimşek Veske](#) , [Seda Tural Önür](#)

Sağlık Bilimleri Üniversitesi, Yedikule Göğüs Hastalıkları ve Göğüs Cerrahisi Eğitim Ve Araştırma Hastanesi

Amaç: Sigara dumanından kaynaklanan sağlık riskleri, sigarayı bırakarak belirli bir yoksunluk sürecinden sonra azaltılabilir. Sigarayı bırakmayı tetikleyen faktörlerin cinsiyet ve yaşa göre farklılıklarını tespit etmeyi amaçladık.

Gereç ve Yöntemler: Bu kesitsel çalışma Mayıs-Eylül 2022 tarihleri arasında göğüs hastalıkları polikliniğine başvuran, sigarayı bir yıldan fazla süredir bırakan ve anketi yanıtlamayı kabul eden 185 hasta üzerinde gerçekleştirildi. Ankette katılımcıların demografik özellikleri, sigara içme ve sigarayı bırakma özellikleri, bırakmaya yönelik destekler ve sigarayı bırakmayı tetikleyen bildirilmiş 25 faktör (psikososyal, ekonomik ve tıbbi faktörler) değerlendirildi. Sigarayı bırakmayı tetikleyen faktörler kadın ve erkekler arasında ve ayrıca yaş grupları (≤ 50 yaş ve > 50 yaş) arasında karşılaştırıldı.

Bulgular: Katılımcıların yaş ortalaması 58 ± 13 'ü (yıl) olup %23'ü kadındı. Katılımcıların sigarayı bırakmalarını sağlayan en önemli faktör sigaranın sağlığa zararlarının bilinmesi (%65) oldu. Kadınlar erkeklerden daha gençti ve yılda daha az paket sigara içiyordu (sırasıyla $P < 0.001$, $P = 0.008$). Erkeklerle göre kadınlarda ortalama sigaraya başlama yaşı daha geç, ortalama sigarayı bırakma yaşı ise daha erkendi (sırasıyla $P = 0.036$, $P = 0.00$). Sigara içmenin sağlığa zararları konusundaki farkındalık, erken ölüm kaygısı ve Kronik Obstrüktif Akciğer Hastalığı (KOAH) tanısı erkekler üzerinde daha fazla etkiliyken (sırasıyla $P = 0,026$, $P = 0,01$, $P = 0,009$) astım tanısı kadınlar üzerinde daha fazla etkiliydi ($P = 0.01$). Ailede akciğer kanseri öyküsü genç katılımcılar için sigarayı

bırakma konusunda daha cesaret vericiyken, kronik hastalıklar (KOAH ve kalp hastalıkları) yaşlılar için daha cesaret vericiydi (sırasıyla $P=0.003$, $P=0.001$).

Sonuç: Sigarayı bırakmayı tetikleyen faktörler cinsiyete ve yaşa göre değişmektedir. Hekimler bu farklı özellikleri dikkate alarak bireysel tedavi yaklaşımları geliştirebilirler. Hekimler polikliniklerde ve halka açık sigara karşıtı kampanyalarda sigara içmenin sağlık risklerini vurgulamalıdır.

Anahtar Kelimeler: Sigara bırakma, sağlık bilgisi, kronik hastalık, cinsiyet, yaş

Türkçe Kısa Başlık: Sigara Bırakmayı Teşvik Eden Faktörler

Objective: Health risks due to cigarette smoke can be reduced after a period of abstinence by quitting smoking. We intended to detect trigger factors and variations in triggering factors for smoking cessation regarding gender and age.

Material and Methods: This cross-sectional study was performed on 185 patients applied to the pulmonology polyclinics, who quit smoking for more than one year and who accepted to answer the questionnaire between May and Sep 2022. The questionnaire evaluated the participants' demographic, smoking, and quitting smoking characteristics, supports for quitting, and the reported 25 trigger factors for smoking cessation (psychosocial, economic, and medical factors). The triggers for smoking cessation were compared between females and males, and also age groups (≤ 50 years and > 50 years).

Results: Of the participants, the mean age was 58 ± 13 yrs, and 23% were female. The most seen factor enabling participants to quit smoking was awareness of smoking harms to health (65%). Females were younger and smoked fewer packs/year cigarettes than males ($P < 0.001$, $P = 0.008$, respectively). The mean initiation age of smoking was later, and the mean age of smoking cessation was earlier in females compared to males ($P = 0.036$, $P = 0.00$, respectively). The awareness of health harms of smoking, anxiety about premature death, and diagnosis of Chronic Obstructive Pulmonary Disease (COPD) had higher effects on males ($P = 0.026$, $P = 0.01$, $P = 0.009$, respectively), and diagnosis of asthma had a higher impact on females ($P = 0.01$). While a family history of lung cancer was more encouraging to younger participants for smoking cessation, chronic diseases (COPD and cardiac diseases) were more encouraging to elders ($P = 0.003$, $P = 0.001$, respectively).

Conclusion: The trigger factors for smoking cessation vary according to gender and age. Physicians may develop individual treatment approaches by taking these different characteristics into account. Physicians should emphasize the health risks of smoking in polyclinics and public anti-smoking campaigns.

Keywords: Smoking cessation, health knowledge, chronic disease, gender, age

İngilizce Kısa Başlık: Promoting factors of smoking cessation

INTRODUCTION

Smoking is one of the avoidable causes of premature mortality. Every year, more than 8 million people pass away as a result of smoking-related health issues. The smoking rate has been increasing in low- and middle-income nations since they are the marketing target of the tobacco industry.^[1] It is estimated that there are more than 1 billion consumers of cigarettes worldwide, which negatively affects public health by its use and has an economic cost and social burden due to its consequences.^[2] Türkiye ranks 29th in worldwide cigarette consumption.^[3] In the current literature, there is ample evidence of the health risks associated with cigarette smoke that may reverse after an adequate term of smoking abstinence. Therefore, achieving lifelong smoking cessation is a vital target of public health in developed and developing nations.^[4,5]

Smoking cessation is an important part of tobacco control policies, and evidence-based advices suggest that quitting smoking is useful for smokers. The Framework Convention of Tobacco Control, which was declared by the World Health Organization (WHO) in 2003 with the aim of “protecting the present and future generations against tobacco use and smoke exposure”, was accepted and entered into force in 2004.^[6] Türkiye was the first country to implement all MPOWER (M: Monitor tobacco use and prevention policies; P: Protect people from tobacco smoke; O: Offer help to quit tobacco use; W: Warn about the dangers of tobacco; E: Enforce bans on tobacco advertising, promotion and sponsorship; R: Raise taxes on tobacco) strategies in 2008.^[7] While the rate of individuals aged ≥ 15 yrs who used tobacco products in Türkiye was 28% in 2019, it increased to 28.3% in 2022. Of these individuals, 41.3% were males, and 15.5% were females in 2022.^[8] Despite serious tobacco control policy

implementations in Türkiye, increasing the smoking rate should guide health professionals to focus on smoking cessation. The spectrum of smoking cessation initiatives ranges from an easy advice to pharmacological treatment and intensive behavioral support.^[9]

The three experimental methods (intravenous and oral nicotine self-administration, as well as nicotine-induced conditioned place preference) that were utilized to demonstrate nicotine preference in animal models were outlined in a previous review. All of these methods emphasized the significance of the individuals' gender and age; females and adolescents seem to have stronger preferences than males and adults, respectively. ^[10] Therefore, we aimed to assess the differences in the triggering factors for smoking cessation regarding gender and age, about which there are few studies in the literature. The outcomes of this study may guide physicians to develop individual intervention strategies for smoking cessation.

MATERIAL AND METHODS

Ethical approval of the study was maintained from the Ethics Committee of the reference chest diseases hospital (approval date:12.05.2022, approval no: 2022-239). The study was conducted on 185 patients admitted to the outpatient clinics of the researchers between May and Sep 2022.

Participants recruitment

The inclusion criteria of the participants were as follows: i) to be over 18 yrs old, ii) to have quit smoking for more than one year, and not to initiate again iii) accepting to answer the questionnaire.

The exclusion criteria of the participants were as follows: i) to be under 18 yrs old, ii) current smokers, iii) to have quit smoking for less than one yr.

Data collection

Data of age, gender, education degree, number of smoked cigarettes/day, number of smoked cigarettes (pack/yr), age of initiating/quitting smoking, count of attempts to quit smoking, therapy for quitting (pharmacological, psychological, and pharmacological+psychological), and

psychosocial, economic, and medical factors that promote smoking cessation were registered via participants' answers to the questionnaires.

The triggers for quitting smoking were compared according to gender and age groups. Age was grouped as ≤ 50 years and > 50 years. ^[11]

Statistical Analysis

A statistical software program (SPSS for Windows, version 16.0; SPSS Inc.; Chicago, IL, USA) was used to analyze the data. Quantitative data were expressed as mean \pm standard deviation (SD) (min-max) for the cohort's general characteristics, while qualitative data were expressed as frequencies. Comparisons between groups were performed by the independent two-sample t test for variables with normal distribution, and with the Mann-Whitney U test for variables with non-normal distribution. Chi square test was used to compare categorical variables. $P < 0.05$ was deemed as significant.

RESULTS

A total of 185 adult participants were evaluated by the questionnaires. The average age of the participants was 58 ± 13 yrs, and the number of males was 143 (77%). The participants were mostly primary school graduates (n:94, 51%). It was determined that the majority of the participants were able to quit smoking without any therapy support (n:166 vs n:19) (Table 1).

It was observed that females were younger (51 ± 11 vs. 60 ± 12 , $P < 0.001$) and had fewer average cigarette packs per year (19 ± 10 vs. 25 ± 12 , $P = 0.008$) than males. Compared with males, the average age of females initiating smoking was two years later ($P = 0.036$), and the average age of females quitting smoking was five years earlier ($P = 0.001$) (Table 1).

Among the factors, the awareness of smoking harms to health and diagnosis of chronic diseases were the most common, respectively. When the factors that enabled smoking cessation were reviewed, the awareness of smoking health harms and anxiety about premature death had higher impacts on males compared to females ($P = 0.026$, $P = 0.01$, respectively). Among the triggers for smoking cessation, the diagnosis of COPD was seen

more frequently in males ($P=0.009$), while the diagnosis of asthma was seen more frequently in females ($P=0.01$) (Table 2).

The trigger factors for smoking cessation were compared between the age groups. While having a family history of lung cancer had more impact on younger participants (15% vs. 2%, $P=0.003$), the diagnosis of chronic diseases [COPD and cardiac diseases ($P<0.0001$, $P=0.041$, respectively)] had more impact on older ones (57% vs. 31%, $P=0.001$) (Table 2).

Participants generally reported multifactorial (>3) triggers for smoking cessation. The count of smoking cessation triggers is shown in Table 3.

DISCUSSION

Among the triggering factors for smoking cessation, the most common was awareness of the smoking harms to health by 65% in our study. Similarly, Argüder E et al^[12] reported that the primary factor leading Turkish adults to smoking cessation among the other factors was awareness of the health harms of smoking at a rate of 44%. Moreover, previous studies reported that concern about future health problems due to this awareness was the most effective factor.^[13,14] In developing nations (Malaysia, Thailand, Indonesia, and Vietnam) that implement MPOWER measures like ours, awareness of health harms of smoking appears as the major factor in smoking cessation.^[15] Levy DT et al^[16] proved this result by showing that health warnings were the MPOWER approach that had the greatest impact on reducing smoking-related deaths (13.3 million as opposed to <1 million for each of the other MPOWER initiatives).

The second most common factor was the diagnosis of chronic diseases such as COPD, asthma, cardiac diseases, and hypertension (HT). Similar to our study, Wang R et al^[17] found that smokers with tobacco related chronic diseases had higher prevalence of quit smoking. On the contrary, Holm M et al^[18] reported that the presence of diabetes, chronic bronchitis, asthma, and HT were not associated with smoking cessation. Similar to this study, Patel K et al^[19] detected no significant associations between smoking cessation and diabetes, HT, and high cholesterol. We consider that the differences between our study and these studies may

be due to the fact that the relationship between chronic diseases and smoking cessation was not explained in detail when informing patients. Patients may wise up that smoking is only a risk factor for lung diseases. Because patients are unaware of the smoking harms to the other organ systems, their chronic diseases, except lung diseases, may not have triggered them to quit smoking.

When smoking cessation triggers were compared according to age, the presence of a chronic disease was more effective in the elderly. Among these chronic diseases, the presence of COPD or cardiac diseases enabled older participants to quit smoking significantly more than younger ones. This outcome was expected due to the parallel increase in chronic diseases as aging. In younger people, a family history of lung cancer, which could lead to future health concerns, was prominent. In agreement with our findings, a previous study reported that current health problems enabled elders to quit smoking, while future health concerns enabled younger ones to quit smoking.^[20]

In this study, compared with males, females initiated smoking later and quit smoking earlier. As expected, females smoked fewer packs of cigarettes per year. Males are more exposed to environmental factors that encourage smoking (such as coffeehouse habits, work environment at an early age) compared to females in Türkiye. Thus, males may be under more sociocultural pressure to smoke and continue smoking for more extended periods than females.

We found that the awareness of smoking health harms and anxiety about premature death had more effects on males for smoking cessation. Similarly, many studies declared that males quit smoking more due to health concerns than females.^[21-23] In contrast, females were more triggered to quit by concerns of health problems from smoking than males in a previous study.^[24] The variation between the mentioned above studies may be due to the participants' different sociocultural characteristics and educational levels.

The leading health problems associated with smoking are respiratory diseases such as COPD and asthma. Having a diagnosis of COPD or asthma is an important factor enabling quitting smoking.^[25,26] The diagnosis of COPD had a significant impact on males compared to females for smoking cessation in our study. Because males smoked longer and more packs/year of

cigarettes, and as expected, they were diagnosed with COPD more often. In line with a previous study, having a diagnosis of asthma was observed to trigger smoking cessation mostly in females compared to males.^[27]

Study Limitations

Since the hospital serves adult patients over 18 yrs, only adult patients could be evaluated in this study. Such a survey study should be performed in a larger population in order to reflect the general population.

CONCLUSION

Effective interventions are valuable for successful smoking cessation. Physicians should target separate approaches to men and women, as well as young people and the elderly, for smoking cessation. Among the triggers, awareness of the health harms of smoking, anxiety about premature death, and diagnosis of COPD were more prominent in males, and the diagnosis of asthma was more prominent in females. A family history of lung cancer appeared to enable younger participants to quit smoking, while chronic diseases appeared to enable the older ones. Physicians should explain the appropriate triggers to the patients according to their gender and age. Also, these triggers should be emphasized in public anti-smoking campaigns.

Conflict of Interest: The authors declared no conflicts of interest concerning with the study.

Financial Disclosure: The authors declared that the study received no financial support.

REFERENCES

1. Center For Disease Control and Prevention, 2022. Global Tobacco Control. Available at: <https://www.cdc.gov/tobacco/global/index.htm>. Accessed April 26, 2023.
2. World Health Organization Report on the Global Tobacco Epidemic 2021b:69.
3. World Health Organization Report on the Global Tobacco Epidemic 2021a:148.
4. Wang L, Kong L, Wu F, et al. Preventing chronic diseases in China. *Lancet* 2005;366: 1821–4.
5. US Department of Health and Human Services. The health benefits of smoking cessation. US Department of Health and Human Services, Public Health Service, Centers for Disease Control, Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 1990 (DHHS Publication No. (CDC) 90–8516).
6. Bilir N. Tütün Kontrolü Çerçeve Sözleşmesi ve Türkiye: Bir Başarı Örneği. *Güncel Göğüs Hastalıkları Serisi* 2016;4: 7–12.
7. T. C. Sağlık Bakanlığı Tütün ve Madde Bağımlılığı ile Mücadele Dairesi Başkanlığı, 2019.
8. Türkiye health research, 2022-Türkiye statistical agency-data portal. Available at: <https://data.tuik.gov.tr>. Accessed June 1, 2023.
9. Fiore MC, Bailey WC, Cohen SJ, Dorfman SF, Goldstein MG, Gritz EF, et al. Treating tobacco use and dependence: clinical practice guidelines. U.S. Department of Health and Human Services. Public Health Service, Rockville, MD2000.
10. Pogun S, Yararbas G, Nesil T, Kanit L. Sex differences in nicotine preference. *J Neurosci Res.* 2017;95: 148–162.
11. Chen D and Wu LT. Smoking Cessation Interventions for Adults Aged 50 or Older: A Systematic Review and Meta-analysis. *Drug Alcohol Depend* 2015;154: 14–24.
12. Argüder E, Hasanoğlu H C, Karalezli A, Kılıç H. [Factors that increasing tendency to smoking cessation] [Article in Turkish]. *Tuberk Toraks* 2012;60: 129–35.
13. Bui TC, Xangsayarath P, Douangvichith D, Siengsounthone L, Phandouangsy K, Tran LT, et al. Factors Associated with Cigarette Smoking Cessation in Lao People’s Democratic Republic: Results from the 2015 National Adult Tobacco Survey. *International Journal of Environmental Research and Public Health* 2020;17: 4953.
14. Vangeli E, West E. Sociodemographic differences in triggers to quit smoking: findings from a national survey. *Tob Control* 2008;17: 410–5
15. Shang C, Huang J, Cheng KW, He Y, Chaloupka F J. The association between warning label requirements and cigarette smoking prevalence by education-findings from the Global Adult Tobacco Survey (GATS). *Int J Environ Res Public Health* 2017;14.
16. Levy DT, Li Y, Yuan Z. Impact of nations meeting the MPOWER targets between 2014 and 2016: An update. *Tob Control* 2020;29: 231–3.
17. Wang R, Jiang Y, Yao C, Zhu M, Zhao Q, Huang L, et al. Prevalence of tobacco related chronic diseases and its role in smoking cessation among smokers in a rural area of Shanghai, China: a cross sectional study. *BMC Public Health* 2019;19: 753.
18. Holm M, Schiöler L, Andersson E, Forsberg B, Gislason T, Janson C, et al. Predictors of smoking cessation: A longitudinal study in a large cohort of smokers. *Respir Med* 2017;132: 164–9.

19. Patel K, Schlundt D, Larson C, Wang H, Brown A, Hargreaves M. Chronic illness and smoking cessation. *Nicotine Tob Res* 2009;11: 933–9.
20. Gallus S, Muttarak R, Franchi M, Pacifici R, Colombo P, Boffetta P, et al. Why do smokers quit? *Eur J Cancer Prev* 2013;22: 96–101.
21. Curry SJ, Grothaus L, McBride C. Reasons for quitting: intrinsic and extrinsic motivation for smoking cessation in a population-based sample of smokers. *Addict Behav* 1997;22: 727–39.
22. McKee SA, Maciejewski PK, Falba T, Mazure CM. Sex differences in the effects of stressful life events on changes in smoking status. *Addiction* 2003;98: 847–55.
23. Grøtvedt L, Stavem K. Association between age, gender and reasons for smoking cessation. *Scand J Public Health* 2005;33: 72–76.
24. Chinwong D, Mookmanee N, Chongpornchai J, Chinwong S. A Comparison of Gender Differences in Smoking Behaviors, Intention to Quit, and Nicotine Dependence among Thai University Students. *J Addict* 2018: 8081670.
25. Lindsay H G, Wamboldt F S, Holm K E, Make B J, Hokanson J, Crapo J D, et al. Impact of a Medical Diagnosis on Decision to Stop Smoking and Successful Smoking Cessation. *Chronic Obstr Pulm Dis* 2021;8: 360–70.
26. Danielsen S E, Løchen M-L, Medbø E, Vold M L, Melbye H. A new diagnosis of asthma or COPD is linked to smoking cessation- the Tromsø study. *Int J Chronic Obstr Pulm Dis* 2016;11: 1453–8.
27. Chen Z, Wasti B, Shang Y, Ouyang R, Yuan Y, He Y, et al. Different clinical characteristics of current smokers and former smokers with asthma: a cross-sectional study of adult asthma patients in China. *Sci Rep* 2023;13: 1035.

Table 1. Demographic, smoking, and quitting smoking characteristics of the participants and comparison of participants' characteristics by gender and age

	Number of participants (%)	Female n=42 (%)	Male n=143 (%)	P	≤50 yr n=55 (%)	>50 yr n=130 (%)	P
<i>Mean age(yr)±SD</i>	58±13	51±11	60±12	<0.001			
<i>Gender</i>							
Male	143 (77)				35 (24)	108 (76)	0.007
Female	42 (23)				22 (52)	20 (48)	
<i>Education degree</i>							
Primary	94 (51)	6 (14)	21 (15)	0.523	21 (38)	73 (56)	0.066
Secondary	44 (24)	7 (17)	13 (9)		14 (26)	30 (23)	
High school	20 (11)	8 (19)	36 (25)		7 (12)	13 (10)	
University	27 (14)	21 (50)	73 (51)		13 (24)	14 (11)	
<i>Mean cigarette number/d ±SD (min-max)</i>	24±12 (3-60)	19±10	25 ± 12	0.008	22±10	24±13	0.277
<i>Cigarette number (pack/yr)</i>							
0-20	57 (31)	19 (45)	38 (27)	0.03	38	19	>0.0001
20-40	90 (49)	19 (45)	71 (50)		17	73	
>40	38 (20)	4 (10)	34 (24)		0	38	
<i>Mean age of smoking initiation (yr)±SD (min-max)</i>	18±5.5 (6-41)	20±6	17.9±5	0.036	18±4	18±6	0.905
<i>Mean age of smoking cessation ± SD (min-max)</i>	45.5±12 (17-71)	41.3±12	46.8±12	0.01	34±8	50±10	N/A
<i>Number of attempts to quit smoking</i>							
1	39 (21)	10	29	0.665	16	23	0.193
2	62 (34)	12	50		14	48	
3	32 (17)	7	25		11	21	
>3 (4-6)	52 (28)	18	39		14	38	
<i>Support for smoking cessation</i>	19 (10)	7 (12)	7 (5)	0.19	6 (11)	8 (6)	0.360
<i>Pharmacotherapy</i>							
Varenicline	7	4	3	0.179	4	3	0.131
Nicotine patch	2	0	2		2	0	
Nicotine gum	1	1	0		1	0	
Bupropion	1	1	0		1	0	
<i>Psychotherapy</i>	5	2	3		1	4	
<i>Pharmacotherapy+ Psychotherapy</i>	3	1	2	0.285	1.2	2	0.548

SD, standard deviation; N/A, nonavailable

Table 2. Comparison of trigger factors for smoking cessation by gender and age

Triggers	Number of participants (%)	Gender statistics			Age statistics		
		Female n=42 (%)	Male n=143 (%)	P	≤50 yr n=55 (%)	>50 yr n=130 (%)	P
<i>Awareness of smoking harms to health</i>	121 (65)	21 (50)	100 (70)	0.026	33 (60)	88 (68)	0.317
<i>Diagnosis of Chronic diseases*</i>	91 (49)	20 (48)	71 (50)	0.862	17 (31)	74 (57)	0.001
<i>Diagnosis of Chronic Obstructive Pulmonary Disease</i>	63 (34)	7 (17)	56 (39)	0.009	4 (7)	59 (45)	<0.0001
<i>Desire not to be an inappropriate role model for children</i>	61 (33)	13 (31)	48 (34)	0.853	20 (36)	41 (32)	0.608
<i>Recommendation of a healthcare professional</i>	58 (31)	11 (26)	47 (33)	0.455	12 (22)	46 (35)	0.083
<i>Anxiety about requiring care in old age</i>	51 (28)	8 (19)	43 (30)	0.175	14 (26)	37 (29)	0.722
<i>Request of family members</i>	40 (22)	6 (14)	34 (24)	0.209	12 (22)	28 (22)	0.998
<i>Anxiety about premature death</i>	39 (21)	3 (7)	36 (25)	0.01	8 (15)	31 (24)	0.173
<i>Bad breath smell</i>	37 (20)	10 (24)	27 (19)	0.513	14 (26)	23 (18)	0.234
<i>Hypertension</i>	33 (18)	9 (20)	24 (20)	0.884	5 (9)	28 (22)	0.057
<i>Decision after acute diseases[§]</i>	31 (17)	7 (17)	24 (17)	0.998	8(15)	23 (18)	0.672
<i>Diagnosis of Asthma</i>	25 (14)	11 (26)	14 (10)	0.010	12 (22)	13 (10)	0.057
<i>Financial problems</i>	25 (14)	4 (10)	21 (15)	0.454	11 (20)	14 (11)	0.104
<i>Diagnosis of cardiac diseases[¶]</i>	20 (11)	3 (7)	17 (12)	0.573	2 (4)	18 (14)	0.041
<i>Heart attack</i>	19 (10)	5 (12)	14 (10)	0.773	4 (7)	15 (12)	0.441
<i>Smoking ban at home</i>	16 (9)	4 (10)	12 (8)	0.762	2 (4)	14 (11)	0.155
<i>Effect of public service</i>	15 (8)	1 (2)	14 (10)	0.197	6 (11)	9 (7)	0.384

<i>announcements in the Media</i>							
<i>Other triggers</i>	14 (8)	4 (10)	10 (6)	0.236	5 (9)	9 (6)	0.415
<i>Anxiety about beauty loss</i>	13 (7)	4 (10)	9 (6)	0.498	6 (11)	7 (5)	0.213
<i>A family history of lung cancer</i>	11 (6)	4 810)	7 (5)	0.274	8 (15)	3 (2)	0.003
<i>Diagnosis of malignancy</i>	7 (4)	1 (2)	6 (4)	0.998	3 (6)	4 (3)	0.426
<i>Sexual intercourse disorders</i>	6 (3)	1 (2)	5 (4)	0,998	1 (2)	5 (4)	0.671
<i>Stroke</i>	5 (3)	2 85)	3 (2)	0.318	2 (4)	3 82)	0.635
<i>Infertility</i>	3 (2)	2 (5)	1 (1)	0.130	2 (4)	1 (1)	0.21
<i>Pregnancy</i>	2 (1)	2 (5)			2		

* Chronic diseases: Chronic obstructive pulmonary disease, asthma, cardiac diseases, hypertension;

§Acute diseases: Heart attack, stroke, and others; ¶ Cardiac diseases: Coronary artery disease, cardiac arrhythmias, heart valve diseases, heart failure. Triggers have more than one option.

Table 3. Count of smoking cessation triggers

<i>Count of smoking cessation triggers</i>	Number of participants (%)	Female (n=42)	Male (n=143)	P	≤50 yr (n=55)	>50 yr (n=130)	P
1	29 (16)	9 (21)	20 (14)	0.19	16 (29)	19 (15)	0.071
2-3	53 (29)	10 (24)	43 (30)		15 (27)	42 (32)	
>3	103 (65)	23 (55)	80 (56)		24 (44)	69 (53)	
<i>Mean count of smoking cessation triggers±SD</i>	3.7±2 (1-9)	3.6	3.8	0.312	3.4	3.9	0.224

SD, standard deviation