

Striction Lines of Non-developable Ruled Surfaces in Euclidean 3-Space*

Ali ÇAKMAK

ABSTRACT: The ruled surfaces, one of the areas of interest of differential geometry, have been one of the surface types studied by many mathematicians from the past to the present day. Similarly, some special curves which helix, slant helix, Bertrand curve, etc. are also the curve types discussed often by mathematicians. In this paper, it will be shown that striction lines of non-developable ruled surfaces are helix, slant helix, Bertrand or Mannheim curve in some special cases.

Keywords: Non-developable ruled surface, line of striction, helix, Bertrand curve, Mannheim curve.

Öklid-3 Uzayında Açılabilir Olmayan Regle Yüzeylerin Striksiyon Çizgileri

ÖZET: Diferansiyel geometrinin ilgi alanlarından olan regle yüzeyler, geçmişten günümüze bir çok matematikçi tarafından çalışılan yüzey tiplerinden birisidir. Benzer şekilde helis, slant helis, Bertrand eğrisi gibi bazı özel eğriler de matematikçiler tarafından sıklıkla tartışılan eğri tipleridir. Bu makalede, bazı özel durumlarda açılabilir olmayan regle yüzeylerin striksiyon çizgilerinin helis, slant helis, Bertrand eğrisi ya da Mannheim eğrisi olduğu gösterilecektir.

Anahtar Kelimeler: Açılabilir olmayan regle yüzey, striksiyon çizgisi, helis, Bertrand eğrisi, Mannheim eğrisi.

¹ Ali ÇAKMAK (0000-0002-2783-9311), Bitlis Eren Üniversitesi, Fen Edebiyat Fak., Matematik, Bitlis, Türkiye
Sorumlu yazar/Corresponding Author: Ali ÇAKMAK, acakmak@beu.edu.tr

* Bu makale 22-24 Ağustos 2017 tarihinde Van ilinde Caucasian Mathematics Conference II (CMC - II)'de sunulmuştur.

INTRODUCTION

It is known that if a surface is formed by the movement of a line, this surface is called the ruled surface. A ruled surface is the locus of a line depending on a parameter. We assume that this line does not have an envelope; hence the surface is

parameters u and v in E^3 by $X(u, v)$. Let

$$X(u, v) = a(u) + vb(u) \quad (1)$$

be a non-developable ruled surface in E^3 with $b^2(u) = 1$ and the parameter u is the arc length parameter of $b(u)$ as a unit spherical curve in E^3 (Liu et al., 2014; Yu et al., 2014). If $a'(u) \cdot b'(u) = 0$, base curve $a(u)$ is striction line of ruled surface.

MATERIAL AND METHOD

Definition 2.1. A curve β with $\kappa(s) \neq 0$ is defined as a cylindrical helix if the tangent lines of the curve

$$\left(\frac{\tau}{\kappa} \right) (s) = \text{constant}. \quad (2)$$

Here, if $\kappa(s)$ and $\tau(s)$ are constants we call a circular helix.

Definition 2.2. A curve β with $\kappa(s) \neq 0$ is

non-developable (Kühnel, 2006). In other words, it is mean that a non-developable surface free of points of vanishing Gaussian curvature in a 3-dimensional Euclidean space (Yoon, 2007). We denote the Euclidean 3-space by E^3 and a regular parameter surface with the

Some special curves which are helix, slant helix, Bertrand and Mannheim curves are examined by the some authors. (Izumiya and Takeuchi, 2002; Liu and Wang, 2008; Yaylı and Saraçoğlu, 2012).

β creative a constant angle with a fixed direction.

Proposition 2.1. A curve β with $\kappa(s) \neq 0$ is a cylindrical helix if and only if

defined as a slant helix if the principal normal lines of β creative a constant angle with a fixed direction.

Proposition 2.2. Let β be an unit speed space curve with $\kappa(s) \neq 0$. Then, β is a slant helix if and only if

$$\sigma(s) = \left(\frac{\kappa^2}{(\kappa^2 + \tau^2)^{\frac{3}{2}}} \left(\frac{\tau}{\kappa} \right)' \right) (s) \tag{3}$$

is a constant function.

Definition 2.3. A curve $\beta : I \rightarrow E^3$ with $\kappa(s) \neq 0$ is defined as a Bertrand curve if there is a curve $\bar{\beta} : I \rightarrow E^3$ under the condition that the principal

normal lines of β and $\bar{\beta}$ at $\forall s \in I$ are equal.

Proposition 2.3. Let β be a space curve with $\kappa(s) \neq 0$ and $\tau(s)$. Then, β is a Bertrand curve if and only if

$$A(\tau'(s)\kappa(s) - \kappa'(s)\tau(s)) - \tau'(s) = 0 \tag{4}$$

such that $A \neq 0$ and $A \in R$. Here, $\bar{\beta} = \beta(s) + An(s)$ is Bertrand mate of β .

of β with the binormal lines of $\bar{\beta}$ at $\forall s \in I$ are equal.

Definition 2.4. A curve $\beta : I \rightarrow E^3$ with $\kappa(s) \neq 0$ is defined as a Mannheim curve if there is a curve $\bar{\beta} : I \rightarrow E^3$ such that the principal normal lines

Proposition 2.4. A space curve in E^3 is a Mannheim curve if and only if

$$\kappa = c(\kappa^2 + \tau^2), \tag{5}$$

where c is a nonzero constant and κ and τ are curvature and torsion of curve, respectively.

Definition 2.5. A ruled surface is defined by transformation

$$X(a,b) : I \times E \rightarrow E^3$$

$$(u,v) \rightarrow X(a,b)(u,v) = a(u) + vb(u), \tag{6}$$

where $a: I \rightarrow E^3, b: I \rightarrow E^3$ are differentiable transformations and I is an open interval. $a(u)$ and $b(u)$ are called base curve and director curve, respectively. If $b(u)$ is a constant, ruled surface is called as a cylinder.

It is known that if Gaussian curvature is zero, the ruled surface is developable ruled surface. In addition, it is mean that a non-developable ruled surface free of points of vanishing Gaussian curvature.

$$x'(u) = \alpha(u), \quad (7)$$

$$\alpha'(u) = -x(u) + k_g(u)y(u), \quad (8)$$

$$y'(u) = -k_g(u)\alpha(u), \quad (9)$$

where $k_g(u)$ is called the spherical curvature function and $\{x(u), \alpha(u), y(u)\}$ is called the spherical Frenet frame of the spherical curve $b(u)$ (Liu et al., 2014; Yu et al., 2014).

Definition 2.7. Suppose that $X(u, v) = a(u) + vb(u)$ is a non-developable ruled surface and $a(u)$ is the striction line of $X(u, v)$ under the conditions $d'(u) = \lambda(u)x(u) + \mu(u)y(u)$ and u is the arc length parameter of $b(u)$. Let $k_g(u)$ be the spherical curvature function of $b(u)$, then $X(u, v)$ can be given by $\{k_g(u), \lambda(u), \mu(u)\}$ up to a transformation in E^3 . Here, the functions $k_g(u)$,

Definition 2.6. Suppose that $X(u, v) = a(u) + vb(u)$ is a non-developable ruled surface in E^3 . We assume that $b(u)$ is a unit spherical curve in E^3 such that $b^2(u) = 1$ and the parameter u is the arc length parameter of $b(u)$ and base curve $a(u)$ is the striction line of the non-developable ruled surface $X(u, v)$. In this case, if we write as $x(u) = b(u)$, $x'(u) = \alpha(u)$ and $y(u) = \alpha(u)' x(u)$, the spherical Frenet formulas of the curve $b(u)$ can be given by

$\lambda(u)$ and $\mu(u)$ are defined as structure functions of $X(u, v)$ in E^3 (Liu et al., 2014; Yu et al., 2014).

Definition 2.8. Suppose that $X(u, v) = a(u) + vb(u)$ is a non-developable ruled surface in E^3 and $a(u)$ is the striction line of $X(u, v)$ under the condition $a'(u) = \lambda(u)x(u) + \mu(u)y(u)$. Here, $\{\alpha(u), x(u) = b(u), y(u)\}$ is the spherical Frenet frame of the spherical curve $b(u)$ and the parameter u is the arc length parameter of $b(u)$.

We assume that $\lambda(u) \neq 0$, then $X(u, v)$ is described as pitched ruled surface. (Liu et al., 2014; Yu et al., 2014).

RESULTS AND DISCUSSION

Let $X(u, v) = a(u) + vb(u)$ be a non-developable ruled surface with the structure functions $\lambda(u)$, $\mu(u)$ and $k_g(u)$. Then, the curvature \mathbf{K} and torsion \mathbf{T} of striction line $a(u)$ are, respectively, given by

$$\kappa^2 = \frac{(\lambda - k_g \mu)^2 (\lambda^2 + \mu^2) + (\lambda' \mu - \lambda \mu')^2}{(\lambda^2 + \mu^2)^3} \tag{10}$$

and

$$\tau = \frac{(\lambda - \mu k_g)(\lambda \mu'' - \lambda'' \mu + (\lambda k_g + \mu)(\lambda - \mu k_g)) + (\lambda' \mu - \lambda \mu')(2\lambda' - 2k_g \mu' - k_g' \mu)}{(\lambda - k_g \mu)^2 (\lambda^2 + \mu^2) + (\lambda' \mu - \lambda \mu')^2} \tag{11}$$

(Liu et al., 2014; Yu et al., 2014).

From ‘‘Eq. 10.’’ and ‘‘Eq. 11.’’, we obtain

$$\frac{\tau}{\kappa} = \frac{(\lambda - \mu k_g)(\lambda \mu'' - \lambda'' \mu + (\lambda k_g + \mu)(\lambda - \mu k_g)) + (\lambda' \mu - \lambda \mu')(2\lambda' - 2k_g \mu' - k_g' \mu)}{\sqrt{\frac{(\lambda - k_g \mu)^2 (\lambda^2 + \mu^2) + (\lambda' \mu - \lambda \mu')^2}{(\lambda^2 + \mu^2)^3}}} \tag{12}$$

By definition of helix, the last equation should be constant. In this case, if we choose

$$\frac{\lambda}{\mu} = constant, \text{ we get}$$

$$\lambda' \mu - \lambda \mu' = 0 \tag{13}$$

and

$$\lambda \mu'' - \lambda'' \mu = 0 \tag{14}$$

Then, we have

$$\frac{\tau}{\kappa} = \frac{(\lambda - \mu k_g)(\lambda k_g + \mu)(\lambda - \mu k_g)}{(\lambda - \mu k_g)^2 (\lambda^2 + \mu^2)} \frac{(\lambda^2 + \mu^2) \sqrt{\lambda^2 + \mu^2}}{(\lambda - \mu k_g) \sqrt{(\lambda^2 + \mu^2)}} \tag{15}$$

$$\frac{\tau}{\kappa} = \frac{\lambda k_g + \mu}{\lambda - k_g \mu} \tag{16}$$

If striction line is helix ‘‘Eq. 16.’’ is constant and so, we can write

$$\frac{\lambda k_g + \mu}{\lambda - k_g \mu} = c, \quad (c \in \mathbb{R}) \quad (17)$$

and we obtain

$$k_g = \frac{c\lambda - \mu}{\lambda + c\mu}. \quad (18)$$

From here, we can express the following theorem:

Theorem 3.1. Suppose that $X(u, v) = a(u) + vb(u)$ is any non-developable ruled surface in E^3 and $a(u)$ is the striction line of $X(u, v)$ under the condition $a' = \lambda x(u) + \mu y(u)$.

$a(u)$ is a helix if and only if $\frac{\lambda}{\mu} = \text{constant}$ and $k_g = \frac{c\lambda - \mu}{\lambda + c\mu}$, ($c \in \mathbb{R}$).

By the definition of slant helix, ‘‘Eq. 3.’’ should be constant. From ‘‘Eq. 10.’’ and ‘‘Eq.

11.’’, if we choose $\frac{\lambda}{\mu} = \text{constant}$, we have

$$\kappa = \frac{\lambda - k_g \mu}{\lambda^2 + \mu^2} \quad (19)$$

and

$$\tau = \frac{\lambda k_g + \mu}{\lambda^2 + \mu^2}. \quad (20)$$

Then, we obtain

$$\kappa^2 + \tau^2 = \frac{1 + k_g^2}{\lambda^2 + \mu^2}, \quad (21)$$

$$\left(\frac{\tau}{\kappa}\right)' = \left(\frac{\lambda k_g + \mu}{\lambda - k_g \mu}\right)' = \frac{(\lambda' k_g + \lambda k_g' + \mu')(\lambda - k_g \mu) - (\lambda' - k_g' \mu - k_g \mu')(\lambda k_g + \mu)}{(\lambda - k_g \mu)^2} \quad (22)$$

$$\left(\frac{\tau}{\kappa}\right)' = \frac{k_g'(\lambda^2 + \mu^2)}{(\lambda - k_g \mu)^2} \quad (23)$$

$$\frac{(\lambda - k_g \mu)^2}{(\lambda^2 + \mu^2)^2} \frac{k_g' (\lambda^2 + \mu^2)}{\left(\frac{1 + k_g^2}{\lambda^2 + \mu^2}\right)^{\frac{3}{2}} (\lambda - k_g \mu)^2} = \frac{k_g'}{(1 + k_g^2)^{\frac{3}{2}}}. \tag{24}$$

Here, if we choose $\frac{k_g'}{(1 + k_g^2)^{\frac{3}{2}}} = c$ and solve this differential equation, we obtain

$$\frac{k_g}{\sqrt{1 + k_g^2}} = cu + d, (c, d \text{ constant}). \tag{25}$$

Therefore, we can say the following theorem:

Theorem 3.2. Suppose that $X(u, v) = a(u) + vb(u)$ is a non-developable ruled surface in E^3 and $a(u)$ is the striction line of $X(u, v)$ under the condition $a'(u) = \lambda x(u) + \mu y(u)$. $a(u)$ is a slant helix if and only if $\frac{\lambda}{\mu} = \text{constant}$ and

$\frac{k_g}{\sqrt{1 + k_g^2}} = cu + d$, where $k_g(u)$, $\lambda(u)$ and $\mu(u)$ are structure functions of $X(u, v)$ and c, d are constants.

It is know that a space curve is a Bertrand curve under the condition that

$$A(\tau' \kappa - \tau \kappa') = \tau', \tag{26}$$

where A is nonzero a real number. From ‘‘Eq. 26.’’, we can write

$$\left(\frac{\tau}{\kappa}\right)' \kappa^2 A = \tau', \tag{27}$$

Here, if we use value in ‘‘Eq. 23.’’ instead of $\left(\frac{\tau}{\kappa}\right)'$ and consider $\frac{\lambda}{\mu} = \text{constant}$, we get

$$\frac{k_g' (\lambda^2 + \mu^2) (\lambda - \mu k_g)^2}{(\lambda - \mu k_g)^2 (\lambda^2 + \mu^2)^2} A = \lambda' k_g + \lambda k_g' + \mu' \tag{28}$$

$$\frac{k_g'(\lambda^2 + \mu^2)(\lambda - \mu k_g)^2}{(\lambda - \mu k_g)^2(\lambda^2 + \mu^2)^2} A = \lambda' k_g + \lambda k_g' + \mu'$$

$$A = \lambda + \frac{\lambda' k_g + \mu'}{k_g'}, \quad (29)$$

where u is the arc length parameter of $a(u)$. Since A is a non-zero constant, λ and μ are constants.

In this case, we can write the following theorem:

Theorem 3.3. Suppose that $X(u, v) = a(u) + vb(u)$ is any non-developable ruled surface in E^3 and $a(u)$ is the striction line of $X(u, v)$ under the condition $a'(u) = \lambda x(u) + \mu y(u)$. $a(u)$ is a Bertrand curve if and only if λ and μ are constants, where u is arc length parameter of $a(u)$.

Corollary 3.1. The striction line $a(u)$ of $X(u, v)$ is a Bertrand curve if and only if $A = \lambda$.

Proof: If λ and μ are considered constants in “Eq. 29.”, it can be easily obtained $A = \lambda$.

It is just known that a space curve in E^3 is a Mannheim curve if “Eq. 5.” is provided. If we suppose that $\frac{\lambda}{\mu}$ is constant and use “Eq. 19.”, “Eq. 20.” and “Eq. 21.”, we obtain

$$\kappa = c(\kappa^2 + \tau^2) \quad (30)$$

$$c \frac{1 + k_g^2}{\lambda^2 + \mu^2} = \frac{\lambda - k_g \mu}{\lambda^2 + \mu^2} \quad (31)$$

$$c = \frac{\lambda - k_g \mu}{1 + k_g^2}, \quad (32)$$

where $c \neq 0$, $c \in IR$.

Hence, we can give the following theorem:

Theorem 3.4. Suppose that $X(u, v) = a(u) + vb(u)$ is a non-developable ruled surface in E^3 and $a(u)$ is the striction line of $X(u, v)$ under the condition

$a'(u) = \lambda x(u) + \mu y(u)$. $a(u)$ is a Mannheim curve if and only if $\frac{\lambda}{\mu} = \text{constant}$ and

$c = \frac{\lambda - k_g \mu}{1 + k_g^2}$, where $k_g(u)$, $\lambda(u)$ and $\mu(u)$ are structure functions of $X(u, v)$ and c is a

non-zero constant.

CONCLUSION

In this paper, firstly, we gave definitions of helix, slant helix, Bertrand and Mannheim curves. Then, non-developable ruled surfaces and striction lines of non-developable ruled

surfaces was defined by means of structure functions. Finally, we showed that under certain conditions, striction lines of non-developable ruled surfaces are helix, slant helix, Bertrand or Mannheim curve.

REFERENCES

- Izumiya S, Takeuchi N, 2003. Special Curves and Ruled Surfaces. Contributions to Algebra and Geometry, 44: 203-212.
- Kuhnel W, 2006. Differential geometry, Curves-Surfaces-Manifolds. Second Edition, American Mathematical Society, USA. 380p.
- Liu H, Wang F, 2008. Mannheim partner curves in 3-space. Journal of Geometry, 88: 120-126.
- Liu H, Yu Y, Jung SD, 2014. Invariants of non-developable ruled surfaces in Euclidean 3-space. Contrib. Algebra Geom., 55: 189-199.
- Yaylı Y, Saraçoğlu S, 2012. Different Approaches To Ruled Surfaces. SDU Journal of Science, 7 (1): 56-68
- Yoon DW, 2007. On Non-Developable Ruled Surfaces in Euclidean 3- Spaces. Indian J. pure appl. Math., 38(4): 281-290.
- Yu Y, Liu H, Jung SD, 2014. Structure and characterization of ruled surfaces in Euclidean 3-space. Applied Mathematics and Computation, 233: 252259.