






# Leaf indumentum in some Turkish species of *Teucrium* (Lamiaceae)

Gülay Ecevit-Genç<sup>1\*</sup> , Taner Özcan<sup>2</sup> , Tuncay Dirmenci<sup>2</sup> 

<sup>1</sup>Department of Pharmaceutical Botany, Faculty of Pharmacy, İstanbul University, Beyazıt, 34452, İstanbul, Turkey

<sup>2</sup>Department of Biology Education, Faculty of Necatibey Education, Balıkesir University, 10145, Balıkesir, Turkey

ORCID IDs of the authors: G.E.G. 0000-0002-1441-7427, T.Ö. 0000-0003-1683-7297, T.D. 0000-0003-3038-6904.

**Cite this article as:** Ecevit-Genç G, Özcan T, Dirmenci T. Leaf indumentum in some Turkish species of *Teucrium* (Lamiaceae). *Istanbul J Pharm* 48 (1): 6-11.

## ABSTRACT

Trichome micromorphology can be used as a discriminating character in the separation of species and subspecies. The micromorphological characteristics of foliar trichomes from five *Teucrium* taxa (*T. scordium* subsp. *scordium*, *T. sirnakense*, *T. chasmophyticum*, *T. andrusi* and *T. spinosum*) were investigated by scanning electron microscope. Seven types of trichomes were identified, including glandular and non-glandular. The glandular trichomes were recorded in three types; short clavate, long clavate, and sessile. The non-glandular trichomes were identified with four types; unicellular thin-walled, 2(5)-celled thin-walled, 3–7(11)-celled flexuose, elongated thin-walled and 2-celled thick-walled trichomes. Generally, the trichome types were similar on both sides of the leaves, except in *T. spinosum*. Sessile glandular trichomes were the most common type and were occurred in all the investigated species. Also clavate glandular trichomes were found in all the species, except *T. scordium* subsp. *scordium*. All trichomes were distinctively thick-walled, whereas thin-walled trichomes were observed in only *T. chasmophyticum* species. The leaf micromorphology of *T. sirnakense* and *T. chasmophyticum* has also been reported in detail for the first time in this study.

**Keywords:** *Teucrium*, Lamiaceae, trichome, micromorphology, SEM, leaf indumentum

## INTRODUCTION

*Teucrium* L. belong to the Lamiaceae family is a well-known, wide-distributed and also one of the largest genus with more than 260 species distributed all around the world. (Kastner 1989; Abu-Assab and Cantino 1993). Mediterranean region, the center of diversity of the genus, has about 96% of all taxa (Cantino et al. 1992; Navarro and El Oualidi 2000).

*Teucrium* species have been divided into ten sections all over the world regarding their calyx shapes and the inflorescence structures (McClintock and Epling 1946; Tutin and Wood 1972). These are the sections: sect. *Teucriopsis* Benth., sect. *Teucrium*, sect. *Chamaedrys* Miller Schreber, sect. *Polium* Miller Schreber, sect. *Isotriodon* Boissier, sect. *Pycnobotrys* Benth., sect. *Scorodonia* (Hill) Schreber, sect. *Stachyobotrys* Benth., sect. *Scordium* Reichenbach, and sect. *Spinularia* Boissier. The species of the *Teucriopsis* and *Pycnobotrys* sections are not distributed in Turkey. (Ekim 1982). The eight sections of the genus, consist of 49 taxa (36 species) and 18 of them endemic are naturally grown in Turkey (Govaerts 1999; Duman 2000; Dönmez 2006; Parolly and Eren 2007; Dönmez et al. 2010; Dinç et al. 2011, Dirmenci 2012; Özcan et al. 2015; Vural et al. 2015; Dinç and Doğu 2016).

*Teucrium sirnakense* Özcan and Dirmenci (endemic) and *T. scordium* L. subsp. *scordium* (Sect. *Scordium*), *T. Andrusi* Post (endemic) and *T. chasmophyticum* Rech. f. (Sect. *Isotriodon* and *T. spinosum* L. (Sect. *Spinularia*) were examined in this study. Sect.

## Address for Correspondence:

Gülay Ecevit-Genç, e-mail: gulecevit@gmail.com

© Copyright 2018 by İstanbul University Faculty of Pharmacy. Available on-line at [www.dergipark.gov.tr/iujfp](http://www.dergipark.gov.tr/iujfp)

Received: 04.01.2018

Accepted: 05.03.2018

*Scordium* has toothed leaves, rectangular stems, subgibbous calyces and subequal calyx teeth, and Sect. *Isotriodon* has dentate or entire leaves, terete stems, gibbous and bilabiate calyces. Sect. *Spinularia* is quite different than these two sections. *T. spinosum* is the only annual species in the Turkish *Teucrium*, and has resupinate corolla according to Flora of Turkey (Ekim 1982).

*Teucrium* species has traditionally been used in Turkey for abdominal pain, antidiabetic, antipyretic, stomachache, common cold, high fever and rheumatic pain (Aksoy-Sagirli et al. 2015).

Plant trichomes are important to descriptive and experimental botanists and data about them and their indumenta are routinely included in many studies. Many authors, such as Behnke (1984); Navarro and El Oualidi (2000); Beyrouthy et al. (2009); Moon et al. (2009); Kaya et al. 2012; Khalik and Hassan (2012); Osman (2012); Ecevit-Genç et al. (2017); Genç et al. (2017); Zareh et al. (2017) emphasize the importance of trichomes in taxonomy.

In many genera of Lamiaceae, the trichome morphology is very useful for the classification of all taxonomic levels (Marin et al. 1994; Navarro and El Oualidi 2000; Moon et al. 2009; Salmaki et al. 2009; Ecevit-Genç et al. 2015, 2017).

Trichomes are widely distributed over the different parts of the Lamiaceae genus and they are generally distinguished as glandular and non-glandular trichomes. Micromorphological features, especially trichomes, are available taxonomic characters in *Teucrium*. Trichomes have an important role in the infrageneric classification of the genus. There have been many studies on the trichomes of *Teucrium* species in recent years (Navarro and El Oualidi 2000; Grubescic et al. 2007; Dinç et al. 2011; Eshratifar et al. 2011; Doğu et al. 2013, Ecevit et al. 2015, 2017).

The main purposes of this paper are to provide a detailed description of the leaf micromorphology of five *Teucrium* species belonging to three different sections.

## MATERIALS AND METHODS

The material was collected from different localities in Turkey by the authors. Voucher specimens have been deposited in the ISTE. A list of taxa included in the study was given in Table 1.

For the micromorphological study, the materials were obtained from collected specimens and micromorphological investigations were conducted using scanning electron microscope (SEM). For SEM analysis, leaves parts were mounted on the stubs and coated with gold layer. They were studied

**Table 1. Collection data of *Teucrium* taxa studied.**

Taxon	Collection data
<i>T. scordium</i> subsp. <i>scordium</i>	Edirne, İpsala border gate, 01.viii.2014, T.Dirmenci, ISTE 101 691
<i>T. sirmakense</i>	Şırnak, Taşdelen village, rocky slopes, 10.vi.2013, T.Dirmenci, E.Akçiçek, Ö.Güner, ISTE 101 694
<i>T. chasmophyticum</i>	Siirt, Between Eruh-Gölgelikonak village, 11.vi.2013, T.Dirmenci, E.Akçiçek, Ö.Güner, ISTE 101 711
<i>T. andrusi</i>	Mardin, Bakırkırı hill, cliffs, 23.vi.2013, T.Özcan, M.Açar, ISTE 101 712
<i>T. spinosum</i>	Diyarbakır, Diyarbakır-Ergani roadsides, 09.vi.2014, T.Dirmenci, E.Akçiçek, Ö.Güner, ISTE 101 719

ISTE: Herbarium of the Faculty of Pharmacy of İstanbul University

**Table 2. Trichome types (Navarro and El Oualidi 2000)**

Glandular trichomes	
A	Clavate glandular trichomes
A1	Short clavate glandular trichomes. Generally with two, large and thin stalk cells.
A2	Long clavate glandular trichomes. Generally with long 3-5 stalk cells.
B	Subsessile glandular trichomes, peltate trichomes
Non-glandular trichomes	
C	Thin-walled trichomes
C1	Triangular, large and very thin -walled unicellular hairs.
C2	Large, thin-walled, 2[5]-celled trichomes which are acute apical cell. Internodes have ridges or marked.
C3	Flexuose and elongated, 3-7(11)-celled trichomes, with internodes distinct, the apical cell acute with micro-papillae.
D	Short and slightly conical, generally 2-celled thick-walled trichomes which are pointed short or elongated apical cell, erect or sometimes slightly curved.

with a scanning electron microscope (FEI Quanta 450 FEG-EDS). All leaves were scanned from adaxial and abaxial surfaces. Terminology of the indumenta on leaves were based on Navarro and El Oualidi (2000), (Table 2).

## RESULTS

The micromorphological characteristics and distribution of the trichomes on leaves of five *Teucrium* taxa were examined in this study. Different type of indumentum shows considerable among species (Table 3). SEM micrographs of all indumentum types are presented in Figure 1.

In the sect. *Scordium*, *T. scordium* subsp. *scordium* leaves have sparsely B, C1 and C2 trichome types on both surface, although *T. sirnakense* leaves have A1, A2, B type glandular trichomes and C2 and C3 non-glandular trichomes on the adaxial and abaxial surface. Dense indumentum appears on the abaxial surface than the adaxial surface of this species leaves (Figure 1).

In the Sect. *Isotriodon*, *T. chasmophyticum* has A1, B, D trichome types on both surface of the leaves. The lower surface of the leaf has a much denser indumentum than the upper surface (Figure 1). *T. andrusi* has A1, A2, B, C2, C3 trichome types adaxial and abaxial surface of the leaves. Dense indumentum appears on both surface of the leaves (Figure 1). *Teucrium spinosum* belongs the sect. *Spinularia*, A2, B, C2 trichome types are observed at the abaxial surface of leaves; A2, B, C3 trichome types were found on the adaxial surface of leaves (Figure 1).

## DISCUSSION

Trichome morphology is the useful taxonomic markers in some genera of Lamiaceae. Their absence or presence can be used as taxonomic characters in the infrageneric classification of some genera (Metcalf and Chalk 1950; Navarro and El Oualidi 2000; Moon et al. 2009). For the classification of trichome in *Teucrium*, distinction into thin and thick-walled provides taxonomic support to the delimitation of the species, could be regarded as a valid taxonomic character (Navarro

and El Oualidi; Eshratifar et al. 2011; Ecevit-Genç et al. 2015, Ecevit-Genç et al. 2017).

As a result of our work, seven trichome types were observed in five species belong to three different section of *Teucrium* (Table 2). The trichome types located on the adaxial and abaxial sides of leaves are same except *T. spinosum*. Among them subsessile glandular trichomes are most widespread in all taxa examined. Clavate glandular trichomes are generally found all of the species except *T. scordium* subsp. *scordium*. 2-celled thick-wall non-glandular trichomes are common trichome types but only *T. chasmophyticum* have thin-wall non-glandular trichome. In this work, the leaves trichome micro-morphology of *T. sirnakense*, *T. chasmophytum* are reported in detail by SEM for the first time.

Many studies have been presented about leaves indumentum of *Teucrium* species. For example, 56 *Teucrium* species related to the nine sections were investigated by Navarro and El Oualidi (2000). The authors analyzed five species of the sect. *Spinularia* including *T. spinosum* and they found different trichome types between species. According to their results, *T. spinosum* has flexuose and elongated thin-walled non-glandular trichomes and subsessile glandular trichomes. In addition to this species, long clavate glandular and large thin-walled non-glandular trichome types were identified in this study. The author have investigated four species of sect. *Isotriodon*. Thick-walled non-glandular and long clavate glandular trichomes are most common trichomes and flexuose and elongated thin-walled trichome types are less frequent in this section according to Navarro and El Oualidi (2000). But elongated 5-7(8)-celled thick-wall non-glandular trichomes was not observed on any taxa of studied in this study. Also long clavate glandular trichomes were not observed on *T. chasmophytum*.

We have observed subsessile glandular trichomes and large thin-walled non-glandular trichomes on *T. scordium* subsp. *scordium* leaves. The other subspecies of *T. scordium* subsp. *scordioides* leaves indumentum was investigated many previous studies and long clavate glandular trichomes, subsessile glandular trichomes and flexuose and elongated thin-walled

**Table 3. Trichome types and distribution on the adaxial-abaxial leaf surfaces of studied *Teucrium* taxa**

Section	Taxon	Leaves trichomes (According to Navarro & El Oualidi 2000)	
		Adaxial surface	Abaxial surface
Scordium	<i>T. scordium</i> subsp. <i>scordium</i>	B, C1, C2	B, C1, C2
	<i>T. sirnakense</i>	A1, A2, B, C2, C3	A1, A2, B, C2, C3
Isotriodon	<i>T. chasmophyticum</i>	A1, B, D	A1, B, D
	<i>T. andrusi</i>	A1, A2, B, C2, C3	A1, A2, B, C2, C3
Spinularia	<i>T. spinosum</i>	A2, B, C2	A2, B, C2




non-glandular trichomes were found by Navarro and El Oualidi (2000) and Ecevit-Genç et al. (2017). Peltate, capitate acicular and flagelliform trichome types were found by Jurišić Grubešić et al. (2007). According to all these findings trichome types are show some differences between two sub-species.

*T. scordium* investigated morpho-anatomically by Lakusic et al. (2010) from Balkan peninsula and they found glandular (peltate, unicellular capitate, multicellular capitate), and non-

glandular (unicellular unbranched, multicellular unbranched) trichomes on the leaves of this species. But the unicellular trichomes were not observed on the leaves of *T. scordium* subsp. *scordium* in this study.

Clavate glandular trichomes are ordinarily observed on the leaves of taxa of sect. *Isotriodon* by our team's previous paper (Ecevit-Genç et al. 2017). The results of this study about the species of the sect. *Isotriodon* are overlapped with the previous study of our team. *T. andrusi* leaves were examined by



**Figure 1.** a-d. SEM micrographs of leaves of *Teucrium*. Adaxial surface (a, b), abaxial surface (c, d). *T. scordium* subsp. *scordium* (1), *T. sirnakense* (2), *T. chasmophytum* (3), *T. andrusi* (4), *T. spinosum* (5). (scale bars: a, c =500 micrometer (µm); b, d =200 µm).

Dinç et al. (2011) and their results are corresponding to our results.

This paper gives detailed information on the micromorphological features of the *T. spinosum*, *T. scordium* subsp. *scordium*, *T. sirnakense*, *T. chasmophytum* and *T. andrusi* species. We concluded that trichome types are useful for specific delimitation of *Teucrium* species. However, micromorphological features must be supported by other morphological, molecular, biogeographical characters.

### Acknowledgements

This work was supported by the Research Fund of Istanbul University (Project number 31081) and Research Fund of Balıkesir University (Project number 2012/8).

### REFERENCES

- Abu-Assab MS, Cantino PD, (1993). Phylogenetic implications of pollen morphology in tribe Ajugeae (Labiatae). *Syst Bot* **18**: 100–122.
- Aksoy-Sagirlı, Ozsoy N, Ecevit-Genç G, Melikoglu G (2015). *In vitro* antioxidant activity, cyclooxygenase-2, thioredoxin reductase inhibition and DNA protection properties of *Teucrium sandracicum* L. *Ind Crop Prod* **74**: 545–550.
- Behnke HD (1984). Plant trichomes, structure and ultrastructure. In: Rodriguez E, Healey PL, Metha I (eds.). *Biology and chemistry of plant trichomes*. Plenum Press, New York. pp. 1–21.
- Beyrouthy EM, Arnold-Apostolides N, Dupont F (2009). Trichomes morphology of six Lebanese species of *Stachys* (Lamiaceae). *Flora Medit* **19**: 129–139.
- Cantino PD, Harley RM, Wagstaff SJ (1992). Genera of Labiatae: Status and Classification. In: Harley RM, Reynolds T (eds.). *Advances in Labiate Science. Royal Botanic Gardens, Kew*, pp. 511–522.
- Dinç M, Doğu S, (2016). *Teucrium pruinosum* var. *aksarayense* var. nov. (Lamiaceae) from Central Anatolia, Turkey. *Modern Phytomorphology* **9**: 13–17.
- Dinç M, Doğu S, Bağcı Y (2011). Taxonomic reinstatement of *Teucrium andrusi* from *T. paederotoides* based on morphological and anatomical evidences. *Nord J Bot* **29**: 148–158.
- Dinç M, Doğu S, Bilgili B, Duran A (2009). Comparative anatomical and micromorphological studies on *Teucrium creticum* and *Teucrium orientale* var. *orientale* (*Teucrium* sect. *Teucrium*, Lamiaceae). *Nord J Bot* **27**: 251–256.
- Dirmenci T (2012). *Teucrium* L. In: Güner A, Aslan S, Ekim T, Vural M, Babaç MT (eds.). *Türkiye Bitkileri Listesi (Damarlı Bitkiler)*. Nezahat Gökyiğit Botanik Bahçesi ve Flora Araştırmaları Derneği Yayını, İstanbul, pp. 595–598.
- Doğu S, Dinç M, Kaya A, Demirci B (2013). Taxonomic status of the subspecies of *Teucrium lamiifolium* in Turkey: Reevaluation based on macro-and micro-morphology, anatomy and chemistry. *Nord J Bot* **31**: 198–207.
- Dönmez AA (2006). *Teucrium chasmophyticum* Rech. f. (Lamiaceae): A new record for the flora of Turkey. *Turk J Bot* **30**: 317–320.
- Dönmez AA, Mutlu B, Özçelik AD (2010). *Teucrium melissoides* Boiss. and Hauskn. ex Boiss. (Lamiaceae): A new record for Flora of Turkey. *Hacettepe J Biology and Chemistry* **38**: 291–294.
- Duman H (2000). *Teucrium* L. In: Güner A, Özhatay N, Ekim T, Başer KHC (eds.). *Flora of Turkey and the east Aegean islands*, Vol 11, Edinburg University Press, Edinburgh, pp. 197–198.
- Ecevit-Genç G, Özcan T, Dirmenci T (2015). Micromorphological characters on nutlet and leaf indumentum of *Teucrium* sect. *Teucrium* (Lamiaceae) in Turkey. *Turk J Bot* **39**: 439–448.
- Ecevit-Genç G, Özcan T, Dirmenci T (2017). Nutlet and leaf micromorphology in some Turkish species of *Teucrium* L. (Lamiaceae). *Phytotaxa* **321**: 71–82.
- Ekim T (1982). *Teucrium* L. In: Davis, PH, (eds.). *Flora of Turkey and the East Aegean Islands islands*, Vol 7, Edinburg University Press, Edinburgh, pp. 53–75.
- Eshratifar M, Attar F, Mahdigholi K (2011). Micromorphological studies on nutlet and leaf indumentum of genus *Teucrium* L. (Lamiaceae) in Iran. *Turk J Bot* **35**: 25–35.
- Genç İ, Kültür Ş, Ecevit-Genç G (2017). Capsule, leaf and gland morphology of Turkish perennial taxa of *Euphorbia* L. section *Pithyusa* (raf.) Lazoro. *Trakya J Natural Sciences* **19**: 11–19. Govaerts R, (1999). *World Checklist Seed Plants 3*. Continental Publishing, Deurne.
- Grubescic RJ, Vladimir-Knezevic S, Kremer D, Kalodera Z, Vukovic J (2007). Trichome micromorphology in *Teucrium* (Lamiaceae) species growing in Croatia. *Biologia* **62**: 148–156.
- Jurišić Grubescic RJ, Vladimir-Knezevic S, Kremer D, Kalodera Z, Vukovic J (2007). Trichome micromorphology in *Teucrium* (Lamiaceae) species growing in Croatia. *Biologia* **62**: 148–156.
- Kastner A (1989). Übersicht zur systematischen gliederung der gattung *Teucrium* L. *Biocosme Mésogéen* **6**: 63–77.
- Kaya A, Satil F, Dirmenci T, Selvi S (2012). Trichome micromorphology in Turkish species of *Ziziphora* (Lamiaceae) *Nord J Bot* **30**: 1–8.
- Khalik KA, Hassan NMS (2012). Seed and trichome morphology of the Egyptian *Fagonia* (Zygophyllaceae) with emphasis on their systematic implications. *Nord J Bot* **30**: 116–126.
- Lakusic B, Stevanovic B, Jancic R, Lakusic D (2010). Habitat-related adaptations in morphology and anatomy of *Teucrium* (Lamiaceae) species from the Balkan peninsula. *Flora* **205**: 633–646.
- Marin PD, Petkovic B, Duletic S (1994). Nutlet sculpturing of selected *Teucrium* species (Lamiaceae): a character of taxonomic significance. *Plant Syst Evol* **192**: 199–214.
- McClintock E, Epling C (1946). A revision of *Teucrium* in the new world, with observations on its variation, geographical distribution and history. *Brittonia* **5**: 491–510.
- Metcalfe CR, Chalk L (1950). *Anatomy of the Dicotyledons*, Vol. 2, Oxford Press, UK:
- Moon HK, Hong SP, Smets E, Huysmans S (2009). Phylogenetic significance of leaf micromorphology and anatomy in the tribe Mentheae (Nepetoideae: Lamiaceae). *Flora* **205**: 633–264631.
- Navarro T, El Oualidi J (2000). Trichome morphology in *Teucrium* L. (Labiatae) a taxonomic review. *An Jardin Bot Madrid* **57**: 277–297.
- Osman AK (2012). Trichome micromorphology of Egyptian *Bal-lota* (Lamiaceae) with emphasis on its systematic implication. *Pak J Bot* **44**: 33–46.
- Özcan T, Dirmenci T, Coşkun F, Akçiçek E, Güner O (2015). A new species of *Teucrium* sect. *Scordium* (Lamiaceae) from SE of Turkey. *Turk J Bot* **39**: 310–317.
- Parolly G, Eren O (eds.) (2007). Contributions to the flora of Turkey, 2. *Willdenowia* **37**: 243–271.

- Salmaki Y, Zarre S, Jamzad Z, Bräuchler C (2009). Trichome micromorphology of Iranian *Stachys* (Lamiaceae) with emphasis on its systematic implication *Flora* **204**: 371–381
- Tutin TG, Wood D (1972). *Teucrium* L. In: Heywood, VH, Burges, NA, Moore, DM, Valentine, DH, Walters, SM, Webb, DA (eds.). *Flora Europaea*, Vol. 3, Cambridge University Press, Cambridge.
- Vural M, Duman H, Dirmenci T, Özcan T (2015). A new species of *Teucrium* sect. *Stachyobotrys* (Lamiaceae) from the south of Turkey. *Turk J Bot* **30**: 318–324.
- Zareh M, Faried A, Farghaly N (2017). Micromorphological studies on the genus *Lotus* L. (Fabaceae: Loteae) from Egypt. *Turk J Bot* **41**: 273-288.