

## Seizure worsening caused by low serum valproate levels from an interaction between valproate and meropenem

Valproat ve meropenem arasındaki etkileşime bağlı olarak serum valproat düzeyinin düşmesinin ortaya çıkardığı nöbetler

Suat BİÇER, Gülay ÇİLER ERDAĞ, Canan KOCAMAN YILDIRIM, Tuba GİRAY, Defne ÇÖL, Öznur KÜÇÜK, Zerrin YALVAÇ, Ayça VİTRİNEL

### ABSTRACT

We present an interaction of meropenem with valproate in an epileptic child, leading to seizure exacerbations owing to the rapid lowering of serum valproate concentration. An increase of seizure frequency and somnolence were observed in the patient after the addition of meropenem to the treatment, and a rapid decline of valproate serum concentrations was observed after two doses of meropenem. This decline was the most likely cause of the increase in seizure frequency. The dosage of valproate was raised and meropenem was stopped. Two days later, the seizures stopped. Five days later, the serum valproate concentrations raised to three fold, and they rose to therapeutic levels four weeks later. To avoid drug interaction that reduces the serum concentration of valproate, meropenem in epileptic patients using valproate for the treatment of epilepsy should be administered cautiously. If concomitant administration is essential, close observation of serum concentration of valproate and clinical course of the patient are necessary.

**Keywords:** Carbapenems, Drug interactions, Meropenem, Seizures, Valproate

### ÖZET

Bu yazıda, meropenem ve valproat etkileşimine sekonder olarak serum valproat düzeyinin hızlı düşmesi sonucu nöbetleri alevlenen epileptik bir çocuk sunulmuştur. Hastanın tedavisine meropenem eklendikten ve iki doz kullanıldıktan sonra nöbet sıklığı ve somnolans durumunda artış ortaya çıkan hastanın serum valproat düzeyinde ani düşme görüldü. Nöbet sıklığındaki artışın valproat serum düzeyinin düşüşüne bağlı olduğu düşünüldü. Hastanın valproat dozu artırılarak meropenem tedavisi kesildikten iki gün sonra nöbetler sona erdi. Beş gün sonra serum valproat düzeyi üç katına, dört hafta sonra ise terapötik düzeylere yükseldi. Serum valproat düzeyinin düşmesine neden olan ilaç etkileşimine neden olmamak için, valproat kullanan epileptik hastalarda meropenem dikkatle uygulanmalı, iki ilacın birlikte kullanımı mutlaka gerekliyse hastanın klinik durumu ve serum valproat düzeyi yakından izlenmelidir.

**Anahtar kelimeler:** İlaç etkileşimi, Karbapenemler, Meropenem, Nöbetler, Valproat

### Introduction

Meropenem, is a carbapenem antibiotic, and has a broad spectrum of antimicrobial activity. Valproate is widely used in the treatment of epileptic seizures in children. In recent years, low serum concentrations of valproate were reported in children and adults receiving concomitant treatment with meropenem [1]. We present an interaction of meropenem with valproate in an epileptic child, leading to seizure exacerbations owing to the rapid lowering of serum valproate concentration.

### Case report

This report describes a fourteen-year-old epileptic inpatient girl who had received concurrent treatment with meropenem for the treatment of lobar pneumonia and pleuresia, and

Suat Biçer (✉), Gülay Çiler Erdağ, Tuba Giray, Defne Çöl, Öznur Küçük, Zerrin Yalvaç, Ayça Vitrinel  
Department of Child Health and Pediatrics, School of Medicine, Yeditepe University, Istanbul, Turkey  
e-mail: suat.bicer@yeditepe.edu.tr

Canan Kocaman Yıldırım  
Pediatric Neurology Clinic, Erdem Private Hospital, Istanbul, Turkey

Submitted/Gönderilme: 25.01.2015 Accepted/Kabul: 27.03.2015



and partial seizures [2]. This drug is heavily attached to plasma proteins and largely metabolized in the liver by glucuronidation (70%). Beta-oxidation, omega-oxidation and hydroxylation are minor pathways in the metabolism of valproate. All carbapenems, including meropenem, have lowered the serum concentration of valproate in epileptic patients to a sub-therapeutic level. Because serum valproate concentration was not decreased by carbapenems in hepatectomized rats, the liver is considered as the key organ for the decrease of valproate concentration by carbapenems. Valproate-glucuronidase in human liver microsomes, and especially in cytosol, was inhibited by carbapenems [5]. The interaction between valproate and a carbapenem (panipenem/betamipron) was first reported by Nagai et al [6] in two children in 1997. Other cases have been reported since then; as in our case, seizures have occurred during concomitant therapy of valproate and meropenem [1,2,7,8].

Decline in valproate serum concentration and seizures were observed within 24 hours after concomitant administration of valproate and meropenem in our patient. The most rapid decrease of serum valproate within 24 [7] and 36 hours [6] were reported in other studies. In some other reports, this decline was observed within three to seven days after the initiation of meropenem to patients receiving valproate therapy [1]. This rapid decline in serum concentration of valproate is not explained with enzyme induction because this mechanism usually requires several days [9]. Inhibition of plasma protein binding of valproate by carbapenems has been suggested for the interaction between valproate and carbapenems [5,7]. Valproate has a high protein-binding ratio (90 - 95%), whereas, meropenem is a low (2%) protein bound drug. For this reason, these rapid declines may not be explained by the mechanism of competition in protein binding [10]. The most comprehensive research study on this topic was undertaken by Yokogawa et al [10]. They studied the effects of meropenem on valproate metabolism in rabbits, because metabolism of this drug is similar to that in humans. This study indicated that urinary excretion of valproate-glucuronide, and suppression of valproate-glucuronide hydrolysis in the liver were increased at the time of concomitant administration of valproate and meropenem [10]. Furthermore, suppression of valproate-glucuronide hydrolysis in the gastrointestinal tract was also increased in this concomitance [10]. The mechanisms of rapid decline of valproate serum concentration has been elucidated by recent studies. Some studies have indicated that the clearance of valproate is increased by meropenem by increasing the urinary excretion of valproate glucuronide [7,8,10]. Other mechanisms in which decreases occur in the serum levels of valproate by carbapenems are: inhibition

of the hydrolysis of valproate glucuronide, increase in the glucuronidation of valproate, suppression of enterohepatic recirculation of valproate, and increase in the erythrocyte distribution of valproate, which induce the decrease in serum levels of valproate [11]. The reduced blood levels of valproate at a fairly short time period in our patient may be caused by decreased hydrolysis of valproate-glucuronide conjugation resulting from eliminating of microorganism with beta-glucuronidase activity owing to antibacterial drugs. For these reasons, reabsorption of free valproate decreases.

Carbapenems may induce seizures during acute neurologic diseases as intracranial infections, hemorrhage, stroke, trauma, and tumors [3,9]. None of these have occurred in our patient during concomitant therapy of valproate and meropenem.

In the reported cases to date, none of the antibiotics (i.e. ceftazidime, cefotaxime, amikacin, vancomycin, ampicillin, tobramycin, and clindamycin) other than carbapenems have caused a decline of the serum concentration of valproate. In addition, serum valproate concentrations increased after the termination of carbapenems according to these reports [1]. Rapidly changed serum concentrations of valproate is not consistent with the induction of enzymes participated in valproate metabolism process.

Serum valproate concentration was decreased while under meropenem treatment and gradually returned to normal levels after cessation of meropenem in a reported 14-year old girl with cystic fibrosis and multidrug resistant pulmonary infection with intravenous antibiotics [12]. Her serum valproate concentration kept low despite elevated daily dosing of valproate and a loading dose. Finally, on the 18th day of cessation of meropenem the blood valproate level reached its normal value. The daily doses were increased, however, the loading dose was not used in our case. The low blood levels of valproate until the 36th day may be related with inadequate dose of valproate without the loading dose. Conversely, seizures were controlled after two days and the serum valproate concentration rose to three-fold after the discontinuation of meropenem and increased daily valproate doses. Therefore, the loading valproate dose with increased daily doses may be used to increase the blood valproate levels.

Clinicians should be aware of the interaction between valproate and carbapenems and of the possible clinical consequences like increasing the frequency of epileptic seizures and/or status epilepticus. To avoid drug interaction that reduces the serum concentration of valproate, care should be taken when administering meropenem in epileptic

patients using valproate for the treatment of epilepsy. If possible, another antibiotic, except carbapenems, should be used in patients using valproate or the antiepileptic therapy should be switched to another drug other than valproate. However, this second option may not be feasible in many patients if their seizures are controlled only with valproate. In addition, if the antiepileptic drug regime has changed, at least two – four weeks will be required for sufficient levels of the new antiepileptic drug to be attained [1]. If concomitant administration is essential, close observation of serum concentration of valproate and clinical course of patient are necessary.

There were not plasma levels of other antiepileptic drugs excluding valproate.

#### Acknowledgement

We are grateful to Cahit Zereyçan (writing consultant of Writing Center at Yeditepe University) for the revision of this paper in English.

**Financial disclosure:** None

**Funding:** This study was not funded.

**Author contributions:** Suat Biçer developed the idea of the case report. He abstracted analysed, prepared, wrote and drafted it. Gülay Çiler Erdağ, Canan Kocaman, Tuba Giray, Defne Çöl, Öznur Küçük, Zerrin Yalvaç, and Ayça Vitrinel contributed the case report. Ayça Vitrinel and Canan Kocaman revised the case report for important intellectual content. Ayça Vitrinel revised it for English.

#### References

- Nacarkucuk E, Saglam H, Okan M. Meropenem decreases serum level of valproic acid. *Pediatr Neurol* 2004; 31: 232-4. <http://dx.doi.org/10.1016/j.pediatrneurol.2004.03.014>
- De Turck BJ, Diltoer MW, Cornelis PJ, et al. Lowering of plasma valproic acid concentrations during concomitant therapy with meropenem and amikacin. *J Antimicrob Chemother* 1998; 42: 563-4. <http://dx.doi.org/10.1093/jac/42.4.563>
- Calandra G, Lydick E, Carrigan J, Weiss L, Guess H. Factors predisposing to seizures in seriously ill infected patients receiving antibiotics: Experience with imipenem/cilastatin. *Am J Med* 1988; 84: 911-8.
- Cunha BA. Meropenem in elderly and renally impaired patients (corrected and republished in *Int J Antimicrob Agents* 1999; 11: 167-77). *Int J Antimicrob Agents* 1998; 10: 107-17. [http://dx.doi.org/10.1016/S0924-8579\(98\)00031-4](http://dx.doi.org/10.1016/S0924-8579(98)00031-4)
- Nakamura Y, Nakahira K, Mizutani T. Decreased valproate level caused by VPA-glucuronidase inhibition by carbapenem antibiotics. *Drug Metab Lett* 2008; 2: 280-5. <http://dx.doi.org/10.2174/187231208786734049>
- Nagai K, Shimizu T, Togo A, et al. Decrease in serum levels of valproic acid during treatment with a new carbapenem, panipenem/betamipron. *J Antimicrob Chemother* 1997; 39: 295-6. <http://dx.doi.org/10.1093/jac/39.2.295>
- Spriet I, Meersseman W, De Troy E, Wilmer A, Casteels M, Willems L. Meropenem-valproic acid interaction in patients with cefepime-associated status epilepticus. *Am J Health Syst Pharm* 2007; 64: 54-8. <http://dx.doi.org/10.2146/ajhp050512>
- Yoon H, Kim DH. Unusual drug reaction between valproate sodium and meropenem. *Int J Clin Pharm* 2013; 35: 316-8. <http://dx.doi:10.1007/s11096-013-9763-2>.
- Mann HI. Drug-associated disease: Cytochrome P450 interactions. *Crit Care Clin* 2006; 22: 329-45. <http://dx.doi.org/10.1016/j.ccc.2006.02.004>
- Yokogawa K, Iwashita S, Kubota A, et al. Effect of meropenem on disposition kinetics of valproate and its metabolites in rabbits. *Pharm Res* 2001; 18: 1320-6.
- Ogawa K, Yumoto R, Hamada N, Nagai J, Takano M. Interaction of valproic acid and carbapenem antibiotics with multidrug resistance-associated proteins in rat erythrocyte membranes. *Epilepsy Res* 2006; 71: 76-87. <http://dx.doi.org/10.1016/j.eplepsyres.2006.05.016>
- Taha FA, Hammond DN, Sheth RD. Seizures from valproate-carbapenem interaction. *Pediatric Neurology* 2013; 49: 279-81.