



Effect of Conception Time and Maternal Age Upon Fetal Sex Ratio in Çukurova Population

Çukurova Populasyonunda Gebelik Zamanı ve Maternal Yaşın Fetal Cinsiyet Oranı Üzerine Olan Etkisi

Erdal Tunç¹, Osman Demirhan¹, Ayfer Pazarbaşı¹, Deniz Taştemiş², Ali İrfan Güzel³, Mülkiye Kasap¹, Fatma Tuncay Özgünen⁴, Bertan Yılmaz¹, Sabriye Kocatürk-Sel¹

¹Çukurova University Faculty of Medicine, Department of Medical Biology and Genetics and ⁴Gynecology and Obstetrics Department. ADANA

²Adiyaman University, Vocational School of Health Sciences, ADIYAMAN

³Recep Tayyip Erdogan University, Faculty of Medicine. Department of Medical Biology and Genetics, RİZE

Cukurova Medical Journal 2014 39(2):248-255.

ABSTRACT

Purpose: This investigation was conducted to determine whether the month of conception, parental age, and environmental factors such as seasonal variation, consumption of alcohol and cigarette by parents, play a role on sex ratio at pregnancy in Çukurova region and surrounding area population (Turkey).

Materials and Methods: We have analyzed the data of patients who were referred to our prenatal diagnosis laboratory for amniocentesis with the purpose of karyotyping between the years of 2005-2007. We used Pearson Chi-square, Fisher's exact and T-tests for statistical analyses.

Results: No association was established between the primary sex ratio and time for pregnancy. Differences in primary sex ratio among the months were not statistically significant, but the difference for the month of September was noted. For September, we have found statistically significant distortion of sex ratio toward male, (males:females ratio=1.8).

Conclusion: In conclusion, our speculation is that the climate change in September may affect the primary sex ratio under the summer/autumn photoperiod.

Key Words: Karyotyping, month of conception, primary sex ratio, prenatal diagnosis

ÖZET

Amaç: Bu çalışmanın amacı, Çukurova ve çevresindeki populasyonda konsepsiyon ayı, ebevyenlerin yaşları, mevsimsel değişimler, ebevyenlerin alkol ve sigara kullanım durumları gibi çevresel değişkenlerin fetal cinsiyet oranını etkileyip etkilemediklerini tespit etmektir.

Materyal ve Metod: Bu çalışmada, 2005-2007 yılları arasında amniyosentez maksadıyla prenatal tanı laboratuvarımıza yönlendirilen hastaların verilerini analiz ettik. İstatistiksel analiz için ki-kare, Fisher's exact ve T testlerini uyguladık.

Bulgular: Konsepsiyon zamanı ile primer cinsiyet oranı arasında bir ilişki tespit edilemedi. Aylara göre primer cinsiyet oranındaki değişimlerin istatistiksel olarak anlamlı olmadığı gözlemlendi. İstisna olarak eylül ayındaki değişim kaydedildi. Eylül ayında istatistiksel olarak anlamlı bir şekilde cinsiyet oranının erkeğe doğru değiştiği saptandı (erkek:kadın oranı=1.8).

Sonuç: Yaz/sonbahar fotoperiyodunda bulunan eylül ayındaki iklimsel değişimin, cinsiyet oranındaki bu farklılaşmayı etkilediğini düşünmekteyiz.

Anahtar Kelimeler: Karyotipleme, konsepsiyon ayı, primer cinsiyet oranı, prenatal tanı

INTRODUCTION

Is there any relation between the maternal age and the month of conception, and the fetal sex ratio? This question draws attention of many people, specially couples who want to have a child with a specific gender (boy or girl). For answering this question, there is a method developed by chinese, chinese calender method. According to this method, firstly, the maternal age and the date (month) of conception must be determined, and the date of conception must be converted into their lunar equivalents. So, a predictive value for having a boy or girl can be calculated. The properly applied Chinese Gender Chart gives couples at least a 75% chance of having a child of the desired sex. Some researchers have reported success rates up to 90%. Therefore, it's considered that there is a relation between offspring sex ratio and time to pregnancy¹.

The sex ratio at birth (SRB) is rarely 1.0 (males:females) for most vertebrates and humans². In humans, the SRB is shifted towards boys by some 2–3%³. For most human populations the sex ratio at birth, defined as the number of male births per 100 female births, is around 105⁴⁻⁷. However, notable variations from this mean value have been observed in certain human populations^{6,8,9}. In recent years, some published reports focused on whether some environmental factors affecting before and during periconceptional period, have an impact on sex ratio. At the same time, several biological mechanisms have been suggested to explain the remarkable constancy of this ratio. As a practical matter, the difficulties of measuring these biological factors have persuaded most researchers to rely on less direct, and more easily observable factors underlying the sex ratio¹⁰. This is the reason why variables such as the ages of

the parents and the birth order have been widely studied.

A seasonal variation in human SRB values has been identified for the German population¹¹. This investigation analyzed birth records and found a highly significant, albeit low-amplitude, annual rhythm, with the highest values (relatively the most boys born) between April and June and the lowest values in October. Seasonality and period effects are among the other factors that have drawn attention. Significant variations in the sex ratio by month of birth have been observed in certain populations^{8,12,13,14} but these variations are generally very small.

Our investigation was conducted to determine whether the month of conception, maternal age, environmental factors such as seasonal variation, consumption of alcohol and cigarette by parents, and parental age play a role on sex ratio at pregnancy in Çukurova region, Adana and surrounding area population (Turkey).

MATERIALS and METHODS

We have analyzed the data of patients who were directed from Gynecology and Obstetrics Department of Çukurova University to our Genetic Laboratory, at the Medical Biology and Genetics Department of Çukurova University, for amniosynthesis with purpose of karyotyping between the years of 2005-2007. Birth records (number of boys and girls born per month) were obtained from our prenatal laboratory records. We asked parents or the closest relative of parents about the alcohol and cigarette consumption of parents, ages of parents, and the time of conception. We also asked the mothers about where conception had been occurred. Since our hospital is a reference hospital for a large region, including subregions with different climate

conditions, we classified our patients according to subregions where conception had been occurred to understand the climate's effect on sex ratio. This study was conducted on 2155 cases. Monthly average temperature, photoperiod and rainfall values were obtained from the National Climatic Data Center in Adana-Turkey for the period 1975–2006 (Table 1). Correlations were assessed for

possible effects of temperature on the sex ratio at pregnancy (SRP). We have analysed the effects of all forementioned parameters on primary sex ratio (SRP), employing statistical methods. We used Pearsons Chi-square, Fisher's exact and t-tests for statistical analyses. This study has been approved by the local institutional review board.

Table 1. Monthly weather values for Adana city between years of 1975-2006 are presented www.meteor.gov.tr (June-2009).

Adana city-TURKEY	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec
Average values for years between 1975-2006												
Average temperature(°C)	9.7	10.4	13.3	17.5	21.7	25.6	28.3	28.4	26.1	21.6	15.3	11.1
Average time with daylight (hour)	4.6	5.2	5.8	6.8	9.0	10.4	10.5	10.2	8.6	7.1	5.7	4.3
Average of days with rainfall	10.5	9.8	10.0	10.4	6.9	3.5	1.9	1.5	3.2	5.7	7.6	10.6
Highest and lowest values for years between 1975-2006												
Highest temperature(°C)	23.0	25.0	32.0	36.8	40.6	41.3	44.0	43.8	43.2	39.4	33.3	27.4
Lowest temperature(°C)	-4.2	-6.4	-3.6	-1.3	5.6	13.7	16.8	16.8	10.9	4.8	-1.0	-3.5


RESULTS

In this study, we analyzed the effects of parameters; month of conception, maternal age, environmental factors such as seasonal variation, alcohol consumption and cigarette smoking on sex ratio at pregnancy in population of Çukurova region, Adana and surrounding area. In sum, we studied 2155 cases. But only for 893 cases we were able to obtain information including all mentioned parameters by directly interviewing with mothers or their close relatives. For other 1262

cases we evaluated their data retrospectively, so we could only get information about maternal age during pregnancy, month of pregnancy and subregion where pregnancy might occur. The mean age of the mother and father screened was 32.5 (range 17-48) and 36.1 years (range 20-65), respectively. In total, we calculated the value of sex ratio at pregnancy as male:female ratio=1.02 (49,5% of cases were recorded as female whereas 50,5% of them were male) (Table 2).

Table 2. Primary sex ratio by month

Month	male (%)	female (%)	Total	M/F ratio
1	76 (45,5)	91 (54,5)	167	0.84
2	92 (55,4)	74 (44,6)	166	1.24
3	83 (45,6)	99 (54,4)	182	0.84
4	95 (51,9)	88 (48,1)	183	1.08
5	77 (45,6)	92 (54,4)	169	0.84
6	87 (46,8)	99 (53,2)	186	0.88
7	88 (48,1)	95 (51,9)	183	0.93
8	101 (53,7)	87 (46,3)	188	1.16
9	115 (64,2)	64 (35,8)	179	1.80
10	123 (52,1)	113 (47,9)	236	1.09
11	80 (49,1)	83 (50,9)	163	0.96
12	72 (47,1)	81 (52,9)	153	0.89
Total	1089 (50,5)	1066 (49,5)	2155	1.02

**Figure 1.** Primary sex ratio at pregnancy by month

We analyzed the effects of per months on primary sex ratio at pregnancy. Monthly values for the sex ratio of 2155 cases are shown in Figure 1 and Table 2. Differences among the months were not statistically significant, except marked

difference for the month of September. The sex ratio is significantly higher than 105 for September (M/F ratio=1.80) (Table 2 and Figure 1). We statistically tested the effect of getting pregnant naturally and getting pregnant with treatment

categories on sex ratio at pregnancy. We performed non-parametric chi-square test for mentioned categories and we did not find any statistically significant difference from normal value of primary sex ratio at birth ($p>0,05$) (Tablo 3). For this analysis we utilized answers of question how and of which location pregnancy had occurred. In this regard, we classified our patients into three categories; getting pregnant in hometown, in seaside residence and in high altitude residence. For statistical analyses of these categories we also employed non-parametric chi-square test however we did not find any statistically significant difference from normal value of sex ratio at birth ($p>0,05$) for all categories (Tablo 3). Our patients came from three different subregions which have different climate conditions (Central anatolia region, southeastern region and mediterranean region). Therefore, we wanted to test whether there is an effect of different climate conditions on sex ratio at

pregnancy. However, we did not find any statistically significant difference from normal value of primary sex ratio at pregnancy for all three subregions ($p>0,05$) (Tablo 3). We also wanted to test the effects of parents' alcohol and cigarette consumption on sex ratio at pregnancy. According to our analyses parental alcohol and cigarette consumption also did not have any effect on primary sex ratio at pregnancy (non-parametric chi-square test, $p>0,05$) (Tablo 3). Finally, we tested the effect of parental age on primary sex ratio, employing T test. We did not find statistically significant distortion of sex ratio at pregnancy from normal value of 105 under the effect of parental age (Tablo 3). In this study, for all parameters we tested with chi-square test, we also performed Fisher's Exact test. As a result, we did not find any statistically significant difference from normal value (105) of primary sex ratio at pregnancy for all mentioned parameters .

Table 3. Male and female distributions for constituted categories

How pregnancy had occurred	Male	Female	P value
Naturally	439	413	0,338
With treatment	22	14	
Where pregnancy had occurred			
Home	399	393	0,353
Seaside home	48	34	
Plateau home	9	10	
Subregion patients came from			
Mediterranean	1039	1030	0,445
Southeastern	41	30	
Central anatolia	7	6	
Mothers' cigarette consumption			
No	392	370	0,650
yes	64	67	
Mothers' alcohol consumption			
No	455	435	0,617
Yes	1	2	
Fathers' cigarette consumption			
No	217	191	0,259
yes	238	246	
Fathers' alcohol consumption			
No	382	357	0,411
Yes	73	80	

DISCUSSION

We tested the effects of several candidate variables that could play roles upon sex ratio values independently, for the Çukurova and surrounding area population, Adana-Turkey.

For most human populations the sex ratio at birth, defined as the number of male births per 100 female births, is around 105⁵⁻⁷ and the ratio is rarely 1.0 (males:females) for most vertebrates and humans¹⁵. We calculated the value of primary sex ratio at pregnancy as 1.02 (males:females) (50,5% of cases was recorded as male versus 49,5% of female, Table 2). It's well known that human semen contains equal amounts of X and Y bearing sperms. But, as expressed above more boys than girls are born. The reasonable explanations for such distortion of sex ratio at birth toward male has been argued. It's hypothesised that Y bearing sperms swim faster than X bearing sperms in the viscous surrounding of cervical mucus¹⁶. Therefore, swimming capability in viscous surrounding must be accepted as an phenomenon leading to distortion of sex ratio at birth¹⁷.

Some variations in the sex ratio by month of birth have been observed in certain populations^{8,12,13,14} but these variations are generally very small. Several biological mechanisms have been suggested to explain the remarkable constancy of this ratio. In the present study, we did not find any association between time to pregnancy and primary sex ratio and we did not find any statistically significant distortion from normal value (105 males for 100 females) of sex ratio at pregnancy for any month except for September. But, according to Chinese calendar, it had been reported that there is a relation between the mothers' date of birth and the month of conception and sex ratio at birth. According to this method, firstly, the maternal age and the date (month) of conception must be determined, and the date of conception must be converted into their lunar equivalents. But, we used the sun calendar (the calendar now we are using) in our study. So, used calendar difference might have caused this

different result. We found an apparent and statistically significant distortion of sex ratio toward male, the highest values, rate of a boy is two times more than a girl (males:females ratio=1.8) in September. Our results supports those observed by Nonaka et al. (1999)¹⁴ in their study of the sex ratio in the French-Canadian population. Their study showed slightly lower values for the February-April and May-July quarters, and slightly higher values for the August-October and November-January quarters. Lerchl (1998)¹² observed significantly higher values for the months of May and December, and lower values for March and October in a study of postwar Germany. In the United States, it appears that the ratio may be slightly higher at the beginning of summer, and lower at the end of autumn⁸.

We observed significantly higher and lower values of climatic changes (temperatures, photoperiod and humidity) for September in our study (Table 1). We speculated that approximate climatic changes in this month can affect the primary sex ratio under the summer/autumn photoperiod. Photoperiodism and daylength affect hormonal mechanisms of living organisms and so, causes physiological and behavioral changes. A seasonal rhythm of human sex ratio has been described as inconsistent¹⁸⁻²⁰, its observed amplitude has been reported to be small (<0.5%)^{12,18,19}. Relative humidity is another environmental factor that may affect offspring sex ratio (Table 1). Its effect varies among species, and the effect may change with temperature²¹. Computerized literature searches have yielded no publication concerning temperature effects on the sex ratio at birth (SRB) in humans. However, there are only two relevant reports in the literature with respect to temperature effects on sex ratios in mammals: a study in rats showed decreased SRB values (i.e., an excess of female offspring) when animals were held and bred at 57 C for 20 generations²². The other investigation in wild bat colonies revealed relatively high SRB values at high environmental temperatures. The underlying

mechanism responsible for these phenomena, however, remains unclear. It may be speculated that the effects are due to processes within the testes: it has been shown²³, for example, that increased temperatures affect the dissociation of X and Y chromosomes during meiosis in rats and mice. Since scrotal skin temperatures in men show seasonal variations caused by the environmental temperatures²⁴, it is very likely that intratesticular temperatures also vary seasonally. Variations in temperature, humidity, and photoperiod may affect offspring sex ratio because they are used as seasonal cues. It can only be speculated that effects of these factors (temperature, photoperiod and humidity) are direct or indirect.

We have not found any significant effect of parental age. As elsewhere, in the Çukurova region the parental age does not show any effect upon sex ratio. It has been reported that the paternal age have a greater influence than maternal age, older fathers having a tendency to produce fewer sons than younger fathers⁹. However, Tremblay et al.(2003)²⁵ note a slight positive influence of paternal age (between 35 and 45) and a negative influence of maternal age (between 30 and 37) in the Saguenay population when birth order and parental age difference are taken into account. Since birth order is closely tied to parental ages, some researchers have attempted to separate their relative contributions. It appears that the influence of parental age is diminished when birth order is taken into account^{9,26,27}. Manning et al.(1997)²⁸ provide evidence of a positive relationship between parental age gap and the sex ratio for first births. However, other studies have not found a significant effect for spousal age difference²⁹⁻³¹. Alternatively, hormones may play a role in this context as well³². According to our analyses parental (both mother and father) alcohol and cigarette consumption also did not have an effect on sex ratio at birth ($p>0,05$).

In conclusion, our findings showed that there is no any association between time to pregnancy, maternal age and the fetal sex ratio of fetuses. We found an apparent and statistically significant distortion of primer sex ratio toward male, highest values, rate of a boy is two times more than a girl in September. Our speculation is that climate changes in this month can affect the primary sex ratio under the summer/autumn photoperiod.

REFERENCES

1. Weijin Z, Olsen J. Offspring sex ratio as an indicator of reproductive hazards. *Occup Environ Med* 1996; 53:503-4.
2. Clutton Brock TH, Iason GR. Sex ratio variation in mammals. *Q Rev Biol.* 1986;61:339-74.
3. Roenneberg T, Aschoff J. Annual rhythm of human reproduction: II. Environmental correlations. *J Biol Rhythms* 1990;5:217-39.
4. Cavalli-Sforza L, Bodmer WF. *The Genetics of Human Populations*. San Francisco, USA, Freeman, 1971.
5. Henry L, Blum A. *Techniques d'analyse en démographie historique*. Paris/INED: INSTITUT DE LA STATISTIQUE DU QUÉBEC, 1988. <http://www.stat.gouv.qc.ca/donstat/demograp/regiona/l/203.htm>. (page consulted on 8 March 2002).
6. Sieff DF. Explaining biased sex ratios in human populations: a critique of recent studies. *Curr Anthropol.* 1990;31:25-48.
7. Caselli G, Vallin J. Dynamique de la population : mouvement et structure. In: Caselli G, Valin J, Wunsch G, et al, Editors. *Démographie : analyse et synthèse*. Paris/INED, La dynamique des populations, 2001;l:35-79.
8. James WH. The human sex ratio. Part 1: a review of the literature. *Hum Biol.* 1987;59:721-52.
9. Chahnazarian A. Determinants of the sex ratio at birth: review of recent literature. *Soc Biol.* 1988; 35:214-35.
10. Allan BB, Brant R, Seidel JE, Jarrell JF. Declining sex ratios in Canada. *Can Med Assoc J.* 1997; 156:37-41.
11. Lerchl A. Sex ratios at birth and environmental temperatures. *Naturwissenschaften* 1999;86:340-2.
12. Lerchl A. Seasonality of sex ratio in Germany. *Hum Reprod* 1998;13:1401-2.
13. Nonaka K, Desjardins B, Charbonneau H, Legare J, Miura T. Marriage season, promptness of successful pregnancy and first-born sex ratio in a historical natural fertility population: evidence for sex-

- dependent early pregnancy loss?. *Int J Biometeorol* 1998;42:89-92.
14. Nonaka K, Desjardins B, Charbonneau H, Legare J, Miura T. Human sex ratio at birth and mother's birth season: multivariate analysis. *Hum Biol* 1999; 71:875-84.
 15. Clutton-Brock TH, Iason GR. Sex ratio variation in mammals. *Q Rev Biol* 1986;61:339-74.
 16. Keye WR, Chang RJ, Rebar RW, Soules MR. *Infertility; evaluation and treatment*. Philadelphia, USA, WB Saunders, 1995.
 17. Smits LJM, Bie RA, Essed GG and Piet A, Brandt VD. Time to pregnancy and sex of offspring: cohort study. *BMJ* 2005;331:1437-8.
 18. James WH. Male reproductive hazards and occupation. *Lancet* 1996;347:773.
 19. Jongbloet PH, Groenewoud JMM, Zielhuis GA. Further concepts on regulators of the sex ratio in human offspring. Non-optimal maturation of oocytes and the sex ratio. *Hum Reprod* 1996;11:2-9.
 20. Maconochie N and Roman E. Sex ratios: are there natural variations within the human population?. *Br J Obstet Gynaecol* 1997; 104:1050-3.
 21. Legner EF. Temperature, humidity and depth of habitat influencing host destruction and fecundity of muscoid fly parasites. *Entomophaga* 1977; 22:199-206.
 22. Moriya K, Hiroshige T. Sex ratio of offsprings of rats bred at 5 degrees C. *Int J Biometeorol* 1978; 22:312-5.
 23. van Zelst SJ, Zupp JL, Hayman DL, Setchell BP. X-Y chromosome dissociation in mice and rats exposed to increased testicular or environmental temperatures. *Reprod Fertil Dev* 1995; 7:1117-21.
 24. Lerchl A. Impact of environmental temperature on human scrotal temperatures. In: Hocevar A, Crepinsek Z, Kajfez-Bogataj L, editors: 215-221. 14th International Congress of Biometeorology, 1996.
 25. Tremblay pM, Vezina H, Houde eL. Demographic Determinants of the Sex Ratio at Birth in the Saguenay Population, Quebec, Institut National d'Etudes Démographiques. *Population* 2003; 58:383-94.
 26. Teitelbaum MS. Factors associated with the sex ratio in human populations. In: Harrison GA, Boyce AJ, editors: 90-109. *The Structure of Human Populations*, London, Oxford University Press, 1972.
 27. Erickson JD. The secondary sex ratio in the United States 1969-71: association with race, parental ages, birth order, paternal education and legitimacy. *Ann Hum Genet.* 1976; 40:205-12.
 28. Manning JT, Anderton RH, Shutt M. Parental age gap skews child sex ratio. *Nature* 1997; 389:344.
 29. Arnold F, Rutstein S. Sex ratio unaffected by parental age gap. *Nature* 1997; 390:242.
 30. Boklage CE. Sex ratio unaffected by parental age gap. *Nature* 1997; 390:243.
 31. James WH. Sex ratio unaffected by parental age gap. *Nature* 1997b; 390:242-3.
 32. James WH. Hormonal control of the sex ratio. *J Theor Biol* 1986; 118:427-41.

Yazışma Adresi / Address for Correspondence:

Dr. Erdal Tuğç
 Çukurova University Faculty of Medicine,
 Department of Medical Biology and Genetics
 01330 Balcalı-ADANA
 Tel. 90 322 3386060-3498
 Fax. 90 322 3386572
 e-mail: terdal@cu.edu.tr

geliş tarihi/received :29.07.2013

kabul tarihi/accepted:16.09.2013