



### Seroprevalence of Toxocariasis in Children in East-Azerbaijan Province, Iran

İran Doğu Azerbaycan Eyaletinde Çocuklarda Toksokariyazis Seroprevalansı

Garedaghi Yagoob<sup>1</sup>

<sup>1</sup>Department of Parasitology, Tabriz Branch, Islamic Azad University, Tabriz, IRAN

*Cukurova Medical Journal 2013; 38 (4): 581-586.*

#### ABSTRACT

**Purpose:** Toxocariasis is a zoonotic disease caused by the ascarid of dogs and cats, the main representative of which is a *Toxocara canis*. Distribution of the disease is world wide and is more prevalent in children. The present study was carried out in children of East Azerbaijan Province, Iran, to determine the toxocariasis seropositivity.

**Material and Methods:** For the present seroepidemiological study, blood samples were collected at random from children of all the five districts of the East Azerbaijan Province. A total of 336 children, 187 males and 149 females in age group of 0-15 years were selected for the present study. ELISA was used for detection of IgG antibodies against *Toxocara* excretory secretory antigen. A questionnaire interview was conducted to obtain the data concerning their age, sex and habits. The particular points in the questionnaire asked were recorded on the format right on the spot.

**Results:** Gender was found to be a significant risk factor for the *Toxocara* infection in children population. Male children were found more infected (41.71% as compared to females (24.16%). The total seroprevalence of *T. canis* antibodies in children of East Azerbaijan Province was 29.46 %. The risk factors that were found associated with the infection of toxocariasis in children population of East Azerbaijan Province include family back ground, status of living conditions, awareness, etc.

**Conclusion:** The present study reveals high prevalence of *T. canis* infection in children of East Azerbaijan Province. It is important to raise the awareness of health professionals, public and educators to the fact that toxocariasis is a public health problem. Health promotion by means of a school based educational approach, diagnosis and continuous programme of treatment are necessary.

**Key Words:** Seroprevalence, Toxocariasis, Children, East Azerbaijan Province, Iran.

#### ÖZET

**Amaç:** Toksokariyazis; temel temsilcisi *toxocara canis* olan, köpek ve kedi bağırsak solucanlarının neden olduğu, hayvanlardan insanlara bulaşan zoonotik bir hastalıktır. Bu hastalık dünyanın her tarafında görülebilen ve en çok da çocukları etkileyen bir hastalıktır. Bu çalışma İran Doğu Azerbaycan Eyaletindeki çocukların toksokariyazis seropozitifliğinin belirlenmesi amaçlandı.

**Materyal ve Metod:** Çalışma için Doğu Azerbaycan Eyaletinin 5 farklı bölgesinden rastgele seçilmiş çocuklardan kan örnekleri toplandı. Bu çalışmada 0 ile 15 yaşları arasında 187'si erkek, 149'u kız olan 336 çocuk seçilmiştir. *Toxocara* boşaltım salgısı antijenine karşı IgG antikorlarını belirlemek için ELİSA yöntemi kullanılmıştır. Çocukların yaşlarını, cinsiyetlerini ve yaşadıkları ortamları öğrenmek amacıyla anket düzenlendi. Anketteki önemli sorular yerinde kaydedildi.

**Bulgular:** Cinsiyetin çocuk popülasyonlarında toksokara enfeksiyonu için önemli bir risk faktörü olduğu bulundu. Erkek çocukların (%41,7) kız çocuklara (%24,16) nazaran daha çok etkilendiği gözlemlendi. Doğu Azerbaycan Eyaletinde çocuklarda gözlenen *T.canis* antikorun seroprevalansı %29,46 idi. Doğu Azerbaycan Eyaletinde ki çocuk popülasyonlarında gözlenen toksokara enfeksiyonun aile öyküsü, yaşam statüsü ve bilinçli olma gibi parametrelerle ilişkili olduğu tespit edildi.

**Sonuç:** Bu çalışmada Doğu Azerbaycan Eyaleti çocuklarında *T. Canis* enfeksiyonunun yüksek prevalansa sahip olduğu belirlenmiştir. Toksokariyazis bir halk sağlığı problemidir ve bu nedenle sağlık uzmanlarının ve eğitimcilerinin bu konuda bilinçlendirilmesi oldukça önemlidir. Ayrıca bu konudaki okul temelli eğitimsel yaklaşımların, teşhis ve devam eden tedavi programlarının da geliştirilmesi gereklidir.

**Anahtar Kelimeler:** Seroprevalans, toksokariyazis, Çocuklar, Doğu Azerbaycan Eyaleti, İran

## INTRODUCTION

Toxocariasis is a zoonotic disease caused by the ascarid of dogs and cats and the main representative of which is a *Toxocara canis*<sup>1</sup>. The eggs of *T. canis* are unembryonated when passed in the faeces of dogs into the environment. Under optimal temperature and humidity the eggs develop into embryonated eggs that are infective to both final and paratenic hosts. Infective eggs are reported to survive optimal circumstances for at least one year. Human may acquire the infection by oral ingestion of infective *Toxocara* eggs from contaminated soil (sapro-zoonosis) from unwashed hands or consumption of raw vegetables. The disease manifests itself as two distinct forms, visceral larva migrans (VLM) and ocular larva migrans (OLM). The signs and symptoms of VLM vary from an asymptomatic state with mild eosinophilia to a severe and potentially fatal disorder including hepatomegaly, hyperglobulinemia, pneumonitis and neurological disorders<sup>2,31</sup>. The disease has a chronic state and the symptoms can even persist for more than a year. Patients with OLM also show variable clinical signs varying from asymptomatic state to acute lesions including endophthalmitis accompanying loss of vision and mass similar to retinoblastoma [3-5]. Distribution of the disease is world wide and is more prevalent in children. There is no definitive method in diagnosing *Toxocara* infection. As the larvae of *T. canis* are arrested in the paratenic host-larvae during migration and they do not mature into adults, hence a stool examination of the patient will not give any clue about the infection. However numerous studies have shown that immunoassay for detection of antibodies using a purified excretory-secretory antigen from the

larval stage significantly improves sensitivity and specificity compare to assays using crude antigens<sup>6,7</sup>. The most widely used test, because of its high sensitivity and specificity is the ELISA in which antibodies to *T. canis* larval excretory secretory antigens<sup>1,8</sup> or to larval extracts are measured<sup>9,10</sup>. In Iran reports of human toxocariasis have been presented<sup>3,11</sup>, but there are limited studies from East Azerbaijan Province<sup>12,30</sup>. Therefore the present study was conducted to determine the seroprevalence of toxocariasis in children of East Azerbaijan Province.

## MATERIALS and METHODS

For the present seroepidemiological study, samples were randomly selected from all the districts. A total of 336 children, out of which 187 were males and 149 were females in age group of 0-15 years were randomly selected from different districts of the East Azerbaijan Province. A short questionnaire to obtain data concerning their age, sex, habits was filled for each child. Blood samples were collected using disposable syringes and sera were separated and stored in -20 °C until tested. Antibody (IgG) specific to *Toxocara* purified excretory secretory (ES) antigen was detected by ELISA in all serum samples using kit obtained from IVD research Inc. The test was performed as per manufacturer's instructions. Optical density (OD) value was recorded in an automatic ELISA reader at 450 nm. The samples were considered positive if absorbance reading was equal to or greater than 0.3 OD units and negative if absorbance reading was less than 0.3 OD units. Fisher's exact test was used for statistical analysis.

## RESULTS

In the present study out of 336 children, 99 (29.46%) were found *Toxocara* seropositive. The risk factors that were found associated with the infection of toxocariasis in children population of East Azerbaijan Province included family background, status of living conditions, and awareness about the disease. Gender was found a significant risk factor ( $P=0.00$ ). Male children were found more infected (41.71%) as compared to females (24.16%). Paternal education remained an important risk factor for *Toxocara* infection ( $P<0.05$ ). The percentage of infection in children whose fathers were not educated was 42.03% (66/157) compared to 26.81% (48/179) children whose fathers were educated. Maternal education had a great impact on the *Toxocara* infection in children and remained a significant risk factor for the *Toxocara* infection in the children ( $P=0.002$ ). Children whose mothers were educated had significantly lower prevalence (25.19%) than those whose mother's were not educated (39.23%). The prevalence of *Toxocara* infection in families with fenced houses was less 29.84% compared to families with unfenced houses 39.31%. The prevalence of *Toxocara* infection was higher in those having contact with dogs 45.07% or with pets in their house 60.71% (Table). A related factor for the prevalence of *Toxocara* infection in humans was the condition and status of vegetables taken by the subjects. The children who were in the habit of eating raw vegetables were more prone to infection (36.84%) than those who were not in the habit of eating raw vegetables (25.84%). Children who were in the habit of geophagia were also more prone to infection (36.84%). The majority of the study populations were using tap water, but some also used the well water, river water and other sources of water which included water from all these or some sources at a time. Children using the water from rivers and streams were more infected (44.44%) than those using well water (38.70%), followed by those using water from tap (public piped water supply) (27.17%). Water

source was found significant risk factor for the prevalence of *Toxocara* infection ( $P<0.05$ ). Majority of children were using unboiled water. Boiled water was also used by some children in the study population. Prevalence of infection was higher (40.11%) in the children drinking unboiled water than those drinking boiled water (28.40%). Water pretreatment was a significant risk factor for the prevalence of toxocariasis infection.

## DISCUSSION

Serological studies are of immense importance in the detection of infection by *T. canis*, as the clinical symptoms of toxocariasis are variable and non specific. The use of *Toxocara* ES antigen to detect antibodies against *T. canis* does not require the preabsorption of sera with embryonated *Ascaris* egg antigen<sup>9,1,29</sup> and further no cross reaction between purified ES antigen and sera from individuals with *Ascaris lumbricoids*, Hook worms, *E. coli* or *Giardia lamblia* were observed<sup>13,23,28</sup>. The present study reports serological prevalence of human toxocariasis in children of East Azerbaijan Province. In different parts of the world, serological studies have demonstrated a variation in *Toxocara* seroprevalence ranging from 2.3% to 86%<sup>11,14</sup>. However the present study showed a higher rate of infection (29.46 %) than that of (6.4 %) subjects residing in a rural area near Chandigarh<sup>11</sup>, Slovak Republic<sup>15</sup> which may be due to low standards of hygiene, frequent contact with the contaminated soil and less paternal education<sup>23,26,27</sup>. In our study, higher prevalence of infection was found among males than females. The difference among male and female was found significant; a similar result was reported earlier<sup>15</sup>. In previous epidemiological studies association of several risk factors for toxocariasis has been reported such as exposure to dogs[24], socio economic status<sup>1</sup>. In our study the children whose parents were illiterate were more positive indicating the effect of economic situation on seropositivity.

**Table 1: Epidemiological analysis for *Toxocara* seroprevalence in the children of East Azerbaijan Province, Iran.**

Particulars/Age P-value	No. of samples analyzed	Positive (%)	Negative (%)
Upto 15 yr	336	99 (29.46)	237 (70.53)
<b>Sex</b>			
Male	187	78 (41.71)	109 (58.28)
Female	149	36 (24.16)	113 (75.83)
0.000			
<b>Father's education</b>			
Yes	179	48 (26.81)	131(73.18)
No	157	66(42.03)	91(57.96)
0.001			
<b>Mother's education</b>			
Yes	127	32(25.19)	95(74.80)
No	209	82(39.23)	127(60.76)
0.002			
<b>House fenced</b>			
Yes	191	57 (29.84)	134 (70.15)
No	145	57 (39.31)	88 (60.76)
0.005			
<b>Pet in house</b>			
Yes	28	17(60.71)	11(39.28)
No	308	102(33.11)	206(66.88)
0.42			
<b>Contact with dogs (neighbourhood/semi domesticated)</b>			
Yes	71	32(45.07)	39(54.92)
No	265	82(30.94)	183(69.05)
0.01			
<b>Eating raw vegetables/geophagia</b>			
Yes	247	91(36.84)	156(63.15)
No	89	23(25.84)	66(74.15)
0.015			
<b>Source of drinking water</b>			
Stream/river/Ponds	90	40(44.44)	50(55.55)
Well	62	24(38.70)	38(61.29)
Public piped water supply	184	50(27.17)	134(72.82)
0.01			
<b>Condition of drinking water</b>			
Boiled	169	48(28.40)	121(71.59)
Unboiled	167	67(40.11)	100(59.88)
0.01			

Contact with dogs or presence of pet in house was the high risk for *Toxocara* prevalence and was found significant factor for toxocariasis ( $P<0.05$ ). Also significant association was found between presence of dogs in houses or in neighborhood and prevalence of toxocariasis in humans. A higher frequency of infection in individuals who maintained contact with dogs has been reported by many workers in different parts of the world<sup>16,19</sup>.

During this study, the persons having the habit of eating raw vegetables or having a habit of geophagia were found having high seroprevalence of *Toxocara* infection. Holland *et al.* found a significant association between *Toxocara* seropositivity in children and a history of geophagia<sup>20,25</sup>. In the current study the prevalence of *Toxocara* infection was more in people using water from streams, rivers, ponds and wells than

those using water from public piped water supply and the difference was found statistically significant. Similar results were found by Hayashi *et al.*<sup>21,24</sup>. In our study it was observed that individuals whose houses were fenced were less likely to be infected with *Toxocara* relative to those whose houses were not fenced. Similar results were reported by Abe and Yasukawa<sup>22</sup>. Thus reveals the high percentage of *T. canis* infection in toxocariasis in humans.

### CONCLUSION

In conclusion, the high prevalence of *Toxocara* in East Azerbaijan Province could be due to high prevalence of infection in large untreated and unconstrained dog population, low standards of hygiene and geophagic behavior among children[23]. Health promotion by means of a school based programme of treatment improving standards of hygiene and control of infection in dogs are necessary for control and prevention of the disease.

### Acknowledgement

The researchers wish to thanks the Islamic Azad University, Tabriz Branch, Tabriz, Iran for the financial supports and all laboratory technicians for technical aids in this project.

### REFERENCES

- Glickman LT, Schantz PM. Epidemiology and pathogenesis of zoonotic toxocariasis. *Epi Rev.* 1981; 3:230-50.
- Gillespie SH. The clinical spectrum of human toxocariasis. In: Lewis JW and Maizel RM, Editoris. *Toxocara and toxocariasis, clinical, epidemiological and molecular perspectives: British society of parasitology*, London. 1993; 55-61.
- Mirdha BR, Khokar SK. Ocular toxocariasis in a north Indian population. *J of Trop Ped.* 2002; 48:328-330.
- Shimizu Y, Imai M, Fukasawa A, Iijima H. Premacular membrane peeling without removal of sub retinal granuloma in an eye with ocular toxocariasis. *Acta Ophthalmologica Scandinavica.* 2005;14:395-96.
- Fomda BA, Ahmad Z, Khan NN, Tanveer S, Wani SA. Ocular toxocariasis in a child: A case report from Kashmir, North India. *Indian Journal of Medical Microbiology.* 2007; 25:411-2.
- Fenoy S1 Cuellar C, Guillen JL. Seroprevalence of toxocariasis in children and adults in Madrid and Tenerife. Spain *J Helm.* 1996; 70:109-13.
- Kenny JV, Maccabe RJ, Smith HV, Holland C. Serological evidence for the presence of toxocariasis in Turkara district of Kenya. *Trans Roy Soc Trop Med Hyg.* 1995; 89:377-78.
- De Savigny DH, Voller A, Woodreff AW. Toxocariasis: Serological diagnosis by enzyme immunoassay. *J S Pathol.* 1979; 32:284-88.
- Fan CK, SUKE. Cross Reactions with *Ascaris suum* antigens of sera from mice infected with *A. suum*, *Toxocara canis* and *Angiostrongylus contonensis*. *Parasitology International.* 2004; 53:263-271.
- Glickman LT, Grieve RB, Lauria SS, Jones DL. Serodiagnosis of Ocular toxocariasis; a comparison of two antigens. *J Clin Pathol.* 1985; 38:103-07.
- Malla N, Aggarwal AK, Mahajan RC. A serological study of human toxocariasis in north India. *The National Medical Journal of India.* 2002; 15:145-47.
- Ahmad B, Bhatti G, Thokar MA, Malla N. Human toxocariasis and Ascariasis: concomitant parasitism in Srinagar, Kashmir, India. *Indian J Pathol Microbiol.* 2002; 45:315-318.
- Hakim SL, Mak JW, Lam PLM, Nazma S and Normaznah H. Seroprevalence of toxocariasis canis antibodies among Orang Asli (Aborigines) in Peninsular Malaysia. *SE Asian J Trop Med Publ Hlth.* 1992; 23:493-496.
- Thompson DS, Budy DAP, Cooper ES, Schantz PM. Epidemiological characteristics of *Toxocara canis* zoonotic infection of children in Caribbean community. *Bull WHO.* 1986; 64:283-90.
- Havasiova K, Dubinsky P, Stefancikova A. A seroepidemiological study of human *Toxocara* infection in the Slovak Republic. *Journal of Helminthology.* 2006; 67:291-6.
- Fan CK, Hung CC, DU Wen-Yuan, Liau Chen-Wei, Su-Kua-Euyre. Seroepidemiology of *Toxocara canis* infection among mountain aboriginal school children

- living in contaminated districts in Eastern Taiwan. *Tropical Medicine and International Health*. 2004; 9:1312-8.
17. Fan CK, Liano CW, Kao TC, Li MH, Du WY, Su KE. Seroepidemiology of *Toxocara canis* infection among aboriginal school children in the mountainous areas of north-eastern Taiwan. *Ann Trop Med Parasitol*; 2005; 99: 593-600.
  18. Sadjjadi SM, Khosravi M, Mehraabani D, Oryan A. Seroprevalence of *Toxocara* infection in school children of Shiraz, Southern Iran. *J of Trop Ped*. 2000; 46:327-30.
  19. Figueiredo SDP, Taddei JAAC, Menezes JJC, Novo NF, Silva EOM, Crislova HLG, Cury MCFS. Clinical-epidemiological study of *Toxocariasis* in a pediatric population. *J Pediatrics*. 2009; 81: 126-32.
  20. Holland CV, O'Loracin P, Taylor MRH, Kelly A. Sero-epidemiology of *Toxocariasis* in school children. *Parasitology*. 1995; 110: 535-45.
  21. Hayashi E, Tuda J, Imada M, Akao N, Fujita K. The high prevalence of asymptomatic *Toxocara* infection among school children in Manado, Indonesia. *South East Asian J Trop Med Public Health*. 2005; 36:1399-1406.
  22. Abe N, Yasukawa A. Prevalence of *Toxocara* spp. eggs in sand pits in Osaka city, Japan, with notes on the prevention of egg contamination by fence construction. *Vet Med Sci*. 1996; 59:79--80.
  23. Dar ZA, Tanveer S, yatoo GN, sofi BA, Wani SA, Dar PA, fomda BA. Seroprevalence of *Toxocariasis* in children in Kashmir, J&K state, India. *Iranian J parasitol*. 2008; 3:45-50.
  24. Garedaghi Y, safar mashaei S. Prevalence of gastrointestinal helminthic infestation in pet and stray dogs in Tabriz, Iran. *Journal of animal and veterinary advances*. 2011; 10: 1477-9.
  25. Ishiyama, S., K. Ono, S.K. Rai and S. Uga. Method for detecting circulating *Toxocara canis* antigen and its application in human serum samples. *Nepal Med. Coll. J*. 2009; 11: 9-13. PMID: 19769229.
  26. Neelam, T. and S. Meenu. Enhancement of plant growth and decontamination of nickel-spiked soil using PGPR. *Journal of Basic Microbiology*. 2009; 49: 195-204.
  27. Shane, H.L., J.R. Verani, B. Abudho, S.P. Montgomery and A.J. Blackstock et al. Evaluation of urine CCA assays for detection of *Schistosoma mansoni* infection in Western Kenya. *Plos Negl. Trop. Dis*. 2011; e951-e951. DOI: 10.1371/journal.pntd.0000951.
  28. Shoiab, M. and M. Irshadulla. Protein profile of male and female *Haemonchus contortus*. *J. Vet. Parasitol*., 2009; 23: 147-9.
  29. Zibaei, M., S.M. Sadjjadi, S. Ishiyama, B. Sarkari and S. Uga. Production of monoclonal antibody against *Toxocara cati* second-stage larvae and its application for the detection of circulating antigens. *Hybridoma*. 2010; 29: 217-20. DOI: 10.1089/hyb.2009.0108.
  30. Walker, J.M. Gradient SDS Polyacrylamide Gel Electrophoresis of Proteins. In: *The Protein Protocols Handbook*, Walker, J.M. (Ed.), Springer, Totowa, NJ., ISBN-10: 0896039404. 2012; 69-72.
  31. Woodward T.M. Update on CAA (aka-Acantho-Pathology, 40: 93–106. tous Epulis). *Animal Dental Care*. Web: [www.wellpets.com](http://www.wellpets.com). Retrieved on 03.04.2012. 2011; 40:93-106.

#### Yazışma Adresi / Address for Correspondence:

Dr. Garedaghi Yagoob  
 Islamic Azad University  
 Department of Parasitology, Tabriz Branch  
 Tabriz, Iran.  
 e-mail: Y\_garedaghi@yahoo.com

geliş tarihi/received :18.02.2013

kabul tarihi/accepted:20.03.2013