

ELEMENTAL ANALYSIS OF BEYBAĞ-MUĞLA (TURKEY) BYZANTINE SKELETONS

BEYBAĞ-MUĞLA (TÜRKİYE) BİZANS DÖNEMİ İSKELETLERİNİN ELEMENT ANALİZİ

Seda KARAÖZ ARIHAN* - Ali Akın AKYOL**

İsmail ÖZER*** - Okan ARIHAN****

Keywords: Trace Element, Bone Analysis, Paleopathology, Nutrition, Byzantine Period

Anahtar Kelimeler: İz Element, Kemik Analizi, Paleopatoloji, Beslenme, Bizans Dönemi

Makale Bilgisi

Başvuru: 21 Eylül 2015

Hakem Değerlendirmesi: 9 Aralık 2015

Kabul: 8 Eylül 2017

DOI Numarası: 10.22520/tubaar.2017.21.008

Article Info

Received: September 21, 2015

Peer Review: December 9, 2015

Accepted: September 8, 2017

DOI Number: 10.22520/tubaar.2017.21.008

ABSTRACT

Skeletal material forms the basis of paleoanthropological studies and they are also valuable sources of information for the assessment of elemental analysis which may provide clues about nutritional status of ancient populations. Study material comprises skeletons from the necropolis of Beybağ-Muğla (Turkey) region belonging to Byzantine period and excavated during 2008 period. 52 femoral samples (19 men, 18 women, 15 children) and 3 soil samples were studied with the aid of XRF spectroscopy. Macro elements such as calcium, iron and phosphorus as well as trace elements such as arsenic, barium, lead, strontium and zinc were evaluated in order to figure out nutritional status of this Byzantine population with the help of skeletal elemental analysis. In addition it was also aimed to assess different parts of femoral bone within the study in order to determine the effect of sampling area in this population. Results were evaluated with Kruskal-Wallis and Tukey tests. Strontium values for groups were found significantly higher than soil samples. Similarly zinc and phosphorus values for groups were also found significantly higher than soil. However barium, iron and iodine values were found similar to soil samples and affected by diagenetic rather than a biogenetic source. Phosphorus content of bone material was high and Ca/P rate was found lower than observed values in literature which is possibly due to diagenesis. Log (Ba/Sr) rate was

* Yüzüncü Yıl University, Faculty of Letters, Department of Anthropology, Van, Turkey E-mail: sedaarihan@yyu.edu.tr

** Gazi University, Faculty of Fine Arts, Department of Conservation & Restoration of Cultural Properties, Ankara, Turkey E-mail: aliakinakyol@gmail.com

*** Ankara University, Faculty of Language, History and Geography, Department of Anthropology, Ankara, Turkey E-mail: iozer@ankara.edu.tr

**** Yüzüncü Yıl University, Faculty of Medicine, Department of Physiology, Van, Turkey Corresponding Author E-mail: sedaarihan@yahoo.com E-mail: okanarihan@yyu.edu.tr

altered because of possible diagenesis for Ba. Since those two ratios are used as indicators of nutrition of the individual, presence of potent diagenesis prevents conclusion on nutrition and diet choices such as marine/ terrestrial for Beybağ-Muğla population. Since aim of this study was to assess bone material of this high anemia and other paleopathological disease bearing population in elemental analysis a malnutritional situation was expected. However observation of high zinc and phosphorus compared to literature suggests no lack of protein in diet. In addition low lead and low arsenic values suggest no adverse impact of those toxic elements on physiological state of Beybağ people. According to these findings diet of Beybağ population can be considered within omnivorous limits and without apparent insufficiency of nutrition. However presence of anemia in all of the sample material suggests infections diseases and low hygiene conditions in this population and needs further assessment on burial locations and burial types with less influence of diagenetic processes in order to figure out nutritional status of this population.

ÖZET

İskelet materyali paleoantropolojik çalışmaların temel materyalini oluşturur ve element analizinden elde edilen bilgilerle antik toplumların beslenme durumu hakkında ipuçları sağlar. Çalışma materyali Beybağ-Muğla (Türkiye) bölgesi nekropolünden 2008 yılındaki kazıda elde edilmiş Bizans dönemine ait iskeletleri içermektedir. 52 femur örneği (19 erkek, 18 kadın, 15 çocuk) ve 3 toprak örneği XRF spektroskopisi aracılığıyla araştırıldı. Bizans toplumunun beslenme durumunun iskelet element analizi yardımıyla ortaya konulabilmesi için kalsiyum, demir ve fosfor gibi makro elementlerin yanı sıra arsenik, baryum, kurşun, stronsiyum ve çinko da söz konusu çalışmada incelendi. Buna ek olarak femurun farklı kısımlarının örneklenmesinin bu topluma ait eser element analizindeki etkisini araştırmak da hedeflendi. Sonuçlar Kruskal-Wallis ve Tukey testleri ile değerlendirildi. Gruplar için stronsiyum değerleri toprak örneklerine göre anlamlı derecede yüksek bulundu. Benzer şekilde grupların çinko ve fosfor değerleri de toprağa göre anlamlı derecede yüksek tespit edildi. Ancak baryum, demir ve iyot değerleri toprak örnekleri ile benzer bulundu ve biyogenetik bir kaynaktan çok diyagenetik olaylardan etkilenmiş olarak tespit edildi. Kemik materyalinin fosfor içeriği yüksekti ve Ca/P oranı muhtemelen diyagenezle bağlı olarak literatürden yüksek bulundu. Log (Ba/Sr) oranı muhtemelen baryumun diyagenezden etkilenmesine bağlı olarak farklı bulundu. Bu iki oran bireyin beslenmesinin belirteçleri olarak kullanıldığı için muhtemel diyagenezin güçlü etkisi Beybağ-Muğla toplumunun beslenmesi ve denizel/karasal gibi diyet tercihleri hakkında yargıya varmayı engellemektedir. Bu çalışmanın amacı yüksek oranda anemi ve diğer paleopatolojik hastalıkları taşıyan bu toplumun kemik materyalinin element analiziyle değerlendirilmesi olduğu için bir malnütrisyon durumu beklenmekteydi. Ancak çinko ve fosforun literatüre göre yüksek düzeyde gözlenmesi diyetle protein açısından bir eksikliğin olmadığını düşündürmektedir. Buna ek olarak düşük kurşun ve düşük arsenik değerleri bu toksik elementlerin Beybağ toplumunun fizyolojik durumuna olumsuz bir etkisi olmadığını düşündürmektedir. Bu bulgulara göre Beybağ toplumunun diyetinin omnivor limitler içinde olduğu ve belirgin bir beslenme bozukluğunun gözlenmediği yargısına varılabilir. Ancak incelenen materyallerin tümünde aneminin varlığı bu toplumda enfeksiyon hastalıklarını ve düşük hijyen koşullarını düşündürmektedir. Bu toplumun beslenme durumunun ortaya konulabilmesi için diyagenetik süreçlerden daha az etkilenen gömü alanlarının ve gömü tiplerinin değerlendirilmesine ihtiyaç duyulmaktadır.

1. INTRODUCTION

Human body is composed of elements dominated by carbon, hydrogen and oxygen but also encompasses more than 80 elements classified in the periodic table. Many of them can be incorporated into the bone tissue¹. Amount consumed in foods defines the term macro and microelements in diet. Macro elements which are consumed in large quantities are carbon, hydrogen, nitrogen, oxygen, calcium, potassium, phosphorus (P) and sodium (Na). Whereas strontium (Sr), barium (Ba), selenium (Se), lead (Pb) are classified as micro or trace elements. These are found in very low quantities in bone tissue. This quantity may be as low as parts per million (ppm) or billion (ppb)². However this minute contribution into bone tissue does not lessen the physiological importance of trace elements. In various parts of the world alterations in the trace elements in soil is correlated with certain diseases such as Kashin-Beck disease (osteoarthritis deformans endemica) found in Northern China, Northern Korea and Siberia. This disease is found to be related by low selenium content of soil³.

Use of physicochemical analysis techniques in anthropological studies brought a new area to obtain more data from the skeletal materials in this field. Presence of many of the diseases which existed during historical periods was found with the help of science of paleopathology which deals with the remnant skeleton material of humans. Metabolic diseases which form an important portion of those diseases have direct relation with nutrition. Stable isotopes of some elements and trace elements in the periodic table are among the mostly investigated substances. Both techniques base on the analysis of bone and teeth which are the most durable tissues in the body that can withstand the disintegration occurred during postmortem period. Amount of those elements in those tissues and their ratios to one another gives valuable information in the field of nutrition and related paleopathological conditions⁴. Since elemental compositions of soils from different parts of the world are vastly dissimilar, this situation avoids comparison between anthropological materials from different studies. Nevertheless comparisons of elemental content of skeleton materials from the same burial site give valuable information⁵. Another important detail in this subject is the dissimilarity of data obtained from teeth and skeletal material. Teeth are almost completely formed at the early stages of life. On the contrary bone tissue is a

dynamic structure in which osteoclastic and osteoblastic cells continue to break down and build up bone tissue throughout lifetime. This disparity in the exposure of teeth and bone tissues to different environmental and nutritional impacts give way to assess different conditions faced in distinct phases of lifetime⁶. Elements widely used for clarifying nutritional status and nutrition of past humans are Ba, Sr, Iron (Fe), Copper (Cu), Cadmium (Cd), Silicon (Si), Pb, Se and Iodine (I)⁷. Accumulation of those elements in the bone tissue may be affected from various factors. Food choice of humans may include more animal or plant based food or a balanced diet. This variation is reflected in the bone elemental composition. Calcium (Ca), Ba and Sr which are all members of 2A group of the periodical table are found in soil, plant and animals within the food web. Close physicochemical properties of those elements cause similar accumulation in bone tissue.

Animal body can incorporate all three elements however they favour Ca when it is available. Since absorption of Ca and excretion of Sr is high in the kidneys, Ca accumulation in the organism is much higher than Sr in animals compared to plants and in carnivores compared to herbivores⁸. Sr and Ba are high in plant food resources and so in bones of individuals which have higher plant sources in the daily nutrition. Similarly silicon can be found in higher amounts when plant based food dominates the diet⁹.

Pb is accepted as a toxic element for organisms and its physiological significance is not known yet. Although its toxicity and related symptoms are most visible at central nervous system its accumulation occurs mostly at bone tissue. Copper is another element which can cause toxicity however its adequate amounts are necessary for enzymes of antioxidant system in the body. It also accumulates in the bones. Those elements can occur in food as well as they can be found as a result of contamination from the storage cups to the food¹⁰. Similarly water pipes can contaminate the water inside them. Apart from the accumulation of such elements in the skeleton due to the food or water consumed by the people, diagenesis can be a source of augmented copper or zinc (Zn) content¹¹.

Although Pb and mercury (Hg) can cause very distinct intoxications when they are found in very low quantities, all elements may cause toxicity above certain levels. Fe is an important element for oxygen transport via hemoglobin as well as for oxidant/antioxidant enzyme systems. Its decrease in the body leads to anemia whereas its over accumulation

¹ Smrcka 2005, Bowen 1979, Emsley 1998

² Smrcka 2005.

³ Sokoloff 1988.

⁴ Trimble/Macko 1997; Lee-Thorp/Sponheimer/Vander Merwe 2003.

⁵ Katzenberg/Saunders 2008.

⁶ Grupe 1995.

⁷ Smrcka 2005.

⁸ Smrcka 2005.

⁹ Aksoy 2007.

¹⁰ Smrcka 2005.

¹¹ Zapata/Perez-Sirvent/Martinez-Sanchez/Tovar 2006.

in the body may lead to pathological conditions¹². Under normal circumstances over accumulation of a trace element in the body is rare but it may occur due to geological conditions of the inhabited region, food preparation techniques and used materials for food and water.

Various techniques are applied for the analysis of elements from bone or earth material. Gas chromatography, atomic absorption spectroscopy, atomic emission spectroscopy XRF (X-Ray fluorescence), ICP-MS are examples for such techniques¹³. Such techniques depend on the fact that properties of atoms in absorbing and emitting energy are different. By their different properties elements can be discriminated from each other. The principles of X-Ray Fluorescence (XRF) analysis is related to optical dispersion spectrograph principles. However in the X-ray fluorescence method excitation of atom happens through X-rays (instead of heat). As the atom numbers increase in the elements, the amount of electrons they have and thus their number of orbitals on which the electrons move increase. If any atom is excited with a high energy radiation like X-Ray, this high energy input brings the electrons in the near orbitals to a higher energy level; when the excited electrons go back to the initial energy level, they discharge the extra energy they had gained in the form of X-Rays. This secondary X-Ray emission is called "fluorescence radiation". The wavelength of these radiations that an element emits is characteristic for that element. In other words, these radiations are like fingerprints of the related elements. By identifying the wave-length of the radiation, the type of the element and by measuring the density of this identified ray, the concentration of the element in that substance can be determined¹⁴.

Porous structure which gives bone tissue its strength and flexibility also causes its close interference with blood elemental values such as Ca and P¹⁵. This porous structure is the key for exchange of elements between bone and blood during lifetime and between bone and soil after death. Following the death of individual this close interrelation of bone with blood ends and another interaction starts with soil and burial environment. Some elements in the bone are lost whereas some others may start to accumulate to the bone from the soil such as manganese, Ca and Fe¹⁶. Ba is more affected from diagenesis although it is a better indicator than Sr for plant based food¹⁷. Therefore Ba

content of soil should be assessed prior to the use of Ba as a plant based food in order to discriminate whether this Ba is originated from soil diagenesis or the food consumed by the individual. More abundance of an element in soil rather than bone material suggests a diagenetic process. Since Sr and Zn are less affected from diagenesis unlike the Ba it is much safer to use them in nutrition assessment studies¹⁸. But still soil elemental analysis is essential for confirmation.

In addition to soil analysis, Ca/P ratio in bone can be used as an indicator of diagenesis. Ca/P rate is found at a constant rate in humans. This ratio is 2.16¹⁹. An attenuation of this value or its increase can be considered as a diagenetic process to the skeletal material. Diagenesis is known to effect bone and dental tissue differently. Teeth are more resistant to diagenesis²⁰. Also skeletal tissue is not uniformly affected by diagenesis. Cortical bone is less affected than trabecular bone and surface is more affected than deeper parts of bone²¹. In order to lessen the effect of diagenesis cortical bone surface obtained from areas such as femoral neck is scraped with a suitable material and less affected parts are used for analysis²². There are reports stating the impact of sampling area on the bone on the results. Femoral head (spongiosa part of the femoral head) is used but also compact bone extracted from the proximal part (femoral neck) is also assessed²³ since it is known that compact bone is less influenced from diagenesis²⁴. Even spatial accumulation of elements is also observed in same site of bone²⁵. Therefore additional samplings may aid evaluation of accumulation of elements. Generally bone which have more surface/volume ratios such as ribs are more affected from diagenesis. Due to same physical surface/volume ratio adult bones are less affected from diagenesis compared to children bones²⁶.

Another ratio frequently used is the Sr and Ca ratio. High Sr/Ca ratio is an indicator of plant based diet and vice versa²⁷. Higher levels in food pyramid causes less Sr/more Ca²⁸ since animal tissues prefer Ca in favor and Sr levels attenuate compared to lower levels of this pyramid²⁹. Humans place somewhere in between carnivorous and herbivorous animals because of their omnivorous

¹² Allen/Gurrin/Constantine/Usborne/Delatycki/Nicoll/McLaren/Bahlo/Nisselle/Vulpe/Anderson/Southey/Giles/English/Hopper/Olynyk/Powell/Gertig 2008.

¹³ Katzenberg/Saunders 2008; Smrcka 2005; Johansson/Campbell 1988.

¹⁴ Shackley 2011.

¹⁵ Katzenberg/Saunders 2008.

¹⁶ Sillen/Sealy/vander Merwe 1989

¹⁷ Farnum/Sandford 1997.

¹⁸ Ezzo 1994.

¹⁹ Posner 1969.

²⁰ Kyle 1986.

²¹ Lambert/Weydert-Homeyer 1993.

²² Lambert/Weydert-Homeyer 1993.

²³ Smrcka 2005.

²⁴ Katzenberg/Saunders 2008.

²⁵ Pemmer 2013.

²⁶ Mays 1998.

²⁷ Sillen 1992.

²⁸ Smrcka 2005.

²⁹ Katzenberg/Saunders 2008.



Figure 1: Muğla - Beybağ Region / *Muğla Beybağ Bölgesi*

property³⁰. Also augmented Log Ba/Sr rate in the skeletal material may indicate an augmented marine food in the diet of this human³¹.

Turkey with its long human settlement history starting from prehistoric times provides a vast study area for elemental analysis from burial sites. There are increasing number of studies on the subject presenting different nutritional status of individuals and different burial conditions in the scope of diagenetic processes which alters elemental content of skeletons. Among them Beybağ-Muğla Byzantine population has been studied by different researchers in archaeological and anthropological view. In this population various paleopathological conditions has been found (PhD thesis by S.K. Arıhan). This reminds a possible malnutrition or intoxication case. This study aims to assess nutritional status of this population with the aid of certain chemical elements found in bone material in order to figure out any possible interrelation between those pathologies and nutritional state.

2. MATERIALS AND METHODS

2.1. Skeletal Material

Research material consists of samples from an excavation study conducted at Beybağ Region (Muğla-Turkey-Byzantine Period) by Prof Dr. Ahmet Tırpan from Konya Selçuk University Department of Archaeology in 2008

³⁰ Lambert/Vlasak-Simpson/Szpunar/Buikstra 1984.

³¹ Smrcka 2005; Burton/Price 2002.

(Fig. 1). This site had been inhabited between 10th century AD and 13th century AD³² (Fig. 2). Skeletons belonging to 52 individuals were studied in this research. Chronologic determination of skeletons was performed according to archaeological findings. Burials were in different types but all were found with close contact with soil material.

Different settlement periods are shown in different colors as: Bronze age, Byzantine period and Ottoman period

2.2. Age And Sex Determination

Determination of age and sex are important subjects in anthropological studies. Following cleaning and repair on skeletal material age and sex determination were performed. Methods accepted for this purpose were applied according to literature records. Morphological features on pelvis and skull of adult individuals were chosen for sex determination. For sex determination Brothwell³³, Buikstra and Ubelaker³⁴, Krogman and İşcan³⁵, Olivier³⁶, Steele and Bramblett³⁷ and criteria determined at Workshop of European Anthropologists (1980) were used.

³² Tırpan/Sögüt/Büyüközer 2009.

³³ Brothwell 1981.

³⁴ Buikstra/Ubelaker 1994.

³⁵ Krogman/İşcan 1986.

³⁶ Olivier 1969.

³⁷ Steele/Bramblett 1988.




Figure 2: Beybağ Region Excavation Site (Beybağ Excavation Archive) Different Settlement Periods are Shown in Different Colors as: Bronze Age, Byzantine Period and Ottoman period / *Beybağ Bölgesi Kazı Alanı (Beybağ Kazı Arşivi). Farklı Yerleşim Dönemleri Farklı Renklerle Gösterilmiştir: Bronz Çağı, Bizans Dönemi ve Osmanlı dönemi*

For age determination of adults symphysis pubis found on pelvis³⁸ and auricular surface morphology³⁹, degree of sutural closure on skull⁴⁰, cross section of clavícula body⁴¹, teeth erosion⁴², changes of costae on sternal end⁴³ proximal cross section of humerus and femur⁴⁴ were preferred.

For age determination of young adults methods such as epiphyseal aging and symphyseal aging⁴⁵, dental erosion⁴⁶, complex aging⁴⁷ were used.

Epiphyseal aging in young adults⁴⁸ and tooth eruption as well as root length were chosen for children⁴⁹. For aging

of childrens and babies measurement of maximal long bones⁵⁰, and tooth eruption times were⁵¹ chosen.

Mean age of males, females and childrens were 39±12, 41±9 and 5±4 respectively. At least one of the pathological signs of anemia (cribra orbitalia, porotic hyperostosis and thickening of skull bones) were observed in all of the individuals, studied in this research (PhD thesis by S.K.Arihan)

2.3. Elemental Analysis

A total of 52 skeletal (19 male, 18 female and 15 childrens) and 3 soil samples were investigated for this population. 3 samples from man and woman are from the same individuals sampled from two different areas of the femoral bone; femoral head (spongiosa part of the femoral head) and femoral neck (at the opposite position of lesser trochanter). Therefore we aimed to assess different parts of the femoral bone to determine the effect of sampling area.

³⁸ McKern/Stewart, 1957.

³⁹ Lovejoy/Richard/Meindl/Thomas/Pryzbeck/Robert 1985.

⁴⁰ Olivier 1969.

⁴¹ Kaur/Jit 1990.

⁴² Brothwell 1981.

⁴³ Krogman/İşcan 1986; Loth/İşcan, 1989.

⁴⁴ Szilavssy/Kritscher 1990.

⁴⁵ White/Black/Folkons 2012.

⁴⁶ Brothwell 1981.

⁴⁷ Workshop of European Anthropologist, 1980.

⁴⁸ Brothwell 1981.

⁴⁹ Ubelaker 1978.

⁵⁰ Workshop of European Anthropologist 1980.

⁵¹ Ubelaker 1978.

ELEMENTAL ANALYSIS OF BEYBAĞ-MUĞLA (TURKEY) BYZANTINE SKELETONS



Figure 3: Burning Samples in Stove / *Örneklerin Fırında Yakılması*



Figure 4: Sample Crushing in Agate Mortar / *Örneklerin Agat Havanda Ezilmesi*

Samples were prepared for analysis at Historical Material Research and Conservation Laboratory-MAKLAB at Gazi University (Faculty of Fine Arts, Department of Conservation and Restoration of Cultural Properties). By the suggestions made by Katzenberg and Saunders⁵² in sample preparation, our bone samples were processed to make them ready for analysis. Samples were mostly chosen from femoral and humerus head. Contact of samples with metal substances was avoided. All samples taken were washed at ultrasonic bath (Transsonic 470/H) for 10 minutes for each sample with distilled water

⁵² Katzenberg/Saunders 2008.



Figure 5: Prepared Samples of Males, Females and Children (Left to Right Respectively) / *Hazırlanmış Erkek, Kadın ve Çocuk Örnekleri (Soldan sağa doğru)*

without any hand contact. Ultrasonic device was cleaned with distilled water following cleansing of each sample in order to avoid any cross contamination between samples.

Following washing procedure drying had been performed in stove (Raypa) for 12-15 hours at 100 - 110°C. Dried samples were placed in numbered crucible and burned at stove (Nuve MF 120) for 3 hours at 550°C (Fig. 3). Samples were crushed until they are fine as dust in agate mortar, sieved (Retsch 250 µm) and placed in locked plastic bags. In order to avoid contamination between samples, agate mortar and sieve was cleaned with air pump and alcohol after each sample preparation (Fig. 4, 5).

The element contents of the skeleton/bone samples from Beybağ-Muğla were determined by using X-Ray Fluorescence (XRF) Analysis Method (PED-XRF). After pulverizing the analysis samples selected for the analysis in an agate mortar, 32 mm discs were formed, each disc was mixed with a chemical (wacks) used in the XRF analysis and placed on the sample stage of the device and was analysed. In this study, X-Lab 2000 model PED-XRF (Polarized Energy Dispersive-XRF) brand spectrometer was used. X-Lab 2000 PED-XRF spectrometer has a property of being capable of analysing the elements from sodium (Na) with atom number 11 to uranium (U) with atom number

Element	Man femoral head	Man femoral neck	Woman femoral head	Woman femoral neck	Children	Soil	P
Arsenic	7.4±0.7 ^b	5.8±1.9 ^b	9.0±1.2 ^b	3.2±0.9 ^b	7.5±0.6 ^b	26.5±9.3 ^a	*
Barium	250.7±14.4 ^a	117.5±23.0 ^b	260.1±20.2 ^a	179.3±22.0 ^{ab}	292.4±22.2 ^a	237.9±8.9 ^{ab}	*
Calcium	41.8±1.6% ^{cd}	54.4±0.8% ^{ab}	43.3±1.5% ^{bc}	55.3±0.9% ^a	42.9±1.6% ^c	31.5±4.1% ^d	*
Copper	17.4±1.3 ^{ab}	1.5±0.2 ^b	18.5±1.9 ^{ab}	1.53±0.1 ^b	27.7±5.9 ^a	24.17±4.37 ^{ab}	*
Iodine	10.1±1.2 ^a	3.2±0.7 ^b	9.1±1.1 ^a	2.4±0.0 ^b	7.3±0.9 ^a	6.1±2.0 ^a	*
Iron	2.4±0.3% ^a	0.1±0.1% ^b	2.5±0.3% ^a	0.1±0.0% ^b	2.7±0.3% ^a	3.7±0.3% ^a	*
Lead	13.8±1.1 ^a	4.9±0.8 ^b	14.5±1.0 ^a	4.6±1.1 ^b	16.9±1.2 ^a	17.6±1.5 ^a	*
Phosphorus	27.5±0.9% ^{bc}	36.3±1.4% ^a	25.6±1.2% ^c	34.1±0.9% ^{ab}	25.6±1.2% ^c	2.3±1.3% ^d	*
Strontium	131.2±6.7 ^a	133.0±19.0 ^a	144.6±7.1 ^a	138.4±16.1 ^a	131.9±6.7 ^a	92.9±8.0 ^b	*
Zinc	177.5±31.9 ^a	189.4±65.6 ^a	195.7±45.6 ^a	184.7±74.2 ^a	182.1±47.2 ^a	69.4±23.7 ^b	*

Table 1: Results of Analysis from Skeletal Samples from Femoral Head from Man, Woman and Children and Soil Samples. Data were Statistically Evaluated with Kruskal-Wallis and Tukey Tests. Statistically Homogenous Subsets for Groups are Shown in a, b, c and d Letters. Statistical Significance was Accepted as *P<0.05. Data are Presented as Mean ± Standard Error of Mean. (n=19 for Man, n=18 for Woman, n=3 for Soil). Iron was Given in the Form of % Fe203, Calcium in the Form of % Ca0 and Phosphorus in the Form of P205. / Erkek, Kadın ve Çocuk Femur İskelet Örnekleri ve Toprak Numuneleri için Analiz Sonuçları. Veriler Kruskal-Wallis ve Tukey İstatistik Testleriyle Değerlendirildi. İstatistik Açısından Homojen Gruplar a, b, c ve d Harfleri ile Gösterildi. İstatistik Anlamlılık *P<0.05 Olarak Kabul Edildi. Veriler Ortalama ± Standart Hata Olarak Gösterildi. (Erkekler için n=19, Kadınlar için n=18, Toprak için n=3). Demir % Fe203 Cinsinden Verildi. Kalsiyum % Ca0 Biçiminde ve Fosfor P205 Cinsinden Verildi.

92. The sensitivity threshold of the device is 0.5 ppm for heavy elements and about 10 ppm for light elements. In this study approximately 50 elements could be identified. Because of the weight loss (LOI) at high temperature (950°C) lithium, boron and fluorine cannot be detected. In the analysis major and minor elements are given in the form of oxide percentages (%) and trace elements are given in one in a million (ppm) concentration. In the analysis the USGS (U.S. Geological Survey) standards and as reference GEOL, GBW-7109, and GBW-7309 were used.

2.4. Statistical Analysis

Data were statistically evaluated with Kruskal-Wallis and Tukey test. P<0.05 was accepted as statistically significant. Data is presented as mean ± standard error of mean.

3. RESULTS

Sr values were found as 131.2 ppm±6.7 for man femoral head, 144.6 ppm ±7.1 for woman femoral head, 131.9

ppm±6.7 for children and 92.9 ppm±8.0 for soil samples. Values of all three groups were found significantly higher than soil samples (p<0.05). However, no significance were found between man, woman and children groups. Femoral neck values were found as 133.0 ppm ±19.0 in man and 138.4 ppm ±16.1 in woman (Table 1).

Ca values were found as 41.8 % ±1.6 for man femoral head, 43.3 %±1.5 for woman femoral head, 42.9 % ±1.6 for children and 31.5 % ±4.1 for soil samples. Values of all three groups were found significantly higher than soil samples (p<0.05). Femoral neck values were 54.4 % ±0.8 for man and 55.3 % ±0.9 for woman (Table 1).

Ba values were found as 250.7 ppm ±14.4 for man femoral head, 260.1 ppm ±20.2 for woman femoral head, 292.4 ppm ±22.2 for children and 237.9 ppm ±8.9 for soil samples. No difference was observed with groups and soil (p>0.05). Femoral neck values were 117.5 ppm ±23.0 for man, 179.3 ppm ±22.0 for woman (Table 1).

Zn values were found as 177.5 ppm ±7.2 for man femoral head, 195.7±11.86 for woman femoral head, 182.1 ppm ±12.2 for children and 69.4 ppm ±13.7 for soil samples. All values of groups were found significantly higher than

soil samples ($p < 0.05$) (Femoral neck value was found as 189.4 ppm ± 37.9 for man and 184.7 ppm ± 42.8 for woman. Table 1).

P values were found as, 27.5 % ± 0.9 for man femoral head, 25.6 % ± 1.2 for woman femoral head, 25.6 % ± 1.2 for children and 2.3 % ± 1.3 for soil samples. Values found for P for man, woman and children groups were significantly higher than soil samples ($p < 0.05$). No significant difference was found between man, woman and children. Femoral neck values were found as 36.3 % ± 1.4 for man and 34.1 % ± 0.9 for woman (Table 1).

I values were found as 10.1 ppm ± 1.2 for man femoral head, 9.1 ppm ± 1.1 for woman femoral head, 7.3 ± 0.9 for children and 6.1 ppm ± 2.0 for soil samples. Femoral neck values were 3.2 ppm ± 0.7 for man and 2.4 ± 0.0 for woman. Femoral neck values were found significantly lower than those for femoral head and soil sample values ($p < 0.05$).

Fe were found as 2.4 % ± 0.3 for man femoral head, 2.5 % ± 0.3 for woman femoral head, 2.7 % ± 0.3 for children and 3.7 % ± 0.3 for soil samples. Femoral neck values were found as 0.1 % ± 0.1 for man and 0.1 % ± 0.0 for woman. Femoral neck values were found significantly lower than femoral head samples ($p < 0.05$) (Table 1).

Cu were found as 17.4 ppm ± 1.3 for man femoral head, 18.5 ppm ± 1.9 for woman femoral head, 27.7 ppm ± 5.9 for children and 24.2 ppm ± 4.4 for soil samples ($p > 0.05$). Femoral neck values were found as 1.5 ppm ± 0.2 for man and 1.5 ppm ± 0.1 for woman. Femoral neck values were found significantly lower than femoral head samples ($p < 0.05$) (Table 1).

Pb were found as 13.8 ppm ± 1.1 for man femoral head, 14.5 ppm ± 1.0 for woman femoral head, 16.9 ppm ± 1.2 for children and 17.6 ppm ± 1.5 for soil samples ($p > 0.05$). Femoral neck values were found as 4.9 ppm ± 0.8 for man and 4.6 ppm ± 1.1 for woman. These values found for man and woman were significantly lower other groups ($p < 0.05$) (Table 1).

As were found as 7.4 ppm ± 0.7 for man femoral head, 9.0 ppm ± 1.2 for woman femoral head, 7.5 ppm ± 0.6 for children and 26.5 ppm ± 9.3 for soil samples. Femoral neck values were found as 5.8 ppm ± 1.9 for man and 3.2 ppm ± 0.9 for woman. Values of all groups were found significantly lower compared to soil samples ($p < 0.05$) (Table 1).

Sr/Ca rate was found as 3.14 for man femoral head, 3.34 for woman femoral head and 3.07 for children samples. Same ratio was found as 2.44 for man femoral neck and 2.5 for woman femoral neck samples.

Log (Ba/Sr) rate was found as 0.28 for man femoral head, 0.25 for woman femoral head and 0.35 for children. Same rate was found as -0.05 for man femoral neck and 0.11 for woman femoral neck samples.

Ca/P rates were found as 1.52 for man femoral head, 1.69 for woman femoral head and 1.68 for children. Same rate

was found as 1.50 for man femoral neck and 1.62 for woman femoral neck samples.

4. DISCUSSION

In this study elemental analysis of 52 samples from Beybağ-Muğla (Turkey) Byzantine population was conducted. On the bone samples 3 macro (Ca, P and Fe) and 6 trace elements (As, Ba, Pb, Sr, Zn and I) analyses were performed. For the analyses, femoral head samples were used. In order to determine any possible differences in different parts of bone material, femoral neck from 3 samples of man and woman were compared with samples obtained from femoral head.

Sr values of man, woman and children are significantly higher than soil samples from the graves, suggests that this Sr content reflects food origin rather than diagenesis. Comparison of trace element content of bone material and soil samples are a common method for determining the impact of diagenesis⁵³. Similarly close values observed in femoral neck and femoral head samples for Sr supports the possibility that diagenetic processes did not have a significant impact for this element. No significant difference between man and woman were observed (Table 1). Beybağ values (man 131.2 ppm, woman 144.6 ppm) were similar with an ancient Kelenderis population (man 198.2, woman 179 ppm)⁵⁴ whereas much lower than found in ancient İkiştepe population where values for adults vary between 674 to 757 ppm. Soil Sr for İkiştepe population (139 ppm)⁵⁵ was also higher than Beybağ population (92.9 ppm).

Ba is a good indicator of plant based diet however it was shown that it is affected from diagenesis⁵⁶. In Beybağ samples diagenesis is also apparent therefore together with Sr values it avoids conclusion on plant or animal based nutrition for man or woman.

Log (Ba/Sr) rate which is an indicator of marine/terrestrial feeding⁵⁷ is found at similar rates in man (0.28) and in woman (0.25) compared to childrens (0.35). It was expected to observe negative values for this ratio whereas we observed altered values compared to literature. Findings for man and woman are at the limit of terrestrial feeding however strong diagenetic impact on Ba avoids such conclusions. Although higher Ba content in children can be related with breast feeding however high diagenesis obscures such conclusions.

⁵³ Smrcka 2005.

⁵⁴ Çirak 2010a.

⁵⁵ Özdemir/Erdal 2009.

⁵⁶ Katzenberg/Saunders 2008.

⁵⁷ Smrcka 2005.

Ca in Beybağ for adults was found higher (adults 41 to 43% and soil 31%) than İkiztepe population population (37 to 41% and soil 10%) (Özdemir and Erdal 2009). Accumulation of Ca and Sr may show different patterns according to their presence in the diet and fiber content of the food. Increased plant fiber may decrease Ca but enhance Sr. In addition Ca content in bone tissue may be subject to diagenesis. According to our study results, Ca was found significantly higher than soil samples except males.

Sr/Ca rate is observed higher in woman (3.34) than man (3.14) and children (3.07). These values are higher compared to İkiztepe bronze age population (Özdemir and Erdal 2009).

Phosphorus was observed in similar quantities in man (27.5±0.9%), in woman (25.6±1.2%) and in children (25.6±1.2%). Although P content was significantly high compared to soil samples (2.3±1.3%) ($p<0.05$), values observed in Beybağ bone samples were much higher compared to literature. This situation reminds strong diagenesis for this element⁵⁸. were found P as 18 % for adults and 4.3 % for soil. Excessive phosphorus due to agricultural practices interferes with biological content of bone post-mortem. In addition to that in a study by Nielsen and Kristiansen⁵⁹ a prehistoric Celtic field system in Denmark possessed augmented levels in Na, K, P, Ca, Sr and Mn, whereas an attenuation in Co compared to a reference soil. Beybağ site has been inhabited at least since Bronze age therefore such enhancements in soil is also expected which in turn alter chemical composition of buried bones.

One of the most important indicators of diagenesis is the Ca/P rate⁶⁰. Ca/P rates are 1.52 for man, 1.69 for woman, 1.68 for children at femoral bone samples. This rate is lower than 2.16 which is found as a standart value in other studies⁶¹. Ca/P values in our study are much lower than other studies performed in Turkey such as, Özdemir 2008, Çırak 2010a and Özdemir and Erdal 2009. This finding seems related with high P content of sampled bone material due to diagenesis since Ca ratios are in parallel with other elemental analysis studies from Turkey. However there are studies which determine Ca/P rate as 5.3 which suggest that this ratio can change and can not be used as a constant for every site⁶².

Zn was found as 177.5±7.2 ppm for man, 195.7±11.86 ppm for woman and 182.1±12.2 ppm for children and

found statistically higher than soil which is 69.4 ppm ±13.7 ppm ($p<0.05$). Zapata⁶³ found a mean of 133 ppm for a late ancient period, Martinez-Garcia et al (2005) found a mean of 131 ppm in Catagena, Spain. From Anatolian settlements, Ezci⁶⁴ found 102.2 ppm for man, 76.5 ppm for woman and 32.3 ppm for soil and Çırak⁶⁵ found 154.3 for man and 158.5 ppm for woman. Higher values were found by Güner⁶⁶ as 194.3 ppm for man, 156.9 for woman and 72.2 ppm for soil in Adramytteion population. Zn values found for man, woman and childrens in Beybağ population are within the range (150-400 ppm) of omnivors in which the humans are categorized⁶⁷. Conditions where Sr was low whereas Zn was high were related with a diet high in protein with meat, kazein (milk protein), peanut and other protein sources⁶⁸. Çırak⁶⁹ concluded on a vegetation dominant diet in ancient Kelenderis population due to findings on Sr (198.2 ppm for males and 179 ppm for females) and Zn (154.4 ppm for males and 158.7 ppm for females). In Beybağ population Sr values were lower and Zn values were higher than values found by Çırak⁷⁰. Therefore, higher protein intake can be considered for Beybağ population compared to this population. Observation of high Zn in our study suggests a diet high in protein. This protein richness is not just related with meat consumption⁷¹. Therefore according to our results it is not possible to state high meat consumption in this population. Zn is known to incorporate to bone tissue differently due to seasonality⁷². However it is not possible to make such correlations since it is not known the exact burial season of Beybağ samples.

Fe is an important element for nutrition and its low intake may result in diseases such as anemia. Fe was also assessed in our study. Although femoral head has a higher blood circulation compared to femoral neck cortical bone and it's the site of medullary hematopoiesis (blood production) higher Fe content is expected but when high Fe content in the soil samples are considered this discrepancy in both sampling sites reminds an effective diagenesis. In relevant scientific literature Fe in soil was also found in high quantity⁷³. In Beybağ population iron was found as 2.4 % for man, 2.5 % for woman, 2.7 % for childrens and 3.7 % for soil ($p>0.05$). This situation prevents evaluation of this important element in this aspect. However femoral

⁵⁸ Özdemir/Erdal 2009.

⁵⁹ Nielsen/Kristiansen 2014.

⁶⁰ Zapata/Perez-Sirvent/Martinez-Sanchez/ Tover 2006.

⁶¹ Posner 1969.

⁶² Klepinger/Kurn/Williams 1986.

⁶³ Zapata/Perez-Sirvent/Martinez-Sanchez/ Tover 2006.

⁶⁴ Ezci/Kaya/Erdem/Akay/Kural/Soykut/Boçoğlu/Fenyurt/Kılıç/Temiz 2013.

⁶⁵ Çınar 2010a.

⁶⁶ Güner/Türksoy/Atamtürk/Duyar 2012.

⁶⁷ Lambert/Vlasak-Simpson/Szpunar/Buikstra 1984.

⁶⁸ Lambert/Weydert-Homeyer 1993.

⁶⁹ Çırak 2010a.

⁷⁰ Çırak 2010a.

⁷¹ Katzenberg/Saunders 2008.

⁷² Lambert/Weydert-Homeyer 1993.

⁷³ Larsen 2008.

neck samples present much less iron content showing impact of diagenesis on femoral heads. Ezci⁷⁴ found 230 ppm for males, 295 ppm for females and 1278 pp for soil in Camihoyuk population.

Woman was expected to have more I than man because of pregnancy and breastfeeding however results for man (10.1 ppm) and woman (9.1 ppm) are similar and also statistical insignificance with soil shows effect of diagenesis. Nevertheless although different from soil, similar results were observed for man (3.2 ppm) and woman (2.4 ppm) for femoral neck samples.

Insignificant results in copper content of man, woman and soil samples in Beybağ population state high diagenesis for this element. In literature records mean Cu value for ancient populations varies as 12 ppm⁷⁵, 13.4 ppm⁷⁶, 28.5 in İkiztepe population⁷⁷, 38.4 in Resuloğlu population⁷⁸ and 52.5 ppm for a Jordanian population⁷⁹. In ancient Tios population Cu was found as 28.2 ppm for males and 35.3 for females. In Adramytteion population Cu values were found as 40.6 ppm for man, 48.3 ppm for woman and 30.6 ppm for soil⁸⁰. Tios population Cu levels are also high (28 ppm for man and 35 ppm for woman)⁸¹ as Adramytteion population. Cu values were 27.3 ppm for males and 29.8 ppm for females in İkiztepe population⁸² which show similar values with Tios population. Since augmented grain consumption attenuates Zn absorption and leading high Cu level in bones⁸³ it can be interpreted as an indicator of animal based diet. In this aspect low Cu content in femoral neck samples (1.5 ppm for males and females) of Beybağ population suggests a low plant based and high animal food based diet. Compared to other literature records, Beybağ population Cu values are similar to Camihoyuk population (1.2 ppm for males and 0.47 ppm for females)⁸⁴.

As values in Beybağ population presents low arsenic content in bones (man 7.4 ppm, woman 9 ppm and children 7.5 ppm) however soil As content was high (26.5 ppm). In another Anatolian settlement Adramytteion population As values were found higher as 26.6 ppm for man, 35.5 ppm for woman and 7.5 ppm for soil

in⁸⁵. Values of As for another population in Anatolia (İkiztepe Early Bronze Age population) were found in between Adramytteion and Beybağ populations. In this population As values were 16.9 ppm for males, 15.0 ppm for females, 14.1 ppm for childrens and 6.0 ppm for soil samples⁸⁶. No metal object findings within graves in Beybağ population eliminates such an interaction with burial objects however significantly lower As content observed in femoral neck samples (5.8 ppm for man and 3.2 ppm for woman) suggests partial impact of diagenesis on this burial site for this element.

Lead is a toxic element with well known physiological outcomes. It accumulates in bones. It has a mean of 5 ppm in ancient populations⁸⁷. Among most potent reasons for lead accumulation in bones, use of lead containing cookware is being accused. In İkiztepe population in Northern Turkey high lead values were related with the use of lead made cookware⁸⁸. In Beybağ population information about lead containing cookware is not present however high lead content of soil compared to finding in Minnetpınarı population (male 5.78 ppm, female 5.67 ppm, children 4.63 ppm)⁸⁹ reminds a strong diagenetic process. Pb values in ancient populations may vary vastly. Zapata et al⁹⁰ found mean Pb for population as 225.0 ppm whereas Gonzalez-Reimers et al⁹¹ found mean Pb for population as 4.06 for a Bronze age population in Spain. Another low Pb finding was from Camihoyuk, Turkey as 3.43 ppm for males, 2.71 ppm for females and 0 ppm for soil samples⁹².

5. CONCLUSION

Results of elemental analysis of Beybağ samples present a high diagenetic impact of burial environment on Ba, I, Fe and P. This situation hampers evaluation of Log (Ba/Sr) and assessment of diet choices of Beybağ population such as animal/plant or terrestrial/marine based diet. Although sample sizes are small for femoral neck samples they show great difference with femoral head samples for the elements Ba, Cu, I, Fe and Pb.

High Ba content in all of the groups, observation of Ca/P rate altered from the standart limits in humans and dissimilarity of femoral neck with femoral head for certain

⁷⁴ Ezci/Kaya/Erdem/Akay/Kural/Soykut/Boçoğlu/Fenyurt/Kılıç/Temiz 2013.

⁷⁵ Zapata/Perez-Sirvent/Martinez-Sanchez/ Tover 2006.

⁷⁶ Martinez-Garcia 2005.

⁷⁷ Özdemir 2010.

⁷⁸ Güner 2011.

⁷⁹ Grattan/Abu Karaki/Hine/Poland/Gilbartian/Al-Saad/Pyatt 2005.

⁸⁰ Güner/Türksoy/Atamtürk/Duyar 2012.

⁸¹ Çırak/Akyol 2015.

⁸² Özdemir/Erdal/Demirci 2010.

⁸³ Ezzo 1994.

⁸⁴ Ezci/Kaya/Erdem/Akay/Kural/Soykut/Boçoğlu/Fenyurt/Kılıç/Temiz 2013.

⁸⁵ Güner/Türksoy/Atamtürk/Duyar 2012.

⁸⁶ Özdemir/Erdal/Demirci 2010.

⁸⁷ Becker/Spadaro/Berk 1968.

⁸⁸ Özdemir 2008.

⁸⁹ Çırak/Akyol 2014.

⁹⁰ Zapata/Perez-Sirvent/Martinez-Sönehez/Tovar 2006.

⁹¹ Gonzales-Reimers/Velasco-Vázquez/Arnay/de/la/Rosa/Santolaria/Fernández/Galindo/Martin 2001.

⁹² Ezci/Kaya/Erdem/Akay/Kural/Soykut/Başoğlu/Fenyurt/Kılıç/Temiz 2013.

elements demonstrate a strong impact of diagenesis for this burial site. When Zn is considered as a non affected element from diagenetic processes it is possible to state that man and woman in this population had similar protein intake. Similar content of Sr in both genders suggests a similar plant source was incorporated into diet. However Sr/Ca rate observed in woman (3.34) is higher than man (3.14) and childrens (3.07). More animal food in the diet was mentioned in past populations such as İkiztepe population⁹³.

Low As, Cu and Pb levels in Beybağ samples show no sign of toxication by those elements. This can be concluded as no potential impact of those elements on the observed paleopathological signs on this population samples.

Although sample sizes are small for femoral necks, comparison of femoral head to femoral neck bone values for certain elements reveals importance of sampling area from bone tissue. In areas such as Beybağ region where burial types are simple earth grave, agricultural practices are frequent and contamination with diagenetic processes are augmented, sampling from femoral neck may prevent loss of information from elemental analysis.

As a result it can be concluded from the data obtained from this study, correlation of reported paleopathological findings in Beybağ Byzantine population with any toxication from assessed toxic elements or any apparent lack of macro or micro nutrients is not evident. However presence of anemia signs in all of the sample material suggests infectious diseases and low hygiene conditions in this population and needs further assessment on burial locations and burial types with less influence of diagenetic processes in order to figure out nutritional status of this population.

This study presents an example for assessment of nutritional status of individuals belonging to a high paleopathological disease bearing ancient population with the help of elemental analysis.

5. ACKNOWLEDGEMENTS

Authors would like to thank to Prof. Dr. Ahmet Adil Tırpan for providing research material and Prof. Dr. Erksin Güleç for scientific contribution to the study. Furthermore, the authors owe a great dept of gratitude to Prof. Dr. Yusuf Kağan Kadioğlu (Ankara University, Earth Science Application and Research Center-YEBİM) for the XRF analyses and project laboratory assistants Gülşen Albuz (Gazi University) and Kıymet Deniz (Ankara University) for sample preparations.

6. REFERENCES

- AKSOY, M. 2007.
Ansiklopedik Beslenme Diyet ve Gıda Sözlüğü. Ankara.
- ALLEN, K.J. / GURRIN, L.C. / CONSTANTINE, C.C. / OSBORNE, N.J. / DELATYCKI, M.B. / NICOLL, A.J. / MCLAREN, C.E. / BAHLO, M. / NISSELLE, A.E., / VULPE, C.D. / ANDERSON, G.J. / SOUTHEY, M.C. / GILES, G.G. / ENGLISH, D.R. / HOPPER, J.L. / OLYNYK, J.K. / POWELL, L.W. / GERTIG, D.M. 2008
“Iron-Overload-Related Disease in HFE Hereditary Hemochromatosis”, *N Engl J Med.* 358: 221-30.
- ARIHAN, S.K. 2013.
Beybağ Mevkii (Muğla) Bizans Dönemi Toplumunda Beslenmeye Bağlı Gelişen Paleopatolojik Rahatsızlıklar (Ankara Üniversitesi, Sosyal Bilimler Enstitüsü, Antropoloji (Paleoantropoloji) Anabilim Dalı). Ankara.
- BECKER, R.O. / SPADARO, J.A. / BERK, E.W. 1968
“The Trace Elements of Human Bone”, *Journal of Bone and Joint Surgery* 50: 326-336.
- BOWEN, H.J.M. 1979.
Environmental Chemistry of the Elements. London.
- BUIKSTRA, J.E. / UBELAKER, D.H. 1994.
Standards: For Data Collection From Human Skeletal Remains, *Arkansas Archeological Survey Research Series No: 44.*
- BROTHWELL, D.R. 1981.
Digging up Bones. Oxford.
- BURTON, J. / PRICE, T.D. 2002.
The Use and Abuse of Trace Elements for Paleodietary Research p. Biogeochemical Approaches to Paleodietary Analysis. (Eds. S.H.Abrose/M.A.Katzenberg). New York: 159-171.
- ÇIRAK, M.T. 2010A
“Eski Anadolu Toplumunda Eser Element Analiziyle Beslenmenin Belirlenmesi”, *ZfWT* 2: 191-200.

⁹³ Özdemir 2008.

ÇIRAK, M.T. 2010B.

Minnetpınarı Ortaçağ Toplumunda Eser Element Analizi ile Paleodiyetin Belirlenmesi (Ankara Üniversitesi Sosyal Bilimler Enstitüsü Yayınlanmamış Doktora Tezi). Ankara.

ÇIRAK, M.T. / ÇIRAK, A. / AKYOL, A.A.

“2015 Bizans Dönemi Tios Toplumunu İskeletlerinde Bakır (Cu) Birikimi. 2006-2012 Arkeolojik Çalışmaları ve Genel Değerlendirme”, (Eds. S. Atasoy/Ş. Yıldırım) Zonguldak.

ÇIRAK, M.T. / AKYOL, A.A. 2014.

“Kilikya Toplumunu İskeletlerinde Kurşun (Pb) Düzeyleri”, Hitit Üniversitesi Sosyal Bilimler Enstitüsü Dergisi 7/1: 36-47.

EMSLEY, J. 1998.

The Elements, 3rd edition. Oxford: 1998.

FARNUM, J.F. / SANDFORD, M.K. 1997.

“Trace Element Analysis of the Tutu Skeletal Remains”, Report Submitted to the Office of Planning and Natural Resources, Division of Archaeology and Historic Preservation, Government of the United States Virgin Islands.

EZCI, Y. / KAYA, S. / ERDEM, O. / AKAY, C. / KURAL, C. / SOYKUT, B. / BAŞOĞLU, O. / FENYURT, Y. / KILIÇ, S. / TEMİZ, Ç. 2013.

“Paleodietary Analysis of Human Remains from A Hellenistic-Roman Cemetery at Camihöyük, Turkey”, Journal of Anthropology 2: 1-7.

EZZO, J.A. 1994.

“Putting The ‘Chemistry’ Back Into Archeological Chemistry Analysis: Modeling Potential Paleodietary Indicators”, Journal of Archeological Science 13: 1-34.

GONZALEZ-REIMERS, E. / VELASCO-VÁZQUEZ, J., / ARNAY-DE-LA-ROSA, M. / SANTOLARIA-FERNÁNDEZ, F. / GALINDO-MARTÍN, L. 2001.

“Paleonutritional analysis of the Pre-Hispanic population from Fuerteventura (Canary Islands)”, Science of the Total Environment 264: 215-220.

GRATTAN, J. / ABU KARAKI, L., / HINE, D. / TOLAND, H. / GILBERTSON, D. / AL-SAAD, Z. / PYATT, B. 2005.

“Analyses of Patterns of Copper and Lead Mineralization in Human Skeletons Excavated from An Ancient Mining and Smelting Centre in the Jordanian Desert: A Reconnaissance study”, Mineralogical Magazine 69: 653-666.

GRUPE, G. 1995.

“Reconstructing Migration in the Bell Beaker Period By $87\text{Sr}/86\text{Sr}$ Isotope Ratios in Teeth and Bones”, Proceedings Of The 10th International Symposium On Dental Morphology. Berlin: 338-342.

GÜNER, C. / ALİYEYEV, V. / ATAMTÜRK, D. / DUYAR, İ. / SÖYLEMEZOĞLU, T. 2011.

“Retention of Zn, Cu, Cd, Pb, and As on Human Bones Unearthed at a Central Anatolian Early Bronze Age excavation site (Resuloğlu, Turkey)”, Eurasian Journal of Anthropology 2: 27-39.

GÜNER, C. / TÜRKSOY, V.A. / ATAMTÜRK, D. / DUYAR, İ. 2012.

“Adramytteion (Örentepe, Balıkesir) Erken Bizans Dönemi İnsan İskeletlerinin Kimyasal Analizi”, İnsanbil Dergi: 1/2: 81 -93.

JOHANSSON, S.A.E. / CAMPBELL, J.L. 1988.

PIXE: A Novel Technique for Elemental Analysis. Chichester.

KATZENBERG, M.A. / SAUNDERS, S.R. 2008.

Biological Anthropology of the Human Skeleton. London.

KAUR, H. / JIT, I. 1990.

“Age Estimation from Cortical Index of The Human Clavicle in Northwest Indians”, American Journal of Physical Anthropology 83/3: 297- 305.

KLEPINGER, L.K. / KURN, L.J. / WILLIAMS, W.S. 1986.

“An Elementary Analysis of Archaeological Bone from Sieily as Test of Predictability of Diagenetic Change”, American Journal Of Physical Anthropology 70: 325-331.

KROGMAN W.M., İŞCAN M.Y. 1986.

The Human Skeleton in Forensic Medicine. Illinois.

KYLE, J.H. 1986.

“Effect Of Post-Burial Contamination on the Concentration of Major And Minor Elements in Human Bones and Teeth-The Implivations for Palaeodietary Research”, Journal of Archeological Science 13: 403-416.

- LARSEN, C.S. 2008.
Bioarchaeology. Interpreting Behaviour from the Human Skeleton. Cambridge.
- LAMBERT, J.B. / VLASAK-SIMPSON, S. / SZPUNAR, C.B. / BUIKSTRA, J.E. 1984.
“Ancient Human Diet from Inorganic Analysis of Bone”, *Acc. Chem.* 17: 298-305.
- LAMBERT, J.B. / WEYDERT-HOMEYER, J.M. 1993.
“Dietary Inferences from Element Analyses of bone”, *Prehistoric Human Bone-Archaeology at the Molecular Level* (Eds. J.B.Lambert/G. Grupe): 226-227.
- LEE-THORP, J.A. / SPONHEIMER, M. / VAN DER MERWE, N.J. 2003.
“What do Stable Isotopes Tell Us About Hominid Dietary and Ecological Niches in the Pliocene?” *International Journal of Osteoarchaeology* 13: 104-113.
- LOTH, S.R. / İŞCAN, M.Y. 1989.
Osteological Manifestations of Age in the Adult. Reconstruction of Life From the Skeleton. New York.
- MAYS, S. 1998.
The Archaeology of Human Bones. London and New York. Routledge.
- MARTÍNEZ-GARCÍA, M.J. / MORENO, J.M. / MORENO-CLAVEL, J. / VERGARA, N. / GARCÍA-SÁNCHEZ, A. / GUILLAMÓN, A. / PORTÍ, M. / MORENO-GRAU, S. 2005.
“Heavy Metals in human Bones in Different Historical Epochs”, *Science of the Total Environment* 348: 51-72.
- MCKERN, T.W. / STEWART, J.H. 1957.
Skeletal Age Changes in Young American Males, Analyzed from the Standpoint of Identification. Headque QM Res and Dev Command. Tech Rep EP-45. Natick, Mass.
- NIELSEN, N.H. / KRISTIENSEN, S.M. 2014.
“Identifying Ancient Manuring: Traditional Phosphate vs Multi Element Analysis of archaeological soil”, *Journal of Archaeological Science.* 42: 390-398.
- OLIVIER, G. 1969.
Practical Anthropology. Illinois.
- ÖZDEMİR, K. 2008.
İkiztepe Tunç Çağı Topluluğunda Element Analiziyle Beslenme Yapısının Belirlenmesi (Hacettepe Üniversitesi, Sosyal Bilimler Enstitüsü Yayınlanmamış Doktora Tezi). Ankara.
- ÖZDEMİR, K. / ERDAL, Y.S. 2009.
“Erken Tunç Çağı İkiztepe Topluluğunda Stronsiyum-kalsiyum Oranı ile Sütten Kesme Yaşının Belirlenmesi” *Çocuk Sağlığı ve Hastalıkları Dergisi* 52: 128-140
- ÖZDEMİR, K. / ERDAL Y.S. / DEMİRCİ, S. 2010.
“Arsenic Accumulation on the Bones in the Early Bronze Age İkiztepe Population, Turkey”, *Journal of Archaeological Science* 37: 1033-1041.
- ÖZDEMİR, K. / ERDAL Y.S., 2012.
“Element Analizleri ile Erken Tunç Çağı İkiztepe Toplumunun Yaşadığı Ekolojik Ortam ve Besin Kaynaklarının Belirlenmesi Üzerine Bir Çalışma”, *Türkiye’de Arkeometrinin Ulu Çınarları*. Prof. Dr. Ay Melek Özer ve Prof. Dr. Şahinde İronci’ye Armağan Kitabı. İstanbul.
- PEARSON, K. 1899.
“Mathematical Contribution on the Theory of Evolution. On the Reconstruction of the Stature of Prehistoric Races”, *Philosophical Transactions of the Royal Society of London. Series A*, 192: 169-244.
- PEMMER, B. / ROSCHGER, A. / WASTL / A. / HOFSTAETTER, J.G. / WOBRAUSCHEK, P. / SIMON, R. / THALER, H.W. / ROSCHGER, P. / KLAUSHOFER, K. / STRELIA, C. 2013.
“Spatial Distribution of the Trace Elements Zinc, Strontium and Lead in Human Bone Tissue”, *Bone* 57/1: 184-193.
- ROSCHGER, P. / KLAUSHOFER, K. / STRELI, C. 2013.
“Spatial Distribution of the Trace Elements Zinc, Strontium and Lead in Human Bone Tissue”, *Bone* 57: 184-193.
- SHACKLEY, M.S. (ED.), 2011.
X-Ray Fluorescence Spectrometry (XRF) in Geoarchaeology, DOI 10.1007/978-1-4419-6886-9-2.

ELEMENTAL ANALYSIS OF BEYBAĞ-MUĞLA (TURKEY) BYZANTINE SKELETONS

- SILLEN, A. / SEALY, J.C. / VAN DER MERWE, N.J. 1989. Workshop of European Anthropologists, 1980. "Recommendations for Age and Sex Diagnoses of Skeletons", *Journal of Human Evolution* 9: 517-549.
- "Chemistry and Paleodietary Research: No More East Answers", *American Antiquity* 54: 504-512.
- SILLEN, A. 1992. Diagenesis of the inorganic Phase of Femoral neck. *The Chemistry of the Prehistoric Bone* (Ed. T.D. Price) Cambridge: 221-223.
- SMRCKA, V. 2005. Trace Element in Bone Tissue. Univerzita Karlova Praga Nakladatelstvi Karolinium.
- SOKOLOFF, L. 1988. "Kashin-beck Disease: Current Status", *Nutrition Reviews* 46/3:113-119.
- STEELE, D.G. / BRAMBLETT, C.A. 1988. *The Anatomy of Biology of the Human Skeleton*. Texas.
- SZILAVSSY, J. / KRITSCHER, H. 1990. "Estimation of Chronological Age in Man Based on the Spongy Structure of Long Bones", *Anthrop. Anz.* 48: 159-164.
- POSNER, A.S. 1969. "Crystal Chemistry of Bone Mineral", *Physical Review* 49: 760-792.
- TIRPAN, A. / SÖĞÜT, B. / BÜYÜKÖZER, A. 2009. "Lagina, Börükçü, Belentepe ve Mengefe 2008 Yılı Çalışmaları", 31. Kazı Sonuçları Toplantısı-3. Ankara: 505-527.
- TRIMBLE, C.C. / MACKO, S.A. 1997. "Stable Isotope Analysis of Human Remains: A Tool for Cave Archaeology," *Journal of Cave and Karst Studies* 59/3: 137-142.
- UBELAKER, D. H. 1978. *Human Skeletal Remains*. Chicago.
- WHITE, T.D. / BLACK, M.T. / FOLKENS, A.P. 2012. *Human Osteology*. Elsevier.
- ZAPATA, J. / PÉREZ-SIRVENT, C. / MARTÍNEZ-SÁNCHEZ, M.J. / TOVAR, P. 2006. "Diagenesis, Not Biogenesis: Two Late Roman Skeletal Examples", *Science of the Total Environment* 369: 357-368.