

Antiplasmodial and cytotoxicity evaluation of *Artocarpus sericarpus* leaves extracts and fractions as a potential source of antimalarial substances

Lidya TUMEWU^{1,2,3} , Hilkatul ILMI² , Rina PUSPITASARI⁴ , Defi Kartika SARI² , Hanifah Khairun NISA² , Adita Ayu PERMANASARI² , Irfan Rayi PAMUNGKAS² , Aty WIDYAWARUYANTI^{2,5} , Achmad Fuad HAFID^{2,5*} 

¹ Doctoral Program Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia.

² Natural Product Medicine Research and Development, Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia.

³ Research Center on Global Emerging and Re-emerging Infectious Diseases, Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia.

⁴ Magister Program of Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia.

⁵ Department of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia.

* Corresponding Author. E-mail: achmadfuad@ff.unair.ac.id (A.F.H); Tel. +62-31-593 31 50.

Received: 11 April 2023 / Revised: 23 June 2023 / Accepted: 01 July 2023

ABSTRACT: Malaria is one of the tropical diseases that become a health problem worldwide. The elimination of this disease faced various burdens, leading to the need for new antimalarial drugs. Based on the previously reported antimalarial activity of *Artocarpus* genus, *Artocarpus sericarpus* has been selected as one of the candidates which are potential to be explored as a source of antimalarial substances. This study aimed to obtain active antimalarial fractions from *A. sericarpus* leaves. Extraction of *A. sericarpus* leaves was conducted by ultrasonic-assisted extraction using n-hexane, dichloromethane, and methanol as solvents. All extracts were tested for their antimalarial activity by Lactate Dehydrogenase (LDH) assay. The most active extract was fractionated by Vacuum Liquid Chromatography (VLC) and fractions were further tested by LDH assay as well. Cytotoxicity test conducted by resazurin assay on several cell lines to determine the active and nontoxic fractions which have the potential to be further purified and identified the active substances. The n-hexane, dichloromethane, and methanol extract showed antimalarial activity with IC₅₀ values of 23.96±0.06 µg/mL, 2.72±0.08 µg/mL, and 23.39±0.05 µg/mL, respectively. Dichloromethane extract was chosen for further separation due to its highest antimalarial activity. The separation was obtained in nine fractions. Fractions 2-9 had an IC₅₀ value of less than 10 µg/mL, indicating an active antimalarial substance. Meanwhile, fraction 1 has moderate antimalarial activity. Cytotoxicity test results considered all fractions to be non-toxic. The TLC profile of dichloromethane extract identified polyphenolic compounds, suggesting the active polyphenolic compounds can be further isolated from the active fractions (F2-F9).

KEYWORDS: *Artocarpus sericarpus*; cytotoxicity; extracts; fraction; malaria.

1. INTRODUCTION

Malaria is one of the tropical diseases that become a health problem worldwide. According to the World Malaria Report 2021, there were 241 million cases of malaria globally, increasing from 227 million in 2019 [1]. The elimination of this disease faced various burdens, leading to the need for new antimalarial drugs.

The history of antimalarial drug discovery from nature especially plants contributed to the search for antimalarial drug sources. The *Artocarpus* genus is reported to have antimalarial properties [2]. New prenylflavone namely Artocarpone A and new dihydrobenzoxanthone namely Artocarpone B, along with seven known prenylated flavones have been isolated from the bark of *A. champeden* and reported to be active as antimalarials [3]. Antimalarial active compounds, class of dihydrochalcone which identified as 1-(2,4-dihydroxy phenyl)-3-[8-hydroxy-2-methyl-2-(4-methyl-3-pentenyl)-2H-1-benzopyran-5-yl]-1-propanone have also been isolated from *A. altilis* leaves with a mechanism of action on inhibition of parasitic stage

How to cite this article: Tumewu L, Ilmi H, Puspitasari R, Sari DK, Nisa HK, Permanasari AA, Pamungkas IR, Widyawaruyanti A, Hafid AF. Antiplasmodial and Cytotoxicity Evaluation of *Artocarpus sericarpus* Leaves Extracts and Fractions as a Potential Source of Antimalarial Substances. J Res Pharm. 2024; 28(1): 126-132.

development. The in silico study predicted strong interaction of the compound with falcipain-2 receptor, as a cysteine protease inhibitor [4].

In this study, the potential of *A. sericarpus* as an antimalarial will be investigated. This plant was selected based on a chemotaxonomic approach with *Artocarpus* species, namely *A. champeden* and *A. altilis*. The approach considers that plants from the same taxa had close kinship relations, especially at the taxonomic level of family, genus, and species. The existence of this close relationship allows for the similarity of the substances or compounds contained in those plants [5].

The preliminary study also suggested *A. sericarpus* as selected species from *Artocarpus* genus to be further explored, apart from the chemotaxonomic approach. In our previous studies, screening of antimalarial activity on several *Artocarpus* species was conducted and the results showed that n-hexane, dichloromethane, and methanol extract from *A. sericarpus* leaves at a test concentration of 100 µg/mL could inhibit *P. falciparum* in vitro with a range of 99-100% average inhibition. Therefore, we aim to continue the investigation of the antiplasmodial activity of *A. sericarpus* leaves. This study was conducted to determine the IC₅₀ value of *A. sericarpus* leaves extract and fractions.

2. RESULTS AND DISCUSSION

This study was part of our continuing study of the development of natural product medicine as antimalarial sources, especially for *Artocarpus* genus. Our preliminary study has screened *A. sericarpus* leaves extracts and obtained data on the inhibition of *Plasmodium* growth but the effectivity level was not yet conducted. Therefore, the determination of IC₅₀ value was important to perform.

In this present study, the extraction of *A. sericarpus* leaves (800 g) yielded three extracts namely n-hexane extract (3.77 g), dichloromethane extract (16.07 g), and methanol extract (57.43 g). Thin Layer Chromatography (TLC) of extracts showed that extracts containing groups of polyphenolic compounds based on the TLC spot indicated brown to orange colors after spraying with sulfuric acid.

The antimalarial activity by LDH assay was conducted to determine the active extract. *Plasmodium falciparum* lactate dehydrogenase enzyme has been considered a potential molecular target for antimalarials due to this parasite's dependence on glycolysis for energy [6]. LDH assay was chosen because this assay has acceptable demand for equipment, labor, technical skills, and affordability and offers a good opportunity for scientists in low-and middle-income countries to participate in the global effort of discovering future antimalarial drugs [7]. The LDH assay of n-hexane, dichloromethane, and methanol extract of *A. sericarpus* leaves revealed dichloromethane extract as the most active extract, with an IC₅₀ value of less than 10 µg/mL. Meanwhile, n-hexane and methanol extract had IC₅₀ values of more than 10 µg/mL (Table 1). These results were in accordance with the previously reported antimalarial activity of *A. sericarpus* stem bark extracts in which dichloromethane had superior activity compared to n-hexane and methanol extract [8]. Dichloromethane extract from *A. sericarpus* stem bark and leaves had a similar IC₅₀ value which is 2.11 µg/mL and 2.72 µg/mL.

The criteria for the antimalarial activity of the extract according to Basco et al., 1994 and Dolabela et al., 2008 were as follows: IC₅₀ <10 µg/mL was the active extract, IC₅₀ 10-50 µg/mL was included in the moderate criteria, IC₅₀ 50-100 µg/mL extract has low activity, and IC₅₀ >100 µg/mL indicates the extract is inactive. Meanwhile, isolated compound(s) were classified as highly active if the IC₅₀ value was ≤1 µg/mL [9, 10]. From the IC₅₀ value of extracts, it is known that the dichloromethane extract was included in the criteria for the active extract while the n-hexane extract and methanol extract were included in the extract with moderate antimalarial activity. The dichloromethane extract showed the lowest IC₅₀ value compared to n-hexane and methanol extract meaning that dichloromethane was the most active among the three extracts. This study conducted bioassay-guided separation and suggested that the most active extract will be chosen for further separation. Therefore, the dichloromethane extract was continued for further fractionation process.

The separation of dichloromethane extract by VLC resulted in nine fractions. The TLC profile showed that chlorophyll is present in all fractions, but mostly in fraction 2 and 3 (F2 and F3). Meanwhile, other compounds detected in fractions were terpenoids, flavonoids, and chalcones which were represented by violet, yellow-brown, and orange spots after spraying with sulfuric acid.

The fractions were further tested for their antimalarial activity by LDH assay. The test results showed that all fractions were active with an IC₅₀ value of less than 10 µg/mL, except for fraction 1 (F1) as shown in Table 1. Fraction 1 (F1) had moderate antimalarial activity, meanwhile, fractions 2-9 (F2-F9) exhibited active antimalarial activity. The most active fraction was F8 with an IC₅₀ value of 1.43±0.06 µg/mL. These results

were similar to the antimalarial activity of dichloromethane fractions from *A. sericarpus* stem bark which reported IC₅₀ value of active fractions at a range of 1.53–3.65 µg/mL [8].

Table 1. Antimalarial activity (IC₅₀) of extracts and fractions

Samples	IC ₅₀ (µg/mL)*	Classification
n-Hexane extract	23.96±0.06	Moderate
Dichloromethane extract	2.72±0.08	Active
Methanol extract	23.39±0.05	Moderate
F1	28.13±0.07	Moderate
F2	7.43±0.04	Active
F3	2.84±0.07	Active
F4	2.39±0.07	Active
F5	2.38±0.08	Active
F6	1.75±0.06	Active
F7	1.97±0.06	Active
F8	1.43±0.06	Active
F9	2.49±0.06	Active
Chloroquine diphosphate	0.008±0.05	Highly Active

F1-F9 were fractions obtained from the separation of dichloromethane extract.

*Data represent Mean ± SD

The cytotoxicity test by Resazurin assay was also conducted for extracts and fractions. Resazurin dye (7-hydroxy-3H-phenoxazin-3-one 10-oxide) has been broadly used as an indicator of cell viability in several types of proliferation and cytotoxicity assays. Cellular viability quantification based on resazurin has advantages, including rapidity, reliability, sensitivity, safety, and cost [11]. The cytotoxicity test was conducted on several cell lines namely two carcinoma cells (Huh7, HepG2), and two normal cells (BHK-21, Vero). Two kinds of cell lines, carcinoma, and normal cells were used in the cytotoxicity assay to assure the safety of extracts and fractions as antimalarial. The active and safe substances normally showed nontoxic criteria on both carcinoma and normal cell lines. If substances showed cytotoxic on normal cell lines then their antimalarial activity should be considered whether the activity appeared due to toxic effect on cells. The cytotoxic effect of substances on carcinoma cells might be considered the substance's potential as an anticancer agent. The results showed that either extracts or fractions had CC₅₀ values higher than 20 µg/mL on all tested cell lines (Table 2). According to the U.S. National Cancer Institute (NCI) plant screening program, plant extracts with IC₅₀ values below 20 µg/mL or 10 µM following incubation between 48 and 72 h are recognized as potential cytotoxic substances [12]. In the present study, all extracts and fractions were considered nontoxic substances based on their CC₅₀ values which are above 20 µg/mL.

Evaluating Selectivity Index (SI) for any research on herbal drugs is crucial for determining whether further work can be continued. SI can be defined as the ratio of the toxic concentration of a sample against its effective bioactive concentration. The ideal drug should have a relatively high toxic concentration but a very low active concentration [13, 14]. The acceptance criterion of SI ≥ 10 was recommended for the selected bioactive sample that is the potential to be further investigated [15, 16]. F8 showed the lowest IC₅₀, high CC₅₀ values, and high SI amongst others. This data suggested that F8 was the potential fraction to be further studied to isolate the active antimalarial compounds.

Table 2. Cytotoxicity of extracts and fractions against several cell lines

Samples	Huh7		HepG2		BHK-21		Vero	
	CC ₅₀ (µg/mL)	SI	CC ₅₀ (µg/mL)	SI	CC ₅₀ (µg/mL)	SI	CC ₅₀ (µg/mL)	SI
n-Hexane extract	>100	4.17	>100	4.17	>100	4.17	>100	4.17
Dichloromethane extract	>100	36.76	>100	36.76	82.74 ± 0.02	30.42	90.00 ± 0.04	33.08
Methanol extract	>100	4.27	>100	4.27	>100	4.27	>100	4.27
F1	>100	3.55	>100	3.55	>100	3.55	>100	3.55
F2	>100	13.45	>100	13.45	>100	13.45	>100	13.45
F3	>100	35.21	93.98 ± 0.04	33.09	59.46 ± 0.04	20.94	58.72 ± 0.03	20.67
F4	>100	41.84	98.82 ± 0.03	41.34	73.99 ± 0.08	30.96	71.85 ± 0.05	30.06

F5	>100	42.02	>100	42.02	81.60 ± 0.02	34.28	86.70 ± 0.04	36.42
F6	77.02 ± 0.04	44.01	>100	57.14	68.79 ± 0.03	39.31	89.50 ± 0.02	51.14
F7	84.33 ± 0.05	42.81	93.98 ± 0.04	47.71	65.73 ± 0.06	33.36	90.95 ± 0.01	46.16
F8	>100	69.93	98.82 ± 0.03	69.10	96.46 ± 0.04	67.45	>100	69.93
F9	89.83 ± 0.03	36.08	>100	40.16	73.73 ± 0.003	29.61	78.72 ± 0.01	31.61
Chloroquine diphosphate	93.73±0.15	>100	>100	>100	90.52±0.24	>100	>100	>100

F1-F9 were fractions obtained from the separation of dichloromethane extract.

*Data represent Mean ± SD. SI = CC₅₀/IC₅₀

F8 was subjected to HPLC analysis using Shimadzu HPLC system. Reverse-phase analytical HPLC experiment was performed on Lichrospher RP-18 column (4.6 × 250 mm, 5 μm). A wavelength ultraviolet-visible detector was set at 254 nm. Gradient mobile phase acetonitrile-water 50%-50% to 80%-20% at flowrate 0.5 mL/min was used as eluent. The HPLC profile of F8 is shown in Figure 1. Several major peaks contained in F8 were observed based on the % area of peaks at Retention time (Rt) of 2.4 min, 3.9 min, 6.0 min, and 14.6 min. The assignment UV spectrum of those major peaks which represent several substances was shown in Figure 2. The UV spectrum exhibited absorption maxima at 282 nm, 276 nm, 255 nm, and 278 nm (band II), which were suggestive of a chalconoid structure [17]. Based on TLC and HPLC data, it could be suggested that F8 possibly contains polyphenolic compounds with a chalconoid structure as one of the active compounds which take a role in its antimalarial activity [18, 19].

This study was focused on the finding of antimalarial substances from *A. sericarpus* leaves extracts based on the bioassay-guided separation and determining the cytotoxicity as well. The results showed several fractions which exhibited antimalarial activity and were nontoxic. The HPLC profile of the most active fraction (F8) was performed, but the identification of substances contained in F8 was not conducted yet. This study was also limited to the active fractions activity and cytotoxicity data. Therefore, further studies need to be conducted to isolate and identify the active antimalarial compounds from active fractions, especially F8.




Figure 1. HPLC profile of F8. The analysis using Shimadzu HPLC system, Lichrospher RP-18 column (4.6 × 250 mm, 5 μm), gradient mobile phase acetonitrile-water 50%-50% to 80%-20%, flowrate 0.5 mL/min and observed at 254 nm.




Figure 2. The UV spectra assignment of several major peaks contained in F8 was detected at Retention time (Rt) of 2.4 min (A), 3.9 min (B), 6.0 min (C), and 14.6 min (D). A chalconoid structure's characteristics were detected from absorption maxima at 282 nm, 276 nm, 255 nm, and 278 nm (band II).

3. CONCLUSION

The study results described that dichloromethane extract and eight fractions (F2-F9), of which the most active fraction is F8 of *Artocarpus sericarpus* leaves contain antimalarial active substances so this plant becomes one of the potential sources of antimalarial compounds. In this study, the active compounds can be further isolated from the active fractions, which are possibly belonging to polyphenolic compounds.

4. MATERIALS AND METHODS

4.1 Plant material

Artocarpus sericarpus leave was obtained from Balikpapan Botanical Garden in East Kalimantan, Indonesia. The plant was identified at Purwodadi Botanical Garden, East Java with determination letter No.0074 IPH.06/HM/XII/2015.

4.2 Extraction and fractionation methods

Artocarpus sericarpus leaves were dried and ground into powder form. The powder (800 g) was extracted successively using n-hexane, dichloromethane, and methanol as solvents by ultrasonic-assisted extraction. The extract was then filtered and evaporated in vacuo to obtain dried extracts. All extracts were tested against *Plasmodium falciparum* 3D7 strain. The most active extract was fractionated by vacuum liquid chromatography (VLC) using silica gel as a stationary phase, gradient n-hexane-ethyl acetate (70%-30% to 60%-40% v/v), and chloroform-methanol (95%-5% to 50%-50% v/v) as mobile phase. The fractions were further tested for their antimalarial activity to determine the active fractions.

4.3 *Plasmodium falciparum* culture

P. falciparum 3D7 strain was cultured in 2% hematocrit type O human red blood cell (RBC) in RPMI-1640 (Gibco, ThermoFisher Scientific, Waltham, MA, USA), supplemented with 25 mM HEPES buffer, 2 g/L sodium hydrogen bicarbonate, 50 µg/mL hypoxanthine, 50 µg/mL gentamicin sulfate, and 0.5% (w/v) Albumax II (Gibco, ThermoFisher Scientific, Waltham, MA, USA) under 5% O₂, 90% N₂, and 5% CO₂ at 37 °C. Human RBCs were received from The Indonesian Red Cross. The medium was replaced daily and parasitemia was maintained below 5% for a routine subculture [20]. Parasitemia was determined by the examination of Giemsa's stained thin blood smears of infected erythrocytes. The percentage of parasitemia was determined by counting infected erythrocytes per 1000 total erythrocytes under a light microscope.

4.4 Antimalarial activity test by LDH assay

Antimalarial activity of extracts and fractions was determined by *lactate dehydrogenase* (LDH) assay against *P. falciparum* 3D7 strain. Parasite culture was synchronized with 5% (w/v) d-sorbitol as previously described [21]. Ring-stage parasites (100 μ L/well) were placed in a 96-well plate at 0.3% parasitemia. The 50% growth inhibition concentrations of extracts and fractions were determined by the addition of 0.4 μ L of serial dilutions of each sample to the wells. As a control, wells containing culture medium and RBCs alone were prepared. After 72 hours of incubation, parasite growth was determined by diaphorase-coupled lactate dehydrogenase (LDH) assay, as previously described [22]. The absorbance of each well was measured at 650 nm using Multiskan Skyhigh microplate spectrophotometer (Thermo Fisher Scientific, USA). The inhibition rate was calculated with the absorbance of uninfected wells defined as 100% inhibition. The IC₅₀ values were analyzed and calculated with GraphPad PRISM 6.0 by applying the “log (inhibitor) vs. response – Variable slope (four parameters)” in the “Dose-response – inhibition” equation family.

4.5 Cytotoxicity test by Resazurin assay

To measure cell viability based on metabolic activity in mammalian culture and primary cells, resazurin can be reduced resorufin by NAD(P)H has been widely used [23]. The carcinoma cell (Huh7, HepG2) and normal cell (BHK-21, Vero) lines were performed on D-MEM (High Glucose) media with L-Glutamine and Phenol Red (Wako, Fujifilm) and NaHCO₃, supplemented with 10% (v/v) Fetal Bovine Serum (Gibco) and 1% (v/v) Penicillium streptomycin (Sigma) as an antibiotic. Plant extracts and fractions were diluted with DMEM media to obtain the serial concentration: 100; 50; 25; 12.5; 6.25 μ g/mL and seeded in a micro-96 well plate. Following the extracts and fractions, the 3x10⁴, 1x10⁴, 5x10³, and 1x10⁴ cells were added into each well for Huh7, HepG2, BHK-21, and Vero cell lines, respectively, and incubated for 44 hours in a 5% CO₂ incubator at 37 °C. After 44h incubation, 10 μ L of 0.5mM Resazurin was added to each well and incubated for 4h. After the completion of incubation, the fluorescence data were measured at wavelengths ex 530 nm and em 595 nm using a Nivo plate reader (PerkinElmer). The CC₅₀ values were analyzed and calculated with GraphPad PRISM 6.0.

Acknowledgements: The authors would like to thank Professor Tomoyoshi Nozaki, The University of Tokyo, for providing Huh7, HepG2, BHK-21, and Vero cell lines. This research was funded by Universitas Airlangga through Penelitian Unggulan Fakultas (PUF) with contract No. 130/UN3.9.4/PT/2022. The authors were grateful to Natural Product Medicine Research and Development, Institute of Tropical Disease, Universitas Airlangga for facilities support.

Author contributions: Concept – A.W., A.F.H.; Design – A.W., A.F.H.; Supervision – A.W., A.F.H.; Resources – A.W., A.F.H., L.T., H.I., D.K.S.; Materials – A.W., A.F.H., L.T., H.I., D.K.S.; Data Collection and/or Processing – L.T., H.I., R.P., D.K.S., H.K.N., A.A.P., I.R.P.; Analysis and/or Interpretation – L.T., H.I., R.P., D.K.S., H.K.N., A.A.P., I.R.P.; Literature Search – L.T., H.I., R.P., D.K.S., H.K.N.; Writing – L.T., H.I., D.K.S.; Critical Reviews – L.T., H.I., R.P., D.K.S., H.K.N., A.A.P., I.R.P., A.W., A.F.H.

Conflict of interest statement: The authors declare that they have no conflict of interest.

REFERENCES

- [1] WHO. World malaria report 2021. <https://www.who.int/publications/i/item/9789240040496>. 2021 (accessed on 07 November 2022).
- [2] Hafid AF, Febrianty N, Septiani RP, Ranggaditya D, Fabriana LH, Widyawaruyanti A. Antimalarial activity of crude extracts of *Artocarpus heterophyllus*, *Artocarpus altilis* and *Artocarpus camansi*. Asian J Pharm Clin Res. 2016;9(1):279–281.
- [3] Widyawaruyanti A, Subehan, Kalauni SK, Awale S, Nindatu M, Zaini NC, Syafruddin D, Asih PBS, Tezuka Y, Kadota S. New prenylated flavones from *Artocarpus champeden*, and their antimalarial activity in vitro. J Nat Med. 2007;61(4):410–413. <https://doi.org/10.1007/s11418-007-0153-8>.
- [4] Hidayati AR, Widyawaruyanti A, Ilmi H, Tanjung M, Widiandani T, Siswandono S S, Syafruddin D, Hafid AF. Antimalarial activity of flavonoid compound isolated from leaves of *Artocarpus altilis*. Pharmacogn J. 2020;12(4):835–842. <https://doi.org/10.5530/pj.2020.12.120>.
- [5] Wang Y. Needs for new plant-derived pharmaceuticals in the post-genome era: An industrial view in drug research and development. Phytochem Rev. 2008;7(3):395–406. <https://doi.org/10.1007/s11101-008-9092-6>.
- [6] Penna-Coutinho J, Cortopassi WA, Oliveira AA, França TCC, Krettli AU. Antimalarial activity of potential inhibitors of *Plasmodium falciparum* lactate dehydrogenase enzyme selected by docking studies. PLoS One. 2011;6(7):e21237. <https://doi.org/10.1371/journal.pone.0021237>
- [7] Fokou PVT, Tali BMT, Dize D, Mbouna CDJ, Ngansop CAN, Keumoe R, Tchokouaha LRY, Tchouankeu JC, Escudie F, Duffy J, Boyom FF. Implementation and continued validation of the malaria *Plasmodium falciparum* lactate

- dehydrogenase-based colorimetric assay for use in antiplasmodial drug screening. *Anal Biochem.* 2022 ;648:114669. <https://doi.org/10.1016/j.ab.2022.114669>
- [8] Tumewu L, A'yun LQ, Ilmi H, Hafid AF, Widyawaruyanti A. *Artocarpus sericarpus* stem bark contains antimalarial substances against *Plasmodium falciparum*. *J Basic Clin Physiol Pharmacol.* 2021;32(4):853–858. <https://doi.org/10.1515/jbcpp-2020-0397>
 - [9] Basco LK, Mitaku S, Skaltsounis AL, Ravelomanantsoa N, Tillequin F, Koch M, Le Bras J. In vitro activities of furoquinoline and acridone alkaloids against *Plasmodium falciparum*. *Antimicrob Agents Chemother.* 1994;38(5):1169–1171. <https://doi.org/10.1128/aac.38.5.1169>
 - [10] Dolabela MF, Oliveira SG, Nascimento JM, Peres JM, Wagner H, Póvoa MM, de Oliveira AB. In vitro antiplasmodial activity of extract and constituents from *Esenbeckia febrifuga*, a plant traditionally used to treat malaria in the Brazilian Amazon. *Phytomedicine.* 2008;15(5):367–372. <https://doi.org/10.1016/j.phymed.2008.02.001>
 - [11] Borra RC, Lotufo MA, Gaglioti SM, Barros F de M, Andrade PM. A simple method to measure cell viability in proliferation and cytotoxicity assays. *Braz Oral Res.* 2009; 23(3):255–262. <https://doi.org/10.1590/S1806-83242009000300006>
 - [12] Boik J. *Natural Compounds in Cancer Therapy: Promising Nontoxic Antitumor Agents From Plants & Others Natural Sources.* Minnesota: Oregon Medical Press; 2001.
 - [13] Indrayanto G, Putra GS, Suhud F. Validation of *in-vitro* bioassay methods: Application in herbal drug research. In: Al-Majed AA, editor. *Profiles of Drug Substances, Excipients and Related Methodology.* Academic Press; 2021. p. 273–307. <https://doi.org/10.1016/bs.podrm.2020.07.005>
 - [14] Pritchett JC, Naesens L, Montoya J. Treating HHV-6 Infections. In: Flamand L, Lautenschlager I, Krueger GRF, Ablashi DV., editors. *Human Herpesviruses HHV-6A, HHV-6B & HHV-7.* 3rd ed. Amsterdam: Elsevier; 2014. p. 311–31. <https://doi.org/10.1016/C2011-0-08453-3>.
 - [15] Peña-Morán O, Villarreal M, Álvarez-Berber L, Meneses-Acosta A, Rodríguez-López V. Cytotoxicity, post-treatment recovery, and selectivity analysis of naturally occurring podophyllotoxins from *Bursera fagaroides* var. *fagaroides* on breast cancer cell lines. *Molecules.* 2016;21(8):1013. <https://doi.org/10.3390/molecules21081013>
 - [16] Awouafack MD, McGaw LJ, Gottfried S, Mbouangouere R, Tane P, Spiteller M, Eloff JN. Antimicrobial activity and cytotoxicity of the ethanol extract, fractions and eight compounds isolated from *Eriosema robustum* (Fabaceae). *BMC Complement Altern Med.* 2013;13(1):289. [10.1186/1472-6882-13-289](https://doi.org/10.1186/1472-6882-13-289)
 - [17] Aksoz BE, Ertan R. Spectral Properties of Chalcones II. *FABAD J Pharm Sci.* 2012;37(4):205–216.
 - [18] Hafid AF, Ariantari NP, Tumewu L, Hidayati AR, Widyawaruyanti A. The active marker compound identification of *Artocarpus champeden* Spreng. stem bark extract, morachalcone A as antimalarial. *Int J Pharm Pharm Sci.* 2012;4(5):246–249.
 - [19] Qin HL, Zhang ZW, Lekkala R, Alsulami H, Rakesh KP. Chalcone hybrids as privileged scaffolds in antimalarial drug discovery: A key review. *Eur J Med Chem.* 2020;193:112215. <https://doi.org/10.1016/j.ejmech.2020.112215>
 - [20] Trager W, Jensen JB. Human malaria parasites in continuous culture. *Science.* 1976;193(4254):673–675.
 - [21] Lambros C, Vanderberg JP. Synchronization of *Plasmodium falciparum* erythrocytic stages in culture. *J Parasitol.* 1979;65(3):418–420.
 - [22] Wang X, Miyazaki Y, Inaoka DK, Hartuti ED, Watanabe YI, Shiba T, Harada S, Saimoto H, Burrows JN, Benito FJG, Nozaki T, Kita K. Identification of *Plasmodium falciparum* mitochondrial malate: Quinone oxidoreductase inhibitors from the pathogen box. *Genes (Basel).* 2019;10(6):471. <https://doi.org/10.3390/genes10060471>
 - [23] Lavogina D, Lust H, Tahk MJ, Laasfeld T, Vellama H, Nasirova N, Vardja M, Eskla KL, Salumets A, Rinken A, Jaal J. Revisiting the resazurin-based sensing of cellular viability: Widening the application horizon. *Biosensors (Basel).* 2022;12(4):196. <https://doi.org/10.3390/bios12040196>