

Endophytic *Bacillus* spp. isolated from *Archidendron pauciflorum*: Pharmacological property and their phytochemical constituents

Jepri Agung PRIYANTO¹ *, Muhammad Eka PRASTYA² , Rika Indri ASTUTI¹ , Minarti² ,
Tjandrawati MOZEF² 

¹ Division of Microbiology, Department of Biology, Faculty of Mathematics and Natural Sciences, IPB University, Bogor, West Java, Indonesia

² Research Center for Pharmaceutical Ingredients and Traditional Medicine, National Research, and Innovation Agency (BRIN), Kawasan Sains dan Teknologi (KST) B.J Habibie (PUSPIPTEK) Serpong, South Tangerang, Banten, Indonesia

* Corresponding Author. E-mail: jepriyanto@apps.ipb.ac.id (J.A.P); Tel. +62-251-8622833.

Received: 22 February 2023 / Revised: 27 March 2023 / Accepted: 03 April 2023

ABSTRACT: Endophytic *Bacillus* spp. associated with *Archidendron pauciflorum* could be the potential source of active compounds attributed to the pharmacological property. The present study aimed to determine the antioxidant, antibacterial, antibiofilm activities, and phytochemical profiles of the crude extracts derived from four endophytic *Bacillus* spp. isolated from *A. pauciflorum*. Among all extracts tested, extract from *Bacillus pseudomycoloides* strain DJ8 displayed the strongest antioxidant activity against DPPH and ABTS with IC₅₀ values of 24.87±1.81 and 9.18±0.10 µg/mL, respectively. As evaluated by the dual culture-agar diffusion and disc diffusion methods, three out of four *Bacillus* spp. extracts also exhibited broad spectra of antibacterial activity against four ATCC strains. Among other extracts, *B. pseudomycoloides* strain DJ8 extract became the most active extract inhibiting *Staphylococcus aureus* strain ATCC 6538, indicated by the lowest MIC (31.25 µg/mL) and MBC (125 µg/mL), which is considered a highly active antibacterial agent. Three extracts attributed to broad spectra antibacterial activity were also capable to inhibit biofilm formation of four ATCC strains, up to 62.59% in 2 MIC of the extract concentration. Phytochemicals (flavonoid, alkaloid, and terpenoid) were present in four endophytic *Bacillus* extracts. Among other extracts, an extract derived from *B. pseudomycoloides* strain DJ8 contains the highest flavonoid content (40.10±1.18 mg QE/g extract) and phenol (58.50±4.56 mg GAE/g extract). In conclusion, these four endophytic *Bacillus* spp. could be the source of pharmaceutically valuable compounds attributed to antioxidant, antibacterial, and antibiofilm properties.

KEYWORDS: Antibacterial; antibiofilm; antioxidant; *Bacillus*; endophyte.

1. INTRODUCTION

Endophytic bacteria are a facultative or obligate endosymbiotic group that inhabit plant tissues intercellularly or intracellularly without causing disease. The bacteria can enhance plant growth and protect the plant from biotic and abiotic stresses [1]. Among the endophytic bacterial group, *Bacillus* species is considered the prolific producer of active compounds, such as bacillomycin D [2], surfactin [3], fengycin [4], bacillibactin [5] and iturin [6]. Metabolites produced by endophytic *Bacillus* also showed various therapeutic potentials, such as antioxidant [7], antibacterial [8], antifungal [9], antiviral [10] and cytotoxic properties [11]. Therefore, this bacterial group is a precious source of medicinally important compounds.

The emergence of degenerative diseases induced by oxidative stresses and infectious diseases caused by antibiotic-resistant pathogens encourages researchers to explore new sources of antioxidant and antibacterial compounds. The main characteristic of antioxidant compounds is the capability to scavenge and stabilize free radicals and prevent or delay free radical production from cellular activities [12]. These properties reduce oxidative damage in human cells and tissues and become important in preventing degenerative diseases, such as cancer and coronary heart disease [13]. While, an antibacterial is characterized by the ability of certain compounds to kill or inhibit particular bacteria growth [14]. The use of available antibiotics (levofloxacin, clarithromycin, and amoxicillin) to control infection can promote many side effects, especially

How to cite this article: Priyanto JA, Prastya ME, Astuti RI, Minarti, Mozef T. Endophytic *Bacillus* spp. isolated from *Archidendron pauciflorum*: pharmacological property and their phytochemical constituents. J Res Pharm. 2023; 27(6): 2511-2521.

increasing reactive oxygen species (ROS) in cellular level which can induce mitochondrial dysfunction and oxydative tissue damage in the human body [15]. Therefore, finding new antibacterial agents with antioxidant property is an urgent need to overcome infectious diseases.

Ethnobotanical knowledge provides a useful approach for selecting plant hosts to discover potential endophytic bacteria. *Archidendron pauciflorum*, a native Southeast Asia plant, was traditionally used to overcome dysentery[16]. Twenty-four endophytic bacteria associated with this plant have been isolated in the earlier study and four isolates identified as *Bacillus* spp. showed promising antibacterial and antibiofilm activities against multidrug-resistant strains[17]. To evaluate their antibacterial spectrum comprehensively, this study aimed to investigate their antibacterial activity against four ATCC (American Type Culture Collection) strains, such as *Pseudomonas aeruginosa* strain ATCC 15442, *Escherichia coli* strain ATCC 8739, *Staphylococcus aureus* strain ATCC 6538, and *Bacillus subtilis* strain ATCC 19659, along with the evaluation of their antibiofilm, and antioxidant activities, and phytochemical content of extracts derived from four endophytic *Bacillus* spp. associated with *A. pauciflorum*.

2. RESULTS

2.1. Antioxidant activity

All extracts derived from four *Bacillus* spp. endophytic bacteria displayed various antioxidant activities against DPPH and ABTS free radicals, as indicated by their IC₅₀ values. *B. pseudomycooides* strain DJ8 exhibited the highest antioxidant activity towards DPPH and ABTS with IC₅₀ values of 24.87±1.81 and 9.18±0.10 µg/mL, respectively. However, the antioxidant activity of four endophytic bacteria extracts is still lower than the positive control, ascorbic acid and trolox (Table 1).

Table 1. Antioxidant activity of endophytic *Bacillus* spp. derived extracts against DPPH and ABTS

Sample	Antioxidant activity*	
	DPPH IC ₅₀ (µg/mL)	ABTS IC ₅₀ (µg/mL)
<i>B. megaterium</i> strain AJ5	98.31±0.94 ^d	56.25±1.87 ^d
<i>B. amyloliquefaciens</i> strain DJ4	419.63±8.88 ^f	103.29±2.04 ^e
<i>B. pseudomycooides</i> strain DJ8	24.87±1.81 ^c	9.18±0.10 ^c
<i>B. velezensis</i> strain DJ9	217.70±9.35 ^e	106.05±5.67 ^f
Ascorbic acid	2.68±0.10 ^b	4.82±0.26 ^b
Trolox	1.77±0.12 ^a	3.17±0.13 ^a

*Note: The IC₅₀ value are expressed as mean ± SEM. Different superscript letters in the same column indicate that the data were significantly different.

2.2. Antibacterial activity

Screening of antibacterial activity from four *Bacillus* spp. endophyte using dual culture agar diffusion test showed all isolates exhibited antibacterial activity against at least three ATCC strains (Table 2). A broad spectrum of antibacterial activity showed by *B. pseudomycooides* strain DJ8, *B. megaterium* strain AJ5, and *B. amyloliquefaciens* strain DJ4 against *P. aeruginosa* strain ATCC 15442, *E. coli* strain ATCC 8739, *B. subtilis* strain ATCC 19659, and *S. aureus* strain ATCC 6538. Consistently, their antibacterial activity was also shown by their metabolite extracts in 1 mg/mL concentration (Figure 1). In general, extracts from *B. pseudomycooides* strain DJ8 showed stronger antibacterial activity than other extracts, with inhibition zone ranging from 8±0.8 to 18±0.8 mm against four ATCC strains (Table 3). Extract from the four endophytic bacteria were more active on the Gram-positive strains than on the Gram-negative strains. Tetracycline (200 µg/mL) as a positive control also exhibited antibacterial activity with inhibition zone diameter ranging from 15±0.8 to 29.3±0.9 mm.

The minimum inhibitory concentration (MIC) of all endophyte extracts was determined. The strong bacterial growth inhibitors are indicated by the lower MIC value. Commonly, all endophyte extracts varied in MIC value (31.25 – 1000 µg/mL) on four strains tested (Table 4). Extract from *B. pseudomycooides* strain DJ8 possessed the lowest MIC value (ranging from 31.25-500 µg/mL) against four ATCC strains. Tetracycline as positive control displayed MIC ranging from 1.95-62.5 µg/mL. The endophytic extracts also have different minimum bactericidal concentration (MBC) values. The most promising MBC value was also exhibited by *B. pseudomycooides* strain DJ8 extract (125-500 µg/mL) against four ATCC strains investigated.

Table 2. Antibacterial activity of *Bacillus* spp. endophyte from *A. pauciflorum* against ATCC strains by dual culture agar diffusion test

Endophytic isolates	Inhibition zone diameter (mm)*			
	<i>E. coli</i> ATCC 8739	<i>P. aeruginosa</i> ATCC 15442	<i>B. subtilis</i> ATCC 19659	<i>S. aureus</i> ATCC 6538
<i>B. megaterium</i> strain AJ5	+	+	++	+
<i>B. amyloliquefaciens</i> strain DJ4	+	+	+	++
<i>B. pseudomycooides</i> strain DJ8	+	+	++	++
<i>B. velezensis</i> strain DJ9	+	-	+	++

*Note: + = less active (clear zone diameter: ≤ 6 mm), ++ = moderately active (clear zone diameter: 7-10 mm), +++ = highly active (clear zone diameter: >10 mm), - =No activity




Figure 1. Antibacterial activity of *B. pseudomycooides* strain DJ8-derived extracts (1 mg/mL) against (A) *Escherichia coli* strain ATCC 8739, (B) *Pseudomonas aeruginosa* strain ATCC 15442, (C) *Bacillus subtilis* strain ATCC 19659, (D) *Staphylococcus aureus* strain ATCC 6538. (1) DJ8 extract, (2) Tetracycline (200 µg/ml), (3) DMSO (1% v/v).

Table 3. Antibacterial activity of endophytic *Bacillus* spp. derived extract against ATCC strains by disc diffusion method

Sample	Inhibition zone (mm)*			
	<i>E. coli</i> ATCC 8739	<i>P. aeruginosa</i> ATCC 15442	<i>B. subtilis</i> ATCC 19659	<i>S. aureus</i> ATCC 6538
<i>B. megaterium</i> strain AJ5	7.3 $\pm$ 0.5 ^b	8 $\pm$ 0 ^b	16 $\pm$ 0.8 ^b	8.3 $\pm$ 0.5 ^c
<i>B. amyloliquefaciens</i> strain DJ4	8 $\pm$ 0.8 ^c	10 $\pm$ 0 ^d	17.7 $\pm$ 0.9 ^c	7.7 $\pm$ 0.5 ^b
<i>B. pseudomycooides</i> strain DJ8	8.7 $\pm$ 0.9 ^c	8 $\pm$ 0.8 ^c	18 $\pm$ 0.8 ^d	13 $\pm$ 0.8 ^d
<i>B. velezensis</i> strain DJ9	9.6 $\pm$ 0.5 ^d	NT	18 $\pm$ 0.8 ^d	13 $\pm$ 0.8 ^d
Tetracycline	28 $\pm$ 0.8 ^e	15 $\pm$ 0.8 ^c	29.3 $\pm$ 0.9 ^e	23 $\pm$ 0 ^e
DMSO	0 $\pm$ 0 ^a	0 $\pm$ 0 ^a	0 $\pm$ 0 ^a	0 $\pm$ 0 ^a

*Note: NT: Not tested.

Table 4. The Minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) of endophytic *Bacillus* spp. derived extracts against ATCC strains

Sample	MIC and MBC ($\mu\text{g/mL}$) against ATCC strains							
	<i>E. coli</i> ATCC 8739		<i>P. aeruginosa</i> ATCC 15442		<i>B. subtilis</i> ATCC 19659		<i>S. aureus</i> ATCC 6538	
	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC
<i>B. megaterium</i> strain AJ5	250	1000	1000	>1000	500	>1000	500	1000
<i>B. amyloliquefaciens</i> strain DJ4	1000	>1000	1000	>1000	500	>1000	125	500
<i>B. pseudomycooides</i> strain DJ8	125	500	62.25	250	125	500	31.25	125
<i>B. velezensis</i> strain DJ9	500	1000	NT*	NT*	62.50	250	62.50	250
Tetracycline	1.95	3.91	15.62	62.5	1.95	3.91	1.95	3.91

*Note: NT: Not tested. Endophytic extract, tetracycline, and DMSO are applied at 1 mg/mL, 200 $\mu\text{g/mL}$, and 1% (v/v), respectively. Different superscript letters in the same column indicate the data were significantly different

2.3. Antibiofilm activity

Three extracts with broad spectra of antibacterial activity were selected for antibiofilm analysis. Inhibition of biofilm formation was correlated to the extract concentration in doses dependent response (Figure 2). In general, all extracts in 2 MIC showed the highest inhibition of biofilm formation of the test strains ranging from 39.51-62.59%. *B. pseudomycooides* strain DJ8-derived extract in 2 MIC showed the strongest inhibition of biofilm formation (62.59%) against *B. subtilis* strain ATCC 19659. This extract also inhibited more than 50% biofilm formation of other ATCC strains in the concentration. However, biofilm formation of Gram-negative strains (*E. coli* strain ATCC 8739 and *P. aeruginosa* strain ATCC 15442) was more persistence in endophyte extracts, as indicated by the lower inhibition percentage compared to Gram-positive strains.




Figure 2. Inhibition of biofilm formation of endophytic *Bacillus* spp. associated with *A. pauciflorum* against ATCC strains

2.4. Phytochemical profile

Based on qualitative assay, all endophytic *Bacillus* spp. extract contain alkaloid, flavonoid, and terpenoid. However, tannin and saponin were absent in all endophyte extracts (Table 5).

Table 5. Phytochemical contents of *Bacillus* spp. endophytes from *A. pauciflorum*

Extract	Phytochemical Contents				
	Alkaloid	Flavonoid	Tannin	Saponin	Terpenoid
<i>B. megaterium</i> strain AJ5	+	+	-	-	+
<i>B. amyloliquefaciens</i> strain DJ4	+	+	-	-	+
<i>B. pseudomycooides</i> strain DJ8	+	+	-	-	+
<i>B. velezensis</i> strain DJ9	+	+	-	-	+

*Note: - : absent; +: present

2.5. Total flavonoid and phenolic compounds

Four endophyte extracts varied in the total flavonoid and phenolic content (Table 6). In general, the total phenolic content was higher than the total flavonoid in all extracts. The highest total flavonoid and phenolic compounds exhibited by *B. pseudomycooides* strain DJ8 extract which is 40.10±1.18 mg QE/g extract, and 58.50±4.56 mg GAE/g extract, respectively.

Table 6. Total phenolic and flavonoid of endophytic *Bacillus*-derived extracts

Extract	Total flavonoid (mg QE/g extract)	Total phenolic (mg GAE/g extract)
<i>B. megaterium</i> strain AJ5	32.37±0.50 ^c	41.18±2.86 ^b
<i>B. amyloliquefaciens</i> strain DJ4	23.96±0.53 ^a	50.34±1.34 ^c
<i>B. pseudomycooides</i> strain DJ8	40.10±1.18 ^d	58.50±4.56 ^d
<i>B. velezensis</i> strain DJ9	30.95±1.72 ^b	38.00±0.78 ^a

*Note: Different superscript letters in the same column indicate the data were significantly different

3. DISCUSSION

Medicinal plants provide a favorable environment for the colonization of endophytic microbes capable of producing many valuable natural products. The current study evaluated antioxidant, antibacterial, and antibiofilm properties of the crude extracts derived from four endophytic *Bacillus* spp. Antioxidant analysis showed that all extracts were active to two different free radicals, including DPPH and ABTS, with median inhibition concentration (IC₅₀) under 500 µg/mL. In this concentration, the antioxidant compounds could reduce 50% of free radicals. *B. pseudomycooides* strain DJ8 extract is considered highly active antioxidant activity due to its IC₅₀ being less than 50 µg/mL, which is 24.87±1.81 µg/mL and 9.18±0.10 µg/mL against DPPH and ABTS, respectively. Extract from *B. megaterium* strain AJ5 classified into active antioxidant activity (IC₅₀ range: 50-100 µg/mL), while extract derived from *B. velezensis* strain DJ9 classified to moderate (IC₅₀ range: 101-250 µg/mL) and *B. amyloliquefaciens* strain DJ4 to weak antioxidant activity (IC₅₀ range: 250-500 µg/mL). Compared to their host, methanolic *A. pauciflorum* fruit extract are still more active than endophyte extract, due to it has a lower value of IC₅₀ (11.28 µg/mL) against DPPH [18]. However, growing and harvesting plant biomass takes more time than culturing bacteria. It takes 40-50 days after flowering to mature fruits. Therefore, the active compound extracted from its endophytes could be the new potential source of an antioxidant agent because culturing bacteria have a shorter culture time than the host plant.

Endophytic *Bacillus* spp. associated with *A. pauciflorum* also has antibacterial activity. Based on the dual culture agar diffusion test, these endophytic colonies showed an antagonism effect against ATCC strains (Table 2). Three isolates (*B. megaterium* strain AJ5, *B. amyloliquefaciens* strain DJ4, and *B. pseudomycooides* strain DJ8) displayed broad spectra of antibacterial activity, except *B. velezensis* strain DJ9 that was not active to *P. aeruginosa* strain ATCC 15442. In addition, secondary metabolites derived from each endophytic isolate were also tested on four test strains using the disc diffusion method. Endophytic *Bacillus* extracts were more active on the Gram-positive bacteria than on the Gram-negative bacteria. The difference in the antibacterial effects of endophytic extracts against a different group of bacteria may cause by the different external structures between Gram-positive and Gram-negative bacteria. Gram-negative bacteria have two layers of the cell membrane. Consequently, the Gram-negative bacteria group is commonly more resistant to antibacterial substances[19]. The potential of antibacterial activity observed in the current study suggests that endophytic *Bacillus* spp. may play a significant role in the protection of *A. pauciflorum* against phytopathogens. Although the host also reported having antibacterial activity against pathogens[20].

The minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) of four endophytic extracts against ATCC strains were investigated. The bacterial growth was decreased by increasing endophytic extracts concentration. In general, all endophytic extracts have MIC ≤ 1000 $\mu\text{g/mL}$. In this concentration, endophytic extract prevents the visible growth of ATCC strains under *in vitro* condition. *B. pseudomycooides* strain DJ8 extract was most effective in inhibiting *S. aureus* strain ATCC 6538, as indicated by the lowest MIC of 31.25 $\mu\text{g/mL}$, considered a highly active antibacterial agent [21]. Furthermore, all extracts also were varied in MBC value (except *B. velezensis* strain DJ9 was not to be tested to *P. aeruginosa* strain ATCC 15442). MBC value was indicated by the absence of bacterial growth on the agar plates after being treated with endophytic extracts. In this concentration the extract was completely killing the test strains. The lowest MBC value (125 $\mu\text{g/mL}$) was also shown by *B. pseudomycooides* strain DJ8 extract against *S. aureus* strain ATCC 6538.

Biofilm-forming bacteria are generally more resistant to antimicrobial agents, host defense systems, and other external factors compared to free-living bacteria [22]. Therefore, exploring an antibacterial agent that could inhibit biofilm formation is globally needed. In the present study, three endophytic extracts with broad spectra of antibacterial activity showed significant inhibitory effects against biofilm formation across four test strains. 2 MIC of each extract was found as the most effective concentration in inhibiting biofilm formation. In this concentration, the inhibitory activity ranges from 39.51 to 62.59%. Several mechanisms could possibly inhibit biofilm formations, such as quorum sensing blockage, inhibition of macromolecules synthesis and adhesion, inhibition of cell division and survival, and disruption of extracellular DNA, lipopolysaccharides, protein, and exopolysaccharides [23].

Investigation on the phytochemical profile of extract derived from four endophytic *Bacillus* associated with *A. pauciflorum* exhibited the presence of alkaloid, flavonoid, and terpenoid. The phytochemical profile of endophyte extract was also similar to the extract derived from their host. All these compounds are also found in the leaf and fruit peel extract of *A. pauciflorum* [20, 24]. Previous studies reported that phytochemicals were also commonly found in the crude extract of endophytic bacteria isolated from medicinal plants [25-26]. In some cases, flavonoid, alkaloid, and terpenoid are responsible for natural products' antioxidant and antibacterial properties [27-29]. The presence of these phytochemicals within endophyte extracts can be a potential source of active compounds for medicinal and biotechnological use.

Extract with the most potent antibacterial and antioxidant activities, *B. pseudomycooides* strain DJ8 extract, contains the higher phenol content (58.50 ± 4.56 mg GAE/g extract) and flavonoid content (40.10 ± 1.18 mg QE/g extract), than other endophyte extracts. The total phenol and flavonoid may correlate with their antibacterial and antioxidant activities. The phenolic content of endophyte extracts is still lower than their host fruit extract (97.50 ± 2.45 mg GAE/g extract). However, all endophyte extracts contain higher flavonoid (ranging from 23.96 ± 0.53 to 40.10 ± 1.18 mg QE/g extract) than host fruit extract (6.67 ± 0.38 mg QE/g extract) [18].

4. CONCLUSION

Four endophytic *Bacillus* spp. associated with *Archidendron pauciflorum* showed different antioxidant, antibacterial, and antibiofilm properties. Among four extracts tested, extract derived from *B. pseudomycooides* strain DJ8 markedly showed a highly active antioxidant effect against DPPH and ABTS. Extract derived from this isolate also showed promising antibacterial and antibiofilm activities, which were more active to *Bacillus subtilis* and *Staphylococcus aureus*. Phytochemicals (flavonoid, alkaloid, and terpenoid) were present in the extract. The extract also contained phenolic and flavonoid compounds.

5. MATERIALS AND METHODS

5.1. Source of bacterial strains

In the recent study, we used 4 *Bacillus* spp. endophytic strains, which isolated from the leaves and roots of *A. pauciflorum* [17]. Of note, the target bacterial strains utilized in this research were Gram-negative, including *P. aeruginosa* strain ATCC 15442 and *E. coli* strain ATCC 8739 along with Gram-positive strains such as *S. aureus* strain ATCC 6538 and *B. subtilis* strain ATCC 19659. All bacterial tested were kindly provided by the Laboratory of Microbiology, Department of Biology, IPB University. The reason of choosing these four strains for antibacterial testing is it due to their resistance to some antibiotics (clindamycin, oxacillin, lincomycin, bacitracin, quinupristin, and teicoplanin) [30-33].

5.2. Endophytic bacterial culture and secondary metabolite extraction

Bacterial culture was carried out in 1 L nutrient broth (NB) (Himedia) made in 3 L erlenmeyer flask. At the same time, each of endophytic bacteria were re-cultured in 100 mL NB for 1 days at 28 °C with 120 rpm shaking. Then, the bacterial culture was put into a 3 L NB flask, and incubated 3 days at 28 °C with shaking at 120 rpm. After incubation period is completed, ethyl acetate was added to the bacterial culture (1 :1 v/v) followed by ± 15 min strong shaking. The upper layer was then collected and evaporated at 50 °C [34].

5.3. Analysis of 2, 2 di phenyl 1 picryl hydrazyl (DPPH) scavenging activity

A solution of 125 μ M DPPH was prepared, and 100 μ L of this radical was added to 96 well microplates followed by the addition of 100 μ L sample. Further, this mixture was homogenized and placed for 30 min before being observed at 517 nm via spectrophotometer (Thermo Scientific Varioskan Flash-ThermoFischer Elisa reader). The inhibition value was measured using the formula: $((\text{Abs. corrector} - \text{Abs. sample}) / \text{Abs. corrector}) \times 100\%$. Abs. Corrector represent absorbance of methanol + DPPH, while Abs. sample is the absorbance of samples + DPPH. Antioxidants activity was shown as capacity to reduce DPPH radicals by 50% (IC_{50}) [35]. Ascorbic acid and trolox were utilized as positive control. The use of ascorbic acid and trolox as a control in this study because the compounds were commonly known for having strong and stable antioxidant activity against DPPH and ABTS, and that is accepted as standard antioxidant compound in *in vitro* study [36-37].

5.4. Analysis of 2,20- azino-bis-(3- ethylbenzothiazoline-6-sulphonic acid) (ABTS) radical scavenging activity

The ABTS radicals were made by mixing potassium persulfate (2.45 mM) and ABTS (7 mM) followed by incubation in the dark to produce a green-colored ABTS radicals solution with an absorbance at 745 nm of about 0.70 ± 0.02 . Next, 30 μ L of the test sample was added with 270 μ L of ABTS radical in 96-well microplate and incubated for 30 min. The percentage inhibition was determined using the formula: $((\text{Abs. corrector} - \text{Abs. sample}) / \text{Abs. corrector}) \times 100\%$. Abs. corrector is the absorbance of ABTS radical + methanol, and Abs. sample is the absorbance of ABTS radical + samples. Trolox and ascorbic acid were used as a positive control. The antioxidant activity of extracts was reflected as IC_{50} value [35].

5.5. Antibacterial activity by dual-culture agar diffusion method

Four bacterial targets were cultured in 50 mL of NB in a shaker condition of 120 rpm for approximately 24 h at 28 °C. Further, 1% (v/v), equivalent to 1×10^8 CFU/mL of bacterial target was inoculated to molten NA media and poured into a sterile petri dish. After the NA media solid, bacterial endophyte colonies on the 24 h-old were streak on round shape into medium and incubated at ± 37 °C for 24 h. After that, the clear zone diameter was determined and reflected in millimeters [38].

5.6. Antibacterial activity by disc diffusion assay

Initial screening for the antibacterial activity was analyzed by the disc diffusion assay. Shortly, the potent extract (1 mg/mL) was prepared in 1% dimethyl sulfoxide (DMSO). Further, 1% (v/v) of bacterial target strain (equivalent to 1×10^8 CFU/mL) was inoculated to molten NA media and poured into sterile petri dish. Then, a sterilized paper disc (diameter: 6 mm) was impregnated with 10 μ L extract and put on the surface of NA media and incubated at 37 °C for 24 h. Discs treated with 1% DMSO and tetracycline (200 μ g/mL) were applied as a negative and positive control, respectively. The inhibition zone to the nearest millimeter was reflected in the antibacterial activity of extracts. The use of tetracycline in this study because four test strains are susceptible to tetracycline [39].

5.7. Analysis of the minimum inhibitory concentration (MIC) and the minimum bactericidal concentration (MBC)

In this assay, we used the serial microdilution method in 96 sterile wellplates. Briefly, each well was filled with 100 μ L MHB. Further, the sample was to the wells in column A and followed by serial dilution from column A to H. After that, each well was added with 100 μ L of target bacterial culture 24-h in a 0.5 McFarland (equivalent to 1×10^8 CFU/mL) and incubated at 37°C for 24 h with 200 rpm shaking. MIC was reflected as the lowest sample concentration at which bacteria showed no visible growth in wells. As for MBC value was obtained as the lowest extract concentration to kill 99.9% of the target bacteria. Positive and negative controls were conducted using tetracycline and 1% DMSO, respectively [40].

5.8. Analysis of antibiofilm activity

Three endophytic bacterial extracts at MIC value concentration (1/4; 1/2; 1; 2 MIC) were evaluated for their antibiofilm activity. Briefly, 100 μ L of brain heart infusion (BHI) media was load onto 96-well sterile microplate. Subsequently, 100 μ L of sample and controls was introduced into well A, followed by serial dilution up to well H. Next, 100 μ L of bacterial culture in the standard of 0.5 McFarland's (equivalent to 1×10^8 CFU/mL) was pipetted to each well and incubated for 24 h at 37 C with 200 rpm shaking. Following incubation, each well were rinsed two times with phosphate buffer saline (PBS) to eliminate loose non-biofilm. Further, the plates were stained with 200 μ L crystal violet (0.1%) and incubated at 37°C for 30 min. The plates were then rinsed with PBS and reconstituted with 200 μ L of 99% DMSO before photometrically detection. Absorbance was determined using a microplate ELISA reader (Thermo Scientific Varioskan Flash-ThermoFischer Elisa reader) at 595 nm. The negative control was applied in the untreated conditions[40].

5.9. Phytochemical analysis

The endophytic bacterial extracts were used for phytochemical analysis for secondary metabolites identification using the qualitative phytochemical test, which was previously described[41]. Briefly, the presence or absence of phytochemical compounds was determined by mixing an appropriate chemical reagent and sample. The absence or presence of targeted compounds, including alkaloids, terpenoids, flavonoids, tannins, and saponins classes, was qualitatively detected.

5.10. Analysis of total flavonoid and phenolic contents

The total phenolic content (TPC) and total flavonoid content (TFC) were assayed following the previous method[42]. As for TPC, distilled water (3.5 mL) was supplemented to the extract (500 μ L, 1000 μ g/mL). The Folin-Ciocalteu reagent (250 μ L) and sodium carbonate (750 μ L, 20%) were then added, followed by incubation at $\pm 27^\circ\text{C}$ for 2 hours. The absorbance was measured via a spectrophotometer (Thermo Scientific Varioskan Flash-Thermo Fischer Elisa reader) at 765 nm. On the other hand, estimation of TFC, in a 10 ml test tube, 500 μ L of extracts, 2.2 mL of distilled water, 150 μ L of NaNO₂ (5%), and 150 μ L of AlCl₃ (10%) were mixed. After 8 min incubation, 2 mL of NaOH (1 M) was subjected, and the absorbance was measured at 506 nm. Of note, as for TPC, the same method was applied to the standard solutions of gallic acid and stated as mg Gallic Acid Equivalents (GAE)/g extract. As for TFC, the total flavonoids were exhibited as milligrams of quercetin (QE)/g of extract.

5.11. Statistical analysis

Each test was determined in triplicate. The results are shown as mean values completed with standard deviation ($\pm$ SD). The 95% and 99% confidence levels obtained from one-way analysis of variance (ANOVA) were utilized. Tukey's test was performed using a p -values < 0.05, which was considered statistically significant in further statistical analysis.

Acknowledgements: All authors thank The Institution of Research and Community Services, IPB University, Indonesia, through "Young Lecturer Research Program" for supporting this study (Grant Number: 2881/IT3.L1/PT.01.03/M/T/2022, awarded to JAP). This work was also partly supported by the Research Health Organization of National Research and Innovation Agency (BRIN) Indonesia through "Medicine and Vaccine program" in year of 2023 (Grant Number: B-231/III.9/PR.03.08/1/2023, to MEP).

Author contributions: Concept – J.A.P., M.E.P., R.I.A.; Design – J.A.P., M.E.P., R.I.A.; Supervision – R.I.A., T.M.; Resources – J.A.P., M.E.P., M.M., T.M.; Materials – J.A.P., M.E.P., M.M., T.M.; Data Collection and/or Processing – J.A.P., M.E.P., M.M., Analysis and/or Interpretation – J.A.P., M.E.P., M.M., R.I.A.; Literature Search – J.A.P., M.E.P.; Writing – J.A.P., M.E.P.; Critical Reviews – R.I.A., T.M.

Conflict of interest statement: The authors declared no conflict of interest.

REFERENCES

- [1] Gouda S, Das G, Sen SK, Shin HS, Patra JK. Endophytes: A treasure house of bioactive compounds of medicinal importance. *Front Microbiol.* 2016;7:1538. <https://doi.org/10.3389/fmicb.2016.01538>
- [2] Jiao R, Cai Y, He P, Munir S, Li X, Wu Y, Wang J, Xia M, He P, Wang G, Yang H, Karunarathna SC, Xie Y, He Y. *Bacillus amyloliquefaciens* YN201732 produces lipopeptides with promising biocontrol activity against fungal

- pathogen *Erysiphe cichoracearum*. Front Cell Infect Microbiol. 2021;11:598999. <https://doi.org/10.3389/fcimb.2021.598999>
- [3] Hazarika DJ, Goswami G, Gautom T, Parveen A, Das P, Barooah M, Boro RC. Lipopeptide mediated biocontrol activity of endophytic *Bacillus subtilis* against fungal phytopathogens. BMC Microbiol. 2019; 19(1):71. <https://doi.org/10.1186/s12866-019-1440-8>
- [4] JasimB, Sreeakshmi S, Mathew J, Radhakrishnan EK. Identification of endophytic *Bacillus mojavensis* with highly specialized broad spectrum antibacterial activity. 3 Biotech.2016; 6 (2):187. <https://doi.org/10.1007/s13205-016-0508-5>
- [5] Nalli Y, Singh S, Gajjar A, Mahizhaveni B, Dusthacker VNA, Shinde PB. Bacillibactin class siderophores produced by the endophyte *Bacillus subtilis* NPROOT3 as antimycobacterial agents. Lett Appl Microbiol. 2023; 76 (2): ovac026. <https://doi.org/10.1093/lambio/ovac026>
- [6] Rasiya KT, Sebastian D. Iturin and surfactin from the endophyte *Bacillus amyloliquefaciens* strain RKEA3 exhibits antagonism against *Staphylococcus aureus*. Biocat Agric Biotech. 2021;36: 102125. <https://doi.org/10.1016/j.bcab.2021.102125>
- [7] Nisa S, Shoukat M, Bibi Y, Ayoubi SA, Shah W, Masood S, Sabir M, Bano SA, Qayyum A. Therapeutic prospects of endophytic *Bacillus* species from *Berberis lycium* against oxidative stress and microbial pathogens. Saudi J Biol Sci. 2022; 29(1):287-295. <https://doi.org/10.1016/j.sjbs.2021.08.099>
- [8] Bhoonobong A, Sodngam S, Boonlue S, Bunyatratchata W, Mongkolthananarukk W. Antibiotics constituents of endophytic *Bacillus amyloliquefaciens* UD25 extracted from a medicinal plant, *Memecylon edule* Roxb. Chiang Mai J Sci. 2017;44(3):788-799.
- [9] Gond SK, Bergen MS, Torres MS, Junior JFW. Endophytic *Bacillus* spp. produce antifungal lipopeptides and induce host defence gene expression in maize. Microbiol Res. 2015;172:79-87. <https://doi.org/10.1016/j.micres.2014.11.004>
- [10] Sorokan A, Cherepanova E, Burkhanova G, Veselova S, Rumyantsev S, Alekseev V, Mardanshin I, Sarvarova E, Khairullin R, Benkovskaya G, Maksimov I. Endophytic *Bacillus* spp. as a prospective biological tool for control of viral diseases and non-vector *Leptinotarsa decemlineata* Say. in *Solanum tuberosum* L. Front Microbiol. 2020;11:569457. <https://doi.org/10.3389/fmicb.2020.569457>
- [11] Romero-Arguelles R, Romo-Sáenz CI, Morán-Santibáñez K, Tamez-Guerra P, Quintanilla-Licea R, Orozco-Flores AA, Ramírez-Villalobos JM, Tamez-Guerra R, Rodríguez-Padilla C, Gomez-Flores R. *In vitro* antitumor activity of endophytic and rhizosphere Gram-positive bacteria from *Ibervillea sonora* (S. Watson) Greene against L5178Y-R lymphoma cells. Int J Environ Res Publ Health. 2022;19(2):894. <https://doi.org/10.3390/ijerph19020894>
- [12] Żymańczyk-Duda E, Szmigie-Merena B, Brzezińska-Rodak M, Klimek-Ochab M. Natural antioxidants-properties and possible applications. J Appl Biotechnol Bioeng. 2018;5(4):251- 258. <https://doi.org/10.15406/jabb.2018.05.00146>
- [13] Rani K. Role of antioxidants in prevention of diseases. J Appl Biotech Bioeng. 2017;4(1):495-496. <https://doi.org/10.15406/jabb.2017.04.00091>
- [14] LiD, Zhou B, Lv B. Antibacterial therapeutic agents composed of functional biological molecules. J Chem. 2021;2020: 6578579. <https://doi.org/10.1155/2020/6578579>
- [15] Kohanski MA, Tharakan A, Lane AP, Ramanathan MJ. Bactericidal antibiotics promote reactive oxygen species formation and inflammation in humans in nasal epithelial cells. Int Forum Allergy Rhinol. 2016;6(2):191-200. <https://doi.org/10.1002/alr.21646>
- [16] Bunawan H, Dusik L, Bunawan SN, Amin NM. Botany, traditional uses, phytochemistry and pharmacology of *Archidendron jiringa*: a review. Global J Pharmacol. 2013;7(4):474-478. <https://doi.org/10.5829/idosi.gjp.2013.7.4.824>
- [17] Priyanto JA, Prastya ME, Astuti RI, Kristiana R. The antibacterial and antibiofilm activities of the endophytic bacteria associated with *Archidendron pauciflorum* against multidrug-resistant strains. Appl Biochem Biotech. 2023;Published online. <https://doi.org/10.1007/s12010-023-04382-4>
- [18] Molla MM, Kamal MM, Sabuz AA, Chowdhury MGF, Khan MHH, Khatun A, Miaruddin M, Zashimuddin M, Islam MM. Chemical composition, bioactive compounds, antioxidants potential and mycotoxin of minor exotic *Archidendron pauciflorum* fruit with the focus to Bangladesh. Biocat Agric Biotech. 2021;34:102039. <https://doi.org/10.1016/j.bcab.2021.102039>
- [19] Breijyeh Z, Jubeh B, Karaman R. Resistance of Gram-negative bacteria to current antibacterial agents and approaches to resolve it. Molecules. 2020;25(6):1340. <https://doi.org/10.3390/molecules25061340>

- [20] Nasution HM, Yuniarti R, Rani Z, Nursyafira A. Phytochemical screening and antibacterial activity test of ethanol extract of jengkol leaves (*Archidendron Pauciflorum* Benth.) IC Nielsen against *Staphylococcus epidermidis* and *Propionibacterium acnes*. Int J Sci Tech Manag. 2022; 3(3): 647-653. <https://doi.org/10.46729/ijstm.v3i3.509>
- [21] Silva ACO, Santana EF, Saraiva AM, Coutinho FN, Castro RHA, Pisciotano MNC, Amorim ILC, Albuquerque UP. Which approach is more effective in the selection of plants with antimicrobial activity?. Evid-Based Complement Alternat Med. 2013;2013:308980. <https://doi.org/10.1155/2013/308980>
- [22] Sharma D, Misba L, Khan AU. Antibiotics versus biofilm: an emerging battleground in microbial communities. Antimicrob Resist Infect Cont. 2019; 8: 76. <https://doi.org/10.1186/s13756-019-0533-3>
- [23] Roy R, Tiwari M, Donelli G, Tiwari V. Strategies for combating bacterial biofilms: a focus on anti-biofilm agents and their mechanisms of action. Virulence. 2018;9(1):522-554. <https://doi.org/10.1080/21505594.2017.1313372>
- [24] Sihombing JR, Dharma A, Chaidir Z, Almahdy, Fachrial E, Munaf E. Phytochemical screening and antioxidant activities of 31 fruit peel extract from Sumatera, Indonesia. J Chem Pharm Res. 2015;7(11): 190-196.
- [25] Rathod ZR, Shah D, Shukla P, Meghani S, Patel C. Citrus limon as an abundant source of phytochemicals their evaluation and antioxidant analysis with bacterial endophytes. Curr Trends Biomed Eng Biosci. 2022;20(4):556045. <https://doi.org/10.19080/CTBEB.2022.20.55604>
- [26] Morare RF, Ubomba-Jaswa E, Serepa-Dlamini MH. Isolation and identification of bacterial endophytes from *Crinum macowanii* Baker. Afr J Biotech. 2018;17(33):10401047. <https://doi.org/10.5897/AJB2017.16350>
- [27] Achika JL, Ayo RG, Oyewale AO, Habila JD. Flavonoids with antibacterial and antioxidant potentials from the stem bark of *Uapaca heudelottii*. Heliyon. 2020; 6(2):e03381. <https://doi.org/10.1016/J.HELIYON.2020.E03381>
- [28] Achika JL, Ayo RG, Habila JD, Oyewale AO. Terpenes with antimicrobial and antioxidant activities from *Lannea humilis* (Oliv.). Sci Afr. 2020;10:e00552. <https://doi.org/10.1016/j.sciaf.2020.e00552>
- [29] Mabhiza D, Chitemerere T, Mukanganyama S. Antibacterial properties of alkaloid extracts from *Callistemon citrinus* and *Vernonia adoensis* against *Staphylococcus aureus* and *Pseudomonas aeruginosa*. Int J Med Chem. 2016;2016:6304163. <https://doi.org/10.1155/2016/6304163>
- [30] Bac Dive. *Escherichia coli* Crooks is a mesophilic human pathogen that has multiple antibiotic resistances and was isolated from faeces. 2021. <https://bacdive.dsmz.de/strain/4433>. Accessed on 22 March 2023.
- [31] Mbaveng AT, Sandjo LP, Tankeo SB, Ndifor AR, Pantaleon A, Nagdjuu BT, Kuete V. Antibacterial activity of nineteen selected natural products against multi-drug resistant Gram-negative phenotypes. Springer Plus. 2015; 4 : 823. <https://doi.org/10.1186/s40064-015-1645-8>
- [32] Osman ME-S, El-Hendawy HH, Abd El-Ail SM, HassanAM, Mahmoud DE. Correlation between resistance of *Pseudomonas aeruginosa* to benzalkonium chloride and expression of efflux pump genes. J Appl Pharm Sci. 2018;8(12): 044-052. <https://doi.org/10.7324/JAPS.2018.81206>
- [33] Speck S, Wenke C, Feßler AT, Kacza J, Geber F, Scholtzek AD, Hanke D, Eichhorn I, Schwarz S, Rosolowski M, Truyen U. Borderline resistance to oxacillin in *Staphylococcus aureus* after treatment with sub-lethal sodium hypochlorite concentrations. Heliyon. 2020;6(6): e04070. <https://doi.org/10.1016/j.heliyon.2020.e04070>
- [34] Priyanto JA, Astuti RI, Nomura J, Wahyudi AT. Bioactive compounds from sponge associated bacteria: Anticancer activity and NRPS-PKS gene expression in different carbon sources. Am J Biochem Biotech. 2017;13(4):148-156. <https://doi.org/10.3844/ajbbsp.2017.148.156>
- [35] Prastya ME, Astuti RI, Batubara I, Takagi H, Wahyudi AT. Natural extract and its fractions isolated from the marine bacterium *Pseudoalteromonas flavipulchra* STILL-33 have antioxidant and antiaging activities in *Schizosaccharomyces pombe*. FEMS Yeast Res. 2020;20(3): foaa014. <https://doi.org/10.1093/femsyr/foaa014>
- [36] Abramovič H, Grobin B, Ulrih NP, Cigić B. Relevance and standardization of in vitro antioxidant assays: ABTS, DPPH, and Folin-Ciocalteu. J Chem. 2018; 2018: 4608405 . <https://doi.org/10.1155/2018/4608405>.
- [37] Wołosiaś R, Drużyńska B, Derewiaka D, Piecyk M, Majewska E, Ciecierska M, Worobiej E, Pakosz P. Verification of the conditions for determination of antioxidant activity by ABTS and DPPH assays – a practical approach. Molecules. 2022; 27(1): 50. <https://doi.org/10.3390/molecules27010050>
- [38] Rini AF, Yuhana M, Wahyudi AT. Potency of sponge-associated bacteria producing bioactive compounds as biological control of vibriosis on shrimp. J Akuakult Ind. 2017;16(1):41-50. <https://doi.org/10.19027/jai.16.1.41-50>
- [39] Diale MO, Aswa EU, Serepa-Dlamini MH. The antibacterial activity of bacterial endophytes isolated from *Combretum mole*. Afr J Biotech. 2018;17(8): 255-262. <https://doi.org/10.5897/AJB2017.16349>

- [40] NCCLS (National Committee for Clinical Laboratory Standard), Performance Standard for Antimicrobial Susceptibility Testing. Ninth informational supplement. 30th ed. NCCLS, Malvern 2020.
- [41] Harbourne JB. Phytochemical Methods: A Guide to Modern Techniques of Plants Analysis. Chapman and Hall, London 1983.
- [42] Prastya ME, Astuti RI, Batubara I, Wahyudi AT. Antioxidant, antiglycation and *in vivo* antiaging effects of metabolite extracts from marine sponge-associated bacteria. Indian J Pharm Sci. 2019;81(2):344-353. <https://doi.org/10.36468/pharmaceutical-sciences.516>