

# Medicinal and aromatic plants traditionally used to treat metabolic diseases in the Rabat region, Morocco

Chaimae EL KOURCHI <sup>1</sup> , Oumayma BELHOUSSEINE <sup>1</sup> , Hicham HARHAR <sup>1</sup> , Abdelhakim BOUYAHYA <sup>2\*</sup> , Vijay KOTRA <sup>3</sup> , Inayat Ur REHMAN <sup>4</sup> , Long Chiau MING <sup>5</sup> , Pei Lin LUA <sup>6</sup> , Mohamed TABYAOUI <sup>1</sup> 

<sup>1</sup> Department of Chemistry, Faculty of Sciences, Mohammed V University, Rabat, Morocco.

<sup>2</sup> Department of Biology, Faculty of Sciences, Mohammed V University, Rabat, Morocco.

<sup>3</sup> Faculty of Pharmacy, Quest International University, Ipoh, Malaysia.

<sup>4</sup> Department of Pharmacy, Abdul Wali Khan University Mardan, Mardan, Pakistan.

<sup>5</sup> School of Medical and Life Sciences, Sunway University, Sunway City, Malaysia.

<sup>6</sup> Faculty of Pharmacy, Universiti Sultan Zainal Abidin, Terengganu, Malaysia.

\* Corresponding author. E-mail: a.bouyahya@um5r.ac.ma (A.B.); Tel. +212 (0) 5 37 77 18 34.

Received: 6 November 2023 / Revised: 26 January 2024 / Accepted: 27 January 2024

**ABSTRACT:** Diabetes and hyperlipidemia are major human health problems due to their high levels of mortality and morbidity. During the last decades, people affected by metabolic diseases are turning more and more to traditional medicine due to the undesirable side effects of hypoglycemic and hypolipidemic drugs. Morocco has a wide floral diversity, which offers a wide range of aromatic and medicinal plants with potential application in the nutraceutical and pharmaceutical market. In this survey, we are going to highlight the medicinal plants used traditionally by the Rabat community for the treatment of metabolic diseases like hypercholesterolemia and diabetes. A total of 475 participants were interviewed in this survey using a semi-structured questionnaire form. The results obtained were interpreted through the ethnopharmacological parameters such as UV, RFC and FUV. The results revealed 83 and 41 floristic species recorded as a remedy against diabetes and hypercholesterolemia, respectively. The most cited species are: *Olea europaea*, *Solanum melongena* and *Linum usitatissimum*. Leaves and seeds are the most used plant parts, while the oral administration of powder is the most common route of administration. It should also be noted that *Moringa oleifera*, *Stevia rebaudiana*, and *Panax ginseng* are not used by citizens but only by herbalists as anti-diabetic plants imported from outside Morocco, which implies that the people of Rabat region are not familiar with imported plants and are using the endemic and condiment species. The results of this study pave the way for phytochemical and pharmacological studies to confirm the activity of the highlighted species, and toxicological studies regarding the duration of treatment, dose and interaction with other drugs.

**KEYWORDS:** Medicinal plants; nutritional plant; plant mixtures; metabolic diseases; ethnopharmacological survey.

## 1. INTRODUCTION

Metabolic diseases (MD) represent one of the main sources of death, affecting millions of people each year in the world (in developed and developing countries), thus constituting a negative burden for the government and the community, besides its impact on the individual [1–3]. However, the development of MD including obesity, diabetes mellitus, and cardiovascular diseases, can be attributed to several environmental, lifestyle and epigenetic elements such as physical inactivity, smoking, lack, overconsumption or imbalance of dietary nutrients [4–7].

Diabetes is a worldwide health problem due to its high mortality and morbidity rates. It is an endocrine metabolic disease characterized by an increase (hyperglycemia) or decrease (hypoglycemia) of the blood glucose level. Diabetes is a chronic disease that can occur from a hereditary or environmental source, as well as it can increase the risk of several diseases in affected patients [8–13]. However, there are several classes of diabetes, including type 1 (insulin-dependent), type 2 (non-insulin-dependent) and gestational diabetes [9,12,14,15]. Type 1 diabetes, also called juvenile diabetes because of its constant onset in children, is characterized by a lack or total absence of insulin production due to the destruction of beta cells producing this hormone in the pancreatic islets of Langerhans [12,14,16]. This type of diabetes represents only 5-10% of

**How to cite this article:** El Kourchi C, Belhoussaine O, Harhar H, Bouyahya A, Kotra V, Ur Rehman I, Ming LC, Lua PL, Tabyaoui M. Medicinal and aromatic plants traditionally used to treat metabolic diseases in the Rabat region, Morocco. J Res Pharm. 2024; 28(4): 1293-1315.

cases while type 2 diabetes affects the majority of diabetics (90-95%). These diabetic patients do not need insulin from an external source, but they just need to follow a balanced diet, moderate physical exercise and take hypoglycemic and hypolipidemic drugs[12,14,17]. Whereas, gestational diabetes (GD) represents hyperglycemia first detected during pregnancy [18]. GD is linked not only to perinatal morbidity[19], but also to an increased risk of diabetes and cardiovascular disease in later life [20,21] and obesity in children[22].

Dyslipidemia is a disorder in the metabolism of lipids [cholesterol, low-density lipoprotein cholesterol (LDL), high-density lipoprotein (HDL) and triglycerides]. This alteration can result from diet, tobacco exposure, autosomal mutations in LDL receptors that result in elevated LDL cholesterol levels, which can conduct to the development of cardiovascular disease [23,24]. Previous studies conducted in Morocco on the prevalence and risk factors of type 2 diabetes (T2) revealed a group of significant risk factors for chronic complications of T2 in Moroccan patients such as increased duration of diabetes, insulin use, lack of regular exercise, hypertension, hypolipidemic treatment, elevated serum creatinine and elevated low-density lipoprotein cholesterol[25,26]. Therefore, because of the difficulty to maintain changes in the way of life, and of the undesirable secondary effects of drugs namely hepatotoxicity and myopathy, people are turning more towards herbal cure as an alternative to manage metabolic diseases. To this end, several studies have reported on the physiological modifications by primary and secondary metabolites contained in functional foods on metabolic diseases, *via* metabolomic analysis [6,27]. These functional foods derive largely from microorganisms, animals, and plants. They encompass a nutrient ratio beneficial to human health and capable of reducing many diseases [28,29]. One example is given by tea polyphenols (*Camellia sinensis* (L.) Kuntze) that reduced level of blood sugar, total cholesterol and LDL-cholesterol [30]. In addition to its rich flavor, coffee (*Coffea arabica* L.) also boasts polyphenols that are known to reduce hyperglycemia, obesity, and the risk of cardiovascular disease [31,32]. Anthocyanins, flavonoids, and resveratrol contained in many fruits and vegetable help to prevent degenerative diseases such as diabetes, hyperlipidemia, hypertension, obesity, and cancer [33–36]. Polyphenols of pomegranate (*Punica granatum* L.) such as ellagitannins and anthocyanins are responsible for hypoglycemic and hypocholesterolemic effects [37].

African health systems have relied on the traditional medicine approach for about two centuries, allowing about 70-80% of patients to be treated with the so-called complementary and alternative medicine [38–40]. According to the statistics yearbook of Morocco published by the High Commission for Planning in 2020, The region has a remarkable floristic richness where natural forest species occupy an area of 355,999 ha, distributed among the different provinces of the region, including natural deciduous species that mark the first place in terms of area (173,357 ha), followed by the matorral type (75,064 ha), then the natural resinous species (66,082 ha). Thus, the natural hardwood species include a specific richness grouping *Quercus suber* L. (133 867 ha), *Quercus ilex* L. (69 918 ha) and *Quercus canariensis* Willd. (239 ha) with the absence of the argan tree (*Argania spinosa* L.) in the region. However, natural resinous species in the Rabat region are limited to *Pinus atlantica* Endl. (65,522 ha) and *Juniperus phoenicea* L. (560 ha). In terms of cultivated area, Morocco has experienced a decline in the cultivation of cereals, where the region of Rabat produces 12,603.3 thousand quintals on an area of 526.4 thousand hectares with the dominance of soft wheat and barley, and occupies the 2nd rank after the region of Fez. In addition, the cultivation of pulses in this region is in continuous evolution, including the cultivation of chickpea and lentils (318.4 and 225.9 Mq). On the other hand, the region has experienced an evolution in the production of sugar and oilseed crops.

Given the enormous popularity of this healing system based on the use of medicinal and aromatic plants, it is important to shed light on the risk from the combination of several plants in a recipe or their interactions with conventional medicines. Therefore, the main objective of this study is to bring together all the achievements and knowledge of citizens and herbalists in the region of Rabat, Morocco, through an ethnopharmacological survey form established in two forms (online and in the field) allowing the recording of the floristic species used by this community for the treatment of metabolic diseases such as diabetes and hyperlipidemia.

## 2. RESULTS

### 2.1. Population based profile

This survey involved interviewing a total of 475 participants, including 20 herbalists and traditional healers who were questioned in their stores about the two treated diseases, and 455 people who contributed to the success of this online survey, with 241 and 214 people interviewed about diabetes and hyperlipidemia, respectively.

### 2.1.1. Herbal practitioners

A total of 20 herbalists were interviewed during the survey in their herbal stores (Table 1).

**Table 1.** Socio-demographic profile of traditional healers.

Traditional healers	Percentages %
<b>Age</b>	
<20	---
[20-40]	40
[40-60]	50
>60	10
<b>Gender</b>	
Female	5
Male	95
<b>Level study</b>	
Illiterate	5
Primary	45
Secondary	50
University	---
<b>City</b>	
Rabat	50
Temara	25
Salé	10
Kénitra	---
<b>Years' experience</b>	
[5-9]	30
[10-20]	60
>20	10
<b>Origin of information</b>	
Hereditary	63.2
Experience	36.8

The socio-demographic profile of the herbalists showed that all the traditional healers were men; this was due to the fact that in this region, only male traditional healers were listed, as well as women who may not have any encouragement to practice this job especially outside their environment (home and family). However, the age of the participants varies from 20 to 60 years old. The most represented age group was 40-60 years old (50%), followed by 20-40 years old (40%), then 60 years old and above (10%) while the age group under 20 years old was absent. On the basis of experience, 60% of the herbalists had between 10 and 20 years of experience, while 30% of the respondents had between 5 and 9 years, while 10% had more than 20 years of experience (Table 1). The majority of the herbalists had a secondary and primary level of education, with 5% being illiterate. In addition, 63.2% of the informants had knowledge inherited from their families, and 36.8% of the participants based their knowledge on experience gained since they started this work. However, the majority of the traditional healers were accompanied by their parents who have more experience in this field and from where their heritage came.

### 2.1.2. Local population

A total of 455 informants were interviewed and classified into different demographic categories, as shown in Table 2.

**Table 2.** Socio-demographic profile of online respondents.

Informant	Diabetes	Hypercholesterolemia
	%	%
<b>Age</b>		
<20	1.7	2.8
[20-40]	62.3	73.8
[40-60]	23.3	19.2
>60	12.7	4.2
<b>Gender</b>		
Men	35.6	31.8
Female	64.4	68.2
<b>Level study</b>		
Illiterate	5.5	3.3
Primary	6.4	1.9
Secondary	11.4	5.1
University	76.7	89.7
<b>City</b>		
Rabat	36	38.3
Témara	12.7	20.1
Salé	36	30.4
Kénitra	5.3	4.2
Khemisset	10	7
<b>Origin of information</b>		
Hereditary	14.4	13.4
Reading	27.2	43.3
Social initiation	52	37.3
Divine revelation	5.6	6
<b>Are you suffering from the disease?</b>		
Yes	19.9	14
No	80.1	86

Generally, the socio-demographic characteristics of the informants appear to be the same in both therapeutic categories. The age group varies from 20 to 60 years. The most representative age range is 20 to 40 years and this may be due to the increased use of online communication by this age group. However, women showed more knowledge in the field of herbal medicine. Thus, the knowledge of the respondents seems to be more argued with their university study levels and the origin of the information and this can be explained by the knowledge of the Moroccan populations regarding the socio-economic level and the knowledge acquired by the ancestors. The majority of the speakers belonged to the following cities: Rabat followed by Salé, then Temara.

In this study, we remarked that women frequently used more (69%) medicinal plants than men (31%). This trend has been observed in other ethnobotanical and ethnopharmacological studies at the regional scale [41-43], as well as at the national scale [40,44-47]. This preponderance of women in the field of herbal medicine can be explained by the fact that they take responsibility within their homes to take care of their families in terms of their health but with means that are available, effective, and less expensive, in line with the knowledge and tricks of Moroccan cooking based on the use of aromatic plants.

## 2.2. Floristic profile of food and medicinal plants

Morocco, has a remarkable floristic richness due to its geographical position and encompasses about 4500 native species which make it among the richest countries floristically at the scale of the Mediterranean basin [48]. Likewise, the studied area offers a wide ecological diversity characterized by a vast subterranean forest which in turn presents a very diverse range in terms of aromatic and medicinal plants which provides the citizen of the region with an awareness all around the world of traditional medicine. However, this tradition must be scientifically approved by further studies.

The results obtained inventoried about 98 medicinal plants for the managing of diabetes and hyperlipidemia in the region of Rabat and are presented in Table 3.

The ethnobotanical survey disclosed the presence of 98 floristic species divided into 43 families listed by citizens and herbalists in the region of Rabat as herbal remedies against diabetes and cholesterol. The most cited family was the Fabaceae followed by the Oleaceae and then the Lamiaceae (Figure 1).

In relation to the number of species used for treatments, the Asteraceae family is the most common, with 12 species, followed by the Lamiaceae (10 species) and Fabaceae (9 species). The Oleaceae family presents a higher number of citations with only one species (*Olea europaea*) that represents a remarkable socio-economic, cultural and agronomic resource for Morocco. Families most commonly cited in this study area are generally the major groups belonging to the nine main Moroccan families. These three families, as well as the Asteraceae, Lamiaceae, and Fabaceae, represent the most important floristic groups with medicinal plants in Rabat [115]. Geographically, the flora of Morocco is typically Mediterranean with 1441 endemic species and subspecies taxonomically distributed in 7 main families (Asteraceae, Fabaceae, Lamiaceae, Brassicaceae, Caryophyllaceae, Poaceae, Apiaceae, Scrophulariaceae) [49].

However, the climate in Morocco is undergoing a significant change, which is proving to be a hindrance for the development of the forestry and agricultural field. This change is accompanied by an increase in temperature and a decline in annual rainfall, with consequent severe drought. These have direct impacts on water resources, agriculture, human health and economy [116]. In addition to climate change, the overexploitation of natural plant resources has a direct impact on the deterioration of natural areas through the uncontrolled harvesting of plant parts.

In our study, 9 parts of plants are used in phytotherapy including leaves, seeds, fruits, aerial parts, roots, stems, flowers, bark and juice. The survey allowed to classify the plants according to their used parts among which the leaves represent the most used by the practitioners against diabetes and hyperlipidemia with a percentage of 48%, followed by the seeds (34%), the fruits (12%) and the roots (9%) (Figure 2).

While seeds and leaves are found to be preferentially used in the treatment of diabetes, roots and fruits register more uses in recipes treating hypercholesterolemia (Figure 3). This trend has been observed in previous studies reporting the increased use of leaves for treatment of diabetes [44,58,59,74,81], hypercholesterolemia [76], and treatment of diseases with medicinal plants in Morocco [40,89,107,117].

**Table 3.** Medicinals and nutritional plants used to manage metabolic diseases in rabat region.

Family	Scientific name	Vernacular name	Used part	Use value	Metabolic diseases**	Moroccan literature relating to Metabolic diseases plants	Literature documented for the uses of metabolic diseases plants outside Morocco
<b>Aloaceae</b>	<i>Aloe barbadensis</i> Mill.	Aloe vera	L	0.061	D-Hy	Nd	[49–56]
<b>Amaranthaceae</b>	<i>Spinacia oleracea</i> L.	Sabanikh	L	0.018	D	[57,58]	
<b>Anacardiaceae</b>	<i>Pistacia lentiscus</i> L.	Drou	Fr	0.009	D	[41,57–65]	Nd
	<i>Ammi visnaga</i> Lam.	Bachnikha	Fr- Ap	0.018	D	[44,45,58,61,62,64,66,67]	Nd
	<i>Coriandrum sativum</i> L.	Kosber	S-L	0.109	D-Hy	[43,47,58,59,64,68–75]	Nd
	<i>Cuminum cyminum</i> L.	Kamoun	S	0.056	D-Hy	[58,59,63,64,74,76]	Nd
	<i>Carum carvi</i> L.	Carwiya	S	0.028	D	[45,57,59,60,64,67,68,71–74,77–80]	Nd
	<i>Foeniculum vulgare</i> Mill.	Nafaa	S	0.028	D	[44,45,57–60,64,67,68,70–81]	Nd
<b>Apiaceae</b>	<i>Petroselinum crispum</i> Mill.	Maadnouss	L- Ap	0.028	Hy	[58,59,70,72,74,78,80]	Nd
	<i>Apium graveolens</i> L.	Krafess	R	0.028	Hy	[58,59,70,72,78]	Nd
<b>Araliaceae</b>	<i>Panax ginseng</i> C.A.Mey.	Ginseng	R	0.033	D	Nd	[82]
	<i>Artemisia herba-alba</i> Asso.	Chih	Ap	0.123	D-Hy	[44–46,48,58,59,61–63,65,67–69,71–79,81,83–89]	[51,56,57,90,91]
	<i>Artemisia absinthium</i> L.	Chiba	L	0.018	D	[45,46,59–62,64,65,67,68,71–73,77–79,81,83,85]	[55,57,90,92,93]
	<i>Arctium lappa</i> L	Badiana	S	0.028	D		[57]
	<i>Anacyclus pyrethrum</i> L.	Babounj	Fl	0.004	Hy	[59,63,65]	[57]
	<i>Dittrichia viscosa</i> L.	Tarahla	L	0.018	D	[59,63,67,74,78,79,94]	[57]
	<i>Stevia rebaudiana</i> Bertoni.	Stevia	L	0.037	D	Nd	Nd
<b>Asteraceae</b>	<i>Silybum marianum</i> L.	Chouka hmar	L - R	0.037	D-Hy	[72]	[95]
	<i>Cynara cardunculus</i> L.	Qoq	L	0.061	Hy	[59,61,65,67,71,72,75–78,83]	[91]
	<i>Taraxacum officinale</i> L.	Pissenlit	L	0.014	Hy	[65,74]	[55,57]
	<i>Helianthus annuus</i> L.	Nawart chams	S	0.023	Hy	[61,65,96]	[57,93]
	<i>Centaurea maroccana</i> Ball.	Tafgha	St	0.028	D	[59]	
	<i>Scolymus hispanicus</i> L.	Guernina	L - R	0.037	D	[59,63,67,78]	[97]
<b>Berberidaceae</b>	<i>Berberis vulgaris</i> L.	Aghris	Fr - B	0.004	D	[59,67,72,78]	[51,57,92,93]
	<i>Raphanus sativus</i> L.	Fjel	R	0.075	D-Hy	[45,58,59,65,67,71,73,78,81]	[55, 57]
	<i>Brassica oleracea</i> L.	Cheflour	L	0.042	Hy	[45,59,63,67,78,96]	[52,57]
<b>Brassicaceae</b>	<i>Eruca vesicaria</i> L.	Jarjir	L	0.004	D	[59,67,78]	Nd
	<i>Lepidium sativum</i> L.	Heb rchad	S	0.042	D	[44–46,48,49,59,60,65,67,72–74,77–79,81,83,98]	[56,57,91]



Cactaceae	<i>Brassica rapa</i> L.	Laft	R	0.004	D	[59,63,65,67,78,87]	Nd
	<i>Opuntia ficus indica</i> Mill.	Handiya	St	0.047	D	[45,58–60,63,65,71,73,74,77,81,85,98,99]	[53,57,91,100]
Capparaceae	<i>Capparis spinosa</i> L.	Kebbar	Fr	0.004	D	[45,46,59–63,65–68,77–79,81,83–85,101]	[51,56,57,94]
Cistaceae	<i>Cistus salvifolius</i> L.	Irgel	Fl	0.009	D	[58–60,65,67]	Nd
Curcubitaceae	<i>Cucurbita pepo</i> L.	Geraa	S	0.033	D-Hy	[59,62,63,96,102]	[55,91]
		khedra					
Cupressaceae	<i>Juniperus phoenicea</i> L.	Araar	L	0.037	D-Hy	[45,46,48,59,63,65,67,71,72,77,78,81,83,86,88,101,102]	[51,56,57]
Euphorbiaceae	<i>Euphorbia officinarum</i> L.	Daghmouss	St-L	0.090	D	[58,63,65,67,72,74,75,78–80,84,85,96,98]	[55,56]
Ericaceae	<i>Vaccinium myrtillus</i> L.	Oleik	Fr	0.004	D	Nd	[103]
	<i>Arbutus unedo</i> L.	Bakhnou	L- Fr	0.014	D	[45,62,63,65,67,68,74,78,79,81]	Nd
	<i>Trigonella foenum-graecum</i> L.	Halba	S	0.374	D-Hy	[42,45,46,48,58–63,65,67,68,70–81,83–86,88,94,98,99]	[51,55,56,90,92,95,104,105]
	<i>Lupinus albus</i> L.	Tirms / Foul gnawa	S	0.099	D	[45,46,48,59,61,65,67,73,74,78,79,83,85,86,102]	[56,57]
	<i>Glycine max</i> (L.) Merr.	Soja	S	0.033	D-Hy	[45,58,65,67,71,73,74,77,78,80,81]	Nd
	<i>Glycyrrhiza glabra</i> L.	Aarq souss	R	0.009	D-Hy	[44,59,60,65,67,76,78]	[56,93]
Fabaceae	<i>Cerantonia siliqua</i> L.	Kharoub	S – L	0.014	D	[44,58–60,65,67,71,73–76,78,81]	[91,106]
	<i>Senna alexandrina</i> Mill.	Senameki	L	0.004	D	Nd	Nd
	<i>Cicer arietanum</i> L.	Homes	S	0.004	D	[59,67,73,78,81]	[105]
	<i>Medicago sativa</i> L.	Fessa	L	0.004	D	[59,63,67,71,75,78,81,88]	Nd
Gentianaceae	<i>Arachis hypogaea</i> L.	Kawkaw	S	0.004	D	[59]	Nd
	<i>Centaurium erythraea</i> Rafn.	Guesset haya	Fl	0.028	D	[45,59,61–65,67,72,74,78,79,94,107]	[57,91,97]
Juglandaceae	<i>Juglans reiga</i> L.	Gargaa	S	0.014	Hy	[45,48,59,60,63,65,67,68,70,72,76,78,86]	[93,95,97,103,108,109]
	<i>Salvia officinalis</i> L.	Salmiya	L – Ap	0.194	D-Hy	[44,46,48,58–60,62,63,65,67–69,71–81,83–86,89,94,96,98,99]	[56,57,91]
	<i>Salvia hispanica</i> L.	Chia	S	0.014	D	Nd	Nd
	<i>Origanum compactum</i> Benth.	Zaatar	L – Ap	0.085	D-Hy	[42,44–46,48,59,61,63–65,67–69,72,73,78,79]	[57]
	<i>Thymus vulgaris</i> L.	Zaitra	L – Ap	0.028	D	[44,46,48,59,63,67,71–73,76–79,84,98]	[56,90]
	<i>Marrubium vulgare</i> L.	Merriwt	L – Ap	0.052	D	[42,44–46,48,58–60,62,63,65,67–75,77–79,81,83–86,89,94,96]	[51,56,57,91]
Lamiaceae	<i>Rosmarinus officinalis</i> L.	Azir	L – Ap	0.085	D-Hy	[44–46,59,60,62–65,67,68,71–74,76–79,83,89,96,98,102]	[56,57,90,91,97]
	<i>Ocimum basilicum</i> L.	Lahbeq	L	0.009	D	[59,63,72]	[53,57,97]
	<i>Mentha officinalis</i> L.	Naanaa	L – Ap	0.018	D-Hy	[46]	Nd
	<i>Lavandula angustifolia</i> L.	Khezama	L	0.018	D	[45,58,59,64,65,71,72,77,78,89,98]	[57]
	<i>Origanum majorana</i> L.	Merdedouch	L	0.014	D	[59,72,75,77,79]	[57]

	<i>Cinnamomum cassia</i> Blume	Qarfa	B	0.175	D-Hy	[46,59,61,63,65,67,73,76,78,83,94]	[56,57,91,95]
<b>Lauraceae</b>	<i>Persea americana</i> Mill.	Avocat	Fr-L	0.009	D	[59,60,65,72,83,85]	[52,54,100,110]
	<i>Laurus nobilis</i> L.	Rand	L	0.037	D-Hy	[59,60,67,78]	[57]
	<i>Nerium oleander</i> L.	Defla	L	0.004	D	[45,46,48,59,61-64,67-69,71,74,75,77-81,83,87,99]	[51,57]
	<i>Allium cepa</i> L.	Basla	R	0.004	D	[44-46,48,59,60,62-65,67,71-73,76,77,81,83,85,98]	[50,52,54-57,95,104,105,111]
<b>Liliaceae</b>	<i>Allium sativum</i> L.	Toma	R	0.085	D-Hy	[45,46,58-60,63,65,67,68,71,72,75,76,78,79,81,83,89,94,98]	[50,53-56,91,95,97,104,105,111,112]
<b>Linaceae</b>	<i>Linum usitatissimum</i> L.	Zariat ketan	S	0.322	D-Hy	[44,46,59,60,63,65,67,76,78,79,96,98,99,102]	Nd
<b>Lithraceae</b>	<i>Punica granatum</i> L.	Roman	Fr-L	0.004	D	[46,58-61,63-65,67,72,73,77,78,80]	[50,56,57,91,97,100]
<b>Malvaceae</b>	<i>Hibiscus sabdariffa</i> L.	Karkadil	L	0.009	D	[59,67,78]	Nd
<b>Myrtaceae</b>	<i>Eucalyptus globulus</i> L.	Eucalyptus	L	0.104	D	[45,46,59,62-65,71,73,74,77,80,81]	[56,57,78,111]
	<i>Eugenia caryophyllata</i>	Qronfel	L-Fl	0.004	D	[44,45,62,65,67,74,78,81,83]	Nd
<b>Moraceae</b>	<i>Morus alba</i> L.	Toute	L	0.009	D	[59,63,65,83,96,102]	[50,54,55,97]
	<i>Ficus carica</i> L.	Kermouss	L	0.014	D-Hy	[45,58-60,63,65,67,68,73,74,77,78,80,81,96,98,102]	[55,57,91,100]
<b>Moringaceae</b>	<i>Moringa oleifera</i> Lam.	Moringa	L	0.014	D	Nd	[54,55,100,104]
<b>Oleaceae</b>	<i>Olea europea</i> L.	Zaytoun	L	0.521	D-Hy	[44,46,48,58-62,64,65,67,68,70-81,83,85-87,89,98,99,113]	[53,55-57,90,91,97,106]
<b>Poaceae</b>	<i>Avena sativa</i> L.	Choufan	S	0.009	D	[59,63,76,78,79]	[95]
	<i>Triticum durum</i> Desf.	Zraa	S	0.004	D	[59,61,65,67,78]	Nd
	<i>Hordeum vulgare</i> L.	Cheir	S	0.037	D	[59-61,64,65,67,71,78,86,98]	[56,105]
	<i>Lolium perenne</i> L.	Zouane	S	0.023	D	[46,59,67,71,72,77,78,81,85]	Nd
	<i>Pennisetum glaucum</i> L.	Ilane / Bachna	S	0.104	D-Hy	[59,71,78]	Nd
	<i>Zea mays</i> L.	Dera	S	0.004	Hy	[59,60,62,65,76,81]	[54,55,104]
<b>Plantaginaceae</b>	<i>Stipa tenacissima</i> L.	Halfa	Ap	0.004	Hy	[61]	Nd
<b>Portulacaceae</b>	<i>Plantago ovata</i> Forsk.	Katona	S	0.004	D	[81]	Nd
<b>Ranunculaceae</b>	<i>Portulaca oleracea</i> L.	Rejla	L	0.075	Hy	[59,65,67,71,78,101]	[93,97,108]
	<i>Nigella sativa</i> L.	Sanouj	S	0.009	D	[44-46,48,59,60,62,63,65,67,68,71-75,77-79,81,83-85,89,98]	[55-57,90,91,109]
<b>Rhamnaceae</b>	<i>Ziziphus lotus</i> Lam.	Nbeg	S-L	0.014	D	[58-60,65,67,68,73-75,77-81,83,86,98]	[55-57]
	<i>Prunus cerasus</i> L.	Heb mlouk	S	0.009	D	Nd	[93,109]
	<i>Prunus amygdalus</i> Stokes	Louza	S	0.028	D	[58-62,65,67,68,71-74,77,78,81,86,98]	[55,57]
<b>Rosaceae</b>	<i>Rosa damascena</i> Mill.	Lward beldi	Fl	0.009	Hy	Nd	[55]
	<i>Coffea arabica</i> L.	Qahwa	S	0.023	D	[76]	[91]



<b>Rutaceae</b>	<i>Citrus limon</i> Burm.	Hamed	Fr	0.014	D-Hy	[59,60,76]	[52,55,94]
	<i>Citrus maxima</i> Burm.	Ranj	Fr	0.018	Hy	[59]	Nd
<b>Salicaceae</b>	<i>Salix alba</i> L.	Safsaf	L	0.004	D	[59,65,74,83,87]	[103]
<b>Sapotaceae</b>	<i>Argania spinosa</i> L.	Argane	S	0.009	D	[44,58–60,63,65,71,72,75,80]	Nd
<b>Solanaceae</b>	<i>Solanum melongena</i> L.	Danjel	Fr	0.227	Hy	[76]	Nd
<b>Theaceae</b>	<i>Camellia sinensis</i> (L.) Kuntze	Atay	L	0.028	D-Hy	[58,59,65,67,71,72,81]	[57,91]
<b>Tiliaceae</b>	<i>Tilia europea</i> L.	Zayzafoun	L	0.014	Hy	Nd	Nd
<b>Urticaceae</b>	<i>Urtica dioica</i> L.	Hariga	St – L	0.014	D	[45,58,59,61–63,65,67,68,74–77,79,81,101]	[57,91,93,95,109,114]
	<i>Zingiber officinale</i>	Skinjbir	R	0.099	D-Hy	[44,59,60,62,67,72,73,75,76,78,79,83,99]	[54,56,57,91,104]
<b>Zingiberaceae</b>	Roscoe						
	<i>Curcuma longa</i> L.	Kharqom	R	0.066	D-Hy	[44,75,76]	[54,104]

\* Used part; L: leaves, S: seeds, Fr: fruits, Ap: aerial parts, R: roots, St: stems, Fl: flowers, B: bark.

\*\* Metabolic Diseases; D: Diabetes, Hy: Hypercholesterolemia




Figure 1. Number of citation of families.




Figure 2. Used part of the listed plants.




Figure 3. Used part of plants according to the use categories.

#### 2.2.1. Floristic diversity according to use category

As previously mentioned, this survey allowed us to identify 98 plant species suggested for healing the two diseases (diabetes and hypercholesterolemia), of which 26 plants are listed to treat both diseases at the same time and 72 species are cited in the two categories of diseases divided between diabetes (57) and hypercholesterolemia (15) (Figure 4).




Figure 4. The plants inventoried in each category of use.


For diabetes, 83 plants were recorded, of which the most cited were: *O. europaea*, *T. foenum-graecum*, *C. cassia*, *A. herba-alba*, *S. officinalis*, *L. albus*, *C. sativum*, *E. officinarum*, *P. glaucum*, and *R. officinalis*. The bibliographic reports show that most of the floristic species have already been cited in other surveys [46,59,77,89].

Regarding hyperlipidemia, 41 floristic species were recorded in this survey with the dominance of *S. melongena*, *A. sativum*, *L. usitatissimum*, *T. foenum-graecum*, *C. cassia*, *A. herba-alba*, *S. officinalis*, *C. sativum*, *P. glaucum*, and *R. officinalis*. Most of these species have been recorded in other surveys [76].

### 2.2.2. Floristic diversity according to respondent's category

The increased use of these plants reflects from one hand a better efficacy of medicinal plants and from the other hand a remarkable food safety in diabetic and hypercholesterolemic patients, in the face of the lack of control of glucose and cholesterol levels and of information concerning the consequences from a long-term use.

Thus, the results obtained were analyzed according to the RFC and divided corresponding to the citation of herbalists (Figure 5) or local informants of the region (Figure 6).




**Figure 5.** The relative frequency of citation of plants cited by herbalists.

Generally, the values of relative frequency of citation show a significant difference ranging from 0.004 to 0.524. However, the olive tree has the highest value, followed by fenugreek with a value of 0.374, then flaxseed with a value of 0.322 in relation to the total number of species recorded.

In total 36 medicinal species were inventoried by the selected herbalists of the region, of which the majority of plants (31) are anti-diabetic and 10 species were listed as anti-hypercholesterolemic. However, the values of the anti-diabetic plants inventoried by the herbalists ranged from 0.05 to 0.7. Among the 31 documented species, only 2 have a significant RFC, i.e., *T. foenum-graecum* (0.7) and *O. europaea* (0.5), followed by *S. hispanicus* and *L. usitatissimum* which occupy the same rank with a value of 0.4. For cholesterol, the first rank is occupied by *L. usitatissimum* (0.5), followed by *S. melongena* (0.3) and *O. europaea* (0.2). *Linum usitatissimum* keeps a strong interest for both diseases being cited in recipes by the totality of herbalists (Figure 5).


Otherwise, the information collected by the citizens of the region revealed 77 and 38 species dedicated to cure diabetes and cholesterol respectively. The RFC values of diabetes and hypercholesterolemia ranged from 0.008 to 0.661 and 0.014 to 0.626, respectively (Figure 6). Out of a total of 77 medicinal species used against diabetes, only 10 were widely cited in this survey such as *E. officinarum* and *L. albus* which scored an RFC of 0.120, similarly *C. sativum* and *A. herba-alba* (0.129), *L. usitatissimum* (0.161), *E. globulus* (0.177), *C. cassia* (0.201), *S. officinalis* (0.241), *T. foenum-graecum* (0.438), and finally *O. europaea* which was cited by a large number of users and recorded an RFC of 0.661. While for hypercholesterolemia, the relative frequency of citation of the different plants ranged from 0.029 to 0.626. On the other hand, 14 plants scored an RFC > 0.1, including *L. nobilis* and *C. pepo* (0.104), *S. marianum*, *B. oleracea*, and *O. compactum* (0.119), *C. cardunculus*

(0.134), *P. glaucum* (0.149), *C. domestica* (0.179), *O. europaea* (0.208), *P. oleracea* (0.223), *A. sativum* and *Z. officinale* (0.238), *L. usitatissimum* (0.447), and *S. melongena* which has the high-est RFC value (0.626).



**Figure 6.** The relative frequency of citation of plants reported by interviewees.

Overall, the results obtained by statistical analysis show the predominance of *O. europaea* and *L. usitatissimum* as MP used in the management of metabolic diseases in the region of Rabat, and this was also demonstrated by the calculation of the family use value (FUV) (Figure 7), This study valued the two floral families Oleaceae (FUV=0.52) and Linaceae (FUV=0.32) presented by their representatives cited above.



**Figure 7.** Diversity familial according FUV.

*Olea europaea*, the only representative of its floristic family and the most cited by the participants in both pathologies, is a species endemic to Morocco, and includes two cultivars: *Olea europaea* L. subsp. *europaea* var. *europaea*, the cultivated variety, and *O. europaea* subsp. *europaea* var. *sylvestris* (Mill.) Lehr., the wild or spontaneous variety [118]. In Morocco, this species is more widespread in the wild state than in the cultivated state of which it occupies an area of 1,070,000 ha in cultivated areas. The surveys established previously in Morocco have all marked the increased use of the leaves of the olive tree as an antidiabetic agent, and some have cited the leaves of this plant as anticholesterolemic under decoction, infusion or


powder [44,46,48,58–62,64,65,67,68,70–81,83,85–87,89,98,99,113]; similar uses are recorded outside Morocco, namely Algeria [56,57,90,91], Kenya [53], Pakistan [55] and Turkey [97,106]. The antidiabetic activity has also been experimentally proven by *in vivo* or *in vitro* studies [119–124].

*Linum usitatissimum* is an herbaceous plant belonging to the Linaceae family, known in Morocco as "Zariat-kettan". Due to its nutritional and phytochemical value (high fiber content, protein, phytoestrogen, flavonoids, linolenic acid), this plant is endowed with therapeutic effects [125,126]. Flaxseed has been widely used in traditional medicine to reduce inflammation and fever [88], to treat lung diseases, severe colds and coughs, as a muscle relaxant, to cure abdominal pain, as an antidiabetic and anticholesterolemic [127]. The results of our study were in agreement with those previously carried out in Morocco [44,46,59,60,63,65,67,76,78,79,96,98,99,102]. The *in vitro* activity of the polyphenolic fraction of *L. usitatissimum* showed a potent anti-diabetic activity when compared with acarbose. Its enzymatic inhibition of  $\alpha$ -amylase is higher than that of  $\alpha$ -glucosidase [126].

### 2.3. Ethnopharmacological profile

#### 2.3.1. Preparation and administration method

In the totality of the answers, the oral administration of the medicinal plants proved to be more effective for the treatment of diabetes and hypercholesterolemia. Data analysis shows that the majority of the recipes are developed by simple grinding of medicinal plants for both pathologies, followed by decoction with a percentage of 25 and 20%, respectively (Figure. 8). A new preparation method added to the list is the use of cooked plants for the treatment of hypercholesterolemia. Otherwise, the participants use mainly the infusion, the mixture of plants with honey, milk or vegetable oil (very often olive oil), and herbal tea in the recipes prescribed for hypercholesterolemia. However, these results prove the increased use of the different modes of preparations cited in this survey, but the majority of the questionnaires conducted in Morocco report that the decoction is the most used method.



**Figure 8.** Preparation mode of herbal recipes.

#### 2.3.2. Formulation of recipes

The information collected during this survey shows that all the prescribed recipes are administered orally, as well as the informants are found having an important background concerning the treatment of diabetes with medicinal plants, even if they are not affected by the disease, and this is perhaps due to the importance that this disease occupies in the Moroccan population (Table.4). Furthermore, diabetes has been estimated to affect more than 15,000 children in Morocco, with more than 2 million people in the 18-25 age group suffering from this disease without any knowledge of its mitigation.



**Table 4.** Traditional recipes of both diseases.

N°	Prescribed recipes
1	A mixture of barks of <i>Centaurea maroccana</i> , seeds of <i>Trigonella foenum-graecum</i> and <i>Ziziphus lotus</i> , flower bud of <i>Glycyrrhiza glabra</i> with equal portions and flavored by the leaves of <i>Origanum compactum</i> are ground into a fine powder and administered orally one teaspoon during each morning.
2	<i>Solanum melongena</i> fruits cut into slices and macerate overnight in warm water. Drinking the solution every morning is beneficial for lowering bad cholesterol.
3	Infusing half a teaspoon of <i>Curcuma longa</i> in a cup of boiling water and drinking one cup every day is beneficial for diabetes.
4	A mixture of <i>Juniperus phoenicea</i> leaves with yogurt is beneficial for diabetes
5	Powder of <i>Coffea arabica</i> flavored with the leaves of <i>Thymus vulgaris</i> administered one teaspoon before each meal.
6	A mixture of fine powder of the seeds of <i>Trigonella foenum-graecum</i> , <i>Coriandrum sativum</i> , <i>Carum carvi</i> , and <i>Foeniculum vulgare</i> with equal proportions is administered in powder form every morning.
7	A mixture of <i>Trigonella foenum-graecum</i> seed, <i>Cinnamomum cassia</i> Blume bark, <i>Zingiber officinale</i> rhizome, in powder form is macerated for one night. The macerate to which is added a few drops of <i>Citrus limon</i> is administered orally before meals.
8	A mixture of <i>Cinnamomum cassia</i> Blume bark and <i>Pennisetum glaucum</i> seeds was ground and allowed to macerate in a cup of lukewarm water for one night. The mixture was administered daily before meals and one time per day.
9	A mixture of the leaves of <i>Salvia officinalis</i> , <i>Olea europaea</i> , <i>Origanum compactum</i> was macerated and administered orally one time daily, before meals.

*Trigonella foenum-graecum* is an annual plant belonging to the Fabaceae family, known as "Halba". The seeds are widely used in Moroccan cuisine as a spice or food additive and even in traditional medicine for its carminative, expectorant, laxative effects and for treatment of digestive and mucosal ailments [128]. Various ethnobotanical and ethnopharmacological studies have identified this plant as an antidiabetic agent in [42,45,46,48,58–63,65,67,68,70–81,83–86,88,94,98,99], as well as outside Morocco [51,55,56,90,92,95,104,105]. Indeed, Inbaraj and Muniappan [129] were able to determine the correlation between *in vitro* and *in vivo* antidiabetic activity of fenugreek, the *in vitro*  $\alpha$ -glucosidase enzyme inhibition activity showed a percentage inhibition of 68% at a concentration of 100  $\mu\text{g/mL}$  near that of acarbose (94%). This result was also demonstrated *in vivo* in albino rats by the oral glucose tolerance test, which marked a significant reduction in blood glucose level after 15 min of administration.

In addition, Marmouzi et al. [130] tested the *in vitro* antidiabetic activity of the aqueous extracts of the leaves and fruits of *Z. lotus* which marked an inhibitory activity against  $\alpha$ -amylase and  $\alpha$ -glucosidase higher than that of conventional drugs. This shrub of the Rhamnaceae family growing in arid and semi-arid, sub-Saharan and sub-humid regions of Morocco [131], is highly cited by the Moroccan population and herbalists as a medicinal plant used in the treatment of metabolic diseases such as diabetes, hypercholesterolemia and cardiovascular diseases [58–60,65,67,68,73–75,77–81,83,86,98], and in ethnobotanical studies carried out outside Morocco [55–57]. Bencheikh et al. [115], studied the effect of an aqueous fruit extract from *Z. lotus* on plasma triglyceride, total cholesterol and HDL levels in mice exposed to a high-fat diet. Indeed, the cholesterol-lowering activity of the aqueous extract was associated with an increase in plasma HDL levels, leading to the elimination of cholesterol in the form of bile acid. Similarly, Pasha et al [132] studied the effect of an ethanolic extract of coriander on lipid parameters in diabetic rats. This study showed that oral administration of extract reduces the level of total cholesterol, triglycerides, LDL, and VLDL, which decreases the risk of cardiovascular diseases [133]. This condiment plant is endowed with several pharmacological activities namely anticancer, anti-inflammatory, neuroprotective, anxiolytic, anticonvulsant, and analgesic as a result of its richness in gallic acid, linalool [124,134–138]. Aqueous coriander extract showed higher inhibition against  $\alpha$ -amylase and  $\alpha$ -glucosidase compared with the standard acarbose [139].

The hexanic and methanolic extracts and essential oil of *J. phoenicea* showed a significant inhibitory activity towards  $\alpha$ -amylase that was comparable to that of acarbose [140]. This plant of the Cupressaceae family growing in the coastal areas of the Mediterranean [141] is well known for its therapeutic effects

against diarrhea, rheumatism, bronchitis, diabetes and has been recently identified as a plant used for the prevention of Covid-19 [142] and for diabetes [45,46,48,59,63,65,67,71,72,77,78,81,83,86,88,101,102]. These surveys also cited *O. compactum* as a medicinal species treating diabetes. This endemic plant having a great history in herbal medicine as an agent treating digestive disorders, antidiabetic, antifungal, anti-inflammatory, antimutagenic and anticancer[143–145].

It is important to underline that most of the recipes prescribed by herbalists are based on a single plant, while those given by local informants are based on a formulation of more than 3 plants, which can give possible toxicity following negative interactions between these plants. Also, Mechchate et al. [146] studied the *in vivo* antidiabetic activity of a mixture of *L. usitatissimum*, *C. sativum* seeds, and the leaves of *O. europaea* var. *sylvestris* which has the effect of reducing the blood glucose level after oral administration of the combination of these three floral species in the same quantity.

### 3. CONCLUSION

This ethnopharmacological survey brings the first step in the branch of phytotherapy and highlights all medicinal plants used for the treatment of diabetes and hypercholesterolemia by the regional population of Rabat. The three planes (socio-demographic, floristic and pharmacological) are interconnected: the use value of each species depends on the informant profile and the category of use. Among 98 species, 57 were listed as anti-diabetic and 15 as anticholesterolemic, with 26 species presenting indications for both pathologies. The relative frequency of use emphasizes *T. foenum-graecum* and *O. europaea* in the diabetic use category, and *S. melongena* and *L. usitatissimum* in the cholesterolemic use category. The leaves represent the most used part along with the dominance of the oral consumption in the form of powder. The patients are well adapted to the use of a mixture of plants. In conclusion, this survey paves the way for phytochemical and pharmacological studies which are necessary to confirm the activity of the highlighted species and toxicological studies regarding their long-term use effects, and possible interaction with other drugs.

### 4. MATERIALS AND METHODS

#### 4.1. Studied region

Rabat - Sale - Kenitra region covers an area of 17,570 km<sup>2</sup>, which represents 2.5% of the total area of the Morocco (RGPH, 2014). It participates in 15.2% of the national wealth. The region of Rabat-Salé-Kénitra extends over two large watersheds, namely the hydraulic basin of Bouregreg and Chaouia and the hydraulic basin of Sebou; they are in turn formed by several sub-watersheds with inputs of rainfall origin. Thus, the region is characterized by a semi-arid Mediterranean weather with a marine or continental oceanic effect (Figure 9).



Figure 9. Study area location: Rabat-Salé-Kénitra.

#### 4.2. Data extraction

The ethnopharmacological survey was carried out from December 2021 to May 2022. During this phase, various locations in the cities of the region were visited. The survey was conducted by using two

methods: one online using the platform "Google Forms" and social networks "Facebook" and "WhatsApp", and the other in the field by conducting interviews with different traditional healers with experience in the management of metabolic diseases types diabetes and hyperlipidemia.

Data extraction was conducted based on informants following the written and online interview form. During the interview, 241 informants participated in the diabetes survey, 214 in the cholesterol survey. As well as, 20 traditional herbalists were randomly selected in the region of Rabat and were invited to answer questionnaire fixed on different axes allowing to collect the maximum of information. This included a socio-demographic section, a floristic section which summarizes the used floristic species, part of the plant, and a pharmacological section detailing the method of plant extraction, the mode of administration, the duration of the treatment and the related illness.

### 4.3. Statistical analysis

The results obtained from the interview form were interpreted using a quantifiable approach combining the use value of each species (UV), the relative frequency of citation (RFC), and the familial use value (FUV). The two parameters UV and RFC were determined according to the method described by [147] and calculation of these parameters was done using the following formulas (equations (1) and (2)).

$$UV = \sum \frac{U}{N} \quad (1)$$

Where  $U$  is the number of referees who mentioned the use of the species, and  $N$  is the number of referees (Totality).

$$RFC = \frac{Fc}{N} \quad (2)$$

Where  $Fc$  is the citation frequency of the species, and  $N$  is the number of referees (Totality).

In addition, FUV was defined according to [41] and calculated according to equation (3):

$$FUV = \frac{UVf}{Nf} \quad (3)$$

Where  $UVf$  is the use value of the families, and  $Nf$  represent the number of species in each family.

**Author contributions:** Concept – A.B., H.H.; Design – A.B., H.H., C.E.K.; Supervision – A.B., H.H., M.T.; Resources – C.E.K.; Materials – H.H., M.T.; Data Collection and/or Processing – C.E.K.; Analysis and/or Interpretation – C.E.K., O.B.; Literature Search – C.E.K.; Writing – C.E.K.; Critical Reviews – H.H., V.K., I.U.R., L.C.M., P.L.L., A.B.

**Conflict of interest statement:** The authors declared no conflict of interest.

### REFERENCES

- [1] Sangouni AA, Taghdir M, Mirahmadi J, Sepandi M, Parastouei K. Effects of curcumin and/or coenzyme Q10 supplementation on metabolic control in subjects with metabolic syndrome: a randomized clinical trial. *Nutr J* 2022;21:62. <https://doi.org/10.1186/s12937-022-00816-7>.
- [2] Hu W, Lazar MA. Modelling metabolic diseases and drug response using stem cells and organoids. *Nat Rev Endocrinol* 2022;18:744–59. <https://doi.org/10.1038/s41574-022-00733-z>.
- [3] Kutbi EH, Sohoul M, Fatahi S, Lari A, Shidfar F, Aljhdali MM, Alhoshan FM, Elahi SS, Almusa HA, Abu-zaid A. The beneficial effects of cinnamon among patients with metabolic diseases: A systematic review and dose-response meta-analysis of randomized-controlled trials. *Crit Rev Food Sci Nutr* 2022;62:6113–31. <https://doi.org/10.1080/10408398.2021.1896473>.
- [4] Afshin A, Sur PJ, Fay KA, Cornaby L, Ferrara G, Salama JS, Mullani EC, Abate KH, Abbafati C, Abebe Z, Afarideh M, Aggarwal A, Agrawal S, Akinyemiju T, Alahdab F, Bacha U, Bachman VF, Badali H, Badawi A, Bensenor IM, Bernabe E, Biadgilign SKK, Biryukov SH, Cahill LE, Carrero JJ, Cercy KM, Dandona L, Dandona R, Dang AK, Degefa MG, El Sayed Zaki M, Esteghamati A, Esteghamati S, Fanzo J, Farinha CS, Farvid MS, Farzadfar F, Feigin VL, Fernandes JC, Flor LS, Foigt NA, Forouzanfar MH, Ganji M, Geleijnse JM, Gillum RF, Goulart AC, Grosso G, Guessous I, Hamidi S, Hankey GJ, Harikrishana S, Hassen HY, Hay SI, Hoang SL, Horini M, Ikeda N, Islami F, Jackson MD, James SL, Johansson L, Jonas JB, Kasaian A, Khader YS, Khalil IA, Khang YH, Kimokoti RW, Kokubo Y, Kumar GA, Lallukka T, Lopez AD, Lorkowski S, Lotufo PA, Lozano R, Malekzadeh R, Marz W, Meier T, Melaku YA, Mendoza W, Mensink GBM, Micha R, Miller TR, Mirarefin M, Mohan V, Mokdad AH, Mozaffarian D, Nagel G, Naghavi M, Nguyen CT, Nixon MR, Ong KL, Pereira DM, Poustchi H, Qorbani M, Rai RK, Razo-Garcia C, Rehm CD, Rivera JA, Rodríguez-Ramírez S, Roshandel G, Roth GA, Sanabria J, Sánchez-Pimienta TG, Sartorius B, Schmidhuber J, Schutte AE, Sepanlou SG, Shin MJ, Sorensen RJD, Springmann M, Szponar L, Thorne-Lyman AL, Thrift AG, Touvier M, Tran BX, Tyrovolas S, Ukwaja KN, Ullah I, Uthman OA, Vaezghasemi M, Vasankari TJ, Vollset SE, Vos T, Vu GT, Vu LG, Weiderpass E, Werdecker A, Wijeratne T, Willett WC, Wu JH, Xu G, Yonemoto N, Yu C, Murray CJL. Health effects of dietary risks in 195 countries, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *The Lancet* 2019;393:1958–72. [https://doi.org/10.1016/S0140-6736\(19\)30041-8](https://doi.org/10.1016/S0140-6736(19)30041-8).

- [5] Li D, Li Y, Yang S, Lu J, Jin X, Wu M. Diet-gut microbiota-epigenetics in metabolic diseases: From mechanisms to therapeutics. *Biomed Pharmacother* 2022;153:113290. <https://doi.org/10.1016/j.biopha.2022.113290>.
- [6] Fu J, Zhang L-L, Li W, Zhang Y, Zhang Y, Liu F, Zou L. Application of metabolomics for revealing the interventional effects of functional foods on metabolic diseases. *Food Chem* 2022;367:130697. <https://doi.org/10.1016/j.foodchem.2021.130697>.
- [7] Lusis AJ, Attie AD, Reue K. Metabolic syndrome: from epidemiology to systems biology. *Nat Rev Genet* 2008;9:819–830. <https://doi.org/10.1038/nrg2468>.
- [8] Liu Q, Chen L, Hu L, Guo Y, Shen X. Small molecules from natural sources, targeting signaling pathways in diabetes. *Biochim Biophys Acta BBA - Gene Regul Mech* 2010;1799:854–865. <https://doi.org/10.1016/j.bbagr.2010.06.004>.
- [9] Hung H-Y, Qian K, Morris-Natschke SL, Hsu C-S, Lee K-H. Recent discovery of plant-derived anti-diabetic natural products. *Nat Prod Rep* 2012;29:580–606. <https://doi.org/10.1039/C2NP00074A>.
- [10] Emran TB, Dutta M, Uddin MMN, Nath AK, Uddin MZ. Antidiabetic potential of the leaf extract of *Centella asiatica* in alloxan induced diabetic rats. *Jahangirnagar Univ J Biol Sci* 2015;4:51–59. <https://doi.org/10.3329/jujbs.v4i1.27785>.
- [11] American Diabetes Association. Introduction: Standards of Medical Care in Diabetes—2022. *Diabetes Care* 2021;45:S1–2. <https://doi.org/10.2337/dc22-Sint>.
- [12] Rahman MM, Islam MR, Shohag S, Hossain ME, Rahaman MS, Islam F, Ahmed M, Mitra S, Khandaker MU, Idris AM, Chidambaram K, Bin Emran T, Cavalu S. The multifunctional role of herbal products in the management of diabetes and obesity: A Comprehensive review. *Molecules*. 2022;27:1713. <https://doi.org/10.3390/molecules27051713>.
- [13] Quazi A, Patwekar M, Patwekar F, Alghamdi S, Rajab BS, Babalghith AO, Islam F. *In Vitro* Alpha-amylase enzyme assay of hydroalcoholic polyherbal extract: Proof of concept for the development of polyherbal teabag formulation for the treatment of diabetes. *Evid Based Complement Alternat Med*. 2022;2022:e1577957. <https://doi.org/10.1155/2022/1577957>.
- [14] Xu L, Li Y, Dai Y, Peng J. Natural products for the treatment of type 2 diabetes mellitus: Pharmacology and mechanisms. *Pharmacol Res*. 2018;130:451–465. <https://doi.org/10.1016/j.phrs.2018.01.015>.
- [15] Bondar AC, Popa AR. Diabetic neuropathy prevalence and its associated risk factors in two representative groups of Type 1 and Type 2 Diabetes Mellitus patients from Bihor County. *Mædica* 2018;13:229–234. <https://doi.org/10.26574/maedica.2018.13.3.229>.
- [16] Rother KI. Diabetes Treatment – Bridging the Divide. *N Engl J Med*. 2007;356:1499–501. <https://doi.org/10.1056/NEJMp078030>.
- [17] American Diabetes Association. 8. Pharmacologic Approaches to Glycemic Treatment. *Diabetes Care* 2016;40:S64–74. <https://doi.org/10.2337/dc17-S011>.
- [18] López Stewart G. Diagnostic criteria and classification of hyperglycaemia first detected in pregnancy: A World Health Organization Guideline. *Diabetes Res Clin Pract*. 2014;103:341–363. <https://doi.org/10.1016/j.diabres.2013.10.012>.
- [19] Lowe LP, Metzger BE, Dyer AR, Lowe J, McCance DR, Lappin TRJ, Trimble ER, Coustan DR, Hadden DR, Hod M, Oats JN, Persson B. Hyperglycemia and adverse pregnancy outcome (HAPO) study: Associations of maternal A1C and glucose with pregnancy outcomes. *Diabetes Care*. 2012;35:574–580. <https://doi.org/10.2337/dc11-1687>.
- [20] Song C, Lyu Y, Li C, Liu P, Li J, Ma RC, Yang X. Long-term risk of diabetes in women at varying durations after gestational diabetes: A systematic review and meta-analysis with more than 2 million women. *Obes Rev*. 2018;19:421–429. <https://doi.org/10.1111/obr.12645>.
- [21] Li N, Yang Y, Cui D, Li C, Ma RCW, Li J, Yang X. Effects of lifestyle intervention on long-term risk of diabetes in women with prior gestational diabetes: A systematic review and meta-analysis of randomized controlled trials. *Obes Rev*. 2021;22:e13122. <https://doi.org/10.1111/obr.13122>.
- [22] Tam WH, Ma RCW, Ozaki R, Li AM, Chan MHM, Yuen LY, Lao TTH, Yang X, Ho CS, Tutino GE, Chan JCN. In utero exposure to maternal hyperglycemia increases childhood cardiometabolic risk in offspring. *Diabetes Care* 2017;40:679–686. <https://doi.org/10.2337/dc16-2397>.
- [23] Defesche JC, Gidding SS, Harada-Shiba M, Hegele RA, Santos RD, Wierzbicki AS. Familial hypercholesterolaemia. *Nat Rev Dis Primer*. 2017;3:1–20. <https://doi.org/10.1038/nrdp.2017.93>.
- [24] Pappan N, Rehman A. Dyslipidemia. StatPearls, Treasure Island (FL): StatPearls Publishing; 2022.
- [25] Chetoui A, Kaoutar K, Kardoudi A, Boutahar K, Chigr F. Epidemiology of diabetes in Morocco: Review of data, analysis and perspectives. *Int J Sci Eng Res*. 2018;9:1310–1316.
- [26] El Alami H, Haddou I, Benaadi G, Lkhider M, El Habchi D, Wakrim L, Nabih N, Abidi O, Khlil N, Maaroufi A, Naamane A, Hamdi S. Prevalence and risk factors of chronic complications among patients with type 2 diabetes mellitus in Morocco: A cross-sectional study. *Pan Afr Med J*. 2022;41:182. <https://doi.org/10.11604/pamj.2022.41.182.25532>.
- [27] Mohamad NE, Abu N, Yeap SK, Lim KL, Romli MF, Sharifuddin SA, Long K, Alitheen NB. Apoptosis and metastasis inhibitory potential of pineapple vinegar against mouse mammary gland cells in vitro and in vivo. *Nutr Metab*. 2019;16:49. <https://doi.org/10.1186/s12986-019-0380-5>.
- [28] Granado-Lorencio F, Hernández-Alvarez E. Functional Foods and health effects: A nutritional biochemistry perspective. *Curr Med Chem* 2016;23:2929–2957. <https://doi.org/10.2174/0929867323666160615105746>.
- [29] Gul K, Singh AK, Jabeen R. Nutraceuticals and functional foods: The foods for the future world. *Crit Rev Food Sci Nutr*. 2016;56:2617–2627. <https://doi.org/10.1080/10408398.2014.903384>.
- [30] Qin B, Polansky MM, Harry D, Anderson RA. Green tea polyphenols improve cardiac muscle mRNA and protein levels of signal pathways related to insulin and lipid metabolism and inflammation in insulin-resistant rats. *Mol Nutr Food Res* 2010;54:S14–23. <https://doi.org/10.1002/mnfr.200900306>.
- [31] Murase T, Misawa K, Minegishi Y, Aoki M, Ominami H, Suzuki Y, Shibuya Y, Hase T. Coffee polyphenols suppress diet-induced body fat accumulation by downregulating SREBP-1c and related molecules in C57BL/6j mice. *Am J Physiol-Endocrinol Metab*. 2011;300:E122–133. <https://doi.org/10.1152/ajpendo.00441.2010>.



- [32] Panchal SK, Poudyal H, Waanders J, Brown L. Coffee extract attenuates changes in cardiovascular and hepatic structure and function without decreasing obesity in high-carbohydrate, high-fat diet-fed male rats. *J Nutr.* 2012;142:690–697. <https://doi.org/10.3945/jn.111.153577>.
- [33] Chen C, Li Y, Xu Z. Chemical principles and bioactivities of blueberry. *Yao Xue Xue Bao.* 2010;45(4):422–429.
- [34] Stull AJ, Cash KC, Johnson WD, Champagne CM, Cefalu WT. Bioactives in blueberries improve insulin sensitivity in obese, insulin-resistant men and women. *J Nutr.* 2010;140:1764–1768. <https://doi.org/10.3945/jn.110.125336>.
- [35] Prior RL, Wilkes S, Rogers T, Khanal RC, Wu X, Hager TJ, Hager A, Howard L. Dietary black raspberry anthocyanins do not alter development of obesity in mice fed an obesogenic high-fat diet. *J Agric Food Chem.* 2010;58:3977–3983. <https://doi.org/10.1021/jf9030772>.
- [36] Roopchand DE, Kuhn P, Rojo LE, Lila MA, Raskin I. Blueberry polyphenol-enriched soybean flour reduces hyperglycemia, body weight gain and serum cholesterol in mice. *Pharmacol Res.* 2013;68:59–67. <https://doi.org/10.1016/j.phrs.2012.11.008>.
- [37] Medjakovic S, Jungbauer A. Pomegranate: a fruit that ameliorates metabolic syndrome. *Food Funct.* 2013;4:19–39. <https://doi.org/10.1039/C2FO30034F>.
- [38] Brahmi SA, M'rabet FE, Benbrahim Z, Akesbi Y, Amine B, Nejari C, El Mesbahi O. Complementary medicine use among Moroccan patients with cancer: A descriptive study. *Pan Afr Med J.* 2011;10. <https://doi.org/10.4314/pamj.v10i0>.
- [39] Gromek K, Drumond N, Simas P. Pharmacovigilance of herbal medicines. *Int J Risk Saf Med.* 2015;27:55–65. <https://doi.org/10.3233/JRS-150643>.
- [40] Bourhia M, Abdelaziz Shahat A, Mohammed Almarfadi O, Ali Naser F, Mostafa Abdelmageed W, Ait Haj Said A, El Gueddari F, Naamane A, Benbacer L, Khilil N. Ethnopharmacological survey of herbal remedies used for the treatment of cancer in the Greater Casablanca-Morocco. *Evid Based Complement Alternat Med.* 2019;2019:e1613457. <https://doi.org/10.1155/2019/1613457>.
- [41] Sreekeesoon DP, Mahomoodally MF. Ethnopharmacological analysis of medicinal plants and animals used in the treatment and management of pain in Mauritius. *J Ethnopharmacol.* 2014;157:181–200. <https://doi.org/10.1016/j.jep.2014.09.030>.
- [42] Lahsissene H, Kahouadji A. Usages thérapeutiques traditionnels des plantes médicinales dans le Maroc occidental: cas de la région de Zaër. *Phytothérapie.* 2010;8:210–217. <https://doi.org/10.1007/s10298-010-0566-5>.
- [43] Alaoui A, Laaribya S. Etude ethnobotanique et floristique dans les communes rurales Sehoul et Sidi-Abderrazak (cas de la Maamora-Maroc Septentrional). *Nat & Tech.* 2017;9(2):15–24.
- [44] Skalli S, Hassikou R, Arahou M. An ethnobotanical survey of medicinal plants used for diabetes treatment in Rabat, Morocco. *Heliyon.* 2019;5:e01421. <https://doi.org/10.1016/j.heliyon.2019.e01421>.
- [45] Jouad H, Haloui M, Rhiauani H, El Hilaly J, Eddouks M. Ethnobotanical survey of medicinal plants used for the treatment of diabetes, cardiac and renal diseases in the North centre region of Morocco (Fez-Boulemane). *J Ethnopharmacol.* 2001;77:175–182. [https://doi.org/10.1016/S0378-8741\(01\)00289-6](https://doi.org/10.1016/S0378-8741(01)00289-6).
- [46] Eddouks M, Maghrani M, Lemhadri A, Ouahidi M-L, Jouad H. Ethnopharmacological survey of medicinal plants used for the treatment of diabetes mellitus, hypertension and cardiac diseases in the south-east region of Morocco (Tafilalet). *J Ethnopharmacol.* 2002;82:97–103. [https://doi.org/10.1016/S0378-8741\(02\)00164-2](https://doi.org/10.1016/S0378-8741(02)00164-2).
- [47] Amrati FE-Z, Bourhia M, Slighoua M, Mohammad Salamatullah A, Alzahrani A, Ullah R, Bari A, Bousta D. Traditional medicinal knowledge of plants used for cancer treatment by communities of mountainous areas of Fez-Meknes-Morocco. *Saudi Pharm J.* 2021;29:1185–1204. <https://doi.org/10.1016/j.jsps.2021.09.005>.
- [48] Benkhiguel O, Chaachouay N, Khamar H, El Azzouzi F, Douira A, Zidane L. Ethnobotanical and ethnopharmacological study of medicinal plants used in the treatment of anemia in the region of Haouz-Rehamna (Morocco). *J Pharm Pharmacogn Res.* 2022;10:279–302. [https://doi.org/10.56499/jppres21.1196\\_10.2.279](https://doi.org/10.56499/jppres21.1196_10.2.279).
- [49] Fennane M. Propositions de zones importantes pour les plantes au Maroc (ZIP Maroc), Institut Scientifique, Rabat, Morocco, 2004.
- [50] Mukherjee PK, Maiti K, Mukherjee K, Houghton PJ. Leads from Indian medicinal plants with hypoglycemic potentials. *J Ethnopharmacol.* 2006;106:1–28. <https://doi.org/10.1016/j.jep.2006.03.021>.
- [51] Allali H, Benmehdi H, Dib M, Boufeldja T, Said G, Benabadji N. Phytotherapy of diabetes in west Algeria. *Asian J Chem.* 2008;20:2701–2710.
- [52] Katemo M, Mpiana PT, Mbala BM, Mihigo SO, Ngbolua KN, Tshibangu DST, Koyange PR. Ethnopharmacological survey of plants used against diabetes in Kisangani city (DR Congo). *J Ethnopharmacol.* 2012;144:39–43. <https://doi.org/10.1016/j.jep.2012.08.022>.
- [53] Keter LK, Mutiso PC. Ethnobotanical studies of medicinal plants used by traditional health practitioners in the management of diabetes in Lower Eastern Province, Kenya. *J Ethnopharmacol.* 2012;139:74–80. <https://doi.org/10.1016/j.jep.2011.10.014>.
- [54] Ezurike UF, Prieto JM. The use of plants in the traditional management of diabetes in Nigeria: Pharmacological and toxicological considerations. *J Ethnopharmacol.* 2014;155:857–924. <https://doi.org/10.1016/j.jep.2014.05.055>.
- [55] Yaseen G, Ahmad M, Zafar M, Sultana S, Kayani S, Cetto AA, Shaheen S. Traditional management of diabetes in Pakistan: Ethnobotanical investigation from Traditional Health Practitioners. *J Ethnopharmacol.* 2015;174:91–117. <https://doi.org/10.1016/j.jep.2015.07.041>.
- [56] Telli A, Esnault M-A, Ould El Hadj Khelil A. An ethnopharmacological survey of plants used in traditional diabetes treatment in south-eastern Algeria (Ouargla province). *J Arid Environ.* 2016;127:82–92. <https://doi.org/10.1016/j.jaridenv.2015.11.005>.
- [57] Amari A, Hachem K, Hassani MM. Ethnopharmacological survey of medicinal plants used to treat cutaneous leishmaniasis in Ain Sekhoune, Saida, Algeria. *Not Sci Biol.* 2021;13:10961–10961. <https://doi.org/10.15835/nsb13210961>.

- [58] Barkaoui M, Katiri A, Boubaker H, Msanda F. Ethnobotanical survey of medicinal plants used in the traditional treatment of diabetes in Chtouka Ait Baha and Tiznit (Western Anti-Atlas), Morocco. *J Ethnopharmacol.* 2017;198:338–350. <https://doi.org/10.1016/j.jep.2017.01.023>.
- [59] Idm'hand E, Msanda F, Cherifi K. Ethnopharmacological review of medicinal plants used to manage diabetes in Morocco. *Clin Phytosci.* 2020;6:18. <https://doi.org/10.1186/s40816-020-00166-z>.
- [60] Saadi B, Msanda F, Boubaker H. Contributions of folk medicine knowledge in Southwestern Morocco: The case of rural communities of Imouzzer Ida Outanane Region. *Int J Med Plant Res.* 2013;2:135–145.
- [61] Jamila F, Mostafa E. Ethnobotanical survey of medicinal plants used by people in Oriental Morocco to manage various ailments. *J Ethnopharmacol.* 2014;154:76–87. <https://doi.org/10.1016/j.jep.2014.03.016>.
- [62] Orch H, Douira A, Zidane L. Étude ethnobotanique des plantes médicinales utilisées dans le traitement du diabète, et des maladies cardiaques dans la région d'Izarène (Nord du Maroc). *J Appl Biosci.* 2015;86:7940–7956. <https://doi.org/10.4314/jab.v86i1>.
- [63] Hachi M, Benkhniq O, Hachi T, El Bouhaddioui M, Bouabadi I, Rochdi A, Zidane L. Contribution to the ethnobotanical study of antidiabetic medicinal plants of the Central Middle Atlas region (Morocco). *Lazaroa.* 2016;37(2016):1–11.
- [64] Bouyahya A, Abrini J, Et-Touys A, Bakri Y, Dakka N. Indigenous knowledge of the use of medicinal plants in the North-West of Morocco and their biological activities. *Eur J Integr Med.* 2017;13:9–25. <https://doi.org/10.1016/j.eujim.2017.06.004>.
- [65] Bouyahya A, El Omari N, Elmeniyi N, Guaouguaou F-E, Balahbib A, Belmehdi O, Salhi N, Imtara H, Mrabti HN, El-Shazely M, Bakri Y. Moroccan antidiabetic medicinal plants: Ethnobotanical studies, phytochemical bioactive compounds, preclinical investigations, toxicological validations and clinical evidences; challenges, guidance and perspectives for future management of diabetes worldwide. *Trends Food Sci Technol.* 2021;115:147–254. <https://doi.org/10.1016/j.tifs.2021.03.032>.
- [66] Kahouadji MS, Hseini S, Jaafar B, Zaid Y, Kahouadji A, Zidane L. Floristic Diversity and evaluation of the potential of spontaneous medicinal plants in the Bigoudine Watershed (Moroccan Western High Atlas). *Int J For Res.* 2022;2022:e2485502. <https://doi.org/10.1155/2022/2485502>.
- [67] Belhaj S, Chaachouay N, Zidane L. Ethnobotanical and toxicology study of medicinal plants used for the treatment of diabetes in the High Atlas Central of Morocco. *J Pharm Pharmacogn Res.* 2021;9:619–662. [https://doi.org/10.56499/jppres21.1014\\_9.5.619](https://doi.org/10.56499/jppres21.1014_9.5.619).
- [68] Ziyat A, Legssyer A, Mekhfi H, Dassouli A, Serhrouchni M, Benjelloun W. Phytotherapy of hypertension and diabetes in oriental Morocco. *J Ethnopharmacol.* 1997;58:45–54. [https://doi.org/10.1016/S0378-8741\(97\)00077-9](https://doi.org/10.1016/S0378-8741(97)00077-9).
- [69] Khabbach A, Libiad M, Ennabili A, Bousta D. Medicinal and cosmetic use of plants from the province of Taza, Northern Morocco. *Bol Latinoam Caribe Plant Med Aromat.* 2012;11(1):46–60.
- [70] Akdime H, Boukhira S, Mansouri LE, El AH, Bousta D. Ethnobotanical study and traditional knowledge of medicinal plants in Ain Leuh Region (Middle-Atlas of Morocco). *Am J Adv Drug Deliv.* 2015;3: 248–263.
- [71] Laadim M, Ouahidi ML, Zidane L, Hessni AE, Ouichou A, Mesfioui A. Ethnopharmacological survey of plants used for the treatment of diabetes in the town of Sidi Slimane (Morocco). *J Pharmacogn Phytother.* 2017;9:101–110. <https://doi.org/10.5897/JPP2016.0437>.
- [72] Hachlafi NE, Chebat A, Bencheikh RS, Fikri-Benbrahim K. Ethnopharmacological study of medicinal plants used for chronic diseases treatment in Rabat-Sale-Kenitra region (Morocco). *Ethnobot Res Appl.* 2020;20:1–23. <http://dx.doi.org/10.32859/era.20.2.1-23>.
- [73] Mechchate H, Es-safi I, Jawhari FZ, Bari A, Grafov A, Bousta D. Ethnobotanical survey about the management of Diabetes with medicinal plants used by diabetic patients in region of Fez-Meknes, Morocco. *J Ethnobot Res Appl.* 2020;19:1–28. <http://dx.doi.org/10.32859/era.19.12.1-28>.
- [74] Naceiri Mrabti H, Bouyahya A, Naceiri Mrabti N, Jaradat N, Doudach L, Faouzi MEA. Ethnobotanical survey of medicinal plants used by traditional healers to treat Diabetes in the Taza Region of Morocco. *Evid Based Complement Alternat Med.* 2021;2021:e5515634. <https://doi.org/10.1155/2021/5515634>.
- [75] Fatiha BA, Souad S, Ouafae B, Jamila D, Allal D, Lahcen Z. Ethnobotanical study of medicinal plants used in the region of middle oum rbia (morocco). *Plant Arch.* 2019;19(2):2005–1017.
- [76] Arrout A, El Ghallab Y, El Otmani IS, Said AAH. Ethnopharmacological survey of plants prescribed by herbalists for traditional treatment of hypercholesterolemia in Casablanca, Morocco. *J Herb Med.* 2022;36:100607. <https://doi.org/10.1016/j.hermed.2022.100607>.
- [77] Tahraoui A, El-Hilaly J, Israili ZH, Lyoussi B. Ethnopharmacological survey of plants used in the traditional treatment of hypertension and diabetes in south-eastern Morocco (Errachidia province). *J Ethnopharmacol.* 2007;110:105–17. <https://doi.org/10.1016/j.jep.2006.09.011>.
- [78] Belhaj S, Dahmani J, Belahbib N, Zidane L. Ethnopharmacological and Ethnobotanical study of Medicinal plants in the Central High Atlas, Morocco. *Ethnobot Res Appl* 2020;20:1–40.
- [79] Benali T, Khabbach A, Ennabili A, Hammani K. Ethnopharmacological prospecting of medicinal plants from the Province of Guercif (NE of Morocco). *Moroccan J Biol.* 2017;14:1–14.
- [80] Abouri M, Mousadik AE, Msanda F, Boubaker H, Saadi B, Cherifi K. An ethnobotanical survey of medicinal plants used in the Tata Province, Morocco. *Int J Med Plants Res.* 2012;1(7):99–123.
- [81] Mrabti HN, Jaradat N, Kachmar MR, Ed-Dra A, Ouahbi A, Cherrah Y, Faouzi MEA. Integrative herbal treatments of diabetes in Beni Mellal region of Morocco. *J Integr Med.* 2019;17(2):93–99. <https://doi.org/10.1016/j.joim.2019.01.001>.
- [82] Damjanovic I, Kitic D, Stefanovic N, Zlatkovic-Guberinic S, Catic-Djordjevic A, Velickovic-Radovanovic R. Herbal self-medication use in patients with diabetes mellitus type 2. *Turk J Med Sci.* 2015;45:964–971. <https://doi.org/10.3906/sag-1410-60>.



- [83] El Amrani F, Rhallab A, Alaoui T, El Badaoui K, Chakir S. Étude ethnopharmacologique de quelques plantes utilisées dans le traitement du diabète dans la région de Meknès-Tafilalet (Maroc). *Phytotherapie*. 2010;8:161–165. <https://doi.org/10.1007/s10298-010-0552-y>.
- [84] Errajaji A, Ouhdouch F, El-Anssari N. Usage des plantes médicinales dans le traitement du diabète de type 2 au Maroc. *Med des Mal Metab*. 2010;4:301–304. [https://doi.org/10.1016/S1957-2557\(10\)70064-X](https://doi.org/10.1016/S1957-2557(10)70064-X).
- [85] Boust D, Boukhira S, Aafi A, Ghanmi M, El Mansouri L. Ethnopharmacological Study of anti-diabetic medicinal plants used in the Middle-Atlas region of Morocco (Sefrou region). *Int J Pharma Res Health Sci*. 2014;2(1):75-79.
- [86] Ghourri M, Zidane L, Douira A. Usage des plantes médicinales dans le traitement du Diabète au Sahara Marocain (Tan-tan). *J Anim Plant Sci (JAPS)*. 2013;17(1):2388-2411. <http://www.m.elewa.org/JAPS>.
- [87] Benlamdini N, Elhafian M, Rochdi A, Zidane L. Étude floristique et ethnobotanique de la flore médicinale du Haut Atlas oriental (Haute Moulouya). *J Appl Biosci*. 2014;78:6771–6787. <https://doi.org/10.4314/jab.v78i1.17>.
- [88] Fadil M, Farah A, Haloui T, Rachiq S. Étude ethnobotanique des plantes exploitées par les coopératives et les associations de la région Meknès-Tafilalet au Maroc. *Phytotherapie*. 2014;13:19–30. <https://doi.org/10.1007/s10298-014-0902-2>.
- [89] Eddouks M, Ajebli M, Hebi M. Ethnopharmacological survey of medicinal plants used in Daraa-Tafilalet region (Province of Errachidia), Morocco. *J Ethnopharmacol*. 2017;198:516–530. <https://doi.org/10.1016/j.jep.2016.12.017>.
- [90] Bouasla A, Bouasla I. Ethnobotanical survey of medicinal plants in northeastern of Algeria. *Int J Phytomed*. 2017;36:68–81. <https://doi.org/10.1016/j.phymed.2017.09.007>.
- [91] Belmouhoub M, Tacherfiout M, Boukhalfa F, Khodja YK, Bachir-Bey M. Traditional medicinal plants used in the treatment of diabetes: Ethnobotanical and ethnopharmacological studies and mechanisms of action. *Int J Plant Based Pharm*. 2022;2(2):145–154.
- [92] Altundag E, Ozturk M. Ethnomedicinal studies on the plant resources of east Anatolia, Turkey. *Procedia - Soc Behav Sci*. 2011;19:756–777. <https://doi.org/10.1016/j.sbspro.2011.05.195>.
- [93] Ozturk M, Altay V, Gucel S, Altundag E. Plant diversity of the drylands in Southeastern Anatolia-Turkey: role in human health and food security. *Plant Biodivers Monit Assess Conserv*. 2017;83–124. <https://doi.org/10.1079/9781780646947.0083>.
- [94] Kachmar MR, Naceiri Mrabti H, Bellahmar M, Ouahbi A, Haloui Z, El Badaoui K, Bouyahya A, Chakir S. Traditional knowledge of medicinal plants used in the northeastern part of Morocco. *Evid Based Complement Alternat Med*. 2021;2021:e6002949. <https://doi.org/10.1155/2021/6002949>.
- [95] Bahmani M, Zargaran A, Rafieian-Kopaei M, Saki K. Ethnobotanical study of medicinal plants used in the management of diabetes mellitus in the Urmia, Northwest Iran. *Asian Pac J Trop Med*. 2014;7:S348–54. [https://doi.org/10.1016/S1995-7645\(14\)60257-1](https://doi.org/10.1016/S1995-7645(14)60257-1).
- [96] Chaachouay N, Douira A, Zidane L. Herbal medicine used in the treatment of human diseases in the Rif, Northern Morocco. *Arab J Sci Eng*. 2022;47:131–153. <https://doi.org/10.1007/s13369-021-05501-1>.
- [97] Sargin SA, Akçicek E, Selvi S. An ethnobotanical study of medicinal plants used by the local people of Alaşehir (Manisa) in Turkey. *J Ethnopharmacol*. 2013;150:860–874. <https://doi.org/10.1016/j.jep.2013.09.040>.
- [98] Boufous H, Marhoume F, Chait A, Bagri A. Ethnopharmacological survey of medicinal plants with hallucinogenic effect and used against pain, inflammatory diseases, diabetes and urinary lithiasis in Zagora “Morocco”. *Galenic J Pharm*. 2017;6(4):342–350. <https://doi.org/10.5455/jice.20170721062527>.
- [99] Bouayyadi L, Hafian ME, Zidane L. Étude floristique et ethnobotanique de la flore médicinale dans la région du Gharb, Maroc. *J Appl Biosci*. 2015;93:8770–8788. <https://doi.org/10.4314/jab.v93i1>.
- [100] Semenya S, Potgieter M, Erasmus L. Ethnobotanical survey of medicinal plants used by Bapedi healers to treat diabetes mellitus in the Limpopo Province, South Africa. *J Ethnopharmacol*. 2012;141:440–445. <https://doi.org/10.1016/j.jep.2012.03.008>.
- [101] El Rhaffari L, Zaid A. Pratique de la phytothérapie dans le sud-est du Maroc (Tafilalet) : Un savoir empirique pour une pharmacopée rénovée. Fleurentin J, Pelt JM, Mazars G (eds.), Lejosne J.C. (trad.), Cabalion Pierre (collab.), 2002.
- [102] Chaachouay N, Benkhniue O, Fadli M, El Ibaoui H, Zidane L. Ethnobotanical and ethnopharmacological studies of medicinal and aromatic plants used in the treatment of metabolic diseases in the Moroccan Rif. *Heliyon*. 2019;5:e02191. <https://doi.org/10.1016/j.heliyon.2019.e02191>.
- [103] Polat R, Cakilcioglu U, Kaltalioglu K, Ulsan MD, Türkmen Z. An ethnobotanical study on medicinal plants in Espiye and its surrounding (Giresun-Turkey). *J Ethnopharmacol*. 2015;163:1–11. <https://doi.org/10.1016/j.jep.2015.01.008>.
- [104] Kadir MF, Bin Sayeed MS, Shams T, Mia MMK. Ethnobotanical survey of medicinal plants used by Bangladeshi traditional health practitioners in the management of diabetes mellitus. *J Ethnopharmacol*. 2012;144:605–611. <https://doi.org/10.1016/j.jep.2012.09.050>.
- [105] Ahmad M, Qureshi R, Arshad M, Khan MA, Zafar M. Traditional herbal remedies used for the treatment of diabetes from district attock (pakistan). *Pak J Bot*. 2009; 41(6): 2777–2782.
- [106] Gürdal B, Kültür Ş. An ethnobotanical study of medicinal plants in Marmaris (Muğla, Turkey). *J Ethnopharmacol*. 2013;146:113–126. <https://doi.org/10.1016/j.jep.2012.12.012>.
- [107] Labiad H, Et-tahir A, Ghanmi M, Satrani B, Aljaiyash A, Chaouch A, Fadli M. Ethnopharmacological survey of aromatic and medicinal plants of the pharmacopoeia of northern Morocco. *Ethnobot Res Appl*. 2020;19:1–16. <http://dx.doi.org/10.32859/era.19.45.1-16>.
- [108] Polat R. Ethnobotanical study on medicinal plants in Bingöl (City center) (Turkey). *J Herb Med*. 2019;16:100211. <https://doi.org/10.1016/j.hermed.2018.01.007>.
- [109] Palabaş Uzun S, Koca C. Ethnobotanical survey of medicinal plants traded in herbal markets of Kahramanmaraş. *Plant Divers*. 2020;42:443–454. <https://doi.org/10.1016/j.pld.2020.12.003>.

- [110] Kpodar MS, Lawson-Evi P, Bakoma B, Ekl-Gadegbeku K, Agbonon A, Aklikokou K, Gbeassor M. Ethnopharmacological survey of plants used in the treatment of diabetes mellitus in south of Togo (Maritime Region). *J Herb Med.* 2015;5:147–152. <https://doi.org/10.1016/j.hermed.2015.06.002>.
- [111] Demoz M, Gachoki K, Mungai K, Negusse B. Ethnobotanical survey and preliminary phytochemical studies of plants traditionally used for diabetes in Eritrea. *Eur J Med Plants.* 2015;9:1–11.
- [112] Gnagne AS, Camara D, Fofie NBY, Bene K, Zirihi GN. Ethnobotanical study of medicinal plants used in the treatment of diabetes in the Department of Zouenoula (Côte d'Ivoire). *J Appl Biosci.* 2017;113:11257–11266. <http://dx.doi.org/10.4314/jab.v113i1.14>.
- [113] Rhattas M, Douira A, Zidane L. Étude ethnobotanique des plantes médicinales dans le Parc National de Talassemtane (Rif occidental du Maroc). *J Appl Biosci.* 2016;97:9187–9211. <https://doi.org/10.4314/jab.v97i1.5>.
- [114] Ozturk M, Altundağ E, Ibadullayeva SJ, Altay V, Aslanipour B. A comparative analysis of medicinal and aromatic plants used in the traditional medicine of iğdir (turkey), nakhchivan (azerbaijan), and tabriz (iran). *Pak J Bot.* 2018; 50(1): 337–343.
- [115] Bencheikh N, Bouhrim M, Merrouni IA, Boutahiri S, Kharchoufa L, Addi M, Tungmunthum D, Hano C, Eto B, Legssyer A, Elachouri M. Antihyperlipidemic and antioxidant activities of flavonoid-rich extract of *Ziziphus lotus* (L.) Lam. fruits. *Appl Sci.* 2021;11:7788. <https://doi.org/10.3390/app11177788>.
- [116] Woillez M-N. Revue de littérature sur le changement climatique au Maroc: Observations, projections et impacts. *Pap Rech.* 2019;1–33. <https://doi.org/10.3917/afd.woill.2019.01.0001>.
- [117] El Hamsas el Youbi A, Ouahidi I, El Mansouri L, Daoudi A, Bousta D. Ethnopharmacological survey of plants used for immunological diseases in four regions of Morocco. *Eur J Med Plants.* 2016;13:1–24. <https://doi.org/10.9734/EJMP/2016/12946>.
- [118] Kassout J, Terral J-F, El Ouahrani A, Houssni M, Ivorra S, Kadaoui K, El Mahroussi M, Paradis L, Ater M. Species distribution based-modelling under climate change: The Case of two native wild *Olea europaea* subspecies in Morocco, O. e. subsp. *europaea* var. *sylvestris* and O. e. subsp. *maroccana*. In: Leal Filho W, Manolas E, editors. *Clim. Change Mediterr. Middle East. Reg.*, Cham: Springer International Publishing; 2022, p. 21–43. [https://doi.org/10.1007/978-3-030-78566-6\\_2](https://doi.org/10.1007/978-3-030-78566-6_2).
- [119] El SN, Karakaya S. Olive tree (*Olea europaea*) leaves: Potential beneficial effects on human health. *Nutr Rev.* 2009;67:632–638. <https://doi.org/10.1111/j.1753-4887.2009.00248.x>.
- [120] Guex CG, Reginato FZ, de Jesus PR, Brondani JC, Lopes GH, Bauermann L de F. Antidiabetic effects of *Olea europaea* L. leaves in diabetic rats induced by high-fat diet and low-dose streptozotocin. *J Ethnopharmacol.* 2019;235:1–7. <https://doi.org/10.1016/j.jep.2019.02.001>.
- [121] Cheurfa M, Abdallah HH, Allem R, Noui A, Picot-Allain CMN, Mahomoodally F. Hypocholesterolaemic and antioxidant properties of *Olea europaea* L. leaves from Chlef province, Algeria using in vitro, in vivo and in silico approaches. *Food Chem Toxicol.* 2019;123:98–105. <https://doi.org/10.1016/j.fct.2018.10.002>.
- [122] Alesci A, Miller A, Tardugno R, Pergolizzi S. Chemical analysis, biological and therapeutic activities of *Olea europaea* L. extracts. *Nat Prod Res.* 2022;36:2932–2945. <https://doi.org/10.1080/14786419.2021.1922404>.
- [123] Amin ME, Virk P, Elobeid MAR, Almarhoon ZM, Hassan ZK, Omer SA, Merghano NM, Daghestani MH, Al Olayan EM. Anti-diabetic effect of *Murraya koenigii* (L) and *Olea europaea* (L) leaf extracts on streptozotocin induced diabetic rats. *Pak J Pharm Sci.* 2013;26(2):359–365.
- [124] Souto-Maior FN, Fonsêca DV da, Salgado PRR, Monte L de O, de Sousa DP, de Almeida RN. Antinociceptive and anticonvulsant effects of the monoterpene linalool oxide. *Pharm Biol.* 2017;55:63–67. <https://doi.org/10.1080/13880209.2016.1228682>.
- [125] Ghule AE, Jadhav SS, Bodhankar SL. Effect of ethanolic extract of seeds of *Linum usitatissimum* (Linn.) in hyperglycaemia associated ROS production in PBMNCs and pancreatic tissue of alloxan induced diabetic rats. *Asian Pac J Trop Dis.* 2012;2:405–410. [https://doi.org/10.1016/S2222-1808\(12\)60088-7](https://doi.org/10.1016/S2222-1808(12)60088-7).
- [126] Mechchate H, Es-safi I, Conte R, Hano C, Amaghnoije A, Jawhari FZ, Radouane N, Bencheikh N, Grafov A, Bousta D. In Vivo and in vitro antidiabetic and anti-inflammatory properties of Flax (*Linum usitatissimum* L.) seed polyphenols. *Nutrients* 2021;13:2759. <https://doi.org/10.3390/nu13082759>.
- [127] Arslanoğlu F, Aytac S. The important in terms of health of Flax (*Linum usitatissimum* L.). *Int J Life Sci Biotechnol.* 2020;3:95–107. <https://doi.org/10.38001/ijlsb.690295>.
- [128] Nagulapalli Venkata KC, Swaroop A, Bagchi D, Bishayee A. A small plant with big benefits: Fenugreek (*Trigonella foenum-graecum* Linn.) for disease prevention and health promotion. *Mol Nutr Food Res.* 2017;61:1600950. <https://doi.org/10.1002/mnfr.201600950>.
- [129] Inbaraj SD, Muniappan M. Correlation between the in-vitro and in-vivo antihyperglycemic effect of *Ocimum sanctum*, *Trigonella Foenum Graecum* and *Curcuma longa*. *Pharmacogn J.* 2020;12:369–76. <https://doi.org/10.5530/pj.2020.12.58>.
- [130] Marmouzi I, Kharbach M, El Jemli M, Bouyahya A, Cherrah Y, Bouklouze A, Heyden YV, Faouzi EA. Antidiabetic, dermatoprotective, antioxidant and chemical functionalities in *Ziziphus lotus* leaves and fruits. *Ind Crops Prod.* 2019;132:134–139. <https://doi.org/10.1016/j.indcrop.2019.02.007>.
- [131] Regehr DL, Brahli AE. Wild jujube (*Ziziphus lotus*) control in Morocco. *Weed Technol.* 1995;9:326–330. <https://doi.org/10.1017/S0890037X00023423>.
- [132] Pasha HH, Jabbar U, Anjum A, Kanwal A, Nasrullah C, Bashir A, Qureshi HJ. Is *Coriandrum sativum* hypolipidemic in alloxan induced diabetic rats? *Proceedings.* 2022;36:35–38. <https://doi.org/10.47489/PSZMC-825361-35-38>.
- [133] Das S, Rajadnya V, Kothari R, Tilak AV, Raveendran S, Deshpande T. Hypolipidemic activity of *Coriandrum sativum* in diabetic dyslipidemic rats. *Int J Basic Clin Pharmacol.* 2019;8(6):1393–1397. <http://dx.doi.org/10.18203/2319-2003.ijbcp20192208>

- [134] Matsubara E, Fukagawa M, Okamoto T, Ohnuki K, Shimizu K, Kondo R. (-)-Bornyl acetate induces autonomic relaxation and reduces arousal level after visual display terminal work without any influences of task performance in low-dose condition. *Biomed Res.* 2011;32:151-157. <https://doi.org/10.2220/biomedres.32.151>.
- [135] Riella KR, Marinho RR, Santos JS, Pereira-Filho RN, Cardoso JC, Albuquerque-Junior RLC, Thomazzi SM. Anti-inflammatory and cicatrizing activities of thymol, a monoterpene of the essential oil from *Lippia gracilis*, in rodents. *J Ethnopharmacol.* 2012;143:656-663. <https://doi.org/10.1016/j.jep.2012.07.028>.
- [136] Sun G, Zhang S, Xie Y, Zhang Z, Zhao W. Gallic acid as a selective anticancer agent that induces apoptosis in SMMC-7721 human hepatocellular carcinoma cells. *Oncol Lett.* 2016;11:150-158. <https://doi.org/10.3892/ol.2015.3845>.
- [137] Souto-Maior FN, Carvalho FL de, Morais LCSL de, Netto SM, de Sousa DP, Almeida RN de. Anxiolytic-like effects of inhaled linalool oxide in experimental mouse anxiety models. *Pharmacol Biochem Behav.* 2011;100:259-263. <https://doi.org/10.1016/j.pbb.2011.08.029>.
- [138] de Lucena JD, Gadelha-Filho CVJ, da Costa RO, de Araújo DP, Lima FAV, Neves KRT, de Barros Viana GS. L-linalool exerts a neuroprotective action on hemiparkinsonian rats. *Naunyn Schmiedebergs Arch Pharmacol.* 2020;393:1077-1088. <https://doi.org/10.1007/s00210-019-01793-1>.
- [139] Hashmi Z, Sarkar D, Mishra S, Mehra V. An in-vitro assessment of anti-inflammatory, antioxidant, and anti-hyperglycemic activities of traditional edible plants - *Murraya koenigii*, *Mentha spicata*, and *Coriandrum sativum*. *J Biomed Ther Sci.* 2022;9:1-10.
- [140] Keskes H, Mnafigui K, Hamden K, Damak M, El Feki A, Allouche N. In vitro anti-diabetic, anti-obesity and antioxidant proprieties of *Juniperus phoenicea* L. leaves from Tunisia. *Asian Pac J Trop Biomed.* 2014;4:S649-55. <https://doi.org/10.12980/APJTB.4.201414B114>.
- [141] Adams DRP. *Junipers of the World: The Genus Juniperus*, 4th Edition. Trafford Publishing; 2014.
- [142] Alami AE, Fattah A, Chait A. Medicinal plants used for the prevention purposes during the Covid-19 pandemic in Morocco. *J Anal Sci Appl Biotechnol.* 2020;2:2-11. <https://doi.org/10.48402/IMIST.PRSM/jasab-v2i1.21056>.
- [143] Aboukhalid K, Lamiri A, Agacka-Mołodoch M, Doroszevska T, Douaik A, Bakha M, Casanova J, Tomi F, Machon N, Al Faiz C. Chemical polymorphism of *Origanum compactum* grown in all natural habitats in Morocco. *Chem Biodivers.* 2016;13:1126-1139. <https://doi.org/10.1002/cbdv.201500511>.
- [144] Bouyahya A, Abrini J, Bakri Y, Dakka N. Screening phytochimique et évaluation de l'activité antioxydante et antibactérienne des extraits d'*Origanum compactum*. *Phytotherapie.* 2017;15:379-83. <https://doi.org/10.1007/s10298-017-1101-8>.
- [145] Zeroual A, Eloutassi N, Chaouch M, Chaqroune A. Antimicrobial, antioxidant activity, and chemical composition of *Origanum compactum* benth from taounate province, north morocco. *Asian J Pharm Clin Res.* 2020:126-131. <https://doi.org/10.22159/ajpcr.2020.v13i3.36319>.
- [146] Mechchate H, Ouedrhiri W, Es-safi I, Amaghnoije A, Jawhari FZ, Bousta D. Optimization of a new antihyperglycemic formulation using a mixture of *Linum usitatissimum* L., *Coriandrum sativum* L., and *Olea europaea* var. *sylvestris* flavonoids: A mixture design approach. *Biologics.* 2021;1:154-163. <https://doi.org/10.3390/biologics1020009>.
- [147] Vijayakumar S, Morvin Yabesh JE, Prabhu S, Manikandan R, Muralidharan B. Quantitative ethnomedicinal study of plants used in the Nelliampathy hills of Kerala, India. *J Ethnopharmacol.* 2015;161:238-254. <https://doi.org/10.1016/j.jep.2014.12.006>.