

# Cytotoxicity screening of some Turkish plants against renal cancer cells

Feyyaz MIHOĞLUGİL<sup>1\*</sup>, Demet AKALGAN<sup>2</sup>, Fatma TOSUN<sup>3</sup>

<sup>1</sup> Faculty of Pharmacy, Cyprus International University, Nicosia, North Cyprus

<sup>2</sup> Istinye University, Faculty of Pharmacy, Department of Pharmacognosy, Istanbul, Türkiye

<sup>3</sup> Department of Pharmacognosy, School of Pharmacy, Istanbul Medipol University, Istanbul, Türkiye

\* Corresponding Author. E-mail: [fmihoglugil@ciu.edu.tr](mailto:fmihoglugil@ciu.edu.tr) (F.M.); Tel. +905353605545

Received: 29 August 2022 / Revised: 02 December 2022 / Accepted: 02 December 2022

**ABSTRACT:** In this contribution, we performed cytotoxicity screening of 51 extracts from 17 plant species, representing 14 families growing in Turkey. Cytotoxic activities of the dichloromethane, ethyl acetate, and methanol extracts of the plants were investigated against the renal (A498 and UO31) cancer cell lines. 28 extracts exhibited more than 50% growth inhibition at a 25 µg/mL concentration on both renal cancer A498 and UO31 cell lines. The extracts showed higher activity against UO31 cells than A498 cells. The dichloromethane extract of the aerial parts of *Scabiosa columbaria* ssp. *columbaria* showed the highest cytotoxic activity with 64% inhibition at a 25 µg/mL concentration on the renal cancer UO31 cell line. The methanol extract of the aerial parts of *Epilobium minutiflorum* showed the highest cytotoxic activity with 63% inhibition at a 25 µg/mL concentration on the renal cancer A498 cell line.

**KEYWORDS:** Cytotoxic activity; renal cancer; Turkish plants; *Scabiosa columbaria* ssp. *columbaria*

## 1. INTRODUCTION

Cancer remains one of the leading causes of death worldwide [1]. Kidney cancer (Renal cell cancer-RCC) is the second most common cancer of the urinary system [2]. In 2020, there were approximately 431,000 new cases (2.2% of all new cases) and 179,000 deaths due to renal cancer (1.8% of all death cases) [3]. The discovery of new compounds from natural sources to fight cancer has become a matter of great interest among researchers [4,5]. Plant extracts and plant-derived natural compounds such as glycosides, alkaloids, tannins, terpenes, coumarins, and flavonoids have been reported to inhibit the growth of cancer cells such as renal, lung, breast, and colorectal cancer cells [6-17]. The prognosis of patients with RCC remains poor, although advances in therapeutic approaches. Since therapeutic opportunities are limited, approximately less than 40% of patients survive five years or more after diagnosis. Besides, since RCC is resistant to chemotherapy and radiotherapy, RCC recurrence principally owes to a lack of routine adjuvant therapy in the clinic. So, it is required to detect new compounds and develop novel targeted therapies for RCC [7].

The present study aimed to evaluate the cytotoxic activities of 17 plant species, representing 14 families growing in Turkey, against the renal (A498 and UO31) cancer cell lines. This paper also introduces new plant sources whose cytotoxic activities have not been reported previously.

## 2. RESULTS AND DISCUSSION

This study is the first cytotoxicity screening performed on these 17 plant species against the renal (A498 and UO31) cancer cell lines.

Cytotoxic activities of 51 extracts obtained from 17 plant species tested against A498 and UO31 renal cancer cell lines are shown in Table 1.

Among 51 extracts tested, 28 extracts (54.9%) exhibited growth inhibition of more than 50% at a 25 µg/mL concentration in both renal cancer (A498 and UO31) cell lines. Extracts showed higher activity against UO31 cells than A498 cells. The most potent activity was found for the CH<sub>2</sub>Cl<sub>2</sub> extract of *Scabiosa columbaria* ssp. *columbaria* with 64% inhibition at a 25 µg/mL concentration against UO31 cell line. The CH<sub>2</sub>Cl<sub>2</sub> extracts of *Campanula alliarifolia*, *Corylus avellana* and *Geranium divaricatum*; the EtOAc extracts of

**How to cite this article:** Mihoglugil F., Akalgan D., Tosun F. Cytotoxicity screening of some Turkish plants against renal cancer cells. J Res Pharm. 2023; 27(2): 636-641.

*Campanula lactiflora* and *Corylus avellana*; the MeOH extracts of *Campanula alliarifolia*, *Campanula lactiflora*, *Epilobium minutiflorum* and *Epilobium angustifolium* were also active with the inhibition of more than 60% at a 25 ug/mL concentration against UO31 cell line. The most active extracts against A498 cell line were the MeOH extract of *Epilobium minutiflorum* and the EtOAc extract of *Melampyrum arvense* with 60% inhibition at a 25 ug/mL concentration. Of all 17 CH<sub>2</sub>Cl<sub>2</sub> extracts, 15 showed more than 50% growth inhibition at a 25 ug/mL concentration in both renal cancer cell lines. 10 MeOH extracts and 3 EtOAc extracts inhibited the growth of both renal cancer cell lines by more than 50% at a 25 ug/mL concentration. All three extracts of *Echium vulgare* and *Melampyrum arvense* exhibited more than 50% growth inhibition at a 25 ug/mL concentration in both renal cancer cell lines.

Previously six of these 17 plants were tested on different cancer cell lines. For example, the extracts of *Clinopodium vulgare* were evaluated for cytotoxic activities against the Caco-2, HepG2, and MCF-7 cancer cell lines [18] and A2058 (human metastatic melanoma), HEp-2 (epidermoid carcinoma, larynx, human) and L5178Y (mouse lymphoma) cell lines [19] and CaOV (human testis cystadenocarcinoma), HeLa (human cervical adenocarcinoma), HT-29 (human colorectal adenocarcinoma) [20]. Previous studies reported that the leaf and stem extracts of *Corylus avellana* significantly reduced the viability of the HeLa (cervical cancer), HepG2 (liver hepatocarcinoma), and MCF-7 (breast cancer) cell lines and contained over 100 compounds as organic acids, triacylglycerols, phytosterols, phenolic acids, diarylheptanoids, flavonoids, tannins, isoflavones, lignans, terpenes, and taxanes. (3R,5R)-3,5-dihydroxy-1,7-bis(4-hydroxy-phenyl) heptane 3-O-β-D-glucopyranoside and quercitrin were identified as contributors to the strong cell viability-reducing activity of *C. avellana* leaf extracts [21]. *Rhododendron ponticum* were tested against human prostate carcinoma (DU145) and human prostate adenocarcinoma (PC3) cell lines [22]; *R. luteum* extracts on A549 breast, colon, prostate, liver and lung cancer cell lines [23, 24]; *Sedum spurium* extracts against the Hep-2 human larynx epidermoid carcinoma cells [25]. *Epilobium angustifolium* extracts have been shown to exhibit potent antiproliferative activity on the LNCap prostate carcinoma cell lines [26] and has been reported to contain ellagitannins, flavonoids, and phenolic acids [27,28] and Urolithin C, metabolites of *Epilobium* ellagitannins, has been explained to responsible for the activity [27]. In this study, the MeOH extract of *Epilobium angustifolium* and EtOAc extract of *Corylus avellana* exhibited more than 60% growth inhibition at a 25 ug/mL concentration against UO31 cell line. The remaining extracts of these six plants showed between 41% and 60% growth inhibition at a 25 ug/mL concentration in both renal cancer (A498 and UO31) cell lines.

No reports in the literature deal with the cytotoxic activities of *E. vulgare*, *S. columbaria* ssp. *columbaria*, *C. alliarifolia*, *C. lactiflora*, *G. divaricatum*, *L. pratensis*, *E. minutiflorum*, *M. arvense*, *G. leiocarpum*, *T. minus* and *U. minor*.

### 3. CONCLUSION

This study is the first cytotoxicity screening performed on these 17 plant species against the renal (A498 and UO31) cancer cell lines. Of all 17 plant species, 11 were investigated for the first time for the cytotoxic activities in this study.

Previous studies on the among the most active plants (*S. columbaria* ssp. *columbaria*, *C. alliarifolia*, *C. lactiflora*, *G. divaricatum*, *C. avellana* and *M. arvense*) *Scabiosa columbaria* ssp. *columbaria* has been reported to contain six phenolic compounds; chlorogenic acid was the major compound and exhibited antioxidant activity [29]. *Campanula* species have been reported to contain polyphenols, steroids, phenylpropanoid derivatives, triterpenes, and polyacetylenes and show antioxidant activity [30,31]. Previous phytochemical studies on the *Geranium* genus have shown the presence of phenolic compounds such as gallic, ellagic, quinic acid, quercetin, geraniin, and corilagin [32-33]. *Melampyrum arvense* has been reported to exhibit antimicrobial and antioxidant activity and contain phenolic compounds [34]. There are no reports in the literature dealing with the cytotoxic activities of the five plants (*S. columbaria* ssp. *columbaria*, *C. alliarifolia*, *C. lactiflora*, *G. divaricatum* and *M. arvense*) of the most active six plants.

In conclusion, bioactivity-guided fractionation of the *S. columbaria* ssp. *columbaria*, *C. alliarifolia*, *C. lactiflora*, *G. divaricatum*, *C. avellana* and *M. arvense* extracts, which exhibited growth inhibition of more than 60% at a 25 ug/mL concentration, are planned to isolate and identify their cytotoxic principles.

### 4. MATERIALS AND METHODS

#### 4.1. Plant materials

Collection sites and voucher numbers of the plants are listed in Table 1. Voucher specimens are deposited in the Herbarium of Faculty of Pharmacy of Istanbul Medipol University, Istanbul, Turkey (IMEF).

Used plant part is leaves for *Corylus avellana* and *Ulmus minor*. Aerial parts are used for the remaining 15 plants.

## 4.2. Preparation of extracts

Air-dried and coarsely powdered leaves/aerial parts of the plants were sequentially extracted at room temperature with dichloromethane, ethyl acetate, and methanol. The extracts were separately concentrated in a rotary evaporator under reduced pressure to dryness. All extracts were stored at  $-20^{\circ}\text{C}$  prior to screening.

**Table 1.** Growth inhibition of the plant extracts at 25 ug/mL concentration on the renal cancer cell lines

Plants	Voucher No	Collection sites	Inhibition %					
			A498			UO31		
			1	2	3	1	2	3
<b>Betulaceae</b>								
<i>Corylusavellana</i> L.	IMEF1175	Giresun-Şebinkarahisar	53	47	50	60	61	58
<b>Boraginaceae</b>								
<i>Echium vulgare</i> L.	IMEF1161	Giresun-Şebinkarahisar	56	52	51	59	54	58
<b>Campanulaceae</b>								
<i>Campanulaalliarifolia</i> Willd.	IMEF1162	Giresun-Şebinkarahisar	53	47	52	60	52	61
<i>Campanulalactiflora</i> M.Bieb.	IMEF1168	Giresun-Şebinkarahisar	53	46	54	56	60	62
<b>Caprifoliaceae</b>								
<i>Scabiosacolumbari</i> sp.	IMEF1141	Trabzon-Zigana pass	51	48	55	64	56	55
<b>Crassulaceae</b>								
<i>Sedumspuri</i> M.Bieb.	IMEF1142	Trabzon-Hamsiköy	56	47	37	57	55	43
<b>Ericaceae</b>								
<i>Rhododendronluteum</i> Sweet	IMEF1150	Trabzon-Altındere	56	44	36	58	55	41
<i>Rhododendronponticum</i> L.	IMEF1152	Trabzon-Tonya	53	48	53	55	54	45
<b>Geraniaceae</b>								
<i>Geraniumdivaricatum</i> Ehrh.	IMEF1131	Trabzon-Zigana pass	53	42	53	60	46	55
<b>Lamiaceae</b>								
<i>Clinopodiumvulgare</i> L.	IMEF1137	Trabzon Zigana pass	41	47	50	40	54	57
<b>Leguminosae</b>								
<i>Lathyruspratensis</i> L.	IMEF1167	Giresun-Şebinkarahisar	52	48	43	56	50	47
<b>Onagraceae</b>								
<i>Epilobiumangustifolium</i> L.	IMEF1147	Trabzon-Tonya	53	47	53	59	49	61
<i>Epilobiumminutiflorum</i> Hausskn.	IMEF1171	Giresun-Şebinkarahisar	58	47	63	56	49	63
<b>Orobanchaceae</b>								
<i>Melampyrumarvense</i> L.	IMEF1055	Trabzon-Tonya	50	61	52	53	54	54
<b>Papaveraceae</b>								
<i>Glauciumleiocarpum</i> Boiss.	IMEF1135	Ankara-Çankaya	39	50	46	42	57	55
<b>Ranunculaceae</b>								
<i>Thalictrumminus</i> L.	IMEF1172	Giresun-Şebinkarahisar	50	48	50	57	52	47
<b>Ulmaceae</b>								
<i>Ulmusminor</i> Mill.	IMEF1166	Giresun-Şebinkarahisar	55	47	46	58	45	58

1.  $\text{CH}_2\text{Cl}_2$  extract; 2. EtOAc extract; 3. MeOH extract

### 4.3. Cytotoxic activity assay

The assay used for this study was a two-day, two cell line XTT bioassay [17], an *in vitro* antitumor colorimetric assay developed by the MTL Assay Development and Screening Section. Renal cancer cell lines used were A498 and U031. Sanguinarine was used as a positive control. The assay was performed as described previously [16]. Cytotoxic activities of extracts are shown in Table 1.

**Acknowledgements:** We thank Dr. John Beutler, Molecular Targets Laboratory, CCR, NCI, Frederick, MD, U.S.A. for the cytotoxic activity testing and Prof. Mecit Vural for the identification of plant material

**Author contributions:** Concept – F.M.; Design – F.M.; Supervision – F.T.; Materials – F.M., D.A., F.T.; Data Collection and/or Processing – F.M., D.A., F.T.; Analysis and/or Interpretation – F.M., D.A., F.T.; Literature Search – F.M., D.A., F.T.; Writing – F.M., D.A., F.T.; Critical Reviews – F.T., F.M., D.A.

**Conflict of interest statement:** The authors declared no conflict of interest

### REFERENCES

1. Siegel R, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2022. *CA Cancer J. Clin.* 2022; 72: 7-33. <https://doi.org/10.3322/caac.21708>
2. Shu X, Zhang Z, Yao Z-Y and Xing X-L. Identification of five ferroptosis-related lncRNAs as novel prognosis and diagnosis signatures for renal cancer. *Front Mol Biosci.* 2022; 8:763697. <https://doi.org/10.3389/fmolb.2021.763697>
3. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global Cancer Statistics 2020: GLOBOCAN Estimates of incidence and mortality Worldwide for 36 cancers in 185 countries. *CA A. Cancer J Clin.* 2021; 71: 209-249. <https://doi.org/10.3322/caac.21660>
4. Newman DJ, Cragg GM. Natural products as sources of new drugs over the nearly four decades from 01/ 1981 to 09/2019. *J Nat Prod.* 2020; 83: 770-803. <https://doi.org/10.1021/acs.jnatprod.9b01285>
5. Majolo F, Delwing LKOB, Marmitt DJ, Bustamante-Filho IC, Goettert M I. Medicinal plants and bioactive natural compounds for cancer treatment: Important advances for drug discovery. *Phytochem Lett.* 2019; 31: 196-207. <https://doi.org/10.1016/j.phytol.2019.04.003>
6. Isyaka SM, Mas-Claret E, Langat MK, Hodges T, Selway B, Mbala BM, Mvingu BK, Mulholland DA. Cytotoxic diterpenoids from the leaves and stem bark of *Croton haumanianus* (Euphorbiaceae). *Phytochemistry.* 2020; 178: 112455. <https://doi.org/10.1016/j.phytochem.2020.112455>
7. Zhu C, Na N, Sheng H, Feng B, Wang H, Zhu P, Zhang W, Zhang M and Deng Z. Ginkgolic acid inhibits the growth of renal cell carcinoma cells via inactivation of the EGFR signaling pathway. *Exp Ther Med.* 2020; 19: 2949-2956. <https://doi.org/10.3892/etm.2020.8570>
8. Jisha N, Vysakh A, Vijeesh V, Latha MS. Ethyl acetate fraction of *Muntingia calabura* L. exerts anti-colorectal cancer potential via regulating apoptotic and inflammatory pathways. *J Ethnopharmacol.* 2020; 261: 113064. <https://doi.org/10.1016/j.jep.2020.113064>
9. Swaffar DS, Holley CJ, Fitch RW, Elkin KR, Zhang C, Sturgill JP, Menachery MD. Phytochemical investigation and *in vitro* cytotoxic evaluation of alkaloids from *Abuta rufescens*. *Planta Med.* 2012; 78: 230-232. <https://doi.org/10.1055/s-0031-1280383>
10. Versiani MA, Kanwal A, Faizi S, and Farooq AD. Cytotoxic cardiac glycoside from the parasitic plant *Cuscuta reflexa*. *Chem Nat Compd.* 2017; 53: 915-922. <https://doi.org/10.1007/s10600-017-2154-5>
11. Raj Kapoor B, Murugesh N, Krishna DR. Cytotoxic activity of a flavanone from the stem of *Bauhinia variegata* Linn. *Nat. Prod. Res.* 2009; 23(15): 1384-1389. <https://doi.org/10.1080/14786410802553752>
12. Beutler JA, Kashman Y, Pannell LK, Cardellina JH, Alexander MRA, Balaschak MS, Prather TR, Shoemaker RH, and Boyda MR. Isolation and characterization of novel cytotoxic saponins from *Archidendron ellipticum*. *Bioorg Med Chem.* 1997; 5(8): 1509-1517. [https://doi.org/10.1016/S0968-0896\(97\)00098-9](https://doi.org/10.1016/S0968-0896(97)00098-9)
13. Banik K, Khatoon E, Harsha, C., Rana V, Parama D, Thakur KK, Bishayee A, Kunnumakkara, AB. Wogonin and its analogs for the prevention and treatment of cancer: A systematic review. *Phytother Res.* 2022; 36:1854-1883. doi: 10.1002/ptr.7386. <https://doi.org/10.1002/ptr.7386>
14. Nelson VK, Sahoo NK, Sahu M, Sudhan HH, Pullaiah CP, Muralikrishna KS. *In vitro* anticancer activity of *Eclipta alba* whole plant extract on colon cancer cell HCT-116. *BMC Complement Med Ther.* 2020; 20:355. <https://doi.org/10.1186/s12906-020-03118-9>



15. Tosun F, Beutler JA, Ransom TT, Miski M. Anadolycin, a highly potent and selective cytotoxic sesquiterpene coumarin from the root extract of *Heptaptera anatolica*. *Molecules*. 2019; 24: 1153-1160. <https://doi.org/10.3390/molecules24061153>
16. Tosun F, Aytaç EC, Beutler JA, Wilson JA, Miski M. Cytotoxic sesquiterpene coumarins from the roots of *Heptaptera cilicica*. *Rec Nat Prod*. 2021; 15(6): 529-536. <https://doi.org/10.25135/rnp.242.21.02.1990>
17. Devkota KP, Covell D, Ransom T, McMahon JB, Beutler JA. Growth inhibition of human colon carcinoma cells by sesquiterpenoids and tetralones of *Zygogynum calothyrsum*. *J Nat Prod*. 2013; 76: 710-714. <https://doi.org/10.1021/np400042q>
18. Badisa RB, Tzakou O, Couladis M and Pilarinou E. Cytotoxic activities of some Greek Labiatae Herbs. *Phytother Res*. 2003; 17: 472-476. <https://doi.org/10.1002/ptr.1175>
19. Dzhambazov B, Daskalova S, Monteveva A, and Popov N. In vitro screening for antitumour activity of *Clinopodium vulgare* L. (Lamiaceae) extracts. *Biol Pharm Bull*. 2002; 25(4): 499-504. <https://doi.org/10.1248/bpb.25.499>
20. Batsalova T, Bardarov K, Bardarov V, Moten D, Dzhambazov B. Cytotoxic properties of *Clinopodium vulgare* L. Extracts on selected human cell lines. *C.R. Acad Sci*. 2017; 70(5): 645-650.
21. Gallego A, Metón I, Baanante IV, Ouazzani J, Adelin E, Palazon J, Bonfill M, Moyano E. Viability-reducing activity of *Coryllusa vellana* L. extracts against human cancer cell lines. *Biomed Pharmacother*. 2017; 89: 565-572. <https://doi.org/10.1016/j.biopha.2017.02.046>
22. Bilir EK, Tutun H, Sevin S, Kismali G, Yarsan E. Cytotoxic Effects of *Rhododendron ponticum* L. Extract on prostate carcinoma and adenocarcinoma cell line (DU145, PC3). *Kafkas Univ Vet Fak Derg*. 2018; 24 (3): 451-457. <https://doi.org/10.9775/kvfd.2017.19219>
23. Demir S, Turan I, Aliyazicioglu Y. Cytotoxic effect of *Rhododendron luteum* leaf extract on human cancer cell lines. *KSÜ Tar Doğa Derg*. 2018; 21(6): 950-956. <https://doi.org/10.18016/ksutarimdog.vi.421283>
24. Mahomoodally MF, Sieniawska E, Sinan KI, Picot-Allaina MCN, Yerlikaya S, Baloglu MC, Altunoglu YC, Senkardes I, Rengasamy KRR, Zengin G. Utilisation of *Rhododendron luteum* sweet bioactive compounds as valuable source of enzymes inhibitors, antioxidant, and anticancer agents. *Food Chem Toxicol*. 2020; 135: 111052. <https://doi.org/10.1016/j.fct.2019.111052>
25. Şöhretoglu D, Genç Y, Harput ÜŞ, Sabuncuoğlu S, Soral M, Gulin Renda G, Liptaj T. Phytochemical Content, Antioxidant and Cytotoxic Activities of *Sedum spurium*. *Nat Prod Commun*. 2016; 11 (11):1693-1696. <https://doi.org/10.1177/1934578X1601101117>
26. Piwowarski JP, Bobrowska-Korczak B, Stanisławska I, Bielecki W, Wrzesien R, Granica S, Krupa K, Kiss AK. Evaluation of the Effect of *Epilobium angustifolium* Aqueous Extract on LNCaP Cell Proliferation in *In Vitro* and *In Vivo* Models. *Planta Med*. 2017. <https://doi.org/10.1055/s-0043-109372>
27. Stolarczyk M, Piwowarski JP, Granica S, Stefańska J, Naruszewicz M and Kiss AK. Extracts from *Epilobium* sp. Herbs, Their Components and Gut Microbiota Metabolites of *Epilobium* Ellagitannins, Urolithins, Inhibit Hormone-Dependent Prostate Cancer Cells (LNCaP) Proliferation and PSA Secretion. *Phytother Res*. 2013; 27: 1842-1848. <https://doi.org/10.1002/ptr.4941>
28. Nowak A, Zielonka-Brzezicka J, Peruzynska M and Klimowicz A. *Epilobium angustifolium* L. as a Potential Herbal Component of Topical Products for Skin Care and Treatment—A Review. *Molecules* 2022; 27: 3536. <https://doi.org/10.3390/molecules27113536>
29. Akar Z. Chemical compositions by using LC-MS/MS and GC-MS and antioxidant activities of methanolic extracts from leaf and flower parts of *Scabiosa columbaria* subsp. *columbaria* var. *columbaria* L. *Saudi J Biol Sci*. 2021; 28: 6639-6644. <https://doi.org/10.1016/j.sjbs.2021.07.039>
30. Dumlu MU, Gurkan E and Tuzlacı E. Chemical composition and antioxidant activity of *Campanula alliariifolia*. *Nat Prod Res*. 2008; 22(6): 477-482. <https://doi.org/10.1080/14786410701640429>
31. Yaylı N, Yıldırım N, Doğan N, Usta A and Altun L. Triterpenes from *Campanula lactiflora*. *J Asian Nat Prod Res*. 2005; 7(5): 771-775. <https://doi.org/10.1080/1028602042000324907>
32. Celikler Ozer O, Erdogan Orhan I, Caliskan B, Senol Deniz FS, Gokbulut A, Gur Maz T, Aysal A, Emerce E, Shekfeh S, Kahraman A, Banoglu E. Exploration of anti-tyrosinase effect of *Geranium glaberrimum* Boiss. & Heldr. with *in silico* approach and survey of 21 *Geranium* species. *J Herb Med*. 2021; 27: 100431. <https://doi.org/10.1016/j.hermed.2021.100431>

33. Bastian F, Ito Y, Ogahara E, Ganeko N, Hatano T and Ito H. Simultaneous Quantification of Ellagitannins and Related Polyphenols in *Geranium thunbergii* Using Quantitative NMR. *Molecules*. 2018; 23: 1346. <https://doi.org/10.3390/molecules23061346>
34. Karadağ AE., Tosun F. *In vitro* Antimicrobial and Antioxidant Evaluation of *Melampyrum arvense* var. *elatius* L. and *Sedum spurium* M. Bieb. Extracts. *Acta Pharm Sci*. 2019; 57(2): 193-201. <https://doi.org/10.23893/1307-2080.APS.05712>

This is an open access article which is publicly available on our journal's website under Institutional Repository at <http://dspace.marmara.edu.tr>.