



Weighted Tribonacci Sums

Kunle Adegoke¹

¹Department of Physics and Engineering Physics, Obafemi Awolowo University, 220005 Ile-Ife, Nigeria

Abstract

We derive various weighted summation identities, including binomial and double binomial identities, for Tribonacci numbers. Our results contain some previously known results as special cases.

Keywords: binomial summation identity; generating function; partial sum, Tribonacci number; weighted sum

2010 Mathematics Subject Classification: 11B37, 11B39, 65B10

1. Introduction

For $m \geq 3$, the Tribonacci numbers are defined by

$$T_m = T_{m-1} + T_{m-2} + T_{m-3}, \quad T_0 = 0, T_1 = T_2 = 1. \tag{1.1}$$

By writing $T_{m-1} = T_{m-2} + T_{m-3} + T_{m-4}$ and eliminating T_{m-2} and T_{m-3} between this recurrence relation and the recurrence relation (1.1), a useful alternative recurrence relation is obtained for $m \geq 4$:

$$T_m = 2T_{m-1} - T_{m-4}, \quad T_0 = 0, \quad T_1 = T_2 = 1, \quad T_3 = 2. \tag{1.2}$$

Extension of the definition of T_m to negative subscripts is provided by writing the recurrence relation (1.2) as

$$T_{-m} = 2T_{-m+3} - T_{-m+4}. \tag{1.3}$$

Anantakitpaisal and Kuhapatanakul [2] proved that

$$T_{-m} = T_{m-1}^2 - T_{m-2}T_m. \tag{1.4}$$

The following identity (Feng [3], equation (3.3); Shah [7], (ii)) is readily established by the principle of mathematical induction:

$$T_{m+r} = T_r T_{m-2} + (T_{r-1} + T_r)T_{m-1} + T_{r+1}T_m. \tag{1.5}$$

Irmak and Alp [5] derived the following identity for Tribonacci numbers with indices in arithmetic progression:

$$T_{tm+r} = \lambda_1(t)T_{t(m-1)+r} + \lambda_2(t)T_{t(m-2)+r} + \lambda_3(t)T_{t(m-3)+r}, \tag{1.6}$$

where,

$$\lambda_1(t) = \alpha^t + \beta^t + \gamma^t, \quad \lambda_2(t) = -(\alpha\beta)^t - (\alpha\gamma)^t - (\beta\gamma)^t, \quad \lambda_3(t) = (\alpha\beta\gamma)^t,$$

where α, β and γ are the roots of the characteristic polynomial of the Tribonacci sequence $x^3 - x^2 - x - 1$. Thus,

$$\alpha = \frac{1}{3} \left(1 + \sqrt[3]{19 + 3\sqrt{33}} + \sqrt[3]{19 - 3\sqrt{33}} \right),$$

$$\beta = \frac{1}{3} \left(1 + \omega \sqrt[3]{19 + 3\sqrt{33}} + \omega^2 \sqrt[3]{19 - 3\sqrt{33}} \right)$$

and

$$\gamma = \frac{1}{3} \left(1 + \omega^2 \sqrt[3]{19 + 3\sqrt{33}} + \omega \sqrt[3]{19 - 3\sqrt{33}} \right),$$

where $\omega = \exp(2i\pi/3)$ is a primitive cube root of unity. Note that $\lambda_1(t), \lambda_2(t)$ and $\lambda_3(t)$ are integers for any positive integer t [5]; in particular, $\lambda_1(1) = 1 = \lambda_2(1) = \lambda_3(1)$.

2. Weighted sums

Lemma 2.1 ([1], Lemma 2). Let $\{X_m\}$ be any arbitrary sequence, where X_m , $m \in \mathbb{Z}$, satisfies a three term recurrence relation $X_m = f_1 X_{m-a} + f_2 X_{m-b}$, where f_1 and f_2 are arbitrary non-vanishing complex functions, not dependent on m , and a and b are integers. Then,

$$f_2 \sum_{j=0}^k \frac{X_{m-ka-b+aj}}{f_1^j} = \frac{X_m}{f_1^k} - f_1 X_{m-(k+1)a}, \quad (2.1)$$

$$f_1 \sum_{j=0}^k \frac{X_{m-kb-a+bj}}{f_2^j} = \frac{X_m}{f_2^k} - f_2 X_{m-(k+1)b} \quad (2.2)$$

and

$$\sum_{j=0}^k \frac{X_{m-(b-a)k+a+(b-a)j}}{(-f_2/f_1)^j} = \frac{f_1 X_m}{(-f_2/f_1)^k} + f_2 X_{m-(k+1)(b-a)}. \quad (2.3)$$

for k a non-negative integer.

Theorem 2.2. The following identities hold for any integers m and k :

$$\sum_{j=0}^k 2^{-j} T_{m-k-4+j} = 2T_{m-k-1} - 2^{-k} T_m, \quad (2.4)$$

$$2 \sum_{j=0}^k (-1)^j T_{m-4k-1+4j} = (-1)^k T_m + T_{m-4k-4} \quad (2.5)$$

and

$$\sum_{j=0}^k 2^j T_{m-3k+1+3j} = 2^{k+1} T_m - T_{m-3k-3}. \quad (2.6)$$

Proof. From the recurrence relation (1.2), make the identifications $f_1 = 2$, $f_2 = -1$, $a = 1$ and $b = 4$ and use these in Lemma 2.1 with $X = T$. \square

Particular instances of identities (2.4)–(2.6) are the following identities:

$$\sum_{j=0}^k 2^{-j} T_j = 4 - 2^{-k} T_{k+4}, \quad (2.7)$$

giving,

$$\sum_{j=0}^{\infty} 2^{-j} T_j = 4, \quad (2.8)$$

and

$$2 \sum_{j=0}^k (-1)^j T_{4j} = (-1)^k T_{4k+1} - 1 \quad (2.9)$$

and

$$\sum_{j=0}^k 2^j T_{3j} = 2^{k+1} T_{3k-1}. \quad (2.10)$$

Lemma 2.3 (Partial sum of an n^{th} order sequence). Let $\{X_j\}$ be any arbitrary sequence, where X_j , $j \in \mathbb{Z}$, satisfies a $n+1$ term recurrence relation $X_j = f_1 X_{j-c_1} + f_2 X_{j-c_2} + \dots + f_n X_{j-c_n} = \sum_{m=1}^n f_m X_{j-c_m}$, where f_1, f_2, \dots, f_n are arbitrary non-vanishing complex functions, not dependent on j , and c_1, c_2, \dots, c_n are fixed integers. Then, the following summation identity holds for arbitrary x and non-negative integer k :

$$\sum_{j=0}^k x^j X_j = \frac{\sum_{m=1}^n x^{c_m} f_m \left(\sum_{j=1}^{c_m} x^{-j} X_{-j} - \sum_{j=k-c_m+1}^k x^j X_j \right)}{1 - \sum_{m=1}^n x^{c_m} f_m}.$$

Proof. Recurrence relation:

$$X_j = \sum_{m=1}^n f_m X_{j-c_m}.$$

We multiply both sides by x^j and sum over j to obtain

$$\sum_{j=0}^k x^j X_j = \sum_{m=1}^n \left(f_m \sum_{j=0}^k x^j X_{j-c_m} \right) = \sum_{m=1}^n \left(x^{c_m} f_m \sum_{j=-c_m}^{k-c_m} x^j X_j \right),$$

after shifting the summation index j . Splitting the inner sum, we can write

$$\sum_{j=0}^k x^j X_j = \sum_{m=1}^n x^{c_m} f_m \left(\sum_{j=-c_m}^{-1} x^j X_j + \sum_{j=0}^k x^j X_j + \sum_{j=k+1}^{k-c_m} x^j X_j \right).$$

Since

$$\sum_{j=-c_m}^{-1} x^j X_j \equiv \sum_{j=1}^{c_m} x^{-j} X_{-j} \text{ and } \sum_{j=k+1}^{k-c_m} x^j X_j \equiv - \sum_{j=k-c_m+1}^k x^j X_j,$$

the preceding identity can be written

$$\sum_{j=0}^k x^j X_j = \sum_{m=1}^n x^{c_m} f_m \left(\sum_{j=1}^{c_m} x^{-j} X_{-j} + \sum_{j=0}^k x^j X_j - \sum_{j=k-c_m+1}^k x^j X_j \right).$$

Thus, we have

$$S = \sum_{m=1}^n x^{c_m} f_m \left(\sum_{j=1}^{c_m} x^{-j} X_{-j} + S - \sum_{j=k-c_m+1}^k x^j X_j \right),$$

where

$$S = S_k(x) = \sum_{j=0}^k x^j X_j.$$

Removing brackets, we have

$$S = \sum_{m=1}^n x^{c_m} f_m \left(\sum_{j=1}^{c_m} x^{-j} X_{-j} - \sum_{j=k-c_m+1}^k x^j X_j \right) + S \sum_{m=1}^n x^{c_m} f_m,$$

from which the result follows by grouping the S terms. □

Lemma 2.4 (Generating function). *Under the conditions of Lemma 2.3, if additionally $x^k X_k$ vanishes in the limit as k approaches infinity, then*

$$S_\infty(x) = \sum_{j=0}^\infty x^j X_j = \frac{\sum_{m=1}^n \left(x^{c_m} f_m \sum_{j=1}^{c_m} x^{-j} X_{-j} \right)}{1 - \sum_{m=1}^n x^{c_m} f_m},$$

so that $S_\infty(x)$ is a generating function for the sequence $\{X_j\}$.

Theorem 2.5 (Sum of Tribonacci numbers with indices in arithmetometric progression). *For arbitrary x , any integers t and r and any non-negative integer k , the following identity holds:*

$$\begin{aligned} \left(1 - \lambda_1(t)x - \lambda_2(t)x^2 - \lambda_3(t)x^3 \right) \sum_{j=0}^k x^j T_{t+j+r} &= T_r + (x\lambda_2(t) + x^2\lambda_3(t))T_{r-t} + x\lambda_3(t)T_{r-2t} - x^{k+1}T_{(k+1)t+r} - x^{k+2}(\lambda_2(t) + x\lambda_3(t))T_{kt+r} \\ &\quad - x^{k+2}\lambda_3(t)T_{(k-1)t+r}, \end{aligned}$$

where,

$$\lambda_1(t) = \alpha^t + \beta^t + \gamma^t, \quad \lambda_2(t) = -(\alpha\beta)^t - (\alpha\gamma)^t - (\beta\gamma)^t, \quad \lambda_3(t) = (\alpha\beta\gamma)^t,$$

where α , β and γ are the roots of the characteristic polynomial of the Tribonacci sequence $x^3 - x^2 - x - 1$.

Proof. Write identity (1.6) as $X_j = f_1 X_{j-1} + f_2 X_{j-2} + f_3 X_{j-3}$ and identify the sequence $\{X_j\} = \{T_{t+j+r}\}$ and the constants $c_1 = 1$, $c_2 = 2$, $c_3 = 3$ and the functions $f_1 = \lambda_1(t)$, $f_2 = \lambda_2(t)$, $f_3 = \lambda_3(t)$, and use these in Lemma 2.3. □

Corollary 2.6 (Generating function of the Tribonacci numbers with indices in arithmetical progression). *For any integers t and r , any non-negative integer k and arbitrary x for which $x^k T_k$ vanishes as k approaches infinity, the following identity holds:*

$$\sum_{j=0}^{\infty} x^j T_{tj+r} = \frac{T_r + (x\lambda_2 + x^2\lambda_3)T_{r-t} + x\lambda_3 T_{r-2t}}{1 - \lambda_1 x - \lambda_2 x^2 - \lambda_3 x^3},$$

where,

$$\lambda_1 = \alpha^t + \beta^t + \gamma^t, \quad \lambda_2 = -(\alpha\beta)^t - (\alpha\gamma)^t - (\beta\gamma)^t, \quad \lambda_3 = (\alpha\beta\gamma)^t,$$

where α, β and γ are the roots of the characteristic polynomial of the Tribonacci sequence $x^3 - x^2 - x - 1$.

Many instances of Theorem 2.5 may be explored. In particular, we have

$$(\lambda_1(t) + \lambda_2(t) + \lambda_3(t) - 1) \sum_{j=0}^k T_{tj+r} = -T_r - (\lambda_2(t) + \lambda_3(t))T_{r-t} - \lambda_3(t)T_{r-2t} + T_{(k+1)t+r} + (\lambda_2(t) + \lambda_3(t))T_{kt+r} + \lambda_3(t)T_{(k-1)t+r}, \tag{2.11}$$

which at $r = 0$ gives

$$(\lambda_1(t) + \lambda_2(t) + \lambda_3(t) - 1) \sum_{j=0}^k T_{tj} = -(\lambda_2(t) + \lambda_3(t))(T_{t-1}^2 - T_{t-2}T_t) - \lambda_3(t)(T_{2t-1}^2 - T_{2t-2}T_{2t}) + T_{(k+1)t} + (\lambda_2(t) + \lambda_3(t))T_{kt} + \lambda_3(t)T_{(k-1)t}; \tag{2.12}$$

and

$$(1 + \lambda_1(t) - \lambda_2(t) + \lambda_3(t)) \sum_{j=0}^k (-1)^j T_{tj+r} = T_r + (\lambda_3(t) - \lambda_2(t))T_{r-t} - \lambda_3(t)T_{r-2t} + (-1)^k T_{(k+1)t+r} + (-1)^k (\lambda_3(t) - \lambda_2(t))T_{kt+r} - (-1)^k \lambda_3(t)T_{(k-1)t+r}, \tag{2.13}$$

which at $r = 0$ gives

$$(1 + \lambda_1(t) - \lambda_2(t) + \lambda_3(t)) \sum_{j=0}^k (-1)^j T_{tj} = (\lambda_3(t) - \lambda_2(t))(T_{t-1}^2 - T_{t-2}T_t) - \lambda_3(t)(T_{2t-1}^2 - T_{2t-2}T_{2t}) + (-1)^k T_{(k+1)t} + (-1)^k (\lambda_3(t) - \lambda_2(t))T_{kt} - (-1)^k \lambda_3(t)T_{(k-1)t}. \tag{2.14}$$

Many previously known results are particular instances of the identities (2.11) and (2.13). For example, Theorem 5 of [6] is obtained from identity (2.12) by setting $t = 4$. Sums of Tribonacci numbers with indices in arithmetical progression are also discussed in references [4, 5, 6] and references therein, using various techniques.

Weighted sums of the form $\sum_{j=0}^k j^p T_{tj+r}$, where p is a non-negative integer, may be evaluated by setting $x = e^y$ in the identity of Theorem 2.5, differentiating both sides p times with respect to y and then setting $y = 0$. The simplest examples in this category are the following:

$$2 \sum_{j=0}^k j T_{tj+r} = -T_{r-2} + 3T_{r+1} + (k-1)T_{k+r-1} + (2k-1)T_{k+r} + (k-2)T_{k+r+1} \tag{2.15}$$

and

$$2 \sum_{j=0}^k j^2 T_{tj+r} = -3T_{r-1} - 5T_r - 6T_{r+1} + (k^2 - 2k + 3)T_{k+r-1} + (2k^2 - 2k + 5)T_{k+r} + (k^2 - 4k + 6)T_{k+r+1}, \tag{2.16}$$

with the particular cases

$$2 \sum_{j=0}^k j T_j = 2 + (k-1)T_{k-1} + (2k-1)T_k + (k-2)T_{k+1} \tag{2.17}$$

and

$$2 \sum_{j=0}^k j^2 T_j = -6 + (k^2 - 2k + 3)T_{k+r-1} + (2k^2 - 2k + 5)T_k + (k^2 - 4k + 6)T_{k+1}. \tag{2.18}$$

3. Weighted binomial sums

Lemma 3.1 ([1], Lemma 3). Let $\{X_m\}$ be any arbitrary sequence. Let X_m , $m \in \mathbb{Z}$, satisfy a three term recurrence relation $X_m = f_1 X_{m-a} + f_2 X_{m-b}$, where f_1 and f_2 are non-vanishing complex functions, not dependent on m , and a and b are integers. Then,

$$\sum_{j=0}^k \binom{k}{j} \left(\frac{f_1}{f_2}\right)^j X_{m-bk+(b-a)j} = \frac{X_m}{f_2^k}, \quad (3.1)$$

$$\sum_{j=0}^k \binom{k}{j} \frac{X_{m+(a-b)k+bj}}{(-f_2)^j} = \left(-\frac{f_1}{f_2}\right)^k X_m \quad (3.2)$$

and

$$\sum_{j=0}^k \binom{k}{j} \frac{X_{m+(b-a)k+aj}}{(-f_1)^j} = \left(-\frac{f_2}{f_1}\right)^k X_m, \quad (3.3)$$

for k a non-negative integer.

Theorem 3.2. The following identities hold for any integer m and any non-negative integer k :

$$\sum_{j=0}^k (-1)^j \binom{k}{j} 2^j T_{m-4k+3j} = (-1)^k T_m, \quad (3.4)$$

$$\sum_{j=0}^k \binom{k}{j} T_{m-3k+4j} = 2^k T_m \quad (3.5)$$

and

$$\sum_{j=0}^k (-1)^j \binom{k}{j} 2^{-j} T_{m+3k+j} = 2^{-k} T_m. \quad (3.6)$$

Proof. Identify $X = T$ in Lemma 3.1 and use the f_1 , f_2 , a and b values found in the proof of Theorem 2.2. □

Particular cases of (3.4), (3.5) and (3.6) are the following identities:

$$\sum_{j=0}^k (-1)^j \binom{k}{j} 2^j T_{3j} = (-1)^k T_{4k}, \quad (3.7)$$

$$\sum_{j=0}^k \binom{k}{j} T_{4j} = 2^k T_{3k} \quad (3.8)$$

and

$$\sum_{j=0}^k (-1)^j \binom{k}{j} 2^{-j} T_j = 2^{-k} (T_{3k-1}^2 - T_{3k-2} T_{3k}). \quad (3.9)$$

4. Weighted double binomial sums

Lemma 4.1. Let $\{X_m\}$ be any arbitrary sequence, X_m satisfying a four term recurrence relation $X_m = f_1 X_{m-a} + f_2 X_{m-b} + f_3 X_{m-c}$, where f_1 , f_2 and f_3 are arbitrary nonvanishing functions and a , b and c are integers. Then, the following identities hold:

$$\sum_{j=0}^k \sum_{s=0}^j \binom{k}{j} \binom{j}{s} \left(\frac{f_2}{f_3}\right)^j \left(\frac{f_1}{f_2}\right)^s X_{m-ck+(c-b)j+(b-a)s} = \frac{X_m}{f_3^k}, \quad (4.1)$$

$$\sum_{j=0}^k \sum_{s=0}^j \binom{k}{j} \binom{j}{s} \left(\frac{f_3}{f_2}\right)^j \left(\frac{f_1}{f_3}\right)^s X_{m-bk+(b-c)j+(c-a)s} = \frac{X_m}{f_2^k}, \quad (4.2)$$

$$\sum_{j=0}^k \sum_{s=0}^j \binom{k}{j} \binom{j}{s} \left(\frac{f_3}{f_1}\right)^j \left(\frac{f_2}{f_3}\right)^s X_{m-ak+(a-c)j+(c-b)s} = \frac{X_m}{f_1^k}, \quad (4.3)$$

$$\sum_{j=0}^k \sum_{s=0}^j \binom{k}{j} \binom{j}{s} \left(\frac{f_2}{f_3}\right)^j \left(-\frac{1}{f_2}\right)^s X_{m-(c-a)k+(c-b)j+bs} = \left(-\frac{f_1}{f_3}\right)^k X_m, \quad (4.4)$$

$$\sum_{j=0}^k \sum_{s=0}^j \binom{k}{j} \binom{j}{s} \left(\frac{f_1}{f_3}\right)^j \left(-\frac{1}{f_1}\right)^s X_{m-(c-b)k+(c-a)j+as} = \left(-\frac{f_2}{f_3}\right)^k X_m, \quad (4.5)$$

and

$$\sum_{j=0}^k \sum_{s=0}^j \binom{k}{j} \binom{j}{s} \left(\frac{f_1}{f_2}\right)^j \left(-\frac{1}{f_1}\right)^s X_{m-(b-c)k+(b-a)j+as} = \left(-\frac{f_3}{f_2}\right)^k X_m. \quad (4.6)$$

Proof. Only identity (4.1) needs to be proved as identities (4.2)–(4.6) are obtained from (4.1) by re-arranging the recurrence relation. The proof of (4.1) is by induction on k , similar to the proof of Lemma 3 of [1]. \square

Theorem 4.2. *The following identities hold for non-negative integer k , integer m and integer $r \notin \{-17, -4, -1, 0\}$:*

$$\sum_{j=0}^k \sum_{s=0}^j \binom{k}{j} \binom{j}{s} (T_{r-1} + T_r)^{j-s} \frac{T_{r+1}^s}{T_r^j} T_{m-(r+2)k+j+s} = \frac{T_m}{T_r^k}, \quad (4.7)$$

$$\sum_{j=0}^k \sum_{s=0}^j \binom{k}{j} \binom{j}{s} \frac{T_r^{j-s} T_{r+1}^s}{(T_{r-1} + T_r)^j} T_{m-(r+1)k-j+2s} = \frac{T_m}{(T_{r-1} + T_r)^k}, \quad (4.8)$$

$$\sum_{j=0}^k \sum_{s=0}^j \binom{k}{j} \binom{j}{s} \frac{T_{r-1}^{j-s} (T_{r-2} + T_{r-1})^s}{T_r^j} T_{m-(r-1)k-2j+s} = \frac{T_m}{T_r^k}, \quad (4.9)$$

$$\sum_{j=0}^k \sum_{s=0}^j (-1)^s \binom{k}{j} \binom{j}{s} \frac{(T_{r-1} + T_r)^{j-s}}{T_r^j} T_{m-2k+j+(r+1)s} = (-1)^k \left(\frac{T_{r+1}}{T_r}\right)^k T_m, \quad (4.10)$$

$$\sum_{j=0}^k \sum_{s=0}^j (-1)^s \binom{k}{j} \binom{j}{s} \frac{T_{r+1}^{j-s}}{T_r^j} T_{m-k+2j+rs} = (-1)^k \left(\frac{T_{r-1} + T_r}{T_r}\right)^k T_m \quad (4.11)$$

and

$$\sum_{j=0}^k \sum_{s=0}^j (-1)^s \binom{k}{j} \binom{j}{s} \frac{T_{r+1}^{j-s}}{(T_{r-1} + T_r)^j} T_{m+k+j+rs} = (-1)^k \left(\frac{T_r}{T_{r-1} + T_r}\right)^k T_m. \quad (4.12)$$

Proof. Write the identity (1.5) as $T_m = T_r T_{m-r-2} + (T_{r-1} + T_r) T_{m-r-1} + T_{r+1} T_{m-r}$, identify $f_1 = T_r$, $f_2 = T_{r-1} + T_r$, $f_3 = T_{r+1}$, $a = r + 2$, $b = r + 1$, $c = r$ and use these in Lemma 4.1 with $X = T$. \square

Acknowledgement

The author thanks the anonymous referee whose comments helped improve the presentation.

References

- [1] K. Adegoke, Weighted sums of some second-order sequences, *arXiv:1803.09054[math.NT]* (2018).
- [2] P. Anantakitpaisal and K. Kuhapatanakul, Reciprocal sums of the Tribonacci numbers, *Journal of Integer sequences* **19** (2016), 1–9.
- [3] J. Feng, More identities on the Tribonacci numbers, *Ars Combinatorial* **C** (2011), 73–78.
- [4] R. Frontczak, Sums of Tribonacci and Tribonacci-Lucas numbers, *International Journal of Mathematical Analysis* **12:1** (2018), 19–24.
- [5] N. Irmak and M. Alp, Tribonacci numbers with indices in arithmetic progression and their sums, *Miskolc Mathematical Notes* **14:1** (2013), 125–133.
- [6] E. Kilic, Tribonacci sequences with certain indices and their sums, *Ars Combinatorial* **86** (2008), 13–22.
- [7] D. V. Shah, Some Tribonacci identities, *Mathematics Today* **27** (2011), 1–9.