



GENERALIZATION OF $(\alpha - F_d)$ -CONTRACTION ON QUASI-METRIC SPACE

ÖZLEM ACAR

ABSTRACT. In this paper, we introduce the concept of generalized $(\alpha - F_d)$ -contraction and give some fixed point results in quasi metric spaces with different types of completeness.

1. INTRODUCTION

The beginning of metrical fixed point theory is related to the Banach's Contraction Principle, published in 1922. Banach's Contraction Principle says that, whenever (X, d) is complete, then any contraction self-map of X has a unique fixed point. This fixed point result is one of the most powerful tools for many existence and uniqueness problems arising in mathematics. Because of its importance, Banach Contraction Principle has been extended and generalized in many ways. Among all these, an interesting generalization was given by Wardowski [11] using a new concept F -contraction. Then many authors gave some results using this concept in different type metric space.

In this paper, we introduce the concept of generalized $(\alpha - F_d)$ -contraction and give some fixed point results in quasi metric spaces with different types of completeness using this concept.

2. PRELIMINARIES

In this section, we give some basic definitions and properties about quasi metric space.

Definition 1. *Let X be any nonempty set. A function $d : X \times X \rightarrow \mathbb{R}_+$ is said to be a quasi-pseudo metric if and only if for all $x, y \in X$ the following conditions are satisfied:*

$$(i) \quad d(x, x) = 0,$$

Received by the editors: May 31, 2017, Accepted: September 25, 2017.

2010 *Mathematics Subject Classification.* 54H25, 47H10.

Key words and phrases. Quasi-metric space, F -contraction, α -admissible.

©2018 Ankara University
Communications Faculty of Sciences University of Ankara-Series A1 Mathematics and Statistics.
Communications de la Faculté des Sciences de l'Université d'Ankara. Série A1. Mathématiques et Statistiques.

$$(ii) \quad d(x, y) \leq d(x, z) + d(z, y).$$

If a quasi-pseudo metric d satisfies:

$$\bullet \quad d(x, y) = d(y, z) = 0 \Rightarrow x = y$$

then d is said to be quasi metric.

Let d be a quasi metric. If d satisfies

$$d(x, y) = 0 \Rightarrow x = y$$

then d is said to be T_1 -quasi metric.

It is clear that, every metric is a T_1 -quasi metric, every T_1 -quasi metric is a quasi metric and every quasi metric is a quasi-pseudo metric.

Let (X, d) be a quasi-pseudo metric space. Given a point $x_0 \in X$ and a real constant $\varepsilon > 0$, the sets

$$B_d(x_0, \varepsilon) = \{y \in X : d(x_0, y) < \varepsilon\}$$

and

$$B_d[x_0, \varepsilon] = \{y \in X : d(x_0, y) \leq \varepsilon\}$$

are called open ball and closed ball with center x_0 and radius ε .

Each quasi-pseudo metric d on X generates a topology τ_d on X which has a base the family of open balls

$$\{B_d(x, \varepsilon) : x \in X \text{ and } \varepsilon > 0\}.$$

The closure and interior of a subset A of X with respect to τ_d is denoted by $cl_d(A)$ and $int_d(A)$, respectively.

If d is a quasi-pseudo metric on X , then τ_d is a T_0 topology and if d is a T_1 -quasi-metric, then τ_d is a T_1 topology on X .

If d is a quasi metric and τ_d is T_1 topology, then d is T_1 -quasi-metric.

If d is a quasi-pseudo metric on X , then the functions d^{-1} , d^s and d_+ defined by

$$d^{-1}(x, y) = d(y, x)$$

$$d^s(x, y) = \max \{d(x, y), d^{-1}(x, y)\}$$

and

$$d_+(x, y) = d(x, y) + d^{-1}(x, y)$$

are also quasi-pseudo metric on X . If d is a quasi metric, then d^s and d_+ are (equivalent) metrics on X .

Definition 2. Let (X, d) be a quasi metric space and $x \in X$. The convergence of a sequence $\{x_n\}$ to x with respect to τ_d called d -convergence and denoted by $x_n \xrightarrow{d} x$, is defined

$$x_n \xrightarrow{d} x \Leftrightarrow d(x, x_n) \rightarrow 0.$$

Similarly the convergence of a sequence $\{x_n\}$ to x with respect to $\tau_{d^{-1}}$ called d^{-1} -convergence and denoted by $x_n \xrightarrow{d^{-1}} x$, is defined

$$x_n \xrightarrow{d^{-1}} x \Leftrightarrow d(x_n, x) \rightarrow 0.$$

Finally the convergence of a sequence $\{x_n\}$ to x with respect to τ_{d^s} called d^s -convergence and denoted by $x_n \xrightarrow{d^s} x$, is defined

$$x_n \xrightarrow{d^s} x \Leftrightarrow d^s(x_n, x) \rightarrow 0.$$

It is clear that $x_n \xrightarrow{d^s} x \Leftrightarrow x_n \xrightarrow{d} x$ and $x_n \xrightarrow{d^{-1}} x$.

Definition 3. [8] Let (X, d) be a quasi metric space. A sequence $\{x_n\}$ in X is called

- Left K -Cauchy (or forward Cauchy) if for every $\varepsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that

$$\forall n, k, n \geq k \geq n_0, d(x_k, x_n) < \varepsilon,$$

- Right K -Cauchy (or backward Cauchy) if for every $\varepsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that

$$\forall n, k, n \geq k \geq n_0, d(x_n, x_k) < \varepsilon,$$

- d^s -Cauchy if for every $\varepsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that

$$\forall n, k \geq n_0, d(x_n, x_k) < \varepsilon.$$

It is clear that $\{x_n\}$ is d^s -Cauchy if and only if it is both left K -Cauchy and right K -Cauchy. If a sequence is left K -Cauchy with respect to d , then it is right K -Cauchy with respect to d^{-1} . If $\{x_n\}$ is a sequence in a quasi metric space (X, d) such that

$$\sum_{n=1}^{\infty} d(x_n, x_{n+1}) < \infty,$$

then it is left K -Cauchy sequence.

It is well known that every convergent sequence is Cauchy sequence in a metric space. In general this situation is not valid in a quasi metric space. That is, d -convergent or d^{-1} -convergent sequences may not be Cauchy (in the sense of d^s , left K and right K) in a quasi metric space.

Definition 4. Let (X, d) be a quasi metric space. Then (X, d) is said to be

- bicomplete if every d^s -Cauchy sequence is d^s -convergent,
- left (right) K -complete if every left (right) K -Cauchy sequence is d -convergent,
- left (right) \mathcal{M} -complete if every left (right) K -Cauchy sequence is d^{-1} -convergent,
- left (right) Smyth complete if every left (right) K -Cauchy sequence is d^s -convergent.

One can find more information about important properties of quasi metric space in [8, 9].

On the other hand, α -admissibility and F -contractivity of a mapping are popular in recent years. The concept of α -admissibility of a mapping on a nonempty set has been introduced by Samet et. al. [10]. Let X be a nonempty set, T be a self mapping of X and $\alpha : X \times X \rightarrow [0, \infty)$ be a function. Then T is said to be α -admissible if

$$\alpha(x, y) \geq 1 \Rightarrow \alpha(Tx, Ty) \geq 1.$$

Using this concept, Samet et al. [10] provided some general fixed point results including many known theorems on complete metric spaces.

In 2012, Wardowski [11] introduced the concept of F -contraction. Let \mathcal{F} be the family of all functions $F : (0, \infty) \rightarrow \mathbb{R}$ satisfying the following conditions:

- (F1) F is strictly increasing, i.e., for all $\alpha, \beta \in (0, \infty)$ such that $\alpha < \beta$, $F(\alpha) < F(\beta)$,
(F2) for each sequence $\{a_n\}$ of positive numbers

$$\lim_{n \rightarrow \infty} a_n = 0 \text{ if and only if } \lim_{n \rightarrow \infty} F(a_n) = -\infty,$$

- (F3) there exists $k \in (0, 1)$ such that $\lim_{\alpha \rightarrow 0^+} \alpha^k F(\alpha) = 0$.

Definition 5. Let (X, d) be a metric space and $T : X \rightarrow X$ be a mapping. Then T is said to be an F -contraction if $F \in \mathcal{F}$ and there exists $\tau > 0$ such that

$$\forall x, y \in X [d(Tx, Ty) > 0 \Rightarrow \tau + F(d(Tx, Ty)) \leq F(d(x, y))].$$

Various fixed point results for α -admissible mappings and F -contractions on complete metric space can be found in [1, 3, 4, 5] and [1, 2, 3, 7, 12], respectively.

3. MAIN RESULTS

Let (X, d) be a quasi metric space, $T : X \rightarrow X$ be a mapping and $\alpha : X \times X \rightarrow [0, \infty)$ be a function. We will consider the following set (see [1]):

$$T_\alpha = \{(x, y) \in X \times X : \alpha(x, y) \geq 1 \text{ and } d(Tx, Ty) > 0\}. \quad (3.1)$$

Definition 6. Let (X, d) be a quasi metric space and $T : X \rightarrow X$ be a mapping satisfying

$$d(x, y) = 0 \Rightarrow d(Tx, Ty) = 0. \quad (3.2)$$

$\alpha : X \times X \rightarrow [0, \infty)$ and $F \in \mathcal{F}$ be functions. Then T is said to be a generalized $(\alpha - F_d)$ -contraction if there exists $\tau > 0$ such that

$$\tau + F(d(Tx, Ty)) \leq F(M(x, y)) \quad (3.3)$$

for all $x, y \in T_\alpha$ where

$$M(x, y) = \max \left\{ d(x, y), d(x, Tx), d(y, Ty), \frac{1}{2} [d(x, Ty) + d(y, Tx)] \right\}.$$

Theorem 1. *Let (X, d) be a Hausdorff left K -complete quasi metric space, $T : X \rightarrow X$ be a generalized $(\alpha - F_d)$ -contraction. Assume that T is α -admissible and τ_d -continuous. If there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$, then T has a fixed point in X .*

Proof. Let $x_0 \in X$ be such that $\alpha(x_0, Tx_0) \geq 1$. Define a sequence $\{x_n\}$ in X by $x_n = Tx_{n-1}$ for all $n \in \mathbb{N}$. Since T is α -admissible, then $\alpha(x_n, x_{n+1}) \geq 1$ for all $n \in \mathbb{N}$. Now let

$$d_n = d(x_n, x_{n+1}) \quad (3.4)$$

for all $n \in \mathbb{N}$. If there exists $k \in \mathbb{N}$ with $d_k = d(x_k, x_{k+1}) = 0$ then x_k is a fixed point of T since d is T_1 quasi metric space. Suppose $d_n > 0$ for all $n \in \mathbb{N}$. Since T is generalized $(\alpha - F_d)$ -contraction, we get

$$\tau + F(d(x_n, x_{n+1})) \leq F(M(x_{n-1}, x_n)) \quad (3.5)$$

where

$$\begin{aligned} & M(x_{n-1}, x_n) \\ = & \max \left\{ d(x_{n-1}, x_n), d(x_{n-1}, x_n), d(x_n, x_{n+1}), \frac{1}{2} [d(x_n, x_{n+2}) + d(x_{n+1}, x_{n+1})] \right\} \\ = & \max \{d(x_{n-1}, x_n), d(x_n, x_{n+1})\}. \end{aligned}$$

So, we have

$$\begin{aligned} \tau + F(d(x_n, x_{n+1})) & \leq F(\max \{d(x_{n-1}, x_n), d(x_n, x_{n+1})\}) \\ & \leq F(\max \{d(x_{n-1}, x_n), d(x_n, x_{n+1})\}). \end{aligned}$$

If $\max \{d(x_{n-1}, x_n), d(x_n, x_{n+1})\} = d(x_n, x_{n+1})$, then

$$\tau + F(d(x_n, x_{n+1})) \leq F(d(x_n, x_{n+1}))$$

which is a contradiction. Hence, $\max \{d(x_{n-1}, x_n), d(x_n, x_{n+1})\} = d(x_{n-1}, x_n)$, then we obtain

$$\tau + F(d_n) \leq F(d_{n-1}).$$

Continuing this process, we get

$$F(d_n) \leq F(d_{n-1}) - \tau \leq F(d_{n-2}) - 2\tau \leq \dots \leq F(d_0) - n\tau. \quad (3.6)$$

From 3.6, we get $\lim_{n \rightarrow \infty} F(d_n) = -\infty$. Thus from (F2), we get

$$\lim_{n \rightarrow \infty} d_n = 0.$$

From (F3) there exists $k \in (0, 1)$ such that

$$\lim_{n \rightarrow \infty} d_n^k F(d_n) = 0.$$

By 3.6, the following holds for all $n \in \mathbb{N}$

$$d_n^k F(d_n) - d_n^k F(d_0) \leq -d_n^k n\tau \leq 0. \quad (3.7)$$

Letting $n \rightarrow \infty$ in 3.7, we obtain that

$$\lim_{n \rightarrow \infty} nd_n^k = 0. \quad (3.8)$$

From 3.8, there exists $n_1 \in \mathbb{N}$ such that $nd_n^k \leq 1$ for all $n \geq n_1$. So we have for all $n \geq n_1$

$$d_n \leq \frac{1}{n^{\frac{1}{k}}}. \quad (3.9)$$

Therefore $\sum_{n=1}^{\infty} d_n < \infty$. Now, let $m, n \in \mathbb{N}$ with $m > n \geq n_1$, then we get

$$\begin{aligned} d(x_n, x_m) &\leq d(x_n, x_{n+1}) + d(x_{n+1}, x_{n+2}) + \cdots + d(x_{m-1}, x_m) \\ &= d_n + d_{n+1} + \cdots + d_{m-1} \\ &\leq \sum_{k=n}^{\infty} d_k. \end{aligned}$$

Since $\sum_{k=1}^{\infty} d_k$ is convergent, then we get $\{x_n\}$ is left K -Cauchy sequence in the quasi metric space (X, d) . Since (X, d) left K -complete, there exists $z \in X$ such that $\{x_n\}$ is d -converges to z , that is, $d(z, x_n) \rightarrow 0$ as $n \rightarrow \infty$. Since T is τ_d -continuous, then

$$d(Tz, Tx_n) = d(Tz, x_{n+1}) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Since X is Hausdorff, we get $z = Tz$. \square

Theorem 2. *Let (X, d) be a Hausdorff left \mathcal{M} -complete quasi metric space, $T : X \rightarrow X$ be a generalized $(\alpha - F_d)$ -contraction. Assume that T is α -admissible and $\tau_{d^{-1}}$ -continuous. If there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$, then T has a fixed point in X .*

Proof. As in the proof of Theorem 1, we can get the iterative sequence $\{x_n\}$ is left K -Cauchy. Since (X, d) is left \mathcal{M} -complete, there exists $z \in X$ such that $\{x_n\}$ is d^{-1} -converges to z , that is, $d(x_n, z) \rightarrow 0$ as $n \rightarrow \infty$. By $\tau_{d^{-1}}$ -continuity of T , we have

$$d(Tx_n, Tz) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Since X is Hausdorff, we get $z = Tz$. \square

Theorem 3. *Let (X, d) be left Smyth complete T_1 -quasi metric space, $T : X \rightarrow X$ be a generalized $(\alpha - F_d)$ -contraction. Assume that T is α -admissible and τ_d or $\tau_{d^{-1}}$ -continuous. If there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$, then T has a fixed point in X .*

Proof. As in the proof of Theorem 1, we can get the iterative sequence $\{x_n\}$ is left K -Cauchy. Since (X, d) is left Smyth complete, there exists $z \in X$ such that $\{x_n\}$ is d^s -converges to z , that is, $d^s(x_n, z) \rightarrow 0$ as $n \rightarrow \infty$. If T is τ_d -continuous, then

$$d(Tz, Tx_n) = d(Tz, x_{n+1}) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Therefore we get

$$d(Tz, z) \leq d(Tz, x_{n+1}) + d(x_{n+1}, z) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

If T is τ_{d-1} -continuous, then

$$d(Tx_n, Tz) = d(x_{n+1}, Tz) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Therefore we get

$$d(z, Tz) \leq d(z, x_{n+1}) + d(x_{n+1}, Tz) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Since T is T_1 -quasi metric space, we obtain $z = Tz$. □

Example 1. Let $X = \{0, 2, 4, \dots\}$ and

$$d(x, y) = \begin{cases} 0 & ; x = y \\ x + y & ; x \neq y \end{cases}$$

for all $x, y \in X$. Now define $T : X \rightarrow X$ as

$$Tx = \begin{cases} 0 & ; x = 0 \\ x - 2 & ; x \neq 0 \end{cases}$$

and $\alpha : X \times X \rightarrow [0, \infty)$ as

$$\alpha(x, y) = \begin{cases} 0; & (x, y) \in \{(0, 2), (2, 0)\} \\ 2; & \text{otherwise} \end{cases}$$

Since (X, d) is complete metric space then it is left Smyth complete T_1 -quasi metric space. Also T is α -admissible. On the other hand

$$\begin{aligned} T_\alpha &= \{(x, y) \in X \times X : \alpha(x, y) \geq 1 \text{ and } d(Tx, Ty) > 0\} \\ &= \{(x, y) \in X \times X : (x, y) \notin \{(0, 2), (2, 0)\} \text{ and } x \neq y\} \end{aligned}$$

Then we get T is a generalized $(\alpha - F_d)$ -contraction with $F(\delta) = \ln \delta$, $\tau = \ln 2$ for all $x, y \in T_\alpha$ and $z = 0$ is a fixed point of T .

REFERENCES

- [1] Altun, I., Al Arifi, N., Jleli M., Lashin, A., Samet, B., A new concept of $(\alpha - F_d)$ -contraction on quasi metric space, J. Nonlinear Sci. Appl., 9 (2016), no. 5, 3354-3361.
- [2] Cosentino, M., Vetro, P., Fixed point results for F -contractive mappings of Hardy-Rogers-type, Filomat, 28 (2014), 715-722.
- [3] Hussain, N., Vetro, C., Vetro, F., Fixed point results for α -implicit contractions with application to integral equations, Nonlinear Anal. Model. Control, 21 (2016), 362-378.
- [4] Karapinar, E., Samet, B., Generalized $\alpha - \psi$ -contractive type mappings and related fixed point theorems with applications, Abstr. Appl. Anal., 2012 (2012), 17 pages.
- [5] Kumam, P., Vetro, C., Vetro, F., Fixed points for weak $\alpha - \psi$ -contractions in partial metric spaces, Abstr. Appl. Anal., 2013 (2013), 9 pages.
- [6] Künzi, H.P. A., Vajner, V., Weighted quasi-metrics, Ann. New York Acad. Sci., 728 (1994), 64-67.
- [7] Minak, G., Olgun, M., Altun, I., A new approach to fixed point theorems for multivalued contractive maps, Carpathian J. Math., 31 (2015), 241-248.

- [8] Reilly, I. L., Subrahmanyam, P. V., Vamanamurthy, M. K., Cauchy sequences in quasi-pseudo-metric spaces, *Monatsh. Math.*, 93 (1982), 127–140. 1.1, 1, 1.2,
- [9] Romaguera, S., Left K -completeness in quasi-metric spaces, *Math. Nachr.*, 157 (1992), 15–23.
- [10] Samet, B., Vetro, C., Vetro, P., Fixed point theorems for $\alpha - \psi$ -contractive type mappings, *Nonlinear Anal.*, 75 (2012), 2154–2165.
- [11] Wardowski, D., Fixed points of a new type of contractive mappings in complete metric spaces, *Fixed Point Theory Appl.*, 2012 (2012), 6 pages.
- [12] Wardowski, D., Van Dung, N., Fixed points of F -weak contractions on complete metric spaces, *Demonstr. Math.*, 47 (2014), 146–155.

Current address: Özlem ACAR: Department of Mathematics, Faculty of Science and Arts, Mersin University, 33343, Yenişehir, Mersin, TURKEY

E-mail address: ozlemacar@mersin.edu.tr

ORCID Address: <http://orcid.org/0000-0001-6052-4357>