

# Intraabdominal Synovial Sarcoma, A Rare Tumour and Rare Localisation

Intraabdominal Sinovyal Sarkom,  
Nadir Lokalizasyonda Nadir Bir Tümör

**Mustafa Yener Uzunoğlu<sup>1</sup>, Fatih Altıntoprak<sup>2</sup>, Enis Dikicier<sup>2</sup>, Zeynep Kahyaoğlu<sup>3</sup>**

<sup>1</sup>Sakarya University Research and Educational Hospital, Department of General Surgery, Sakarya, Turkey

<sup>2</sup>İstinye University, Faculty of Medicine, Department of General Surgery, İstanbul, Turkey

<sup>3</sup> Sakarya Research and Educational Hospital, Department of Pathology, Sakarya, Turkey

Yazışma Adresi / Correspondence:

**Mustafa Yener Uzunoglu**

Sakarya Üniversitesi Eğitim ve Araştırma Hastanesi, Adnan Menderes Cad. Sağlık Sok.No: 195 Adapazarı, 54100, Sakarya / Turkey

T: +90 505 650 33 94 E-mail: [drmyuzunoglu@gmail.com](mailto:drmyuzunoglu@gmail.com)

Geliş Tarihi / Received : 12.03.2018 Kabul Tarihi / Accepted : 07.08.2018

## Abstract

Synovial sarcomas are often seen in the soft tissues, intra-abdominal synovial sarcomas are extremely rare. An intra-abdominal mass starting from the umbilicus and extending into the pelvis was detected in a 53-year-old man admitted with abdominal pain and a palpable mass. Total mass excision surgery was performed and the mass was found to be an intra-abdominal synovial sarcoma arise from small bowel wall histopathologically. Unfortunately, this is a malignancy with a poor prognosis and a short survival despite all therapeutic efforts. ( *Sakarya Med J*, 2018, 8(2):669-673)

Keywords

intra-abdominal mass; sarcoma; small bowel wall; synovial sarcoma

## Öz

Sinovyal sarkomlar sıklıkla yumuşak dokuda gözlenirler, intra abdominal sinovyal sarkomlar oldukça nadirdir. Karın ağrısı ve palpabl kitle ile başvuran 53 yaşında erkek bir hastada umbilikustan başlayıp, pelvise kadar uzanım gösteren karın içi kitle saptandı. Total kitle eksizyonu yapılan hastada, histopatolojik olarak ince barsak duvarından kaynaklanan intraabdominal sinovyal sarkom bulundu. bu malignite zayıf prognoza sahiptir ve yapılan tüm tedavilere rağmen sağkalımı kısadır.( *Sakarya Tıp Dergisi*, 2018, 8(2):669-673 ).

Anahtar  
kelimeler

ince barsak duvarı; intraabdominal kitle; sarkom; sinovyal sarkom

## Introduction

Synovial sarcoma is a rare malignancy accounting for 5~10% of all soft-tissue sarcomas. It has an aggressive course and poor prognosis, with a reported mean 5-year survival of 66%.<sup>1</sup> It is typically encountered in the third to fifth decades of life with a male preponderance. In 95% of the cases, the tumour is localised to the extremities, most commonly around the knee.<sup>2</sup> Rarely, tumours originating from the retroperitoneum, tongue, neck, oesophagus, and abdominal wall have been reported. The recommended treatment is surgical excision followed by chemotherapy and radiation therapy. Here, we report a 53-year-old man who was admitted with a palpable intra-abdominal mass and was ultimately diagnosed with an intra-abdominal synovial sarcoma.

## Case report

A 53-year-old man presented with abdominal pain for 4 months and a palpable abdominal mass (Fig. 1).





Figure 1. Preoperative patient's view; apparent palpable intraabdominal mass

Physical examination detected a mass starting from the periumbilical region with indistinct borders. Abdominal ultrasonography (US) revealed a mass extending from the umbilical level to the pelvis, with heterogeneous cystic and degenerated regions and irregular margins. The mass filled the entire lower abdominal region (Fig. 2). The symptoms of an intestinal obstruction emerged and we performed surgery. After obtaining written informed consent, surgical exploration showed a 280 × 180 mm mass with lobulated contours and some cystic components, although it was mainly solid in nature, extending from the umbilical level to the pelvis, surrounding a 20-cm ileal segment (Fig. 3A-B). The mass was excised totally and a partial ileal resection and terminal ileostomy were performed. An iatrogenic injury to the urinary bladder during the surgery was repaired primarily. The immunohistochemical staining showed that the major cell types were spindle and epithelial cells revealed a synovial sarcoma with CD99 and EMA positive, but were negative for S-100, protein, desmin, CD34. (Fig. 4,5)

In the postoperative course, the patient had acute renal failure, aspiration pneumonia. The overall



condition of the patient worsened in the 8th postoperative day and he died of multi-organ failure on day 10.

Figure 2. Ultrasound image; a mass extending from the umbilical level to the pelvis, with heterogeneous cystic and degenerated regions and irregular margins




Figure 3A. Operative view; intraabdominal mass is huge, lobulated contour and solid-cystic in nature.

Figure 3B. Operative view; intraabdominal mass is huge, lobulated contour and solid-cystic in nature.




Figure 4. Histological microphotograph of synovial sarcoma, stained with CD99 x20




Figure 5. Histological microphotograph of synovial sarcoma, stained with EMA x20

## Discussion

Synovial sarcomas are rare aggressive tumours that typically arise in the periarticular areas<sup>2</sup>. Other rare anatomic locations of this tumour are the thorax, neck, parathyroid, tongue, larynx, mediastinum, oesophagus, heart, lung, abdominal wall, gastrocolic ligament, small bowel mesentery, prostate, kidney, and retroperitoneum.<sup>3,4</sup> In this case the tumor arose from small bowel wall.

Clinically, an intra-abdominal synovial sarcoma often presents as a painful, palpable soft mass, as with tumours in other locations. Non-specific gastrointestinal complaints such as abdominal pain, bloating, weight loss, and vomiting may accompany the clinical presentation, depending on the size of the intra-abdominal mass.<sup>3</sup> In this case, the abdominal pain had been present for 4 months, but was overlooked by the patient, until a palpable mass was detected.

Intra-abdominal localisation is an extremely rare presentation of synovial sarcomas. In a retrospective study of 300 cases, Fisher et al. reported that only 11 (3.6%) of the patients had an intra-abdominal mass.<sup>5</sup> Those 11 cases had a mean age of 49 (range 25–75) years and a mean mass size of 6 × 47 cm. In our patient, the mass measured 28 × 18 cm and the patient's age was within the reported age range for synovial sarcomas.

Histopathologically, a synovial sarcoma is a soft-tissue tumour that shows signs of mesenchymal and epithelial differentiation, which can be detected under a light microscope, immunohistochemically, and by electron microscopy.<sup>6</sup> Although the diagnosis of biphasic synovial sarcomas is often straightforward, a monophasic fibrous synovial sarcoma should be differentiated morphologically from fibrosarcoma, malignant peripheral nerve sheath tumour, and solitary fibrous tumour.<sup>7</sup> Routine use of immunohistochemical methods may facilitate the differential diagnosis of these conditions. Changchien et al. reported primary or metastatic intra-abdominal synovial sarcomas stained with SMA, EMA, BCL-2, CD99, and S100, but not with desmin.<sup>8</sup> The monophasic and biphasic types were differentiated by the Fluorescence In Situ Hybridization (FISH) method using SYT-SSX-1. In a study of 121 cases with a synovial sarcoma, Bergh et al. reported that 60% of the cases were monophasic and 40% the biphasic type.<sup>9</sup> Guillou et al. reported that the monophasic type had a more aggressive course.<sup>10</sup> In our case, staining was positive for CD99 and EMA, histopathological study confirmed the diagnosis of monophasic synovial sarcoma.

The major determinant of survival is the histological grade of the tumour, and the dimensions and depth of the tumour are other prognostic factors.<sup>8</sup> Wide local excision and radiation therapy are mainstays in the treatment of synovial sarcomas. Chemotherapy has been also suggested in the treatment of some histological subtypes.<sup>9</sup> Nevertheless, half of the cases die from distant metastasis despite all treatment modalities.

In conclusion, although synovial sarcomas are rarely seen outside the extremities, they can also present as an intra-abdominal mass, albeit extremely rarely. Wide excision is the first step in the treatment. Unfortunately, this is a malignancy with a poor prognosis and a short survival despite all therapeutic efforts.

**Conflicts Of Interest:** No potential conflict of interest relevant to this article was reported.

1. Sultan I, Rodriguez-Galindo C, Saab R, Yasir S, Casanova M, Ferrari A. Comparing children and adults with synovial sarcoma in the Surveillance, Epidemiology, and End Results program, 1983 to 2005. *Cancer* 2009; 115(15):3537-3547 DOI: 10.1002/encr.24424
2. Indranil G, Divya M. Synovial sarcoma of the omentum: A rare entity. *Indian J Cancer* 2015;52:166-7
3. Carrillo R, Rodriguez-Peralto JL, Batsakis JG. Synovial sarcoma of the head and neck. *Ann Otol Rhinol Laryngol.* 1992 Apr; 101(4):367-70. DOI: 10.1177/000348949210100415
4. Fritsch M, Epstein JI, Perma EJ, Watts JC, Argani P. Molecularly confirmed primary prostatic synovial sarcoma. *Hum Pathol* 2000; 31:246-250
5. Fisher C, Folpe AL, Hashimoto H, Weiss SW. Intra-abdominal synovial sarcoma: a clinicopathological study. *Histopathology* September 2004; 45(3): 245-253
6. Buiga-Potcoav R, Crişan D, Olinici CD. Primary intraabdominal sarcoma, a case report *Romanian Journal Of Gastroenterology* 2005;14(1): 67-69
7. Harsh KK, Kalwar A, Kapoor A, Jakhar SL, Kumar HS. Giant cell variant of malignant fibrous histiocytoma of male breast: A rare case report. *J Can Res Ther* 2015;11:657 DOI: 10.4103/0973-1482.138129
8. Changchien YC, Katalin U, Fillinger J, Fónyad L, Papp G, Salamon F, Sági Z. A Challenging Case of Metastatic Intra-Abdominal Synovial Sarcoma with Unusual Immunophenotype and Its Differential Diagnosis. *Case Rep Pathol.* 2012; 2012: 786083. DOI: 10.1155/2012/786083.
9. Eriksen C, Burns L, Bohlke A, Haque S, Slakey DP. Management of monophasic synovial sarcoma of the small intestine. *JSLs.* 2010;14(3):421-5. DOI: 10.4293/108680810X12924466006846.
10. Guillou L, Coindre JM, Gallagher G, Terrier P, Gebhard S, Somerhausen NSA, et al. Detection of the synovial sarcoma translocation t(X;18) (SYT;SSX) in paraffin-embedded tissues using reverse transcriptase-polymerase chain reaction: a reliable and powerful diagnostic tool for pathologist. A molecular analysis of 221 mesenchymal tumors fixed in different fixatives. *Hum Pathol* 2001; 32:105-112. DOI: <https://doi.org/10.1053/hupa.2001.21130>

REFERENCE

Sakarya Med J.  
2018;8(2):669-673

**UZUNOĞLU et al.**  
Intraabdominal Synovial Sarcoma,  
A Rare Tumour and Rare Localisation