



**SOME CAPUTO k -FRACTIONAL DERIVATIVES OF
 OSTROWSKI TYPE CONCERNING $(n+1)$ -DIFFERENTIABLE
 GENERALIZED RELATIVE SEMI- $(r; m, p, q, h_1, h_2)$ -PREINVEX
 MAPPINGS**

ARTION KASHURI AND ROZANA LIKO

ABSTRACT. In this article, we first presented some integral inequalities for Gauss-Jacobi type quadrature formula involving generalized relative semi- $(r; m, p, q, h_1, h_2)$ -preinvex mappings. And then, a new identity concerning $(n+1)$ -differentiable mappings defined on m -invex set via Caputo k -fractional derivatives is derived. By using the notion of generalized relative semi- $(r; m, p, q, h_1, h_2)$ -preinvexity and the obtained identity as an auxiliary result, some new estimates with respect to Ostrowski type inequalities via Caputo k -fractional derivatives are established. It is pointed out that some new special cases can be deduced from main results of the article.

1. INTRODUCTION

The subsequent double inequality is known as Ostrowski inequality which gives an upper bound for the approximation of the integral average $\frac{1}{b-a} \int_a^b f(t)dt$ by the value $f(x)$ at point $x \in [a, b]$.

Theorem 1.1. *Let $f : I \rightarrow \mathbb{R}$ be a mapping differentiable on I° and let $a, b \in I^\circ$ with $a < b$. If $|f'(x)| \leq M$ for all $x \in [a, b]$, then*

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t)dt \right| \leq M(b-a) \left[\frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right], \quad \forall x \in [a, b]. \quad (1.1)$$

Ostrowski inequality is playing a very important role in all the fields of mathematics, especially in the theory of approximations. Thus such inequalities were

Received by the editors: Jan. 05, 2017, Accepted: Nov. 30, 2017.

2010 *Mathematics Subject Classification.* Primary: 26A51; Secondary: 26A33, 26D07, 26D10, 26D15.

Key words and phrases. Ostrowski type inequality, Hölder's inequality, Minkowski inequality, power mean inequality, Caputo k -fractional derivatives, s -convex function in the second sense, m -invex.

©2018 Ankara University
 Communications Faculty of Sciences University of Ankara-Series A1 Mathematics and Statistics

studied extensively by many researches and numerous generalizations, extensions and variants of them for various kind of functions like bounded variation, synchronous, Lipschitzian, monotonic, absolutely, continuous and n -times differentiable mappings etc. appeared in a number of papers (see [[2]-[4], [11]-[13], [15], [16], [18], [22], [23], [31], [32], [34], [37], [40], [42], [43], [49], [51], [60], [62], [64], [66]]). In recent years, one more dimension has been added to this studies, by introducing a number of integral inequalities involving various fractional operators like Riemann-Liouville, Erdelyi-Kober, Katugampola, conformable fractional integral operators etc. by many authors (see [[1],[47],[52]-[58]]). Riemann-Liouville fractional integral operators are the most central between these fractional operators. In numerical analysis many quadrature rules have been established to approximate the definite integrals (see [[14], [20], [36], [38], [39], [44], [48], [61], [63]]). Ostrowski inequality provides the bounds for many numerical quadrature rules. In recent decades Ostrowski inequality is studied in fractional calculus point of view by many mathematicians (see [[6]-[10], [24]-[30], [35], [45], [50]]).

Let us recall some special functions and evoke some basic definitions as follows.

Definition 1.2. The Euler beta function is defined for $a, b > 0$ as

$$\beta(a, b) = \int_0^1 t^{a-1} (1-t)^{b-1} dt = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}.$$

Definition 1.3. For $k \in \mathbb{R}^+$ and $x \in \mathbb{C}$, the k -gamma function is defined by

$$\Gamma_k(x) = \lim_{n \rightarrow \infty} \frac{n! k^n (nk)^{\frac{x}{k}-1}}{(x)_{n,k}}. \quad (1.2)$$

Its integral representation is given by

$$\Gamma_k(\alpha) = \int_0^\infty t^{\alpha-1} e^{-\frac{t^k}{k}} dt. \quad (1.3)$$

One can note that

$$\Gamma_k(\alpha + k) = \alpha \Gamma_k(\alpha).$$

For $k = 1$, (1.3) gives integral representation of gamma function.

Definition 1.4. For $k \in \mathbb{R}^+$ and $x, y \in \mathbb{C}$, the k -beta function with two parameters x and y is defined as

$$\beta_k(x, y) = \frac{1}{k} \int_0^1 t^{\frac{x}{k}-1} (1-t)^{\frac{y}{k}-1} dt. \quad (1.4)$$

For $k = 1$, (1.4) gives integral representation of beta function.

Theorem 1.5. Let $x, y > 0$. Then for k -gamma and k -beta function the following equality holds:

$$\beta_k(x, y) = \frac{\Gamma_k(x)\Gamma_k(y)}{\Gamma_k(x+y)}. \quad (1.5)$$

Definition 1.6. [30] Let $\alpha > 0$ and $\alpha \notin \{1, 2, 3, \dots\}$, $n = [\alpha] + 1$, $f \in C^n[a, b]$ such that $f^{(n)}$ exists and is continuous on $[a, b]$. The Caputo fractional derivatives of order α are defined as follows:

$${}^cD_{a+}^\alpha f(x) = \frac{1}{\Gamma(n-\alpha)} \int_a^x \frac{f^{(n)}(t)}{(x-t)^{\alpha-n+1}} dt, \quad x > a \quad (1.6)$$

and

$${}^cD_{b-}^\alpha f(x) = \frac{(-1)^n}{\Gamma(n-\alpha)} \int_x^b \frac{f^{(n)}(t)}{(t-x)^{\alpha-n+1}} dt, \quad x < b. \quad (1.7)$$

If $\alpha = n \in \{1, 2, 3, \dots\}$ and usual derivative of order n exists, then Caputo fractional derivative $({}^cD_{a+}^\alpha f)(x)$ coincides with $f^{(n)}(x)$. In particular we have

$$({}^cD_{a+}^0 f)(x) = ({}^cD_{b-}^0 f)(x) = f(x) \quad (1.8)$$

where $n = 1$ and $\alpha = 0$.

Definition 1.7. [19] Let $\alpha > 0$, $k \geq 1$ and $\alpha \notin \{1, 2, 3, \dots\}$, $n = [\alpha] + 1$, $f \in C^n[a, b]$. The Caputo k -fractional derivatives of order α are defined as follows:

$${}^cD_{a+}^{\alpha,k} f(x) = \frac{1}{k\Gamma_k(n-\frac{\alpha}{k})} \int_a^x \frac{f^{(n)}(t)}{(x-t)^{\frac{\alpha}{k}-n+1}} dt, \quad x > a \quad (1.9)$$

and

$${}^cD_{b-}^{\alpha,k} f(x) = \frac{(-1)^n}{k\Gamma_k(n-\frac{\alpha}{k})} \int_x^b \frac{f^{(n)}(t)}{(t-x)^{\frac{\alpha}{k}-n+1}} dt, \quad x < b. \quad (1.10)$$

Definition 1.8. [65] A set $M_\varphi \subseteq \mathbb{R}^n$ is named as a relative convex (φ -convex) set, if and only if, there exists a function $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that,

$$t\varphi(x) + (1-t)\varphi(y) \in M_\varphi, \quad \forall x, y \in \mathbb{R}^n : \varphi(x), \varphi(y) \in M_\varphi, t \in [0, 1]. \quad (1.11)$$

Definition 1.9. [65] A function f is named as a relative convex (φ -convex) function on a relative convex (φ -convex) set M_φ , if and only if, there exists a function $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that,

$$f(t\varphi(x) + (1-t)\varphi(y)) \leq tf(\varphi(x)) + (1-t)f(\varphi(y)), \quad (1.12)$$

$$\forall x, y \in \mathbb{R}^n : \varphi(x), \varphi(y) \in M_\varphi, t \in [0, 1].$$

Definition 1.10. [14] A non-negative function $f : I \subseteq \mathbb{R} \rightarrow [0, +\infty)$ is said to be P -function, if

$$f(tx + (1-t)y) \leq f(x) + f(y), \quad \forall x, y \in I, t \in [0, 1].$$

Definition 1.11. [5] A set $K \subseteq \mathbb{R}^n$ is said to be invex respecting the mapping $\eta : K \times K \rightarrow \mathbb{R}^n$, if $x + t\eta(y, x) \in K$ for every $x, y \in K$ and $t \in [0, 1]$.

Definition 1.12. [36] Let $h : [0, 1] \rightarrow \mathbb{R}$ be a non-negative function and $h \neq 0$. The function f on the invex set K is said to be h -preinvex with respect to η , if

$$f(x + t\eta(y, x)) \leq h(1-t)f(x) + h(t)f(y) \quad (1.13)$$

for each $x, y \in K$ and $t \in [0, 1]$ where $f(\cdot) > 0$.

Clearly, when putting $h(t) = t$ in Definition 1.12, f becomes a preinvex function [46]. If the mapping $\eta(y, x) = y - x$ in Definition(1.12), then the non-negative function f reduces to h -convex mappings [63].

Definition 1.13. [61] Let $f : K \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a non-negative function, a function $f : K \rightarrow \mathbb{R}$ is said to be a *tgs*-convex function on K if the inequality

$$f((1-t)x + ty) \leq t(1-t)[f(x) + f(y)] \quad (1.14)$$

holds for all $x, y \in K$ and $t \in (0, 1)$.

Definition 1.14. [34] A function $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ is said to be *MT*-convex functions, if it is non-negative and $\forall x, y \in I$ and $t \in (0, 1)$ satisfies the subsequent inequality:

$$f(tx + (1-t)y) \leq \frac{\sqrt{t}}{2\sqrt{1-t}}f(x) + \frac{\sqrt{1-t}}{2\sqrt{t}}f(y). \quad (1.15)$$

Definition 1.15. [39] A function: $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ is said to be *m-MT*-convex, if f is positive and for $\forall x, y \in I$, and $t \in (0, 1)$, among $m \in [0, 1]$, satisfies the following inequality

$$f(tx + m(1-t)y) \leq \frac{\sqrt{t}}{2\sqrt{1-t}}f(x) + \frac{m\sqrt{1-t}}{2\sqrt{t}}f(y). \quad (1.16)$$

Definition 1.16. [45] Let $K \subseteq \mathbb{R}$ be an open m -invex set respecting $\eta : K \times K \times (0, 1) \rightarrow \mathbb{R}$ and $h_1, h_2 : [0, 1] \rightarrow [0, +\infty)$. A function $f : K \rightarrow \mathbb{R}$ is said to be generalized (m, h_1, h_2) -preinvex functions with respect to η , if

$$f(mx + t\eta(y, x, m)) \leq mh_1(t)f(x) + h_2(t)f(y) \quad (1.17)$$

is valid for all $x, y \in K$ and $t \in [0, 1]$, for some fixed $m \in (0, 1]$.

Let us recall the Gauss-Jacobi type quadrature formula as follows.

$$\int_a^b (x-a)^p(b-x)^q f(x) dx = \sum_{k=0}^{+\infty} B_{m,k} f(\gamma_k) + R_m^* |f|, \quad (1.18)$$

for certain $B_{m,k}, \gamma_k$ and rest $R_m^* |f|$, [59].

In [33], Liu obtained integral inequalities for P -function related to the left-hand side of (1.18), and in [41], Özdemir et al. also presented several integral inequalities concerning the left-hand side of (1.18) via some kinds of convexity.

Motivated by the above literatures, the main objective of this article is to establish integral inequalities for the left side of Gauss-Jacobi type quadrature

formula and some new estimates on Ostrowski type inequalities via Caputo k -fractional derivatives associated with generalized relative semi- $(r; m, p, q, h_1, h_2)$ -preinvex mappings. It is pointed out that some new special cases will be deduced from main results of the article.

2. MAIN RESULTS INVOLVING GAUSS-JACOBI TYPE QUADRATURE FORMULA

The following definitions will be used in this section.

Definition 2.1. [17] A set $K \subseteq \mathbb{R}$ is named as m -invex with respect to the mapping $\eta : K \times K \rightarrow \mathbb{R}^n$ for some fixed $m \in (0, 1]$, if $mx + t\eta(y, mx) \in K$ holds for each $x, y \in K$ and any $t \in [0, 1]$.

Remark 2.2. In Definition 2.1, under certain conditions, the mapping $\eta(y, mx)$ could reduce to $\eta(y, x)$. For example when $m = 1$, then the m -invex set degenerates an invex set on K .

We next introduce generalized relative semi- $(r; m, p, q, h_1, h_2)$ -preinvex mappings.

Definition 2.3. Let $K \subseteq \mathbb{R}$ be an open m -invex set with respect to the mapping $\eta : K \times K \times (0, 1] \rightarrow \mathbb{R}$, $h_1, h_2 : [0, 1] \rightarrow [0, +\infty)$ and $\varphi : I \rightarrow K$ are continuous functions. A mapping $f : K \rightarrow (0, +\infty)$, is said to be generalized relative semi- $(r; m, p, q, h_1, h_2)$ -preinvex, if

$$f(m\varphi(x) + t\eta(\varphi(y), \varphi(x), m)) \leq M_r(h_1(t), h_2(t); mf(x), f(y), p, q) \quad (2.1)$$

holds for all $x, y \in I$ and $t \in [0, 1]$, for $p, q > -1$ with some fixed $m \in (0, 1]$, where

$$M_r(h_1(t), h_2(t); mf(x), f(y), p, q) := \begin{cases} \left[mh_1^p(t)f^r(x) + h_2^q(t)f^r(y) \right]^{\frac{1}{r}}, & \text{if } r \neq 0; \\ [mf(x)]^{h_1^p(t)} [f(y)]^{h_2^q(t)}, & \text{if } r = 0, \end{cases}$$

is the weighted power mean of order r for positive numbers $f(x)$ and $f(y)$.

Remark 2.4. In Definition 2.3, if we choose $r = p = q = 1$ and $\varphi(x) = x$, then we get Definition 1.16.

Remark 2.5. For $r = p = q = 1$, let us discuss some special cases in Definition 2.3 as follows.

- (I) If taking $h_1(t) = (1-t)^s$, $h_2(t) = t^s$ for $s \in (0, 1]$, then we get generalized relative semi- (m, s) -Breckner-preinvex mappings.
- (II) If taking $h_1(t) = h_2(t) = 1$, then we get generalized relative semi- (m, P) -preinvex mappings.
- (III) If taking $h_1(t) = (1-t)^{-s}$, $h_2(t) = t^{-s}$ for $s \in (0, 1]$, then we get generalized relative semi- (m, s) -Godunova-Levin-Dragomir-preinvex mappings.
- (IV) If taking $h_1(t) = h(1-t)$, $h_2(t) = h(t)$, then we get generalized relative semi- (m, h) -preinvex mappings.

(V) If taking $h_1(t) = h_2(t) = t(1-t)$, then we get generalized relative semi- (m, tgs) -preinvex mappings.

(VI) If taking $h_1(t) = \frac{\sqrt{1-t}}{2\sqrt{t}}$, $h_2(t) = \frac{\sqrt{t}}{2\sqrt{1-t}}$, then we get generalized relative semi- m -MT-preinvex mappings.

It is worth to mention here that to the best of our knowledge all the special cases discussed above are new in the literature.

We claim the following integral identity.

Lemma 2.6. *Let $\varphi : I \rightarrow K$ be a continuous function. Assume that $f : K = [m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)] \rightarrow \mathbb{R}$ is a continuous function on K° with respect to $\eta : K \times K \times (0, 1] \rightarrow \mathbb{R}$, for $\eta(\varphi(b), \varphi(a), m) > 0$. Then for some fixed $m \in (0, 1]$ and $p, q > 0$, we have*

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b), \varphi(a), m)} (x - m\varphi(a))^p (m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x)^q f(x) dx \\ &= \eta^{p+q+1}(\varphi(b), \varphi(a), m) \int_0^1 t^p (1-t)^q f(m\varphi(a) + t\eta(\varphi(b), \varphi(a), m)) dt. \end{aligned}$$

Proof. It is easy to observe that

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b), \varphi(a), m)} (x - m\varphi(a))^p (m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x)^q f(x) dx \\ &= \eta(\varphi(b), \varphi(a), m) \int_0^1 (m\varphi(a) + t\eta(\varphi(b), \varphi(a), m) - m\varphi(a))^p \\ & \quad \times (m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - m\varphi(a) - t\eta(\varphi(b), \varphi(a), m))^q \\ & \quad \times f(m\varphi(a) + t\eta(\varphi(b), \varphi(a), m)) dt \\ &= \eta^{p+q+1}(\varphi(b), \varphi(a), m) \int_0^1 t^p (1-t)^q f(m\varphi(a) + t\eta(\varphi(b), \varphi(a), m)) dt. \end{aligned}$$

This completes the proof of the lemma. \square

With the help of Lemma 2.6, we have the following results.

Theorem 2.7. *Suppose $h_1, h_2 : [0, 1] \rightarrow [0, +\infty)$ and $\varphi : I \rightarrow K$ are continuous functions. Assume that $f : K = [m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)] \rightarrow (0, +\infty)$ is a continuous function on K° with respect to $\eta : K \times K \times (0, 1] \rightarrow \mathbb{R}$, for $\eta(\varphi(b), \varphi(a), m) > 0$. Let $k > 1$ and $0 < r \leq 1$. If $f^{\frac{k}{k-1}}$ is generalized relative semi- $(r; m, \bar{p}, \bar{q}, h_1, h_2)$ -preinvex mappings on an open m -invex set K for some fixed $m \in (0, 1]$, where $\bar{p}, \bar{q} > -1$, then for any fixed $p, q > 0$, we have*

$$\int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b), \varphi(a), m)} (x - m\varphi(a))^p (m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x)^q f(x) dx$$

$$\leq \eta^{p+q+1}(\varphi(b), \varphi(a), m) \beta^{\frac{1}{k}}(kp+1, kq+1) \\ \times \left[mf^{\frac{rk}{k-1}}(a) \Psi^r(h_1(t); r, \bar{p}) + f^{\frac{rk}{k-1}}(b) \Psi^r(h_2(t); r, \bar{q}) \right]^{\frac{k-1}{rk}},$$

where

$$\Psi(h_1(t); r, \bar{p}) := \int_0^1 h_1^{\frac{\bar{p}}{r}}(t) dt, \quad \Psi(h_2(t); r, \bar{q}) := \int_0^1 h_2^{\frac{\bar{q}}{r}}(t) dt.$$

Proof. Let $k > 1$ and $0 < r \leq 1$. Since $f^{\frac{k}{k-1}}$ is generalized relative semi- $(r; m, \bar{p}, \bar{q}, h_1, h_2)$ -preinvex mappings on K , combining with Lemma 2.6, Hölder inequality, Minkowski inequality and property of the modulus for all $t \in [0, 1]$ and for some fixed $m \in (0, 1]$, we get

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b), \varphi(a), m)} (x - m\varphi(a))^p (m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x)^q f(x) dx \\ & \leq \eta^{p+q+1}(\varphi(b), \varphi(a), m) \left[\int_0^1 t^{kp} (1-t)^{kq} dt \right]^{\frac{1}{k}} \\ & \quad \times \left[\int_0^1 f^{\frac{k}{k-1}}(m\varphi(a) + t\eta(\varphi(b), \varphi(a), m)) dt \right]^{\frac{k-1}{k}} \\ & \leq \eta^{p+q+1}(\varphi(b), \varphi(a), m) \beta^{\frac{1}{k}}(kp+1, kq+1) \\ & \quad \times \left[\int_0^1 \left[mh_1^{\bar{p}}(t) f^{\frac{rk}{k-1}}(a) + h_2^{\bar{q}}(t) f^{\frac{rk}{k-1}}(b) \right]^{\frac{1}{r}} dt \right]^{\frac{k-1}{k}} \\ & \leq \eta^{p+q+1}(\varphi(b), \varphi(a), m) \beta^{\frac{1}{k}}(kp+1, kq+1) \\ & \quad \times \left\{ \left(\int_0^1 m^{\frac{1}{r}} h_1^{\frac{\bar{p}}{r}}(t) f^{\frac{k}{k-1}}(a) dt \right)^r + \left(\int_0^1 h_2^{\frac{\bar{q}}{r}}(t) f^{\frac{k}{k-1}}(b) dt \right)^r \right\}^{\frac{k-1}{rk}} \\ & = \eta^{p+q+1}(\varphi(b), \varphi(a), m) \beta^{\frac{1}{k}}(kp+1, kq+1) \\ & \quad \times \left[mf^{\frac{rk}{k-1}}(a) \Psi^r(h_1(t); r, \bar{p}) + f^{\frac{rk}{k-1}}(b) \Psi^r(h_2(t); r, \bar{q}) \right]^{\frac{k-1}{rk}}. \end{aligned}$$

So, the proof of this theorem is completed. \square

We point out some special cases of Theorem 2.7.

Corollary 2.8. *In Theorem 2.7 for $r = \bar{p} = \bar{q} = 1$ and $h_1(t) = h(1-t)$, $h_2(t) = h(t)$, we have the following inequality for generalized relative semi- (m, h) -preinvex mappings*

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b), \varphi(a), m)} (x - m\varphi(a))^p (m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x)^q f(x) dx \\ & \leq \eta^{p+q+1}(\varphi(b), \varphi(a), m) \beta^{\frac{1}{k}}(kp+1, kq+1) \Psi^{\frac{k-1}{k}}(h(t); 1, 1) \left[mf^{\frac{k}{k-1}}(a) + f^{\frac{k}{k-1}}(b) \right]^{\frac{k-1}{k}}. \end{aligned}$$

Corollary 2.9. *In Theorem 2.7 for $r = \bar{p} = \bar{q} = 1$ and $h_1(t) = (1-t)^s$, $h_2(t) = t^s$, we have the following inequality for generalized relative semi- (m, s) -Breckner-preinvex mappings*

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b), \varphi(a), m)} (x - m\varphi(a))^p (m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x)^q f(x) dx \\ & \leq \eta^{p+q+1}(\varphi(b), \varphi(a), m) \beta^{\frac{1}{k}}(kp+1, kq+1) \left[\frac{mf^{\frac{k}{k-1}}(a) + f^{\frac{k}{k-1}}(b)}{s+1} \right]^{\frac{k-1}{k}}. \end{aligned}$$

Corollary 2.10. *In Theorem 2.7 for $r = \bar{p} = \bar{q} = 1$ and $h_1(t) = (1-t)^{-s}$, $h_2(t) = t^{-s}$, we get the following inequality for generalized relative semi- (m, s) -Godunova-Levin-Dragomir preinvex mappings*

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b), \varphi(a), m)} (x - m\varphi(a))^p (m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x)^q f(x) dx \\ & \leq \eta^{p+q+1}(\varphi(b), \varphi(a), m) \beta^{\frac{1}{k}}(kp+1, kq+1) \left[\frac{mf^{\frac{k}{k-1}}(a) + f^{\frac{k}{k-1}}(b)}{1-s} \right]^{\frac{k-1}{k}}. \end{aligned}$$

Corollary 2.11. *In Theorem 2.7 for $r = \bar{p} = \bar{q} = 1$ and $h_1(t) = h_2(t) = t(1-t)$, we obtain the following inequality for generalized relative semi- (m, tgs) -preinvex mappings*

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b), \varphi(a), m)} (x - m\varphi(a))^p (m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x)^q f(x) dx \\ & \leq \eta^{p+q+1}(\varphi(b), \varphi(a), m) \beta^{\frac{1}{k}}(kp+1, kq+1) \left[\frac{mf^{\frac{k}{k-1}}(a) + f^{\frac{k}{k-1}}(b)}{6} \right]^{\frac{k-1}{k}}. \end{aligned}$$

Corollary 2.12. *In Theorem 2.7 for $r = \bar{p} = \bar{q} = 1$ and $h_1(t) = \frac{\sqrt{1-t}}{2\sqrt{t}}$, $h_2(t) = \frac{\sqrt{t}}{2\sqrt{1-t}}$, we deduce the following inequality for generalized relative semi- m -MT-preinvex mappings*

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b), \varphi(a), m)} (x - m\varphi(a))^p (m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x)^q f(x) dx \\ & \leq \left(\frac{\pi}{4} \right)^{\frac{k-1}{k}} \eta^{p+q+1}(\varphi(b), \varphi(a), m) \beta^{\frac{1}{k}}(kp+1, kq+1) \left[mf^{\frac{k}{k-1}}(a) + f^{\frac{k}{k-1}}(b) \right]^{\frac{k-1}{k}}. \end{aligned}$$

Theorem 2.13. *Suppose $h_1, h_2 : [0, 1] \rightarrow [0, +\infty)$ and $\varphi : I \rightarrow K$ are continuous functions. Assume that $f : K = [m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)] \rightarrow (0, +\infty)$ is a continuous function on K° with respect to $\eta : K \times K \times (0, 1) \rightarrow \mathbb{R}$, for*

$\eta(\varphi(b), \varphi(a), m) > 0$. Let $l \geq 1$ and $0 < r \leq 1$. If f^l is generalized relative semi- $(r; m, \bar{p}, \bar{q}, h_1, h_2)$ -preinvex mappings on an open m -invex set K for some fixed $m \in (0, 1]$, where $\bar{p}, \bar{q} > -1$, then for any fixed $p, q > 0$, we have

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b), \varphi(a), m)} (x - m\varphi(a))^p (m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x)^q f(x) dx \\ & \leq \eta^{p+q+1}(\varphi(b), \varphi(a), m) \beta^{\frac{l-1}{l}}(p+1, q+1) \\ & \times \left[mf^{rl}(a) I^r(h_1(t); r, p, q, \bar{p}) + f^{rl}(b) I^r(h_2(t); r, p, q, \bar{q}) \right]^{\frac{1}{rl}}, \end{aligned}$$

where

$$I(h_1(t); r, p, q, \bar{p}) = \int_0^1 t^p (1-t)^q h_1^{\frac{\bar{p}}{r}}(t) dt, \quad I(h_2(t); r, p, q, \bar{q}) = \int_0^1 t^p (1-t)^q h_2^{\frac{\bar{q}}{r}}(t) dt.$$

Proof. Let $l \geq 1$ and $0 < r \leq 1$. Since f^l is generalized relative semi- $(r; m, \bar{p}, \bar{q}, h_1, h_2)$ -preinvex mappings on K , combining with Lemma 2.6, the well-known power mean inequality, Minkowski inequality and property of the modulus for all $t \in [0, 1]$ and for some fixed $m \in (0, 1]$, we get

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b), \varphi(a), m)} (x - m\varphi(a))^p (m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x)^q f(x) dx \\ & = \eta^{p+q+1}(\varphi(b), \varphi(a), m) \int_0^1 \left[t^p (1-t)^q \right]^{\frac{l-1}{l}} \left[t^p (1-t)^q \right]^{\frac{1}{l}} \\ & \quad \times f(m\varphi(a) + t\eta(\varphi(b), \varphi(a), m)) dt \\ & \leq \eta^{p+q+1}(\varphi(b), \varphi(a), m) \left[\int_0^1 t^p (1-t)^q dt \right]^{\frac{l-1}{l}} \\ & \quad \times \left[\int_0^1 t^p (1-t)^q f^l(m\varphi(a) + t\eta(\varphi(b), \varphi(a), m)) dt \right]^{\frac{1}{l}} \\ & \leq \eta^{p+q+1}(\varphi(b), \varphi(a), m) \beta^{\frac{l-1}{l}}(p+1, q+1) \\ & \quad \times \left[\int_0^1 t^p (1-t)^q \left[mh_1^{\frac{\bar{p}}{r}}(t) f^{rl}(a) + h_2^{\frac{\bar{q}}{r}}(t) f^{rl}(b) \right]^{\frac{1}{r}} dt \right]^{\frac{1}{l}} \\ & \leq |\eta(\varphi(b), \varphi(a), m)|^{p+q+1} \beta^{\frac{l-1}{l}}(p+1, q+1) \\ & \times \left\{ \left(\int_0^1 m^{\frac{1}{r}} t^p (1-t)^q h_1^{\frac{\bar{p}}{r}}(t) f^l(a) dt \right)^r + \left(\int_0^1 t^p (1-t)^q h_2^{\frac{\bar{q}}{r}}(t) f^l(b) dt \right)^r \right\}^{\frac{1}{rl}} \\ & = \eta^{p+q+1}(\varphi(b), \varphi(a), m) \beta^{\frac{l-1}{l}}(p+1, q+1) \\ & \times \left[mf^{rl}(a) I^r(h_1(t); r, p, q, \bar{p}) + f^{rl}(b) I^r(h_2(t); r, p, q, \bar{q}) \right]^{\frac{1}{rl}}. \end{aligned}$$

So, the proof of this theorem is completed. \square

Let us discuss some special cases of Theorem 2.13.

Corollary 2.14. *In Theorem 2.13 for $r = \bar{p} = \bar{q} = 1$ and $h_1(t) = h(1-t)$, $h_2(t) = h(t)$, we have the following inequality for generalized relative semi- (m, h) -preinvex mappings*

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b),\varphi(a),m)} (x - m\varphi(a))^p (m\varphi(a) + \eta(\varphi(b),\varphi(a),m) - x)^q f(x) dx \\ & \leq \eta^{p+q+1}(\varphi(b),\varphi(a),m) \beta^{\frac{l-1}{l}}(p+1, q+1) \\ & \quad \times \left[m f^l(a) I(h(t); 1, p, q, 1) + f^l(b) I(h(t); 1, q, p, 1) \right]^{\frac{1}{l}}. \end{aligned}$$

Corollary 2.15. *In Theorem 2.13 for $r = \bar{p} = \bar{q} = 1$ and $h_1(t) = (1-t)^s$, $h_2(t) = t^s$, we have the following inequality for generalized relative semi- (m, s) -Breckner-preinvex mappings*

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b),\varphi(a),m)} (x - m\varphi(a))^p (m\varphi(a) + \eta(\varphi(b),\varphi(a),m) - x)^q f(x) dx \\ & \leq \eta^{p+q+1}(\varphi(b),\varphi(a),m) \beta^{\frac{l-1}{l}}(p+1, q+1) \\ & \quad \times \left[m f^l(a) \beta(p+1, q+s+1) + f^l(b) \beta(q+1, p+s+1) \right]^{\frac{1}{l}}. \end{aligned}$$

Corollary 2.16. *In Theorem 2.13 for $r = \bar{p} = \bar{q} = 1$ and $h_1(t) = (1-t)^{-s}$, $h_2(t) = t^{-s}$, we get the following inequality for generalized relative semi- (m, s) -Godunova-Levin-Dragomir preinvex mappings*

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b),\varphi(a),m)} (x - m\varphi(a))^p (m\varphi(a) + \eta(\varphi(b),\varphi(a),m) - x)^q f(x) dx \\ & \leq \eta^{p+q+1}(\varphi(b),\varphi(a),m) \beta^{\frac{l-1}{l}}(p+1, q+1) \\ & \quad \times \left[m f^l(a) \beta(p+1, q-s+1) + f^l(b) \beta(q+1, p-s+1) \right]^{\frac{1}{l}}. \end{aligned}$$

Corollary 2.17. *In Theorem 2.13 for $r = \bar{p} = \bar{q} = 1$ and $h_1(t) = h_2(t) = t(1-t)$, we obtain the following inequality for generalized relative semi- (m, tgs) -preinvex mappings*

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b),\varphi(a),m)} (x - m\varphi(a))^p (m\varphi(a) + \eta(\varphi(b),\varphi(a),m) - x)^q f(x) dx \\ & \leq \eta^{p+q+1}(\varphi(b),\varphi(a),m) \beta^{\frac{l-1}{l}}(p+1, q+1) \beta^{\frac{1}{l}}(p+2, q+2) \left[m f^l(a) + f^l(b) \right]^{\frac{1}{l}}. \end{aligned}$$

Corollary 2.18. *In Theorem 2.13 for $r = \bar{p} = \bar{q} = 1$ and $h_1(t) = \frac{\sqrt{1-t}}{2\sqrt{t}}$, $h_2(t) = \frac{\sqrt{t}}{2\sqrt{1-t}}$, we deduce the following inequality for generalized relative semi- m -MT-preinvex mappings*

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b),\varphi(a),m)} (x - m\varphi(a))^p (m\varphi(a) + \eta(\varphi(b),\varphi(a),m) - x)^q f(x) dx \\ & \leq \left(\frac{1}{2} \right)^{\frac{1}{l}} \eta^{p+q+1}(\varphi(b),\varphi(a),m) \beta^{\frac{l-1}{l}} (p+1, q+1) \\ & \quad \times \left[m f^l(a) \beta \left(p + \frac{1}{2}, q + \frac{3}{2} \right) + f^l(b) \beta \left(q + \frac{1}{2}, p + \frac{3}{2} \right) \right]^{\frac{1}{l}}. \end{aligned}$$

3. OTHER RESULTS INVOLVING CAPUTO k -FRACTIONAL DERIVATIVES

For establishing our main results regarding some new Ostrowski type integral inequalities associated with generalized relative semi- $(r; m, p, q, h_1, h_2)$ -preinvexity via Caputo k -fractional derivatives, we need the following lemma.

Lemma 3.1. *Let $\alpha > 0$, $k \geq 1$ and $\alpha \notin \{1, 2, 3, \dots\}$, $n = [\alpha] + 1$. Also, let $\varphi : I \rightarrow K$ be a continuous function. Suppose $K \subseteq \mathbb{R}$ be an open m -invex subset with respect to $\eta : K \times K \times (0, 1] \rightarrow \mathbb{R}$ for some fixed $m \in (0, 1]$. Assume that $f : K \rightarrow \mathbb{R}$ is a function on K° such that $f \in C^{n+1}[m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)]$, where $\eta(\varphi(b), \varphi(a), m) > 0$. Then we have the following equality for Caputo k -fractional derivatives:*

$$\begin{aligned} & \frac{\eta^{n-\frac{\alpha}{k}}(\varphi(x), \varphi(a), m) f^{(n)}(m\varphi(a) + \eta(\varphi(x), \varphi(a), m))}{\eta(\varphi(b), \varphi(a), m)} \\ & - \frac{\eta^{n-\frac{\alpha}{k}}(\varphi(x), \varphi(b), m) f^{(n)}(m\varphi(b) + \eta(\varphi(x), \varphi(b), m))}{\eta(\varphi(b), \varphi(a), m)} \\ & + (-1)^{n+1} \frac{(nk - \alpha) \Gamma_k(n - \frac{\alpha}{k})}{\eta(\varphi(b), \varphi(a), m)} \\ & \times \left[{}^c D_{(m\varphi(a) + \eta(\varphi(x), \varphi(a), m))}^{\alpha, k} f(m\varphi(a)) - {}^c D_{(m\varphi(b) + \eta(\varphi(x), \varphi(b), m))}^{\alpha, k} f(m\varphi(b)) \right] \\ & = \frac{\eta^{n-\frac{\alpha}{k}+1}(\varphi(x), \varphi(a), m)}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 t^{n-\frac{\alpha}{k}} f^{(n+1)}(m\varphi(a) + t\eta(\varphi(x), \varphi(a), m)) dt \\ & - \frac{\eta^{n-\frac{\alpha}{k}+1}(\varphi(x), \varphi(b), m)}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 t^{n-\frac{\alpha}{k}} f^{(n+1)}(m\varphi(b) + t\eta(\varphi(x), \varphi(b), m)) dt. \quad (3.1) \end{aligned}$$

We denote

$$\begin{aligned} & I_{f,\eta,\varphi}(x; \alpha, k, n, m, a, b) \\ & = \frac{\eta^{n-\frac{\alpha}{k}+1}(\varphi(x), \varphi(a), m)}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 t^{n-\frac{\alpha}{k}} f^{(n+1)}(m\varphi(a) + t\eta(\varphi(x), \varphi(a), m)) dt \end{aligned}$$

$$-\frac{\eta^{n-\frac{\alpha}{k}+1}(\varphi(x), \varphi(b), m)}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 t^{n-\frac{\alpha}{k}} f^{(n+1)}(m\varphi(b) + t\eta(\varphi(x), \varphi(b), m)) dt. \quad (3.2)$$

Proof. Integrating by parts, we get

$$\begin{aligned} I_{f,\eta,\varphi}(x; \alpha, k, n, m, a, b) &= \frac{\eta^{n-\frac{\alpha}{k}+1}(\varphi(x), \varphi(a), m)}{\eta(\varphi(b), \varphi(a), m)} \\ &\times \left[\frac{t^{n-\frac{\alpha}{k}} f^{(n)}(m\varphi(a) + t\eta(\varphi(x), \varphi(a), m))}{\eta(\varphi(x), \varphi(a), m)} \right]_0^1 \\ &- \frac{n - \frac{\alpha}{k}}{\eta(\varphi(x), \varphi(a), m)} \int_0^1 t^{n-\frac{\alpha}{k}-1} f^{(n)}(m\varphi(a) + t\eta(\varphi(x), \varphi(a), m)) dt \\ &- \frac{\eta^{n-\frac{\alpha}{k}+1}(\varphi(x), \varphi(b), m)}{\eta(\varphi(b), \varphi(a), m)} \\ &\times \left[\frac{t^{n-\frac{\alpha}{k}} f^{(n)}(m\varphi(b) + t\eta(\varphi(x), \varphi(b), m))}{\eta(\varphi(x), \varphi(b), m)} \right]_0^1 \\ &- \frac{n - \frac{\alpha}{k}}{\eta(\varphi(x), \varphi(b), m)} \int_0^1 t^{n-\frac{\alpha}{k}-1} f^{(n)}(m\varphi(b) + t\eta(\varphi(x), \varphi(b), m)) dt \\ &= \frac{\eta^{n-\frac{\alpha}{k}}(\varphi(x), \varphi(a), m) f^{(n)}(m\varphi(a) + \eta(\varphi(x), \varphi(a), m))}{\eta(\varphi(b), \varphi(a), m)} \\ &- \frac{\eta^{n-\frac{\alpha}{k}}(\varphi(x), \varphi(b), m) f^{(n)}(m\varphi(b) + \eta(\varphi(x), \varphi(b), m))}{\eta(\varphi(b), \varphi(a), m)} \\ &+ (-1)^{n+1} \frac{(nk - \alpha)\Gamma_k(n - \frac{\alpha}{k})}{\eta(\varphi(b), \varphi(a), m)} \\ &\times \left[{}^c D_{(m\varphi(a) + \eta(\varphi(x), \varphi(a), m))}^{\alpha, k} f(m\varphi(a)) - {}^c D_{(m\varphi(b) + \eta(\varphi(x), \varphi(b), m))}^{\alpha, k} f(m\varphi(b)) \right]. \end{aligned}$$

This completes the proof of the lemma. \square

Using Lemma 3.1, we now state the following theorems for the corresponding version for power of $(n+1)$ -derivative.

Theorem 3.2. *Let $\alpha > 0$, $k \geq 1$, $0 < r \leq 1$, $p_1, p_2 > -1$ and $\alpha \notin \{1, 2, 3, \dots\}$, $n = [\alpha] + 1$. Suppose $h_1, h_2 : [0, 1] \rightarrow [0, +\infty)$ and $\varphi : I \rightarrow K$ are continuous functions. Suppose $K \subseteq \mathbb{R}$ be an open m -invex subset with respect to $\eta : K \times K \times (0, 1) \rightarrow \mathbb{R}$ for some fixed $m \in (0, 1]$. Assume that $f : K \rightarrow (0, +\infty)$ is a function on K° (the interior of K) such that $f \in C^{n+1}[m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)]$, where*

$\eta(\varphi(b), \varphi(a), m) > 0$. If $(f^{(n+1)})^q$ is generalized relative semi- $(r; m, p_1, p_2, h_1, h_2)$ -preinvex mappings, $q > 1$, $p^{-1} + q^{-1} = 1$, then the following inequality for Caputo k -fractional derivatives holds:

$$\begin{aligned} |I_{f,n,\varphi}(x; \alpha, k, n, m, a, b)| &\leq \left(\frac{1}{p(n - \frac{\alpha}{k}) + 1} \right)^{\frac{1}{p}} \frac{1}{\eta(\varphi(b), \varphi(a), m)} \\ &\times \left\{ |\eta(\varphi(x), \varphi(a), m)|^{n - \frac{\alpha}{k} + 1} \left[m \left(f^{(n+1)}(a) \right)^{rq} I^r(h_1(t); r, p_1) + \left(f^{(n+1)}(x) \right)^{rq} I^r(h_2(t); r, p_2) \right]^{\frac{1}{rq}} \right. \\ &\left. + |\eta(\varphi(x), \varphi(b), m)|^{n - \frac{\alpha}{k} + 1} \left[m \left(f^{(n+1)}(b) \right)^{rq} I^r(h_1(t); r, p_1) + \left(f^{(n+1)}(x) \right)^{rq} I^r(h_2(t); r, p_2) \right]^{\frac{1}{rq}} \right\}, \end{aligned} \quad (3.3)$$

where

$$I(h_i(t); r, p_i) := \int_0^1 h_i^{\frac{p_i}{r}}(t) dt, \quad \forall i = 1, 2.$$

Proof. Suppose that $q > 1$ and $0 < r \leq 1$. From Lemma 3.1, generalized relative semi- $(r; m, p_1, p_2, h_1, h_2)$ -preinvexity of $(f^{(n+1)})^q$, Hölder inequality, Minkowski inequality and properties of the modulus, we have

$$\begin{aligned} &|I_{f,n,\varphi}(x; \alpha, k, n, m, a, b)| \\ &\leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{n - \frac{\alpha}{k} + 1}}{|\eta(\varphi(b), \varphi(a), m)|} \int_0^1 t^{n - \frac{\alpha}{k}} |f^{(n+1)}(m\varphi(a) + t\eta(\varphi(x), \varphi(a), m))| dt \\ &\quad + \frac{|\eta(\varphi(x), \varphi(b), m)|^{n - \frac{\alpha}{k} + 1}}{|\eta(\varphi(b), \varphi(a), m)|} \int_0^1 t^{n - \frac{\alpha}{k}} |f^{(n+1)}(m\varphi(b) + t\eta(\varphi(x), \varphi(b), m))| dt \\ &\leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{n - \frac{\alpha}{k} + 1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{p(n - \frac{\alpha}{k})} dt \right)^{\frac{1}{p}} \\ &\quad \times \left(\int_0^1 \left(f^{(n+1)}(m\varphi(a) + t\eta(\varphi(x), \varphi(a), m)) \right)^q dt \right)^{\frac{1}{q}} \\ &\quad + \frac{|\eta(\varphi(x), \varphi(b), m)|^{n - \frac{\alpha}{k} + 1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{p(n - \frac{\alpha}{k})} dt \right)^{\frac{1}{p}} \\ &\quad \times \left(\int_0^1 \left(f^{(n+1)}(m\varphi(b) + t\eta(\varphi(x), \varphi(b), m)) \right)^q dt \right)^{\frac{1}{q}} \\ &\leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{n - \frac{\alpha}{k} + 1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{p(n - \frac{\alpha}{k})} dt \right)^{\frac{1}{p}} \\ &\quad \times \left(\int_0^1 \left[mh_1^{p_1}(t) \left(f^{(n+1)}(a) \right)^{rq} + h_2^{p_2}(t) \left(f^{(n+1)}(x) \right)^{rq} \right]^{\frac{1}{r}} dt \right)^{\frac{1}{q}} \end{aligned}$$

$$\begin{aligned}
& + \frac{|\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{p(n-\frac{\alpha}{k})} dt \right)^{\frac{1}{p}} \\
& \times \left(\int_0^1 \left[mh_1^{p_1}(t) \left(f^{(n+1)}(b) \right)^{rq} + h_2^{p_2}(t) \left(f^{(n+1)}(x) \right)^{rq} \right]^{\frac{1}{r}} dt \right)^{\frac{1}{q}} \\
& \leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{p(n-\frac{\alpha}{k})} dt \right)^{\frac{1}{p}} \\
& \times \left\{ \left(\int_0^1 m^{\frac{1}{r}} \left(f^{(n+1)}(a) \right)^q h_1^{\frac{p_1}{r}}(t) dt \right)^r + \left(\int_0^1 \left(f^{(n+1)}(x) \right)^q h_2^{\frac{p_2}{r}}(t) dt \right)^r \right\}^{\frac{1}{rq}} \\
& + \frac{|\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{p(n-\frac{\alpha}{k})} dt \right)^{\frac{1}{p}} \\
& \times \left\{ \left(\int_0^1 m^{\frac{1}{r}} \left(f^{(n+1)}(b) \right)^q h_1^{\frac{p_1}{r}}(t) dt \right)^r + \left(\int_0^1 \left(f^{(n+1)}(x) \right)^q h_2^{\frac{p_2}{r}}(t) dt \right)^r \right\}^{\frac{1}{rq}} \\
& = \left(\frac{1}{p(n-\frac{\alpha}{k})+1} \right)^{\frac{1}{p}} \frac{1}{\eta(\varphi(b), \varphi(a), m)} \\
& \times \left\{ |\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1} \left[m \left(f^{(n+1)}(a) \right)^{rq} I^r(h_1(t); r, p_1) + \left(f^{(n+1)}(x) \right)^{rq} I^r(h_2(t); r, p_2) \right]^{\frac{1}{rq}} \right. \\
& \left. + |\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1} \left[m \left(f^{(n+1)}(b) \right)^{rq} I^r(h_1(t); r, p_1) + \left(f^{(n+1)}(x) \right)^{rq} I^r(h_2(t); r, p_2) \right]^{\frac{1}{rq}} \right\}.
\end{aligned}$$

So, the proof of this theorem is completed. \square

We point out some special cases of Theorem 3.2.

Corollary 3.3. *In Theorem 3.2 for $h_1(t) = h_2(t) = h(t)$, $p_1 = p_2 = m = k = r = 1$, $\eta(\varphi(y), \varphi(x), m) = \varphi(y) - m\varphi(x)$, $\varphi(x) = x$, $\forall x \in I$ and $f^{(n+1)} \leq K$, we get the following inequality for Caputo fractional derivatives:*

$$\begin{aligned}
& \left| \left[\frac{(x-a)^{n-\alpha} - (x-b)^{n-\alpha}}{b-a} \right] f^{(n)}(x) + (-1)^{n+1} \frac{\Gamma(n-\alpha+1)}{b-a} \left[{}^c D_{x-}^\alpha f(a) - {}^c D_{x-}^\alpha f(b) \right] \right| \\
& \leq \frac{2^{\frac{1}{q}} K}{[p(n-\alpha)+1]^{\frac{1}{p}}} \left(\int_0^1 h(t) dt \right)^{\frac{1}{q}} \left[\frac{(x-a)^{n-\alpha+1} + (b-x)^{n-\alpha+1}}{b-a} \right]. \quad (3.4)
\end{aligned}$$

Corollary 3.4. *In Theorem 3.2 for $h_1(t) = h(1-t)$ and $h_2(t) = h(t)$, we have the following inequality for generalized relative semi- $(r; m, p_1, p_2, h)$ -preinvex mappings*

$$\begin{aligned} |I_{f,n,\varphi}(x; \alpha, k, n, m, a, b)| &\leq \left(\frac{1}{p(n - \frac{\alpha}{k}) + 1} \right)^{\frac{1}{p}} \frac{1}{\eta(\varphi(b), \varphi(a), m)} \\ &\times \left\{ |\eta(\varphi(x), \varphi(a), m)|^{n - \frac{\alpha}{k} + 1} \left[m \left(f^{(n+1)}(a) \right)^{rq} I^r(h(1-t); r, p_1) + \left(f^{(n+1)}(x) \right)^{rq} I^r(h(t); r, p_2) \right]^{\frac{1}{rq}} \right. \\ &\quad \left. + |\eta(\varphi(x), \varphi(b), m)|^{n - \frac{\alpha}{k} + 1} \left[m \left(f^{(n+1)}(b) \right)^{rq} I^r(h(1-t); r, p_1) + \left(f^{(n+1)}(x) \right)^{rq} I^r(h(t); r, p_2) \right]^{\frac{1}{rq}} \right\}. \end{aligned} \quad (3.5)$$

Corollary 3.5. *In Theorem 3.2 for $h_1(t) = (1-t)^s$, $h_2(t) = t^s$, we have the following inequality for generalized relative semi- $(r; m, p_1, p_2, s)$ -Breckner-preinvex mappings*

$$\begin{aligned} |I_{f,n,\varphi}(x; \alpha, k, n, m, a, b)| &\leq \left(\frac{1}{p(n - \frac{\alpha}{k}) + 1} \right)^{\frac{1}{p}} \frac{1}{\eta(\varphi(b), \varphi(a), m)} \\ &\times \left\{ |\eta(\varphi(x), \varphi(a), m)|^{n - \frac{\alpha}{k} + 1} \left[m \left(f^{(n+1)}(a) \right)^{rq} \left(\frac{r}{r + sp_1} \right)^r + \left(f^{(n+1)}(x) \right)^{rq} \left(\frac{r}{r + sp_2} \right)^r \right]^{\frac{1}{rq}} \right. \\ &\quad \left. + |\eta(\varphi(x), \varphi(b), m)|^{n - \frac{\alpha}{k} + 1} \left[m \left(f^{(n+1)}(b) \right)^{rq} \left(\frac{r}{r + sp_1} \right)^r + \left(f^{(n+1)}(x) \right)^{rq} \left(\frac{r}{r + sp_2} \right)^r \right]^{\frac{1}{rq}} \right\}. \end{aligned} \quad (3.6)$$

Corollary 3.6. *In Theorem 3.2 for $h_1(t) = (1-t)^{-s}$, $h_2(t) = t^{-s}$, we have the following inequality for generalized relative semi- $(r; m, p_1, p_2, s)$ -Godunova-Levin-Dragomir-preinvex mappings*

$$\begin{aligned} |I_{f,n,\varphi}(x; \alpha, k, n, m, a, b)| &\leq \left(\frac{1}{p(n - \frac{\alpha}{k}) + 1} \right)^{\frac{1}{p}} \frac{1}{\eta(\varphi(b), \varphi(a), m)} \\ &\times \left\{ |\eta(\varphi(x), \varphi(a), m)|^{n - \frac{\alpha}{k} + 1} \left[m \left(f^{(n+1)}(a) \right)^{rq} \left(\frac{r}{r - sp_1} \right)^r + \left(f^{(n+1)}(x) \right)^{rq} \left(\frac{r}{r - sp_2} \right)^r \right]^{\frac{1}{rq}} \right. \\ &\quad \left. + |\eta(\varphi(x), \varphi(b), m)|^{n - \frac{\alpha}{k} + 1} \left[m \left(f^{(n+1)}(b) \right)^{rq} \left(\frac{r}{r - sp_1} \right)^r + \left(f^{(n+1)}(x) \right)^{rq} \left(\frac{r}{r - sp_2} \right)^r \right]^{\frac{1}{rq}} \right\}. \end{aligned} \quad (3.7)$$

Corollary 3.7. *In Theorem 3.2 for $h_1(t) = h_2(t) = t(1-t)$, we have the following inequality for generalized relative semi- $(r; m, p_1, p_2, \eta)$ -preinvex mappings*

$$\begin{aligned} |I_{f,\eta,\varphi}(x; \alpha, k, n, m, a, b)| &\leq \left(\frac{1}{p(n - \frac{\alpha}{k}) + 1} \right)^{\frac{1}{p}} \frac{1}{\eta(\varphi(b), \varphi(a), m)} \\ &\times \left\{ |\eta(\varphi(x), \varphi(a), m)|^{n - \frac{\alpha}{k} + 1} \left[m \left(f^{(n+1)}(a) \right)^{rq} \beta^r \left(\frac{p_1}{r} + 1, \frac{p_1}{r} + 1 \right) \right. \right. \\ &\quad \left. \left. + \left(f^{(n+1)}(x) \right)^{rq} \beta^r \left(\frac{p_2}{r} + 1, \frac{p_2}{r} + 1 \right) \right]^{\frac{1}{rq}} \right. \\ &\quad \left. + |\eta(\varphi(x), \varphi(b), m)|^{n - \frac{\alpha}{k} + 1} \left[m \left(f^{(n+1)}(b) \right)^{rq} \beta^r \left(\frac{p_1}{r} + 1, \frac{p_1}{r} + 1 \right) \right. \right. \\ &\quad \left. \left. + \left(f^{(n+1)}(x) \right)^{rq} \beta^r \left(\frac{p_2}{r} + 1, \frac{p_2}{r} + 1 \right) \right]^{\frac{1}{rq}} \right\}. \end{aligned} \quad (3.8)$$

Corollary 3.8. *In Theorem 3.2 for $h_1(t) = \frac{\sqrt{1-t}}{2\sqrt{t}}$, $h_2(t) = \frac{\sqrt{t}}{2\sqrt{1-t}}$, we have the following inequality for generalized relative semi- $(r; m, p_1, p_2)$ -MT-preinvex mappings*

$$\begin{aligned} |I_{f,\eta,\varphi}(x; \alpha, k, n, m, a, b)| &\leq \left(\frac{1}{p(n - \frac{\alpha}{k}) + 1} \right)^{\frac{1}{p}} \frac{1}{\eta(\varphi(b), \varphi(a), m)} \\ &\times \left\{ |\eta(\varphi(x), \varphi(a), m)|^{n - \frac{\alpha}{k} + 1} \left[m \left(f^{(n+1)}(a) \right)^{rq} \left(\frac{1}{2} \right)^{p_1} \beta^r \left(1 - \frac{p_1}{2r}, 1 + \frac{p_1}{2r} \right) \right. \right. \\ &\quad \left. \left. + \left(f^{(n+1)}(x) \right)^{rq} \left(\frac{1}{2} \right)^{p_2} \beta^r \left(1 - \frac{p_2}{2r}, 1 + \frac{p_2}{2r} \right) \right]^{\frac{1}{rq}} \right. \\ &\quad \left. + |\eta(\varphi(x), \varphi(b), m)|^{n - \frac{\alpha}{k} + 1} \left[m \left(f^{(n+1)}(b) \right)^{rq} \left(\frac{1}{2} \right)^{p_1} \beta^r \left(1 - \frac{p_1}{2r}, 1 + \frac{p_1}{2r} \right) \right. \right. \\ &\quad \left. \left. + \left(f^{(n+1)}(x) \right)^{rq} \left(\frac{1}{2} \right)^{p_2} \beta^r \left(1 - \frac{p_2}{2r}, 1 + \frac{p_2}{2r} \right) \right]^{\frac{1}{rq}} \right\}. \end{aligned} \quad (3.9)$$

Theorem 3.9. *Let $\alpha > 0$, $k \geq 1$, $0 < r \leq 1$, $p_1, p_2 > -1$ and $\alpha \notin \{1, 2, 3, \dots\}$, $n = [\alpha] + 1$. Suppose $h_1, h_2 : [0, 1] \rightarrow [0, +\infty)$ and $\varphi : I \rightarrow K$ are continuous functions. Suppose $K \subseteq \mathbb{R}$ be an open m -invex subset with respect to $\eta : K \times K \times (0, 1) \rightarrow \mathbb{R}$ for some fixed $m \in (0, 1]$. Assume that $f : K \rightarrow (0, +\infty)$ is a function on K° (the interior of K) such that $f \in C^{n+1}[m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)]$, where*

$\eta(\varphi(b), \varphi(a), m) > 0$. If $(f^{(n+1)})^q$ is generalized relative semi- $(r; m, p_1, p_2, h_1, h_2)$ -preinvex mappings, $q \geq 1$, then the following inequality for Caputo k -fractional derivatives holds:

$$\begin{aligned} |I_{f,\eta,\varphi}(x; \alpha, k, n, m, a, b)| &\leq \left(\frac{1}{n - \frac{\alpha}{k} + 1} \right)^{1-\frac{1}{q}} \frac{1}{\eta(\varphi(b), \varphi(a), m)} \\ &\times \left\{ |\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1} \left[m \left(f^{(n+1)}(a) \right)^{rq} I^r(h_1(t); r, \alpha, k, n, p_1) \right. \right. \\ &\quad \left. \left. + \left(f^{(n+1)}(x) \right)^{rq} I^r(h_2(t); r, \alpha, k, n, p_2) \right]^{\frac{1}{rq}} \right. \\ &\quad \left. + |\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1} \left[m \left(f^{(n+1)}(b) \right)^{rq} I^r(h_1(t); r, \alpha, k, n, p_1) \right. \right. \\ &\quad \left. \left. + \left(f^{(n+1)}(x) \right)^{rq} I^r(h_2(t); r, \alpha, k, n, p_2) \right]^{\frac{1}{rq}} \right\}, \end{aligned} \quad (3.10)$$

where

$$I(h_i(t); r, \alpha, k, n, p_i) := \int_0^1 t^{n-\frac{\alpha}{k}} h_i^{\frac{p_i}{r}}(t) dt, \quad \forall i = 1, 2.$$

Proof. Suppose that $q \geq 1$ and $0 < r \leq 1$. From Lemma 3.1, generalized relative semi- $(r; m, p_1, p_2, h_1, h_2)$ -preinvexity of $(f^{(n+1)})^q$, the well-known power mean inequality, Minkowski inequality and properties of the modulus, we have

$$\begin{aligned} &|I_{f,\eta,\varphi}(x; \alpha, k, n, m, a, b)| \\ &\leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1}}{|\eta(\varphi(b), \varphi(a), m)|} \int_0^1 t^{n-\frac{\alpha}{k}} |f^{(n+1)}(m\varphi(a) + t\eta(\varphi(x), \varphi(a), m))| dt \\ &\quad + \frac{|\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{|\eta(\varphi(b), \varphi(a), m)|} \int_0^1 t^{n-\frac{\alpha}{k}} |f^{(n+1)}(m\varphi(b) + t\eta(\varphi(x), \varphi(b), m))| dt \\ &\leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{n-\frac{\alpha}{k}} dt \right)^{1-\frac{1}{q}} \\ &\quad \times \left(\int_0^1 t^{n-\frac{\alpha}{k}} \left(f^{(n+1)}(m\varphi(a) + t\eta(\varphi(x), \varphi(a), m)) \right)^q dt \right)^{\frac{1}{q}} \\ &\quad + \frac{|\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{n-\frac{\alpha}{k}} dt \right)^{1-\frac{1}{q}} \\ &\quad \times \left(\int_0^1 t^{n-\frac{\alpha}{k}} \left(f^{(n+1)}(m\varphi(b) + t\eta(\varphi(x), \varphi(b), m)) \right)^q dt \right)^{\frac{1}{q}} \\ &\leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{n-\frac{\alpha}{k}} dt \right)^{1-\frac{1}{q}} \end{aligned}$$

$$\begin{aligned}
& \times \left(\int_0^1 t^{n-\frac{\alpha}{k}} \left[mh_1^{p_1}(t) \left(f^{(n+1)}(a) \right)^{rq} + h_2^{p_2}(t) \left(f^{(n+1)}(x) \right)^{rq} \right]^{\frac{1}{r}} dt \right)^{\frac{1}{q}} \\
& + \frac{|\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{n-\frac{\alpha}{k}} dt \right)^{1-\frac{1}{q}} \\
& \times \left(\int_0^1 t^{n-\frac{\alpha}{k}} \left[mh_1^{p_1}(t) \left(f^{(n+1)}(b) \right)^{rq} + h_2^{p_2}(t) \left(f^{(n+1)}(x) \right)^{rq} \right]^{\frac{1}{r}} dt \right)^{\frac{1}{q}} \\
& \leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{n-\frac{\alpha}{k}} dt \right)^{1-\frac{1}{q}} \\
& \times \left\{ \left(\int_0^1 m^{\frac{1}{r}} \left(f^{(n+1)}(a) \right)^q t^{n-\frac{\alpha}{k}} h_1^{\frac{p_1}{r}}(t) dt \right)^r + \left(\int_0^1 \left(f^{(n+1)}(x) \right)^q t^{n-\frac{\alpha}{k}} h_2^{\frac{p_2}{r}}(t) dt \right)^r \right\}^{\frac{1}{rq}} \\
& + \frac{|\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1}}{\eta(\varphi(b), \varphi(a), m)} \left(\int_0^1 t^{n-\frac{\alpha}{k}} dt \right)^{1-\frac{1}{q}} \\
& \times \left\{ \left(\int_0^1 m^{\frac{1}{r}} \left(f^{(n+1)}(b) \right)^q t^{n-\frac{\alpha}{k}} h_1^{\frac{p_1}{r}}(t) dt \right)^r + \left(\int_0^1 \left(f^{(n+1)}(x) \right)^q t^{n-\frac{\alpha}{k}} h_2^{\frac{p_2}{r}}(t) dt \right)^r \right\}^{\frac{1}{rq}} \\
& = \left(\frac{1}{n - \frac{\alpha}{k} + 1} \right)^{1-\frac{1}{q}} \frac{1}{\eta(\varphi(b), \varphi(a), m)} \\
& \times \left\{ |\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1} \left[m \left(f^{(n+1)}(a) \right)^{rq} I^r(h_1(t); r, \alpha, k, n, p_1) \right. \right. \\
& \quad \left. \left. + \left(f^{(n+1)}(x) \right)^{rq} I^r(h_2(t); r, \alpha, k, n, p_2) \right]^{\frac{1}{rq}} \right. \\
& \quad \left. + |\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1} \left[m \left(f^{(n+1)}(b) \right)^{rq} I^r(h_1(t); r, \alpha, k, n, p_1) \right. \right. \\
& \quad \left. \left. + \left(f^{(n+1)}(x) \right)^{rq} I^r(h_2(t); r, \alpha, k, n, p_2) \right]^{\frac{1}{rq}} \right].
\end{aligned}$$

So, the proof of this theorem is completed. \square

We point out some special cases of Theorem 3.9.

Corollary 3.10. *In Theorem 3.9 for $h_1(t) = h_2(t) = h(t)$, $p_1 = p_2 = m = k = r = 1$, $\eta(\varphi(y), \varphi(x), m) = \varphi(y) - m\varphi(x)$, $\varphi(x) = x$, $\forall x \in I$ and $f^{(n+1)} \leq K$, we get the following inequality for Caputo fractional derivatives:*

$$\left| \left[\frac{(x-a)^{n-\alpha} - (x-b)^{n-\alpha}}{b-a} \right] f^{(n)}(x) + (-1)^{n+1} \frac{\Gamma(n-\alpha+1)}{b-a} \left[{}^c D_{x-}^\alpha f(a) - {}^c D_{x-}^\alpha f(b) \right] \right|$$

$$\leq 2^{\frac{1}{q}} K \left(\frac{1}{n - \alpha + 1} \right)^{1 - \frac{1}{q}} I^{\frac{1}{q}}(h(t); 1, \alpha, k, n, 1) \left[\frac{(x - a)^{n - \alpha + 1} + (b - x)^{n - \alpha + 1}}{b - a} \right]. \quad (3.11)$$

Corollary 3.11. *In Theorem 3.9 for $h_1(t) = h(1 - t)$ and $h_2(t) = h(t)$, we have the following inequality for generalized relative semi- $(r; m, p_1, p_2, h)$ -preinvex mappings*

$$\begin{aligned} |I_{f,\eta,\varphi}(x; \alpha, k, n, m, a, b)| &\leq \left(\frac{1}{n - \frac{\alpha}{k} + 1} \right)^{1 - \frac{1}{q}} \frac{1}{\eta(\varphi(b), \varphi(a), m)} \\ &\times \left\{ |\eta(\varphi(x), \varphi(a), m)|^{n - \frac{\alpha}{k} + 1} \left[m \left(f^{(n+1)}(a) \right)^{rq} I^r(h(1 - t); r, \alpha, k, n, p_1) \right. \right. \\ &\quad + \left(f^{(n+1)}(x) \right)^{rq} I^r(h(t); r, \alpha, k, n, p_2) \left. \right]^{1/rq} \\ &\quad + |\eta(\varphi(x), \varphi(b), m)|^{n - \frac{\alpha}{k} + 1} \left[m \left(f^{(n+1)}(b) \right)^{rq} I^r(h(1 - t); r, \alpha, k, n, p_1) \right. \\ &\quad \left. \left. + \left(f^{(n+1)}(x) \right)^{rq} I^r(h(t); r, \alpha, k, n, p_2) \right]^{1/rq} \right\}. \end{aligned} \quad (3.12)$$

Corollary 3.12. *In Theorem 3.9 for $h_1(t) = (1 - t)^s$, $h_2(t) = t^s$, we have the following inequality for generalized relative semi- $(r; m, p_1, p_2, s)$ -Breckner-preinvex mappings*

$$\begin{aligned} |I_{f,\eta,\varphi}(x; \alpha, k, n, m, a, b)| &\leq \left(\frac{1}{n - \frac{\alpha}{k} + 1} \right)^{1 - \frac{1}{q}} \frac{1}{\eta(\varphi(b), \varphi(a), m)} \\ &\times \left\{ |\eta(\varphi(x), \varphi(a), m)|^{n - \frac{\alpha}{k} + 1} \left[m \left(f^{(n+1)}(a) \right)^{rq} \beta^r \left(n - \frac{\alpha}{k} + 1, \frac{sp_1}{r} + 1 \right) \right. \right. \\ &\quad + \left(f^{(n+1)}(x) \right)^{rq} \left(\frac{1}{n + \frac{sp_2}{r} - \frac{\alpha}{k} + 1} \right)^r \left. \right]^{1/rq} \\ &\quad + |\eta(\varphi(x), \varphi(b), m)|^{n - \frac{\alpha}{k} + 1} \left[m \left(f^{(n+1)}(b) \right)^{rq} \beta^r \left(n - \frac{\alpha}{k} + 1, \frac{sp_1}{r} + 1 \right) \right. \\ &\quad \left. \left. + \left(f^{(n+1)}(x) \right)^{rq} \left(\frac{1}{n + \frac{sp_2}{r} - \frac{\alpha}{k} + 1} \right)^r \right]^{1/rq} \right\}. \end{aligned} \quad (3.13)$$

Corollary 3.13. *In Theorem 3.9 for $h_1(t) = (1 - t)^{-s}$, $h_2(t) = t^{-s}$, we have the following inequality for generalized relative semi- $(r; m, p_1, p_2, s)$ -Godunova-Levin-Dragomir-preinvex mappings*

$$\begin{aligned}
|I_{f,\eta,\varphi}(x; \alpha, k, n, m, a, b)| &\leq \left(\frac{1}{n - \frac{\alpha}{k} + 1} \right)^{1-\frac{1}{q}} \frac{1}{\eta(\varphi(b), \varphi(a), m)} \\
&\times \left\{ |\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1} \left[m \left(f^{(n+1)}(a) \right)^{rq} \beta^r \left(n - \frac{\alpha}{k} + 1, 1 - \frac{sp_1}{r} \right) \right. \right. \\
&\quad + \left(f^{(n+1)}(x) \right)^{rq} \left(\frac{1}{n - \frac{sp_2}{r} - \frac{\alpha}{k} + 1} \right)^r \left. \right]^\frac{1}{rq} \\
&\quad + |\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1} \left[m \left(f^{(n+1)}(b) \right)^{rq} \beta^r \left(n - \frac{\alpha}{k} + 1, 1 - \frac{sp_1}{r} \right) \right. \\
&\quad \left. \left. + \left(f^{(n+1)}(x) \right)^{rq} \left(\frac{1}{n - \frac{sp_2}{r} - \frac{\alpha}{k} + 1} \right)^r \right]^\frac{1}{rq} \right\}. \tag{3.14}
\end{aligned}$$

Corollary 3.14. *In Theorem 3.9 for $h_1(t) = h_2(t) = t(1-t)$, we have the following inequality for generalized relative semi- $(r; m, p_1, p_2, tgs)$ -preinvex mappings*

$$\begin{aligned}
|I_{f,\eta,\varphi}(x; \alpha, k, n, m, a, b)| &\leq \left(\frac{1}{n - \frac{\alpha}{k} + 1} \right)^{1-\frac{1}{q}} \frac{1}{\eta(\varphi(b), \varphi(a), m)} \\
&\times \left\{ |\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1} \left[m \left(f^{(n+1)}(a) \right)^{rq} \beta^r \left(n + \frac{p_1}{r} - \frac{\alpha}{k} + 1, \frac{p_1}{r} + 1 \right) \right. \right. \\
&\quad + \left(f^{(n+1)}(x) \right)^{rq} \beta^r \left(n + \frac{p_2}{r} - \frac{\alpha}{k} + 1, \frac{p_2}{r} + 1 \right) \left. \right]^\frac{1}{rq} \\
&\quad + |\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1} \left[m \left(f^{(n+1)}(b) \right)^{rq} \beta^r \left(n + \frac{p_1}{r} - \frac{\alpha}{k} + 1, \frac{p_1}{r} + 1 \right) \right. \\
&\quad \left. \left. + \left(f^{(n+1)}(x) \right)^{rq} \beta^r \left(n + \frac{p_2}{r} - \frac{\alpha}{k} + 1, \frac{p_2}{r} + 1 \right) \right]^\frac{1}{rq} \right\}. \tag{3.15}
\end{aligned}$$

Corollary 3.15. *In Theorem 3.9 for $h_1(t) = \frac{\sqrt{1-t}}{2\sqrt{t}}$, $h_2(t) = \frac{\sqrt{t}}{2\sqrt{1-t}}$, we have the following inequality for generalized relative semi- $(r; m, p_1, p_2)$ -MT-preinvex mappings*

$$\begin{aligned}
|I_{f,\eta,\varphi}(x; \alpha, k, n, m, a, b)| &\leq \left(\frac{1}{n - \frac{\alpha}{k} + 1} \right)^{1-\frac{1}{q}} \frac{1}{\eta(\varphi(b), \varphi(a), m)} \\
&\times \left\{ |\eta(\varphi(x), \varphi(a), m)|^{n-\frac{\alpha}{k}+1} \left[m \left(f^{(n+1)}(a) \right)^{rq} \left(\frac{1}{2} \right)^{p_1} \beta^r \left(n - \frac{\alpha}{k} - \frac{p_1}{2r} + 1, 1 + \frac{p_1}{2r} \right) \right. \right. \\
&\quad \left. \left. + \left(f^{(n+1)}(x) \right)^{rq} \left(\frac{1}{2} \right)^{p_1} \beta^r \left(n - \frac{\alpha}{k} - \frac{p_1}{2r} + 1, 1 + \frac{p_1}{2r} \right) \right]^\frac{1}{rq} \right\}.
\end{aligned}$$

$$\begin{aligned}
& + \left(f^{(n+1)}(x) \right)^{rq} \left(\frac{1}{2} \right)^{p_2} \beta^r \left(n + \frac{p_2}{2r} - \frac{\alpha}{k} + 1, 1 - \frac{p_2}{2r} \right) \Bigg]^{1/rq} \\
& + |\eta(\varphi(x), \varphi(b), m)|^{n-\frac{\alpha}{k}+1} \left[m \left(f^{(n+1)}(b) \right)^{rq} \left(\frac{1}{2} \right)^{p_1} \beta^r \left(n - \frac{\alpha}{k} - \frac{p_1}{2r} + 1, 1 + \frac{p_1}{2r} \right) \right. \\
& \left. + \left(f^{(n+1)}(x) \right)^{rq} \left(\frac{1}{2} \right)^{p_2} \beta^r \left(n + \frac{p_2}{2r} - \frac{\alpha}{k} + 1, 1 - \frac{p_2}{2r} \right) \right]^{1/rq} \Bigg\}. \quad (3.16)
\end{aligned}$$

Remark 3.16. For $k = 1$, by our Theorems 3.2 and 3.9, we can get some new special Ostrowski type inequalities associated with generalized relative semi- $(r; m, p_1, p_2, h_1, h_2)$ -preinvex mappings via Caputo fractional derivatives of order α .

Remark 3.17. Also, applying our Theorems 3.2 and 3.9, we can deduce some new inequalities using special means associated with generalized relative semi- $(r; m, p_1, p_2, h_1, h_2)$ -preinvex mappings.

4. CONCLUSIONS

In this article, we first presented some integral inequalities for Gauss-Jacobi type quadrature formula involving generalized relative semi- $(r; m, p, q, h_1, h_2)$ -preinvex mappings. And then, a new identity concerning $(n+1)$ -differentiable mappings defined on m -invex set via Caputo k -fractional derivatives is derived. By using the notion of generalized relative semi- $(r; m, p, q, h_1, h_2)$ -preinvexity and the obtained identity as an auxiliary result, some new estimates with respect to Ostrowski type inequalities via Caputo k -fractional derivatives are established. It is pointed out that some new special cases are deduced from main results of the article. Motivated by this new interesting class of generalized relative semi- $(r; m, p, q, h_1, h_2)$ -preinvex mappings we can indeed see to be vital for fellow researchers and scientists working in the same domain. We conclude that our methods considered here may be a stimulant for further investigations concerning Ostrowski, Hermite-Hadamard and Simpson type integral inequalities for various kinds of preinvex functions involving local fractional integrals, fractional integral operators, Caputo k -fractional derivatives, q -calculus, (p, q) -calculus, time scale calculus and conformable fractional integrals.

REFERENCES

- [1] Abdeljawad, T., On conformable fractional calculus, *J. Comput. Appl. Math.*, **279**, (2015), 57-66.
- [2] Agarwal, R. P., Luo, M. J. and Raina, R. K., On Ostrowski type inequalities, *Fasc. Math.*, **204**, (2016), 5-27.
- [3] Ahmадmir, M. and Ullah, R., Some inequalities of Ostrowski and Grüss type for triple integrals on time scales, *Tamkang J. Math.*, **42** (4), (2011), 415-426.

- [4] Alomari, M., Darus, M., Dragomir, S. S. and Cerone, P., Ostrowski type inequalities for functions whose derivatives are s -convex in the second sense, *Appl. Math. Lett.*, **23**, (2010), 1071-1076.
- [5] Antczak, T., Mean value in invexity analysis, *Nonlinear Anal.*, **60**, (2005), 1473-1484.
- [6] Chu, Y.-M., Khan, M. Adil, Ali, T. and Dragomir, S. S., Inequalities for α -fractional differentiable functions, *J. Inequal. Appl.*, (2017) 2017:93, 12 pages.
- [7] Dahmani, Z., On Minkowski and Hermite-Hadamard integral inequalities via fractional integration, *Ann. Funct. Anal.*, **1** (1), (2010), 51-58.
- [8] Dahmani, Z., New inequalities in fractional integrals, *Int. J. Nonlinear Sci.*, **9** (4), (2010), 493-497.
- [9] Dahmani, Z., Tabharit, L. and Taf, S., New generalizations of Grüss inequality using Riemann-Liouville fractional integrals, *Bull. Math. Anal. Appl.*, **2** (3), (2010), 93-99.
- [10] Dahmani, Z., Tabharit, L. and Taf, S., Some fractional integral inequalities, *Nonlinear. Sci. Lett. A*, **1** (2), (2010), 155-160.
- [11] Dragomir, S. S., On the Ostrowski's integral inequality for mappings with bounded variation and applications, *Math. Ineq. & Appl.*, **1** (2), (1998).
- [12] Dragomir, S. S., The Ostrowski integral inequality for Lipschitzian mappings and applications, *Comput. Math. Appl.*, **38**, (1999), 33-37.
- [13] Dragomir, S. S., Ostrowski-type inequalities for Lebesgue integral: A survey of recent results, *Aust. J. Math. Anal. Appl.*, **14** (1), (2017), 1-287.
- [14] Dragomir, S. S., Pečarić, J. and Persson, L. E., Some inequalities of Hadamard type, *Soochow J. Math.*, **21**, (1995), 335-341.
- [15] Dragomir, S. S. and Wang, S., An inequality of Ostrowski-Grüss type and its applications to the estimation of error bounds for some special means and for some numerical quadrature rules, *Comput. Math. Appl.*, **33** (11), (1997), 15-20.
- [16] Dragomir, S. S. and Wang, S., A new inequality of Ostrowski's type in L_1 -norm and applications to some special means and to some numerical quadrature rules, *Tamkang J. Math.*, **28**, (1997), 239-244.
- [17] Du, T. S., Liao, J. G. and Li, Y. J., Properties and integral inequalities of Hadamard-Simpson type for the generalized (s, m) -preinvex functions, *J. Nonlinear Sci. Appl.*, **9**, (2016), 3112-3126.
- [18] Farid, G., Some new Ostrowski type inequalities via fractional integrals, *Int. J. Anal. App.*, **14** (1), (2017), 64-68.
- [19] Farid, G., Javed, A. and Rehman, A. U., On Hadamard inequalities for n -times differentiable functions which are relative convex via Caputo k -fractional derivatives, *Nonlinear Anal. Forum*, to appear.
- [20] Hudzik, H. and Maligranda, L., Some remarks on s -convex functions, *Aequationes Math.*, **48**, (1994), 100-111.
- [21] Kashuri, A., Liko, R., Khan, M. Adil and Chu, Y.-M., Some new Ostrowski type fractional integral inequalities for generalized $(r; s, m, \varphi)$ -preinvex functions via Caputo k -fractional derivatives, *Journal of Fractional Calculus and Applications*, **9** (2), (2018), 163-177.
- [22] Kashuri, A. and Liko, R., Generalizations of Hermite-Hadamard and Ostrowski type inequalities for MT_m -preinvex functions, *Proyecciones*, **36** (1), (2017), 45-80.
- [23] Kashuri, A. and Liko, R., Ostrowski type fractional integral inequalities for generalized (s, m, φ) -preinvex functions, *Aust. J. Math. Anal. Appl.*, **13** (1), (2016), Article 16, 1-11.
- [24] Katugampola, U. N., A new approach to generalized fractional derivatives, *Bulletin Math. Anal. Appl.*, **6** (4), (2014), 1-15.
- [25] Khalil, R., Horani, M. Al, Yousef, A. and Sababheh, M., A new definition of fractional derivative, *J. Comput. Appl. Math.*, **264**, (2014), 65-70.

- [26] Khan, M. Adil, Ali, T., Dragomir, S. S. and Sarikaya, M. Z., Hermite-Hadamard type inequalities for conformable fractional integrals, *RACSAM, Rev. R. Acad. Cienc. Exactas Fis. Nat., Ser. A Mat.*, 2017(2): 1-16 DOI10.1007/s13398-017-0408-5.
- [27] Khan, M. Adil, Chu, Y.-M., Kashuri, A., Liko, R. and Ali, G., New Hermite-Hadamard inequalities for conformable fractional integrals, *J. Funct. Spaces*, In press.
- [28] Khan, M. Adil, Chu, Y.-M., Khan, T. U. and Khan, J., Some new inequalities of Hermite-Hadamard type for s -convex functions with applications, *Open Math.*, **15**, (2017), 1414-1430.
- [29] Khan, M. Adil, Khurshid, Y. and Ali, T., Hermite-Hadamard inequality for fractional integrals via η -convex functions, *Acta Math. Univ. Comenianae*, Vol. LXXXVI (1), (2017), 153-164.
- [30] Kilbas, A. A., Srivastava, H. M. and Trujillo, J. J., Theory and applications of fractional differential equations, *North-Holland Math. Stud.*, **204**, Elsevier, New York-London, (2006).
- [31] Liu, Z., Some Ostrowski-Grüss type inequalities and applications, *Comput. Math. Appl.*, **53**, (2007), 73-79.
- [32] Liu, Z., Some companions of an Ostrowski type inequality and applications, *J. Inequal. in Pure and Appl. Math.*, **10** (2), (2009), Art. 52, 12 pp.
- [33] Liu, W., New integral inequalities involving beta function via P -convexity, *Miskolc Math. Notes*, **15** (2), (2014), 585-591.
- [34] Liu, W., Wen, W. and Park, J., Ostrowski type fractional integral inequalities for MT -convex functions, *Miskolc Math. Notes*, **16** (1), (2015), 249-256.
- [35] Liu, W., Wen, W. and Park, J., Hermite-Hadamard type inequalities for MT -convex functions via classical integrals and fractional integrals, *J. Nonlinear Sci. Appl.*, **9**, (2016), 766-777.
- [36] Matloka, M., Inequalities for h -preinvex functions, *Appl. Math. Comput.*, **234**, 52-57, (2014).
- [37] Matloka, M., Ostrowski type inequalities for functions whose derivatives are h -convex via fractional integrals, *Journal of Scientific Research and Reports*, **3** (12), (2014), 1633-1641.
- [38] Mitrinovic, D. S., Pečarić, J. E. and Fink, A. M., Classical and new inequalities in analysis, *Kluwer Academic Publishers, Dordrecht*, (1993).
- [39] Omotoyinbo, O. and Mogbodemu, A., Some new Hermite-Hadamard integral inequalities for convex functions, *Int. J. Sci. Innovation Tech.*, **1** (1), (2014), 1-12.
- [40] Özdemir, M. E., Kavurmacı, H. and Set, E. Ostrowski's type inequalities for (α, m) -convex functions, *Kyungpook Math. J.*, **50**, (2010), 371-378.
- [41] Özdemir, M. E., Set, E. and Alomari, M., Integral inequalities via several kinds of convexity, *Creat. Math. Inform.*, **20** (1), (2011), 62-73.
- [42] Pachpatte, B. G., On an inequality of Ostrowski type in three independent variables, *J. Math. Anal. Appl.*, **249**, (2000), 583-591.
- [43] Pachpatte, B. G., On a new Ostrowski type inequality in two independent variables, *Tamkang J. Math.*, **32** (1), (2001), 45-49.
- [44] Pachpatte, B. G., On some inequalities for convex functions, *RGMIA Res. Rep. Coll.*, **6**, (2003).
- [45] Peng, C., Zhou, C. and Du, T. S., Riemann-Liouville fractional Simpson's inequalities through generalized (m, h_1, h_2) -preinvexity, *Ital. J. Pure Appl. Math.*, **38**, (2017), 345-367.
- [46] Pini, R., Invexity and generalized convexity, *Optimization*, **22**, (1991), 513-525.
- [47] Purohit, S. D. and Kalla, S. L., Certain inequalities related to the Chebyshev's functional involving Erdelyi-Kober operators, *Scientia Series A: Math. Sci.*, **25**, (2014), 53-63.
- [48] Qi, F. and Xi, B. Y., Some integral inequalities of Simpson type for GA - e -convex functions, *Georgian Math. J.*, **20** (5), (2013), 775-788.
- [49] Rafiq, A., Mir, N. A. and Ahmad, F., Weighted Čebyšev-Ostrowski type inequalities, *Applied Math. Mechanics (English Edition)*, **28** (7), (2007), 901-906.
- [50] Raina, R. K., On generalized Wright's hypergeometric functions and fractional calculus operators, *East Asian Math. J.*, **21** (2), (2005), 191-203.

- [51] Sarikaya, M. Z., On the Ostrowski type integral inequality, *Acta Math. Univ. Comenianae*, **79** (1), (2010), 129-134.
- [52] Set, E., Akdemir, A. O. and Mumcu, I., Ostrowski type inequalities for functions whose derivatives are convex via conformable fractional integrals, Submitted.
- [53] Set, E., Akdemir, A. O. and Mumcu, I., Chebyshev type inequalities for conformable fractional integrals, Submitted.
- [54] Set, E. and Gözpınar, A., A study on Hermite-Hadamard type inequalities for s -convex functions via conformable fractional integrals, Submitted.
- [55] Set, E., Gözpınar, A. and Choi, J., Hermite-Hadamard type inequalities for twice differentiable m -convex functions via conformable fractional integrals, *Far East J. Math. Sci.*, **101** (4), (2017), 873-891.
- [56] Set, E., Karataş, S. S. and Khan, M. Adil, Hermite-Hadamard type inequalities obtained via fractional integral for differentiable m -convex and (α, m) -convex functions, *International Journal of Analysis*, Vol. 2016, Article ID 4765691, 8 pages.
- [57] Set, E. and Mumcu, I., Hermite-Hadamard-Fejér type inequalities for conformable fractional integrals, Submitted.
- [58] Set, E., Sarıkaya, M. Z. and Gözpınar, A., Some Hermite-Hadamard type inequalities for convex functions via conformable fractional integrals and related inequalities, *Creat. Math. Inform.*, Accepted paper.
- [59] Stancu, D. D., Coman, G. and Blaga, P., Analiză numerică și teoria aproximării, *Cluj-Napoca: Presa Universitară Clujeană*, **2**, (2002).
- [60] Tunç, M., Ostrowski type inequalities for functions whose derivatives are MT -convex, *J. Comput. Anal. Appl.*, **17** (4), (2014), 691-696.
- [61] Tunç, M., Göv, E. and Şanal, Ü., On tgs -convex function and their inequalities, *Facta Univ. Ser. Math. Inform.*, **30** (5), (2015), 679-691.
- [62] Ujević, N., Sharp inequalities of Simpson type and Ostrowski type, *Comput. Math. Appl.*, **48**, (2004), 145-151.
- [63] Varošanec, S., On h -convexity, *J. Math. Anal. Appl.*, **326** (1), (2007), 303-311.
- [64] Yıldız, G., Özdemir, M. E. and Sarıkaya, M. Z., New generalizations of Ostrowski-like type inequalities for fractional integrals, *Kyungpook Math. J.*, **56**, (2016), 161-172.
- [65] Youness, E. A., E-convex sets, E-convex functions, and E-convex programming, *J. Optim. Theory Appl.*, **102**, (1999), 439-450.
- [66] Zhongxue, L., On sharp inequalities of Simpson type and Ostrowski type in two independent variables, *Comput. Math. Appl.*, **56**, (2008), 2043-2047.

Current address: Artion Kashuri: Department of Mathematics, Faculty of Technical Science, University "Ismail Qemali", Vlora, Albania

E-mail address: artionkashuri@gmail.com

ORCID Address: <http://orcid.org/0000-0003-0115-3079>

Current address: Rozana Liko: Department of Mathematics, Faculty of Technical Science, University "Ismail Qemali", Vlora, Albania

E-mail address: rozanaliko86@gmail.com

ORCID Address: <http://orcid.org/x0000-0003-2439-8538>