



## -RESEARCH ARTICLE-

### Phylogenetic Relationships of Turbot Species (Scophthalmidae) Inferred from the Mitochondrial COIII Gene and Morphological Characters

Cemal Turan<sup>1\*</sup>, Petya Ivanova<sup>2</sup>, Mevlüt Gürlek<sup>1</sup>, Deniz Yağlıoğlu<sup>3</sup>, Deniz Ergüden<sup>1</sup>, Serpil Karan<sup>1</sup>, Servet A. Dođdu<sup>1</sup>, Ali Uyan<sup>1</sup>, Bayram Öztürk<sup>4</sup>, Venelin Nikolov<sup>2</sup>, Violin Raykov<sup>2</sup>, Ivan Dobrovolo<sup>2</sup>, Antonia Khanaychenko<sup>5</sup>

<sup>1</sup>Iskenderun Technical University, Faculty of Marine Sciences and Technology, Fisheries Genetics and Molecular Ecology Laboratory, Iskenderun, Turkey.

<sup>2</sup>Institute of Oceanology, Bulgarian Academy of Sciences, Department of Marine Biology and Ecology, Varna, Bulgaria.

<sup>3</sup>Duzce University, Biology Department, Duzce, Turkey.

<sup>4</sup>Istanbul University, Fisheries Faculty, Istanbul, Turkey.

<sup>5</sup>Institute of Marine Biological Research, Sevastopol, Crimea, (Ukrania) Russia.

#### Abstract

In this study, the validity, distribution and structure of three turbot species, *Scophthalmus maeoticus*, *S. maximus*, *S. rhombus*, belong to Scophthalmidae family in Turkish, Bulgarian and Russian coastal waters were determined with mtDNA sequencing of Cytochrome c oxidase subunit III (COIII). The sequencing of the COIII region revealed 8 bp variable and 6 bp parsimony informative sites between all turbot species. The overall genetic and haplotype diversities among all turbot species were found to be 0.004109 and 0.7655, respectively. Genetic distance analysis showed that the highest nucleotide differences was observed between *S. maximus* and *S. rhombus* species with a value of 0.09620 and, the lowest value (0.02482) was observed between *S. maximus* and *S. maeoticus* species. Neighbor Joining and Maximum Parsimony phylogenetic approaches resulted in the similar tree topologies that *S. maximus* and *S. maeoticus* were found as sister group, whereas *S. rhombus* was more divergent from this group. The mtDNA COIII gene is a useful genetic marker for species specific identification of the genus *Scophthalmus* due to its inter-specific heterogeneity producing a species-specific pattern. In morphological analyses, *S. rhombus* was most differentiated from *S. maximus* and *S. maeoticus*. The genetic data was supported by the detected morphometric variations among the turbot species.

\* Corresponding Author: Cemal TURAN, e-mail: cemal.turan@iste.edu.tr

**Keywords:** Turbot, phylogeny, mtDNA sequencing, morphology, the Black Sea, the Marmara Sea.

**Article history:** Received 05 February 2018, Accepted 24 January 2019, Available online 31 January 2019

## Introduction

The Scophthalmidae are a family of sinistral flatfish found in the North Atlantic Ocean, Baltic Sea, Mediterranean Sea, Black Sea and Marmara Sea are important species in the world due to their economic and ecological value (Chanet, 2003). Fishes of this family are known commonly as turbot which may exceed 1 m of length and 20 kg of weight (Froese & Pauly, 2018). Colors of these species are variable and they can adapt at colors of their environment. Scophthalmid fishes are benthic marine species, living on sandy and muddy bottoms, and younger individuals tend to live in shallower areas. These species generally feed on demersal and pelagic fish species, crustaceans and molluscs; the feeding ratio decreases during the spawning period and becomes more intensive in autumn and winter (Slastenenko, 1956; Karapetkova, 1980; Ivanov & Beverton, 1985).

In the world, the Scophthalmidae family is represented by 9 species (*Scophthalmus maximus*, *S. maeoticus*, *S. rhombus*, *S. aquosus*, *Lepidorhombus whiffiagonis*, *L. boscii*, *Zeugopterus punctatus*, *Z. regius*, *Phrynorhombus norvegicus*) within 4 genera, *Scophthalmus*, *Lepidorhombus*, *Zeugopterus*, *Phrynorhombus* (Nelson, 1994). Scophthalmidae family is represented with three species (*Scophthalmus maximus* (Linnaeus, 1758), *S. maeoticus* (Pallas, 1814), *S. rhombus* (Linnaeus, 1758) in the Black and Marmara Seas (Nelson, 1994; Turan, 2007). *Scophthalmus maximus*, *S. maeoticus* and *S. rhombus* are closely related congeneric species (Pardo et al., 2005; Turan, 2007; Azevedo et al., 2008) which show a similar distributional range (Blanquer et al., 1992; Pardo et al., 2001) and generally considered and excepted that there is one species, *Scophthalmus maximus*, exist in the Black Sea and Marmara Sea (Muus & Dahlström, 1978; Suziki et al., 2004). Unfortunately, the argument on the existence of the number of the species in these waters are still ongoing issue and need clarification.

In the Black and Marmara Seas there is a dilemma on the existence of the number of turbot species (Evseenko, 2003; Voronina, 2010; Froese & Pauly, 2018). The turbot, *Scophthalmus maximus* (Linnaeus, 1758), formerly recognized as *Psetta maxima* (Bailly & Chanet, 2010), distributed in the Northeast Atlantic, throughout the Mediterranean and along the European coasts to Arctic Circle as well as most of the Baltic Sea (Prado et al. 2018). The Black Sea brill, *Scophthalmus maeoticus* (Pallas, 1814), distributed in the Black Sea, was considered as a synonym of *S. maximus* by Bailly & Chanet (2010). Eschmeyer (2011) spelled a specific epithet as *maeotica* and accepted as *S. maeoticus*. The brill, *Scophthalmus rhombus* (Linnaeus, 1758), distributed in the Eastern Atlantic, Norway to Morocco and reported from Iceland (Jonsson, 1992), throughout the Mediterranean and Black Sea. Turan et al. (2016) reported that *S. rhombus* populations were seriously declined and rarely found in the Black Sea and Marmara Sea due to overfishing, habitat degradation and pollution and should be taken into consideration to be Near Threatened (NT) in the IUCN Red List of Threatened Species.

Molecular genetic studies on mtDNA have demonstrated useful in order to determine hypotheses about the phylogeny and phylogeography of marine species (Meyer, 1993; Avise, 1994; Turan et al. 2015a). The pattern of maternal inheritance and rapid rate of evolutionary change of mtDNA compared to nuclear DNA make it a suitable tool for genetic studies among taxa of several fish groups at multiple taxonomic levels (Kocher & Stepien, 1997; Zardoya et al., 1999; Durand et al., 2002). Sequence analysis of mtDNA regions may be a quick tool to reveal phylogenetic relationships of marine species (Avise, 1994; Tabata & Taniguchi, 2000; Turan et al. 2008). Since different regions of mtDNA evolve at different rates, specific mtDNA regions have been targeted for inter and intra specific variation (Hauser et al. 2001; Mohindra et al. 2007; Turan et al. 2015b).

In the present study, the validity, distribution and phylogenetic relationships of three turbot species, *Scophthalmus maeoticus*, *S. maximus*, *S. rhombus*, belong to Scophthalmidae family in the Black and Marmara Seas comprising Turkish, Bulgarian and Russian coasts were investigated with mitochondrial Cytochrome c oxidase subunit III (COIII) gene sequence data.

## Materials and Methods

### Sampling

*S. maximus* specimens were collected from three locations (at fishing ports) throughout the species' distributional range, comprising Trabzon and Düzce and Marmara Sea from Turkish Coasts, Varna Coast from Bulgaria and Sevastopol Coast from Russia. *S. maeoticus* specimens were collected from Trabzon, Düzce and Marmara Sea from Turkish Coasts. *S. rhombus* specimens were only found and collected from only the Marmara Sea (Fig. 1). All the specimens were sampled in winter season of 2013-2014. All samples were put in plastic bags individually and frozen at -20 °C until they were transported to the laboratory.




Figure 1. Sampling locations of turbot species analysed.

### *Morphology*

Meristic characters commonly used to describe Scopthalmidae species such as dorsal fin, caudal fin, anal fin, ventral fin, pectoral fin, back pectoral fin, gill rakers number, vertebra numbers were examined. Moreover, morphological characters that differentiate the species were also used as illustrated in Figure 2.




Figure 2. Morphometric distances between the numbers were taken as morphometric characters. Fish drawings is from Froese & Pauly (2018)

Most of the variability in morphological characters is due to size differences (Junquera & Perez-Gandaras, 1993, Turan, 1999). Therefore, shape analysis should be free from the effect of size to avoid misinterpretation of the results (Strauss, 1985). Therefore allometric size correction method (Elliott et al., 1995) was used to remove the effect of size variation. Principal component analysis (PCA) were used for detecting morphological characters which from differences among populations. Discriminant function analysis (DFA) were used for distinguishing morphological differences between populations (Turan, 1999). Univariate analysis of variance (ANOVA) was used to compare the variation between size-adjusted characters. SPSSv23 and SYSTATv13 were used for all statistical analysis.

### *Genetic analysis*

All tissue samples were stored at -20 °C and 95 % ethanol till the analysis. Total DNA from all fish was extracted from the muscle using the standard phenol: chloroform: isoamyl alcohol procedure

(Sambrook et al., 1989). PCR amplification of the mitochondrial COIII gene was carried out using the universal primers (Valles-Jiménez, 2005):

COIII-F: 5'- AGC CCA TGA CCT TTA ACA GG -3'

COIII-R: 5'- GAC TAC ATC AAC AAA ATG TCA GTA TCA -3'

Polymerase chain reactions were conducted by using a reaction volume of 50 µl containing 5 units of *Taq* polymerase (Thermo scientific), 2 mM of each primer, 10 mM dNTPs (Thermo scientific), 25 mM MgCl<sub>2</sub> (Thermo scientific), 10 mM Tris-HCl pH 8.8, 50 mM KCl and 1 µl template DNA (~10–25 ng). The amplification was performed with pre-denaturation at 95°C for 1 min followed by 5 cycles of denaturation at 94°C for 30 s, 50°C for 30 s, and 72°C for 45 s for 30 cycles and followed by a final extension for 7 min at 72 °C. After PCR amplification, 4 µl sample of each PCR product was controlled in 1.5% agarose gels. Quantitation of the PCR product was completed using spectrophotometer. The DNA sequencing was attempted to determine the order of the nucleotides of a gene. The chain termination method by Sanger et al. (1977) was applied with Bigdye Cycle Sequencing Kit V3.1 and ABI 3130 XL genetic analyzer.

#### *Sequence alignment and genetic analysis*

The initial alignments of partial COIII sequences were performed with Clustal W program (Thompson et al. 1994) and final alignment was completed manually with BioEdit (Hall, 1999). MtDNA sequence data were analyzed to assess levels of genetic variation and to determine nucleotide composition for each taxon using MEGA 6 (Tamura et al. 2011). Modeltest (Posada & Crandall, 1998) was used to determine the best-fit model of DNA evolution. Molecular phylogenetic analyses and molecular phylogenetic trees were conducted using MEGA version 6 (Tamura et al. 2011). A distance-based method as neighbor joining (NJ) (Saitou & Nei, 1987) and a cladistic phylogenetic tree as maximum parsimony (MP) criterion were used. The robustness of the internal branches of trees was assessed by bootstrapping (Felsenstein, 1985) with 1000 replicates. One species from the Solidae family was included in the molecular phylogenetic trees and rooted as an outgroup species *Solea senegalensis* from published sequences in GenBank under accession number, AB270760.

## **Results**

### *Genetic results*

There were 8 variable and 557 conservative nucleotides of which 6 were parsimony informative over 565 bp sequences. The average nucleotide composition was 23.01% for A, 28.80% for T, 19.77% for G and 28.41% for C. The Jukes-Cantor (Jukes-Cantor, 1969) model was determined to be the best model for our dataset. The mean genetic diversity between and within species was calculated as 0.004109 and 0.000446, respectively. The sequence analysis of COIII revealed 6 different haplotypes which were not shared between any species (Table 1). Average haplotype diversity between populations was found to be 0.7655. The minimum spanning tree showing the phylogenetic relationships among the COIII haplotypes was given in Figure 3. Ancestral haplotype (Hap\_6) was only found in *S. rhombus*. The distribution of the haplotypes in the tree generally reflects the species-specific distribution of the samples.




Figure 3. Minimum spanning tree that shows the relationships among the haplotypes.

The matrix of pairwise genetic distances within species and between species are presented in Table 2. Intra-specific genetic diversity was not observed within *S. rhombus* specimens while highest (0.000709) was found within *S. maeoticus* specimens. The lowest genetic distance (0.002482) was observed between *S. maximus* and *S. maeoticus* while the highest (0.009620) was observed between *S. rhombus* and *S. maximus*.

Table 1. The number of haplotype and its distribution among the species.

| Haplotypes | <i>S. maeoticus</i> | <i>S. maximus</i> | <i>S. rhombus</i> | Total |
|------------|---------------------|-------------------|-------------------|-------|
| Hap 1      | 8                   |                   |                   | 8     |
| Hap 2      | 1                   |                   |                   | 1     |
| Hap 3      | 1                   |                   |                   | 1     |
| Hap 4      |                     | 8                 |                   | 8     |
| Hap 5      |                     | 2                 |                   | 2     |
| Hap 6      |                     |                   | 10                | 10    |
| Total      | 10                  | 10                | 10                | 30    |

Table 2. The matrix of intra-specific genetic divergence given in bold (transversal diagonal) and pairwise genetic distances between species with statistical significance. \*\*\*, P<0.001.

|                     | <i>S.maeoticus</i> | <i>S.maximus</i> | <i>S.rhombus</i> |
|---------------------|--------------------|------------------|------------------|
| <i>S. maeoticus</i> | <b>0.000709</b>    |                  |                  |
| <i>S. maximus</i>   | 0.002482***        | <b>0.000630</b>  |                  |
| <i>S. rhombus</i>   | 0.007471***        | 0.009620***      | <b>0.000000</b>  |

Neighbor Joining and Maximum Parsimony phylogenetic approaches resulted in the similar tree topologies. In the both Neighbor Joining and Maximum Parsimony phylogenetic trees (Figure 4, 5), two phylogenetic nodes were detected; in the first node, *S. rhombus* was grouped separately. In the second node two branches were detected; *S. maximus* and *S. maeoticus* were grouped together.




Figure 4. Neighbor joining phylogenetic tree based on COIII sequences. The tree was rooted using the outgroup species *S. senegalensis*. Numbers on nodes indicate the bootstrap values. Fish drawings: *S. maximus*, *S. rhombus* and *S. senegalensis* is from Froese & Pauly (2018) and *S. maeoticus* is from Zaharia (2002).




Figure 5. Maximum parsimony tree phylogenetic tree based on COIII sequences. The tree was rooted using the outgroup species *S. senegalensis*. Numbers on nodes indicate the bootstrap values. Fish drawings: *S. maximus*, *S. rhombus* and *S. senegalensis* is from Froese & Pauly (2018) and *S. maeoticus* is from Zaharia (2002).

*Morphological results*

The range of the meristic characters used to differentiate turbot species is given in Table 3. Dorsal fin ray number varied between 60 to 69 for *S. maximus*, 59 to 76 for *S. maeoticus* and 72 to 81 for *S. rhombus*. Anal fin ray number varied between 41 to 52 for *S. maximus*, 42 to 51 for *S. maeoticus* and 53 to 61 *S. rhombus*. Pectoral fin ray number varied between 9 to 14 for all species. Ventral fin ray number varied between 5 to 7 for all species. The gill raker number and vertebra number of all species were found different ranges from each other (Table 3).

Table 3. The range of the meristic characters observed in turbot species

| Species             | N   | DFR   | PFR   | AFR   | VFR | BPFR  | CFR   | GRN   | VN    |
|---------------------|-----|-------|-------|-------|-----|-------|-------|-------|-------|
| <i>S. maximus</i>   | 158 | 60-69 | 10-14 | 41-52 | 6-7 | 10-14 | 15-22 | 13-17 | 30-32 |
| <i>S. maeoticus</i> | 162 | 59-76 | 10-13 | 42-51 | 6   | 10-13 | 15-19 | 14-17 | 30-31 |
| <i>S. rhombus</i>   | 25  | 72-81 | 10-12 | 53-61 | 6   | 10-12 | 12-18 | 15-18 | 36-38 |

DFR: Dorsal fin ray, PFR: Pectoral fin ray, AFR: Anal fin ray, VFR: Ventral fin ray, BPFR: Back pectoral fin ray, CFR: Caudal fin ray, GRN: Gill raker number VN: Vertebra number, N: number of individuals analyzed.

In PCA, 27 principal components (PCs), which contain the percentage of total variance of all variables, were produced, and 52.10 and 18.79 % of the total variation was presented in the first and second PCs. Morphometric characters generally highly contributed in differentiation, and meristic characters which are anal fin ray, back pectoral fin rays, vertebra numbers, dorsal fin rays were more efficient to differentiate turbot species (Table 4). In discriminant function analysis, the overall random assignment of individuals into their original group was 100%, showing a clear differentiation of species from each other.

Table 4. Contribution of each morphological character on the main principal components that played a role in discriminating each species.

| Variable | Component |        |        |        |
|----------|-----------|--------|--------|--------|
|          | 1         | 2      | 3      | 4      |
| 1_14     | 0.986     | -0.152 | 0.015  | -0.019 |
| 1_15     | 0.985     | -0.154 | 0.015  | -0.005 |
| 3_8      | 0.985     | -0.144 | -0.020 | -0.015 |
| 7_9      | 0.985     | -0.161 | 0.025  | -0.009 |
| 8_14     | 0.984     | -0.148 | -0.014 | -0.013 |
| 1_13     | 0.984     | -0.155 | 0.017  | -0.003 |
| 2_4      | 0.984     | -0.164 | 0.027  | -0.009 |
| 2_14     | 0.983     | -0.157 | 0.017  | -0.018 |
| 3_14     | 0.983     | -0.150 | -0.006 | -0.007 |
| 1_12     | 0.981     | -0.158 | -0.010 | 0.003  |
| 5_6      | 0.980     | -0.156 | -0.019 | -0.013 |
| 10_11    | 0.980     | -0.142 | -0.008 | -0.013 |
| Linea L. | 0.970     | -0.151 | 0.037  | -0.011 |
| 1_2      | 0.970     | -0.132 | -0.010 | -0.026 |
| BVFR     | 0.412     | 0.740  | -0.417 | 0.006  |
| D_O      | 0.277     | 0.726  | 0.495  | 0.057  |
| VN       | -0.383    | -0.717 | 0.404  | -0.062 |
| PFL      | 0.209     | 0.712  | 0.255  | -0.045 |
| DFR      | -0.360    | -0.699 | 0.354  | -0.073 |
| ID       | 0.227     | 0.683  | 0.378  | 0.087  |
| HL       | 0.404     | 0.670  | 0.495  | 0.018  |
| AFR      | -0.457    | -0.630 | 0.462  | -0.080 |
| PFR      | 0.189     | 0.486  | -0.364 | -0.006 |
| ED       | 0.203     | 0.469  | 0.219  | -0.276 |
| M_E      | 0.114     | 0.566  | 0.623  | -0.135 |

|     |       |        |       |       |
|-----|-------|--------|-------|-------|
| GRN | 0.123 | -0.486 | 0.536 | 0.096 |
| VFR | 0.080 | -0.095 | 0.104 | 0.791 |
| CFR | 0.048 | 0.132  | 0.071 | 0.577 |

In phylogenetic tree showed morphological relationship among turbot species, *S. maximus* and *S. maeoticus* were branched together in the same nod. On the other hand, *S. rhombus* was found to be separated from other two turbot species (Figure 6).




Figure 6. Systematic relationship of *Scophthalmus maximus*, *S. maeoticus* and *S. rhombus* based on morphometric and meristic characters.

## Discussion

The phylogenetic relationships of three commercially important turbot species belongs to Scophthalmidae family (*Scophthalmus maximus*, *S. maeoticus* and *S. rhombus*) from Turkish and Bulgarian coasts were investigated based on mtDNA COIII gene sequencing data in the present study. The mean intra-specific genetic diversity was found to be 0.000446 and the mean inter-specific genetic diversity was found to be 0.004109. The mean genetic divergence was low (0.00456). Feng et al. (2005) analyzed phylogenetic relationships of 10 flatfish species (*Psetta maxima*, *Platichthys flesus*, *Solea solea*, *Solea senegalensis*, *Microchirus variegatus*, *Monochirus hispidus*, *Synaptura kleini*, *Paralichthys olivaceus*, *Pseudorhombus cinnamomeus* and *Kareius bicoloratus*) belongs to the Pleuronectiformes order on the basis of 16S rRNA sequencing data and found genetic divergence values between these species ranged from 0.0141 to 0.2639. Karan & Turan (2019) reported significant genetic divergence ( $F_{ST}$ , 0.001593) based on COIII region of mitochondrial DNA between populations of *S. maeoticus* collected from the Black Sea and Marmara Sea. Feng et al. (2005) also reported that the turbot *Psetta maxima* and the thickback sole *Microchirus variegatus* has the genetically closer relationship (0.0395). Additionally, Feng et al. (2005) found genetic divergence of *P. maxima* from the other flatfish species, The European flounder *Platichthys flesus*, the common sole *Solea solea*, the whiskered sole *Monochirus hispidus*, the Klein's sole *Synaptura kleinii* and the Senegalese sole *Solea senegalensis*, to be 0.1492, 0.1798, 0.1969, 0.1734 and 0.1785, respectively. When we compare the mean genetic divergence finding with other flatfish species reported by Feng et al. (2005), genetic divergence is lower at turbot in the present study. It was thought that these turbot species may be distributed in close locations to each other in Turkish marine waters and may happen limited gene flow between them. On the other hand, similar biological and ecological features of the species together with their low mean genetic divergence may support the production of natural  $F_1$  hybrids from inter-specific mating events in

the areas where they are sympatric. However, owing to the maternal inheritance of mtDNA in vertebrates, mitochondrial markers can reveal only the maternal genome sequence. These markers cannot elucidate inter-specific hybridization occurrences that may happen between species with covering reproductive, biological, and ecological features (Turan, 2008).

The genetic differentiation of *S. rhombus* was found to be relatively higher than *S. maximus* and *S. maeoticus* that was also found statistically significant ( $P < 0.001$ ). Moreover, our results indicated that the relationship between *S. maeoticus* and *S. maximus* was closer than that both between *S. rhombus* and *S. maeoticus*, and *S. rhombus* and *S. maximus*, and confirmed the presence of three turbot species in Turkish marine waters. Pardo et al. (2005), investigated phylogenetic relationships of a total of 30 species belongs to 7 flatfish family including Scophthalmidae, Pleuronectidae, Paralichthyidae, Cynoglossidae, Soleidae, Bothidae and Achiridae from the North East Atlantic, Southwest Atlantic and Indian Ocean inferred from 16S rRNA gene and found the close relationship between *S. maximus* and *S. rhombus*. Azevedo et al. (2008) studied phylogenetic analysis of 19 species belongs to the families of the Pleuronectiformes order (Achiridae, Bothidae, Cynoglossidae, Paralichthyidae, Pleuronectidae, Scophthalmidae and Soleidae) from the South America coasts based on 12S and 16S rRNA combined sequence data and found the close relationship between *S. rhombus* and *S. maximus*, as well. As a result, our findings about the relationship between *S. rhombus* and *S. maximus* were not supported by above mentioned studies. This case may be due to the fact that above mentioned studies were conducted by different gene, such as 16S rRNA, in different seas far away from Turkish marine waters and disadvantages of 16S rRNA in discriminating closely related species (Seyhan & Turan 2016).

There was congruent between genetic and morphological analyses that *S. rhombus* was genetically and morphologically differentiated from *S. maximus* and *S. maeoticus*. Phylogenetic studies of molecular genetic variation are increasingly being used to test evolutionary hypotheses about morphological, behavioral, and life-history evolution. In particular, the reconstruction of the evolutionary history of ecological interactions is most robust when phylogenies are based on characters independent of phenotypic characters involved in, or affected by, the history of these interaction (Brooks & McLennan, 2012; Hansen, 2014). Although suitable morphological characters may be found that resolve relationships and provide an independent template for testing evolutionary hypotheses about other phenotypic characters, mitochondrial genes provide a wealth of variation almost certainly independent of changes in ecologically relevant characters. This variation may be particularly useful in generating phylogenies in groups of taxa in which most variation in morphological characters is associated with apparent adaptation (such as in groups that have undergone recent adaptive radiation), or where excessive homoplasy in such morphological characters is apparent (Brown, 1994).

In conclusion, as a result of phylogenetic analysis for turbot species, the presence of three different turbot species is determined as *S. maximus*, *S. maeoticus* and *S. rhombus* in the Black Sea and the Marmara Sea. *S. rhombus* is genetically found as the most different species. In this study, it has been proven that *S. maeoticus*, which has not been reported from the Marmara Sea and Black Sea coasts of Turkey so far, is a distinct species from *S. maximus*. Consequently, the Scophthalmidae family has been began to be represented by six species with the inclusion of *S. maeoticus* in fish fauna of Turkey through this comprehensive study. The present study also suggest that COIII could be adopted as rational approach for resolving unambiguous identification of scophthalmid species in Turkish marine waters with applications in its management and

conservation. Moreover, further analyses on a larger data set and additional molecular markers, such as nuclear genes, could improve the findings and address for more explanation.

### Acknowledgements

Thanks to the Scientific and Technological Research Council of Turkey (TUBITAK-112O920) for financial support.

### References

- Avise, J. C. (Ed.) (1994). *Molecular Markers, Natural History and Evolution*. Chapman and Hall, New York, NY. pp. 511.
- Azevedo, M. F. C., Oliveira, C., Pardo, B. G., Martinez, P., & Foresti, F. (2008). Phylogenetic analysis of the order Pleuronectiformes (Teleostei) based on sequences of 12S and 16S mitochondrial genes. *Genetics and Molecular Biology*, 31, 284-292.
- Bailly, N. & Chanet, B. (2010). Scophthalmus Rafinesque, 1810: the valid generic name for the turbot, *S. maximus* (Linnaeus, 1758) [Pleuronectiformes: Scophthalmidae]. *Cybium*, 34(3), 257-261.
- Blanquer, A., Alayse, J. P., Berrada-Rkhami, O., & Berrebi, P. (1992). Allozyme variation in turbot (*Psetta maxima*) and brill (*Scophthalmus rhombus*) (Osteichthyes, Pleuronectiformes, Scophthalmidae) throughout their range in Europe. *Journal of Fish Biology*, 41(5), 725-736.
- Brooks, D. R., & McLennan, D. A. (2012). *The Nature of Diversity: An Evolutionary Voyage of Discovery*. University of Chicago Press.
- Brown, J. M., Pellmyr, O., Thompson, J. N., & Harrison, R. G. (1994). Phylogeny of Greya (Lepidoptera: Prodoxidae), based on nucleotide sequence variation in mitochondrial cytochrome oxidase I and II: congruence with morphological data. *Molecular Biology and Evolution*, 11(1), 128-141.
- Chanet, B. (2003). Interrelationships of scophthalmid fishes (Pleuronectiformes: Scophthalmidae). *Cybium*, 27(4), 275-286.
- Durand, P., Pointier, J. P., Escoubeyrou, K., Arenas, J., Yong, M., Amarista, M., Bargues, M. D., Mas-Coma, S., & Renaud, F. (2002). Occurrence of a sibling species complex within neotropical lymnaeids, snail intermediate hosts of fascioliasis. *Acta Tropica*, 83, 233-240.
- Elliott, N. C., Farrell, J. A., Gutierrez, A. P., van Lenteren, J. C., Walton, M. P., & Wratten, S. (1995). *Integrated pest management*. Springer Science & Business Media.
- Eschmeyer, W. N. (2011). Catalog of Fishes: Genera, Species, References. (<http://research.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>). Accessed on 07 March 2016.
- Evseenko, S. A. (2003). An annotated catalogue of pleuronectiform fishes (order Pleuronectiformes) of the seas of Russia and adjacent countries. *Journal of Ichthyology*, 43(1), 57-74.
- Felsenstein, J. (1985). Confidence limits on phylogenies: An approach using the bootstrap. *Evolution*, 39(4), 783-791.
- Feng, Y., Jing, L., Peijun, Z., & Jianhai, X. (2005). Preliminary study on mitochondrial 16S rRNA gene sequences and phylogeny of flatfishes (Pleuronectiformes). *Chinese Journal of Oceanology and Limnology*, 23(3), 335-339.
- Froese, R., & Pauly, D. (Editors). 2017. FishBase. World Wide Web electronic publication. [www.fishbase.org](http://www.fishbase.org), version (05/2017).

- Hall, T. A. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium*, 41, 95-98.
- Hansen, T. F. (2014). *Use and misuse of comparative methods in the study of adaptation*. In: Modern phylogenetic comparative methods and their application in evolutionary biology (pp. 351-379). Springer, Berlin, Heidelberg.
- Hauser, L., Turan, C., & Carvalho, G. R. (2001). Haplotype frequency distribution and discriminatory power of two mtDNA fragments in a marine pelagic teleost (Atlantic herring, *Clupea harengus*). *Heredity*, 87, 621-630.
- Ivanov, L. & Beverton, R. J. H. (1985). The fisheries resources of the Mediterranean. pt. 2: Black Sea. FAO.
- Jónsson, G. (1992). Íslenskir fiskar. Reykjavík: Fjölvaútgáfan. 568p.
- Jukes, T. H. & Cantor, C. R. (1969). *Evolution of protein molecules*. Pp. 21-132 in H.N. Munro, ed. Mammalian protein metabolism III. Academic Press, New York.
- Junquera, S., & Perez-Gandaras, G. (1993). Population diversity in Bay of Biscay anchovy (*Engraulis encrasicolus* L. 1758) as revealed by multivariate analysis of morphometric and meristic characters. *ICES Journal of Marine Science*, 50(4), 383-391.
- Karan, S. & Turan, C. (2019). Evaluation of Molecular and Phenotypic Markers for Phylogeographic Analysis of Black Sea Turbot *Scophthalmus maeoticus*. *Acta zoologica bulgarica*, in press.
- Karapetkova, M. (1980). Morphological characteristics of the Black Sea Turbot *Scophthalmus maeoticus* (Pallas). *Khidrobiologiya*, 12, 73-78
- Kocher, T.D. & Stepien, C.A. (Eds.) (1997). *Molecular systematics of fishes*. Academic Press.
- Meyer, A. (1993). Evolution of mitochondrial DNA in fishes. In: Biochemistry and Molecular Biology of Fishes. *Elsevier Science Publishers*, 2, 1-38.
- Mohindra, V., Singh, R. K., Palanichamy, M., Ponniah, A. G., & Lal, K. K. (2007). Genetic identification of three species of the genus *Clarias* using allozyme and mitochondrial DNA markers. *Journal of Applied Ichthyology*, 23(1), 104-109.
- Muus, B., & Dahlström, P. (1978). Meeresfische der Ostsee, der Nordsee, des Atlantiks. BLV Verlagsgesellschaft, München. 244 p.
- Nelson, J. S. (1994). *Fishes of the World*, 3rd edn. Wiley, New York.
- Pardo, B. G., Machordom, A., Foresti, F., Porto-Foresti, F., Azevedo, M. F., Bañon, R., Sánchez, L., & Martínez, P. (2005). Phylogenetic analysis of flatfish (Order Pleuronectiformes) based on mitochondrial 16S rRNA sequences. *Scientia Marina*, 531-543.
- Prado Do, F. D., Vera, M., Hermida, M., Bouza, C., Pardo, B. G., Vilas, R., Blanco, A., Fernández, C., Maroso, F., E. Maes, G., Turan, C., A. M. Volckaert, F., B. Taggart, J., Carr, A., Ogden, R., Nielsen, E., The Aquatrace Consortium, Martínez, P. (2018). Parallel evolution and adaptation to environmental factors in a marine flatfish: Implications for fisheries and aquaculture management of the turbot (*Scophthalmus maximus*). *Evolutionary Applications*, 11:1322–1341.
- Pardo, B. G., Bouza, C., Castro, J., Martínez, P., & Sánchez, L. (2001). Localization of ribosomal genes in Pleuronectiformes using Ag-, CMA3-banding and in situ hybridization. *Heredity*, 86(5), 531-536.
- Posada, D., & Crandall, K. A. (1998). Modeltest: Testing the model of DNA substitution. *Bioinformatics*, 14(9), 817-818.
- Sambrook, J., Fritsch, E. F., & Maniatis, T. (1989). *Molecular cloning: a laboratory manual* (No. Ed. 2). Cold spring harbor laboratory press.

- Sanger, F., Nicklen, S., & Coulson, A. R. (1977). DNA Sequencing with chain-terminating inhibitors. *Proceedings of the National Academy of Sciences*, 74, 5463-5467.
- Slastenenko, E. (1956). Karadeniz Havzasi Balıkları (The fishes of the Black Sea basin). E.B.K. Yayını, İstanbul. 711 pp. (in Turkish).
- Suzuki, N., Nishida, M., Yoseda, K., Ustundag, C., Sahin, T., & Amaoka, K., (2004). Phylogeographic relationships within the Mediterranean turbot inferred by mitochondrial DNA haplotype variation. *Journal of Fish Biology*, 65(2), 580- 585.
- Tabata, K., & Taniguchi, N. (2000). Differences between *Pagrus major* and *Pagrus auratus* through mainly mtDNA control region analysis. *Fisheries Science*, 66(1), 9-18.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. (2011). MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution*, 28(10), 2731-2739.
- Thompson, J. D., Higgins, D. G., Gibson, T. J. (1994). CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research*, 22(22), 4673-4680.
- Turan, C., (1999). A note on the examination of morphometric differentiation among fish populations: The truss system. *Turkish Journal of Zoology*, 23, 259-264.
- Turan, C. (Ed.) (2007). *Atlas and Systematics of Marine Bony Fishes of Turkey*. 1st edition, Nobel Publishing House, Adana, Turkey.
- Turan, C. (2008). Molecular systematic analyses of Mediterranean skates (Rajiformes). *Turkish Journal of Zoology*, 32(4), 437-442.
- Turan, C., Gündüz, İ, Gürlek, M., & Yağlıoğlu, D. (2008). Systematics of Scorpaenidae species in the Mediterranean Sea inferred from mitochondrial 16S rDNA sequence and morphological data. *Folia Biologica*, 57, 219-226.
- Turan, C., Ergüden, D., Çevik, C., Gürlek, M., & Turan, F. (2015a). Molecular systematic analysis of shad species (*Alosa* spp.) from Turkish marine waters using mtDNA genes. *Turkish Journal of Fisheries and Aquatic Sciences*, 15 (1), 149-155.
- Turan, C., Gürlek, M., Ergüden, D., Yağlıoğlu, D., Öztürk, B., Uyan, A., Reyhaniye, A. N., Özbalcılar, B., Erdoğan, Z. A., Ivanova, P., & Soldo, A. (2015b). Population Genetic Analysis of Atlantic Bonito *Sarda sarda* (Bloch, 1793) using Sequence Analysis of mtDNA D-Loop Region. *Fresenius Environmental Bulletin*, 45(3), 231-237.
- Turan, C., Yağlıoğlu, D., Ergüden, D., Gürlek, M., Uyan, A., Karan, S., & Doğdu, S. (2016). Threatened brill species in marine waters of Turkey: *Scophthalmus rhombus* (Linnaeus, 1758) (Scophthalmidae). *Natural and Engineering Sciences*, 1(1), 1-6.
- Valles-Jiménez, R. (2005). Estudios sobre la estructura genética del camarón blanco (*Litopenaeus vannamei*), del Pacífico Oriental inferidos del análisis de microsatélites y ADN mitocondrial. PhD thesis. Centro de Investigaciones Biológicas Del Noroeste, S.C. 74 pp.
- Voronina, E. P. (2010). On Morphology and Taxonomy of Scophthalmids. *Journal of Ichthyology*, 50(9), 695-703.
- Zaharia, T. (2002). Researches for elaborating the technology for reproducing and rearing of the flounder and turbot, in order to renew their natural populations. PhD thesis. University, Dunarea de Jos" Galați, 156 pp. [In Romanian].
- Zardoya, R., Economidis, P. S. & Doadrio, I. (1999). Phylogenetic relationships of Greek Cyprinidae: molecular evidence for at least two origins of the Greek cyprinid fauna. *Molecular Phylogenetics and Evolution*, 13(1), 122-131.