



COEFFICIENT ESTIMATES FOR A NEW SUBCLASS OF m -FOLD SYMMETRIC ANALYTIC BI-UNIVALENT FUNCTIONS

SERAP BULUT

ABSTRACT. Considering a new subclass of m -fold symmetric analytic bi-univalent functions, we determine estimates the coefficient bounds for the Taylor-Maclaurin coefficients $|a_{m+1}|$ and $|a_{2m+1}|$ of the functions in this class. In certain cases, our estimates improve some of those existing coefficient bounds.

1. INTRODUCTION

Let \mathcal{A} denote the class of all functions of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (1.1)$$

which are analytic in the open unit disk $\mathbb{U} = \{z : z \in \mathbb{C} \text{ and } |z| < 1\}$. We also denote by \mathcal{S} the class of all functions in the normalized analytic function class \mathcal{A} which are univalent in \mathbb{U} .

For two functions f and Θ , analytic in \mathbb{U} , we say that the function f is subordinate to Θ in \mathbb{U} , and write

$$f(z) \prec \Theta(z) \quad (z \in \mathbb{U}),$$

if there exists a Schwarz function ω , which is analytic in \mathbb{U} with

$$\omega(0) = 0 \quad \text{and} \quad |\omega(z)| < 1 \quad (z \in \mathbb{U})$$

such that

$$f(z) = \Theta(\omega(z)) \quad (z \in \mathbb{U}).$$

It is well known that every function $f \in \mathcal{S}$ has an inverse f^{-1} , which is defined by

$$f^{-1}(f(z)) = z \quad (z \in \mathbb{U})$$

Received by the editors: November 15, 2017; Accepted: September 09, 2018.

2010 *Mathematics Subject Classification.* Primary 30C45.

Key words and phrases. Analytic functions, univalent functions, bi-univalent functions; m -fold symmetric bi-univalent functions, Taylor-Maclaurin series expansion; coefficient bounds, coefficient estimates.

and

$$f(f^{-1}(w)) = w \quad \left(|w| < r_0(f); r_0(f) \geq \frac{1}{4} \right).$$

In fact, the inverse function $g = f^{-1}$ is given by

$$g(w) = f^{-1}(w) = w - a_2 w^2 + (2a_2^2 - a_3) w^3 - (5a_2^3 - 5a_2 a_3 + a_4) w^4 + \dots \quad (1.2)$$

A function $f \in \mathcal{A}$ is said to be bi-univalent in \mathbb{U} if both f and f^{-1} are univalent in \mathbb{U} , in the sense that f^{-1} has a univalent analytic continuation to \mathbb{U} . We denote by Σ the class of all bi-univalent functions in \mathbb{U} given by (1.1).

For a brief history and interesting examples of functions in the class Σ , see [15] (see also [3]). In fact, the aforementioned work of Srivastava *et al.* [15] essentially revived the investigation of various subclasses of the bi-univalent function class Σ in recent years; it was followed by such works as those by Bulut *et al.* [5], Frasin and Aouf [6], Ramachandran *et al.* [9], Srivastava *et al.* [10, 13], Xu *et al.* [18, 19] and the references cited in each of them.

Let $m \in \mathbb{N} = \{1, 2, 3, \dots\}$. A domain D is said to be m -fold symmetric if a rotation of D about the origin through an angle $2\pi/m$ carries D on itself. It follows that, a function $f(z)$ analytic in \mathbb{U} is said to be m -fold symmetric ($m \in \mathbb{N}$) if

$$f(e^{2\pi i/m} z) = e^{2\pi i/m} f(z).$$

In particular, every $f(z)$ is 1-fold symmetric and every odd $f(z)$ is 2-fold symmetric. We denote by \mathcal{S}_m the class of m -fold symmetric univalent functions in \mathbb{U} .

A simple argument shows that $f \in \mathcal{S}_m$ is characterized by having a power series of the form

$$f(z) = z + \sum_{k=1}^{\infty} a_{mk+1} z^{mk+1} \quad (z \in \mathbb{U}, m \in \mathbb{N}). \quad (1.3)$$

Srivastava *et al.* [14] defined m -fold symmetric bi-univalent functions analogues to the concept of m -fold symmetric univalent functions. For normalized form of f given by (1.3), they obtained the series expansion for f^{-1} as following:

$$\begin{aligned} g(w) &= f^{-1}(w) \\ &= w - a_{m+1} w^{m+1} + [(m+1)a_{m+1}^2 - a_{2m+1}] w^{2m+1} \\ &\quad - \left[\frac{1}{2}(m+1)(3m+2)a_{m+1}^3 - (3m+2)a_{m+1}a_{2m+1} + a_{3m+1} \right] w^{3m+1} \\ &\quad + \dots \\ &= : w + \sum_{k=1}^{\infty} A_{mk+1} w^{mk+1}. \end{aligned} \quad (1.4)$$

We denote by Σ_m the class of m -fold symmetric bi-univalent functions in \mathbb{U} given by (1.3). For $m = 1$, the formula (1.4) coincides with the formula (1.2) of the class Σ . For some examples of m -fold symmetric bi-univalent functions, see [14].

We also denote by \mathcal{P} the family of all functions p analytic in \mathbb{U} for which

$$\Re(p(z)) > 0, \quad p(z) = 1 + c_1z + c_2z^2 + \dots \quad (z \in \mathbb{U}).$$

Thus the m -fold symmetric function p in the class \mathcal{P} is of the form (see [8]),

$$p(z) = 1 + c_mz^m + c_{2m}z^{2m} + \dots \quad (z \in \mathbb{U}).$$

The coefficient problem for m -fold symmetric analytic bi-univalent functions is one of the favorite subjects of geometric function theory in these days (see [1, 4, 7, 11, 14, 16]). The object of the present paper is to introduce a new subclass of bi-univalent functions in which both f and f^{-1} are m -fold symmetric analytic functions and obtain coefficient bounds for $|a_{m+1}|$ and $|a_{2m+1}|$ for functions in this new subclass.

2. THE CLASS $\mathcal{N}_{\Sigma,m}(\tau, \lambda, \delta, \varphi)$

Throughout this paper, we assume that φ is an analytic function with positive real part in the unit disk \mathbb{U} , satisfying $\varphi(0) = 1$, $\varphi'(0) > 0$, and $\varphi(\mathbb{U})$ is symmetric with respect to the real axis. Such a function has a series expansion of the form

$$\varphi(z) = 1 + B_1z + B_2z^2 + \dots \quad (B_1 > 0, B_n \in \mathbb{R}, n = 2, 3, \dots). \quad (2.1)$$

With this assumption on φ , we now introduce the following new class of m -fold symmetric analytic bi-univalent functions.

Definition 1. For $\tau \in \mathbb{C} \setminus \{0\}$, $\lambda \geq 1$ and $0 \leq \delta \leq 1$, a function $f \in \Sigma_m$ given by (1.3) is said to be in the class $\mathcal{N}_{\Sigma,m}(\tau, \lambda, \delta, \varphi)$ if the following conditions are satisfied:

$$1 + \frac{1}{\tau} \left\{ (1 - \lambda) \frac{f(z)}{z} + \lambda f'(z) + \delta z f''(z) - 1 \right\} \prec \varphi(z) \quad (2.2)$$

and

$$1 + \frac{1}{\tau} \left\{ (1 - \lambda) \frac{g(w)}{w} + \lambda g'(w) + \delta w g''(w) - 1 \right\} \prec \varphi(w) \quad (2.3)$$

where $m \in \mathbb{N}$; $z, w \in \mathbb{U}$ and $g = f^{-1}$ is defined by (1.4).

Remark 1. (i) If we set $\tau = 1$ and $\delta = 0$, then we have the class

$$\mathcal{N}_{\Sigma,m}(1, \lambda, 0, \varphi) = \mathcal{B}_{\Sigma,m}(\lambda, \varphi)$$

introduced and studied by Tang *et al.* [17].

(ii) There are many choices of the function $\varphi(z)$ which would provide interesting subclasses of the analytic function class \mathcal{A} . For example, if we let

$$\varphi(z) = \left(\frac{1+z}{1-z} \right)^\alpha \quad (0 < \alpha \leq 1, z \in \mathbb{U}),$$

or

$$\varphi(z) = \frac{1 + (1 - 2\beta)z}{1 - z} \quad (0 \leq \beta < 1, z \in \mathbb{U}),$$

it is easy to verify that these functions are of the form (2.1). If $f \in \mathcal{N}_{\Sigma_m}(\tau, \lambda, \delta, \varphi)$, then $f \in \Sigma_m$ and

$$\left| \arg \left(1 + \frac{1}{\tau} \left\{ (1 - \lambda) \frac{f(z)}{z} + \lambda f'(z) + \delta z f''(z) - 1 \right\} \right) \right| < \frac{\alpha\pi}{2}$$

and

$$\left| \arg \left(1 + \frac{1}{\tau} \left\{ (1 - \lambda) \frac{g(w)}{w} + \lambda g'(w) + \delta w g''(w) - 1 \right\} \right) \right| < \frac{\alpha\pi}{2},$$

or

$$\Re \left(1 + \frac{1}{\tau} \left\{ (1 - \lambda) \frac{f(z)}{z} + \lambda f'(z) + \delta z f''(z) - 1 \right\} \right) > \beta$$

and

$$\Re \left(1 + \frac{1}{\tau} \left\{ (1 - \lambda) \frac{g(w)}{w} + \lambda g'(w) + \delta w g''(w) - 1 \right\} \right) > \beta,$$

where the function $g = f^{-1}$ is defined by (1.4). This means that

$$f \in \mathcal{R}_{\Sigma_m}(\tau, \lambda, \delta; \alpha) \quad \text{or} \quad f \in \mathcal{R}_{\Sigma_m}(\tau, \lambda, \delta; \beta),$$

respectively. These classes are introduced and studied by Atshan and Al-Ziadi [2]. In these classes of m -fold symmetric bi-univalent functions, in particular we have

$$\mathcal{R}_{\Sigma_m}(\tau, 1, \delta; \alpha) = \mathcal{H}_{\Sigma_m}(\tau, \delta; \alpha), \quad \mathcal{R}_{\Sigma_m}(\tau, 1, \delta; \beta) = \mathcal{H}_{\Sigma_m}(\tau, \delta; \beta) \quad (\text{see [11]}),$$

$$\mathcal{R}_{\Sigma_m}(\tau, \lambda, 0; \alpha) = \mathcal{B}_{\Sigma_m}(\tau, \lambda; \alpha), \quad \mathcal{R}_{\Sigma_m}(\tau, \lambda, 0; \beta) = \mathcal{B}_{\Sigma_m}^*(\tau, \lambda; \beta) \quad (\text{see [12]}),$$

and

$$\mathcal{R}_{\Sigma_m}(1, \lambda, 0; \alpha) = \mathcal{A}_{\Sigma_m}^{\alpha, \lambda}, \quad \mathcal{R}_{\Sigma_m}(1, \lambda, 0; \beta) = \mathcal{A}_{\Sigma_m}^{\lambda}(\beta) \quad (\text{see [16]}).$$

Here we propose to investigate the m -fold symmetric bi-univalent function class $\mathcal{N}_{\Sigma_m}(\tau, \lambda, \delta, \varphi)$ introduced in Definition 1 and derive coefficient estimates on the first two Taylor-Maclaurin coefficients $|a_{m+1}|$ and $|a_{2m+1}|$ for a function $f \in \mathcal{N}_{\Sigma_m}(\tau, \lambda, \delta, \varphi)$ given by (1.3). Our results for the m -fold symmetric bi-univalent function class $\mathcal{N}_{\Sigma_m}(\tau, \lambda, \delta, \varphi)$ would generalize and improve the related works of Tang *et al.* [17], Srivastava *et al.* [11, 12] and Sümer Eker [16].

3. A SET OF GENERAL COEFFICIENT ESTIMATES

In this section, we state and prove our general results involving the m -fold symmetric bi-univalent function class $\mathcal{N}_{\Sigma_m}(\tau, \lambda, \delta, \varphi)$ given by Definition 1.

Theorem 1. *Let the function f given by (1.3) be in the class $\mathcal{N}_{\Sigma_m}(\tau, \lambda, \delta, \varphi)$, $\tau \in \mathbb{C} \setminus \{0\}$, $\lambda \geq 1$, $0 \leq \delta \leq 1$ and $m \in \mathbb{N}$. Then*

$$|a_{m+1}| \leq \frac{|\tau| B_1 \sqrt{2B_1}}{\sqrt{|\tau(m+1)\psi_2 B_1^2 - 2\psi_1^2 B_2| + 2\psi_1^2 B_1}} \tag{3.1}$$

and

$$|a_{2m+1}| \leq \begin{cases} \Phi & , \quad B_1 \geq \frac{2\psi_1^2}{|\tau|(m+1)\psi_2} \\ \frac{|\tau|^{2(m+1)}B_1^2}{2\psi_1^2} & , \quad B_1 < \frac{2\psi_1^2}{|\tau|(m+1)\psi_2} \end{cases}, \quad (3.2)$$

where

$$\Phi := \left(\frac{m+1}{2} - \frac{\psi_1^2}{|\tau|\psi_2 B_1} \right) \frac{2|\tau|^2 B_1^3}{|\tau(m+1)\psi_2 B_1^2 - 2\psi_1^2 B_2| + 2\psi_1^2 B_1} + \frac{|\tau| B_1}{\psi_2}, \quad (3.3)$$

$$\psi_j := 1 + jm\lambda + jm(jm+1)\delta \quad (j = 1, 2). \quad (3.4)$$

Proof. Let $f \in \mathcal{N}_{\Sigma, m}(\tau, \lambda, \delta, \varphi)$ and $g = f^{-1}$ be defined by (1.4). Then there exist two Schwarz functions $u, v : \mathbb{U} \rightarrow \mathbb{U}$, with $u(0) = v(0) = 0$, such that

$$1 + \frac{1}{\tau} \left\{ (1-\lambda) \frac{f(z)}{z} + \lambda f'(z) + \delta z f''(z) - 1 \right\} = \varphi(u(z)) \quad (3.5)$$

and

$$1 + \frac{1}{\tau} \left\{ (1-\lambda) \frac{g(w)}{w} + \lambda g'(w) + \delta w g''(w) - 1 \right\} = \varphi(v(w)). \quad (3.6)$$

We suppose that

$$u(z) = p_m z^m + p_{2m} z^{2m} + \dots \quad (z \in \mathbb{U}) \quad (3.7)$$

and

$$v(w) = q_m w^m + q_{2m} w^{2m} + \dots \quad (w \in \mathbb{U}). \quad (3.8)$$

We note that

$$|p_m| \leq 1, \quad |p_{2m}| \leq 1 - |p_m|^2, \quad |q_m| \leq 1, \quad |q_{2m}| \leq 1 - |q_m|^2. \quad (3.9)$$

Using (3.7) and (3.8) together with (2.1), it is evident that

$$\varphi(u(z)) = 1 + B_1 p_m z^m + (B_1 p_{2m} + B_2 p_m^2) z^{2m} + \dots \quad (z \in \mathbb{U}) \quad (3.10)$$

and

$$\varphi(v(w)) = 1 + B_1 q_m w^m + (B_1 q_{2m} + B_2 q_m^2) w^{2m} + \dots \quad (w \in \mathbb{U}), \quad (3.11)$$

respectively. Since

$$\begin{aligned} 1 + \frac{1}{\tau} \left\{ (1-\lambda) \frac{f(z)}{z} + \lambda f'(z) + \delta z f''(z) - 1 \right\} \\ = 1 + \frac{1}{\tau} \{1 + m\lambda + m(m+1)\delta\} a_{m+1} z^m \\ + \frac{1}{\tau} \{1 + 2m\lambda + 2m(2m+1)\delta\} a_{2m+1} z^{2m} + \dots \end{aligned}$$

and

$$1 + \frac{1}{\tau} \left\{ (1-\lambda) \frac{g(w)}{w} + \lambda g'(w) + \delta w g''(w) - 1 \right\}$$

$$= 1 + \frac{1}{\tau} \{1 + m\lambda + m(m + 1)\delta\} A_{m+1} w^m + \frac{1}{\tau} \{1 + 2m\lambda + 2m(2m + 1)\delta\} A_{2m+1} w^{2m} + \dots,$$

then (3.5), (3.6), (3.10) and (3.11) together with (1.4) yield

$$\frac{1}{\tau} \{1 + m\lambda + m(m + 1)\delta\} a_{m+1} = B_1 p_m, \tag{3.12}$$

$$\frac{1}{\tau} \{1 + 2m\lambda + 2m(2m + 1)\delta\} a_{2m+1} = B_1 p_{2m} + B_2 p_m^2, \tag{3.13}$$

$$-\frac{1}{\tau} \{1 + m\lambda + m(m + 1)\delta\} a_{m+1} = B_1 q_m \tag{3.14}$$

and

$$\frac{1}{\tau} \{1 + 2m\lambda + 2m(2m + 1)\delta\} \{(m + 1) a_{m+1}^2 - a_{2m+1}\} = B_1 q_{2m} + B_2 q_m^2. \tag{3.15}$$

Now, considering (3.12) and (3.14), we get

$$p_m = -q_m \tag{3.16}$$

and

$$2\frac{1}{\tau^2} \{1 + m\lambda + m(m + 1)\delta\}^2 a_{m+1}^2 = B_1^2 (p_m^2 + q_m^2). \tag{3.17}$$

Now from (3.13), (3.15) and (3.17), we obtain

$$a_{m+1}^2 = \frac{\tau^2 B_1^3 (p_{2m} + q_{2m})}{\tau(m + 1) \{1 + 2m\lambda + 2m(2m + 1)\delta\} B_1^2 - 2 \{1 + m\lambda + m(m + 1)\delta\}^2 B_2}. \tag{3.18}$$

Therefore, by (3.9), (3.16) from (3.18), we get

$$|a_{m+1}|^2 \leq \frac{2|\tau|^2 B_1^3 (1 - |p_m|^2)}{|\tau(m + 1) \{1 + 2m\lambda + 2m(2m + 1)\delta\} B_1^2 - 2 \{1 + m\lambda + m(m + 1)\delta\}^2 B_2|}.$$

The equality (3.12) and the above inequality give

$$|a_{m+1}|^2 \leq \frac{2|\tau|^2 B_1^3}{|\tau(m + 1) \psi_2 B_1^2 - 2\psi_1^2 B_2| + 2\psi_1^2 B_1}, \tag{3.19}$$

where ψ_j ($j = 1, 2$) is defined by (3.4), which is the desired estimate on the coefficient $|a_{m+1}|$ as asserted in (3.1).

Next, in order to find the bound on the coefficient $|a_{2m+1}|$, we subtract (3.15) from (3.13). Observing (3.16) we get

$$a_{2m+1} = \frac{m + 1}{2} a_{m+1}^2 + \frac{\tau B_1}{2 \{1 + 2m\lambda + 2m(2m + 1)\delta\}} (p_{2m} - q_{2m}). \tag{3.20}$$

We obtain from the (3.9), (3.12) and the above equality

$$\begin{aligned}
 |a_{2m+1}| &\leq \frac{m+1}{2} |a_{m+1}|^2 + \frac{|\tau| B_1}{1+2m\lambda+2m(2m+1)\delta} (1-|p_m|^2) \\
 &\leq \left(\frac{m+1}{2} - \frac{\{1+m\lambda+m(m+1)\delta\}^2}{|\tau|\{1+2m\lambda+2m(2m+1)\delta\}B_1} \right) |a_{m+1}|^2 \\
 &\quad + \frac{|\tau| B_1}{1+2m\lambda+2m(2m+1)\delta}. \tag{3.21}
 \end{aligned}$$

Upon substituting the value of a_{m+1}^2 from (3.12) and (3.19) into (3.21), it follows that

$$|a_{2m+1}| \leq \frac{|\tau|^2 (m+1) B_1^2}{2\psi_1^2} \tag{3.22}$$

and

$$|a_{2m+1}| \leq \left(\frac{m+1}{2} - \frac{\psi_1^2}{|\tau|\psi_2 B_1} \right) \frac{2|\tau|^2 B_1^3}{|\tau(m+1)\psi_2 B_1^2 - 2\psi_1^2 B_2| + 2\psi_1^2 B_1} + \frac{|\tau| B_1}{\psi_2}, \tag{3.23}$$

where ψ_j ($j = 1, 2$) is defined by (3.4), respectively. Therefore considering (3.22) and (3.23), we get the desired estimate on the coefficient $|a_{2m+1}|$ as asserted in (3.2). This completes the proof of the Theorem 1. \square

Setting $\tau = 1$ and $\delta = 0$ in Theorem 1, we have the following corollary.

Corollary 1. *Let the function f given by (1.3) be in the class $\mathcal{B}_{\Sigma,m}(\lambda, \varphi)$, $\lambda \geq 1$ and $m \in \mathbb{N}$. Then*

$$|a_{m+1}| \leq \frac{B_1 \sqrt{2B_1}}{\sqrt{|(m+1)(1+2m\lambda)B_1^2 - 2(1+m\lambda)^2 B_2| + 2(1+m\lambda)^2 B_1}}$$

and

$$|a_{2m+1}| \leq \begin{cases} \Phi & B_1 \geq \frac{2(1+m\lambda)^2}{(m+1)(1+2m\lambda)} \\ \frac{(m+1)B_1^2}{2(1+m\lambda)^2} & B_1 < \frac{2(1+m\lambda)^2}{(m+1)(1+2m\lambda)} \end{cases},$$

where

$$\Phi = \left(\frac{m+1}{2} - \frac{(1+m\lambda)^2}{(1+2m\lambda)B_1} \right) \frac{2B_1^3}{|(m+1)(1+2m\lambda)B_1^2 - 2(1+m\lambda)^2 B_2| + 2(1+m\lambda)^2 B_1} + \frac{B_1}{1+2m\lambda}.$$

Remark 2. Note that Corollary 1 is an improvement of the estimates obtained by Tang *et al.* [17, Theorem 7].

Taking the function

$$\varphi(z) = \left(\frac{1+z}{1-z} \right)^\alpha \quad (0 < \alpha \leq 1, z \in \mathbb{U})$$

in Theorem 1, we have the following corollary.

Corollary 2. Let the function f given by (1.3) be in the class $\mathcal{R}_{\Sigma_m}(\tau, \lambda, \delta; \alpha)$, $\tau \in \mathbb{C} \setminus \{0\}$, $\lambda \geq 1$, $0 \leq \delta \leq 1$, $0 < \alpha \leq 1$ and $m \in \mathbb{N}$. Then

$$|a_{m+1}| \leq \frac{2|\tau|\alpha}{\sqrt{|\tau(m+1)\psi_2 - \psi_1^2|\alpha + \psi_1^2}}$$

and

$$|a_{2m+1}| \leq \begin{cases} \Phi & , \quad \alpha \geq \frac{\psi_1^2}{|\tau(m+1)\psi_2} \\ \frac{2|\tau|^2(m+1)\alpha^2}{\psi_1^2} & , \quad \alpha < \frac{\psi_1^2}{|\tau(m+1)\psi_2} \end{cases},$$

where

$$\Phi = \left((m+1) - \frac{\psi_1^2}{|\tau|\psi_2\alpha} \right) \frac{2|\tau|^2\alpha^2}{|\tau(m+1)\psi_2 - \psi_1^2|\alpha + \psi_1^2} + \frac{2|\tau|\alpha}{\psi_2},$$

and ψ_j is defined by (3.4).

Remark 3. Note that Corollary 2 is an improvement of the estimates obtained by Atshan and Al-Ziadi [2, Theorem 2.1], Srivastava *et al.* [11, Theorem 2] for $\lambda = 1$, Srivastava *et al.* [12, Theorem 2.1] for $\delta = 0$, and Sümer Eker [16, Theorem 1] for $\tau = 1$ and $\delta = 0$, respectively.

Taking the function

$$\varphi(z) = \frac{1 + (1 - 2\beta)z}{1 - z} \quad (0 \leq \beta < 1, z \in \mathbb{U})$$

in Theorem 1, we have the following corollary.

Corollary 3. Let the function f given by (1.3) be in the class $\mathcal{R}_{\Sigma_m}(\tau, \lambda, \delta; \beta)$, $\tau \in \mathbb{C} \setminus \{0\}$, $\lambda \geq 1$, $0 \leq \delta \leq 1$, $0 \leq \beta < 1$ and $m \in \mathbb{N}$. Then

$$|a_{m+1}| \leq \frac{\sqrt{2}|\tau|(1-\beta)}{\sqrt{|\tau(m+1)\psi_2(1-\beta) - \psi_1^2| + \psi_1^2}}$$

and

$$|a_{2m+1}| \leq \begin{cases} \Phi & , \quad \beta \leq 1 - \frac{\psi_1^2}{|\tau(m+1)\psi_2} \\ \frac{2|\tau|^2(m+1)(1-\beta)^2}{\psi_1^2} & , \quad \beta > 1 - \frac{\psi_1^2}{|\tau(m+1)\psi_2} \end{cases},$$

where

$$\Phi = \left((m+1) - \frac{\psi_1^2}{|\tau|\psi_2(1-\beta)} \right) \frac{2|\tau|^2(1-\beta)^2}{|\tau(m+1)\psi_2(1-\beta) - \psi_1^2| + \psi_1^2} + \frac{2|\tau|(1-\beta)}{\psi_2},$$

and ψ_1 and ψ_2 are defined by (3.4).

Remark 3. Note that Corollary 3 is an improvement of the estimates obtained by Atshan and Al-Ziadi [2, Theorem 3.1], Srivastava *et al.* [11, Theorem 3] for $\lambda = 1$, Srivastava *et al.* [12, Theorem 3.1] for $\delta = 0$, and Sümer Eker [16, Theorem 2] for $\tau = 1$ and $\delta = 0$, respectively.

REFERENCES

- [1] Altınkaya, Ş. and Yalçın, S., Coefficient bounds for certain subclasses of m -fold symmetric bi-univalent functions, *J. Math.* (2015), Art. ID 241683, 5 pp.
- [2] Atshan, W.G. and Al-Ziadi, N.A.J., Coefficients bounds for a general subclasses of m -fold symmetric bi-univalent functions, *J. Al-Qadisiyah Comput. Sci. Math.* 9 (2) (2017), 33–39.
- [3] Brannan, D.A. and Taha, T.S., On some classes of bi-univalent functions, in *Mathematical Analysis and Its Applications* (S. M. Mazhar, A. Hamoui and N. S. Faour, Editors) (Kuwait; February 18–21, 1985), KFAS Proceedings Series, Vol. 3, Pergamon Press (Elsevier Science Limited), Oxford, 1988, pp. 53–60; see also *Studia Univ. Babeş-Bolyai Math.* 31 (2) (1986), 70–77.
- [4] Bulut, S., Coefficient estimates for general subclasses of m -fold symmetric analytic bi-univalent functions, *Turkish J. Math.* 40 (6) (2016), 1386–1397.
- [5] Bulut, S., Magesh, N. and Balaji, V.K., Faber polynomial coefficient estimates for certain subclasses of meromorphic bi-univalent functions, *C. R. Acad. Sci. Paris, Ser. I*, 353 (2015), 113–116.
- [6] Frasin, B.A. and Aouf, M.K., New subclasses of bi-univalent functions, *Appl. Math. Lett.* 24 (2011) 1569–1573.
- [7] Hamidi, S.G. and Jahangiri, J.M., Unpredictability of the coefficients of m -fold symmetric bi-starlike functions, *Internat. J. Math.* 25 (7) (2014), 1450064, 8 pp.
- [8] Pommerenke, Ch., On the coefficients of close-to-convex functions, *Michigan. Math. J.* 9 (1962), 259–269.
- [9] Ramachandran, C., Prabhu, Ambrose R. and Magesh, N., Initial coefficient estimates for certain subclasses of bi-univalent functions of Ma-Minda type, *Applied Mathematical Sciences* 9 (47) (2015), 2299–2308.
- [10] Srivastava, H.M., Bulut, S., Çağlar, M. and Yağmur, N., Coefficient estimates for a general subclass of analytic and bi-univalent functions, *Filomat* 27 (5) (2013), 831–842.
- [11] Srivastava, H.M., Gaboury, S. and Ghanim, F., Coefficient estimates for some subclasses of m -fold symmetric bi-univalent functions, *Acta Univ. Apulensis Math. Inform.* 41 (2015), 153–164.
- [12] Srivastava, H.M., Gaboury, S. and Ghanim, F., Initial coefficient estimates for some subclasses of m -fold symmetric bi-univalent functions, *Acta Math. Sci. Ser. B Engl. Ed.* 36 (3) (2016), 863–871.
- [13] Srivastava, H.M., Murugusundaramoorthy, G. and Magesh, N., On certain subclasses of bi-univalent functions associated with Hohlov operator, *Global Journal of Mathematical Analysis* 1 (2) (2013), 67–73.
- [14] Srivastava, H.M., Sivasubramanian, S. and Sivakumar, R., Initial coefficient bounds for a subclass of m -fold symmetric bi-univalent functions, *Tbilisi Math. J.* 7 (2) (2014), 1–10.
- [15] Srivastava, H.M., Mishra, A.K. and Gochhayat, P., Certain subclasses of analytic and bi-univalent functions, *Appl. Math. Lett.* 23 (2010) 1188–1192.
- [16] Sümer Eker, S., Coefficient bounds for subclasses of m -fold symmetric bi-univalent functions, *Turkish J. Math.* 40 (3) (2016), 641–646.
- [17] Tang, H., Srivastava, H.M., Sivasubramanian, S. and Gurusamy, P., The Fekete-Szegő functional problems for some subclasses of m -fold symmetric bi-univalent functions, *J. Math. Inequal.* 10 (4) (2016), 1063–1092.

- [18] Xu, Q.-H., Gui, Y.-C. and Srivastava, H.M., Coefficient estimates for a certain subclass of analytic and bi-univalent functions, *Appl. Math. Lett.* 25 (2012) 990–994.
- [19] Xu, Q.-H., Xiao, H.-G. and Srivastava, H.M., A certain general subclass of analytic and bi-univalent functions and associated coefficient estimate problems, *Appl. Math. Comput.* 218 (2012) 11461–11465.

Current address: Serap Bulut: Kocaeli University, Faculty of Aviation and Space Sciences, Arslanbey Campus, 41285 Kartepe-Kocaeli, TURKEY

E-mail address: serap.bulut@kocaeli.edu.tr

ORCID Address: <http://orcid.org/0000-0002-6506-4588>