



Improved Bounds for the Extremal Non-Trivial Laplacian Eigenvalues

Şerife BÜYÜKKÖSE^{1,♣}, Ercan ALTINIŞIK¹, Feyza YALÇIN²

¹Department of Mathematics, Faculty of Science, Gazi University, 06500 Teknikokullar-Ankara, Turkey

²Department of Mathematics, Faculty of Arts and Sciences, Harran University, 63050 Şanlıurfa, Turkey

Received:12/09/2014 Revised:04/11/2014 Accepted:05/11/2014

ABSTRACT

Let G be a simple connected graph and its Laplacian eigenvalues be $\mu_1 \geq \mu_2 \geq \dots \geq \mu_{n-1} \geq \mu_n = 0$. In this paper, we present an upper bound for the algebraic connectivity μ_{n-1} of G and a lower bound for the largest eigenvalue μ_1 of G in terms of the degree sequence d_1, d_2, \dots, d_n of G and the number $|N_i \cap N_j|$ of common vertices of i and j ($1 \leq i < j \leq n$) and hence we improve bounds of Maden and Büyükköse [14].

Keywords: Laplacian eigenvalues, upper bounds, lower bounds, eigenvalue inequalities.

2010 MSC: 05C50, 15A18.

1. INTRODUCTION

Let $G = (V, E)$ be a simple graph with the vertex set $V = \{v_1, v_2, \dots, v_n\}$ and edge set E . We use $i \sim j$ to denote that $v_i v_j$ is an edge of G and N_i to denote that the set of neighbours of v_i . For $v_i \in V$, the degree of v_i and the average of the degrees of the vertices adjacent to v_i are denoted by d_i and m_i , respectively. We assume that $d_1 \geq d_2 \geq \dots \geq d_n$ without loss of generality and we call d_1, d_2, \dots, d_n the degree sequence of G . Let $A(G)$ be the adjacency matrix of G and let $D(G)$ be the diagonal matrix of vertex degrees. The Laplacian matrix of G is $L(G) = D(G) - A(G)$. For the simplicity of notation, we write $L(G) = L$. Clearly, L is a real symmetric matrix. From this fact and Geršgorin's Theorem, it follows that its eigenvalues are nonnegative real numbers. Moreover, since the sum of rows is 0, it is obvious that 0 is the smallest eigenvalue of L with the all ones vector as an eigenvector. The Laplacian

eigenvalues of G are the eigenvalues of the Laplacian matrix L of G . Throughout this paper, the Laplacian eigenvalues of G are denoted by

$$\mu_1 \geq \mu_2 \geq \dots \geq \mu_{n-1} \geq \mu_n = 0.$$

In addition, by the extremal non-trivial Laplacian eigenvalues, we shall mean μ_{n-1} and μ_1 . It is easy to show that $\mu_{n-1}(G) = 0$ if and only if G is not connected. Thus, μ_{n-1} is called the algebraic connectivity of the graph G [5]. In [1] it is proved that if μ is an eigenvalue of L then $\mu \leq n$ and that the multiplicity of 0 equals the number of components of G . Thus, G is a connected graph if and only if $\mu_{n-1} > 0$.

The Laplacian eigenvalues of a graph are important in the graph theory because they have a relation to numerous graph invariants, including connectivity, expanding property, isoperimetric number, maximum

♣Corresponding author, e-mail: sbuyukkose@gazi.edu.tr

cut, independence number, genus, diameter, mean distance and bandwidth-type parameters of a graph. In many application one needs good lower bound and upper bound of extremal non-trivial Laplacian eigenvalues (see [1], [3], [4], [6], [7], [9], [10], [11], [12], [14]).

In this paper, our aim is to improve the upper bound for the algebraic connectivity μ_{n-1} of G and the lower bound for μ_1 of G given by Maden and Büyükköse [14]. We use Theorem 1 [13] and modify the technique of the proof of Lemma 3 [13], we give an upper bound for the algebraic connectivity μ_{n-1} of G and a lower bound of the largest eigenvalue μ_1 of G in terms of the degree sequence d_1, d_2, \dots, d_n of G and the number $|N_i \cap N_j|$ of common vertices of i and j ($1 \leq i < j \leq n$).

2. THE MAIN RESULT

Firstly we summarize the results of Wolkowicz and Styan on the eigenvalue inequalities which are our fundamental tools in this paper.

Theorem 1. (Theorem 2.1 [13]) *Let A be an $n \times n$ complex matrix with real eigenvalues $\lambda(A)$ and let $m = \frac{\text{tr}A}{n}$, $s = \sqrt{\frac{\text{tr}A^2}{n} - m^2}$. Then*

$$m - s\sqrt{n-1} \leq \lambda_{\min}(A) \leq m - \frac{s}{\sqrt{n-1}} \quad (4)$$

$$m + \frac{s}{\sqrt{n-1}} \leq \lambda_{\max}(A) \leq m + s\sqrt{n-1}. \quad (5)$$

Equality holds on the left (right) of (4) if and only if equality holds on the left (right) of (5) if and only if the $n-1$ largest (smallest) eigenvalues are equal.

In [13] Wolkowicz and Styan proved Theorem 1 by using the following lemmas.

Lemma 2. (Lemma 2.1 [13]) *Let $C = I_n - \frac{ee^T}{n}$, $m = \frac{\lambda^T e}{n}$, $s^2 = \frac{\lambda^T C \lambda}{n}$ where W and $\lambda = (\lambda_j) \in \mathbb{R}^n$ are column vectors, and $e = (1, 1, \dots, 1)^T$. Then*

$$-s\sqrt{nW^T C W} \leq W^T \lambda - mW^T e = W^T C \lambda \leq snW^T C W. \quad (6)$$

Equality holds on the left (right) of (1) if and only if $\lambda = aW + b e$ for some scalars a and b , where $a < 0$ ($a > 0$).

It should be noted that m and s^2 defined in Theorem 1 and Lemma 2 are equivalent [13].

Lemma 3. (Lemma 2.2 [13]) *Let $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$, m and s be defined as in Lemma 2 and $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$. Then*

$$\lambda_n \leq m - \frac{s}{\sqrt{n-1}} \leq m + \frac{s}{\sqrt{n-1}} \leq \lambda_1. \quad (7)$$

Using Theorem 1 Maden and Büyükköse [14] gave upper and lower bounds for μ_{n-1} and μ_1 .

Theorem 4. (Theorem 3 and Corollary 5 [14]) *Let G be a simple graph. Then*

$$\sqrt{m - s \sqrt{\frac{n-2}{2}}} \leq \mu_{n-1} \leq \sqrt{m - s \sqrt{\frac{1}{n-1}}} \quad (8)$$

and

$$\sqrt{m + \frac{s}{\sqrt{n-1}}} \leq \mu_1 \leq \sqrt{m + s\sqrt{n-1}}, \quad (9)$$

where $m = \frac{1}{n} \sum_{i=1}^n d_i (d_i + 1)$ and

$$s^2 = \frac{1}{n} \left(\sum_{i=1}^n (d_i^2 + d_i) + 2 \sum_{\substack{i < j \\ i \sim j}} (d_i + d_j)(d_i + d_j - 2|N_i \cap N_j|) + 2 \sum_{i < j} |N_i \cap N_j|^2 \right) - m^2.$$

We always assume that G is a simple connected graph of order n . The known upper and lower bounds which we used in proof of our main theorem are following:

1. Grone and Merris' bound [15]:

$$\mu_1 \geq d_1 + 1, \quad (1)$$

where d_1 is the largest degree of G .

2. Li and Pan's bound [16]:

$$\mu_2 \geq d_2, \quad (2)$$

where d_2 and μ_2 are the second largest degree and the second largest Laplacian eigenvalue of G , respectively.

3. Fidler's bound [5]: Let G be a graph different from K_n and let d_n be its minimum degree. Then

$$\mu_{n-1} \leq d_n. \quad (3)$$

□

Now, we reprove Lemma 3 for the Laplacian matrix L of G and hence we improve the upper bound for λ_{n-1} in (8) and the lower bound for λ_1 in (9).

Theorem 5. Let G be a simple graph and let m and s be defined as in Theorem 4. Then

$$\mu_{n-1} \leq \left(m - \left(\frac{ns^2 + 2((d_1 + 1)^2 - d_{n-1}^2)(d_2^2 - d_{n-1}^2)}{n^2 - n} \right)^{1/2} \right)^{1/2} \tag{10}$$

and

$$\mu_1 \geq \left(m + \left(\frac{ns^2 + 2(d_1 + 1)^2((d_1 + 1)^2 - d_n^2)}{n^2 - n} \right)^{1/2} \right)^{1/2}. \tag{11}$$

Proof. Let G be a simple graph and let m and s be defined as in Theorem 4. Then we have that

$$\begin{aligned} n^2(m - \mu_{n-1}^2)^2 &= n^2 \left(\frac{1}{n} \sum_{i=1}^n (\mu_i^2 - \mu_{n-1}^2) \right)^2 \\ &= \sum_{i=1}^n (\mu_i^2 - \mu_{n-1}^2)^2 + \sum_{j \neq k} (\mu_j^2 - \mu_{n-1}^2)(\mu_k^2 - \mu_{n-1}^2) \end{aligned}$$

By using (1)-(3) we have that

$$\sum_{j \neq k} (\mu_j^2 - \mu_{n-1}^2)(\mu_k^2 - \mu_{n-1}^2) \geq 2((d_1 + 1)^2 - d_{n-1}^2)(d_2^2 - d_{n-1}^2).$$

On the other hand,

$$\begin{aligned} \sum_{i=1}^n (\mu_i^2 - \mu_{n-1}^2)^2 &= \sum_{i=1}^n (\mu_i^2 - m + m - \mu_{n-1}^2)^2 = \sum_{i=1}^n [(\mu_i^2 - m)(\mu_i^2 + m - 2\mu_{n-1}^2)] + n(m - \mu_{n-1}^2)^2 \\ &= ns^2 + n(m - \mu_{n-1}^2)^2. \end{aligned}$$

Finally, we have that

$$n^2(m - \mu_{n-1}^2)^2 \geq ns^2 + n(m - \mu_{n-1}^2)^2 + 2((d_1 + 1)^2 - d_{n-1}^2)(d_2^2 - d_{n-1}^2).$$

Solving this inequality for μ_{n-1}^2 we obtain the inequality in (10).

Now we similarly expand $n^2(\mu_1^2 - m)$. Then we have

$$n^2(\mu_1^2 - m)^2 = \left(n\mu_1^2 - \sum_{i=1}^n \mu_i^2 \right)^2 = \sum_{i=1}^n (\mu_1^2 - \mu_i^2)^2 + \sum_{j \neq k} (\mu_1^2 - \mu_j^2)(\mu_1^2 - \mu_k^2).$$

By using (1)-(3), we have that

$$\sum_{j \neq k} (\mu_1^2 - \mu_j^2)(\mu_1^2 - \mu_k^2) \geq 2(d_1 + 1)^2((d_1 + 1)^2 - d_n^2).$$

We have that

$$n^2(\mu_1^2 - m)^2 \geq n(\mu_1^2 - m)^2 + ns^2 + 2(d_1 + 1)^2((d_1 + 1)^2 - d_n^2).$$

Solving this inequality for μ_1^2 we obtain the inequality in (11). □

In the proof of Lemma 3 in [13, Lemma 2.2], the second sum is omitted but we consider it to improve the upper bound for μ_{n-1} in (8) and the lower bound for μ_1 in (9). Now we compare our bounds with the bounds of Maden and Büyükköse [14].

Exercise 6. Let $G = (V, E)$ with $V = \{1,2,3,4,5,6,7,8\}$ and

$$E = \{\{1,2\}, \{1,3\}, \{2,3\}, \{2,4\}, \{3,4\}, \{2,5\}, \{2,6\}, \{2,8\}, \{4,5\}, \{4,6\}, \{4,8\}, \{6,7\}, \{7,8\}\}.$$

For this graph $\mu_7 = 1.13$ and $\mu_1 = 7.1$. We present aforesaid upper bounds for μ_7 and lower bounds for μ_1 of the graph G as follows:

	μ_7	(8)	(10)		μ_1	(9)	(11)
G	1.13	4.72	2.76		7,10	3.02	5.17

CONFLICT OF INTEREST

No conflict of interest was declared by the authors.

REFERENCES

- [1] Anderson W.N., Morley T.D., Eigenvalues of the Laplacian of a graph, *Linear Multilinear Algebra* 18 (1985) 141-145.
- [2] Büyükköse Ş., Bounds for singular values using matrix traces, Msc Thesis, *Selçuk University*, (2000).
- [3] Chung F.R.K., Eigenvalues of Graphs, In Proceeding of the International Congress of Mathematician, Zürich, Switzerland, (1994) 1333-1342.
- [4] Das K. Ch., An improved upper bound for Laplacian graph eigenvalues, *Linear Algebra Appl.* 368 (2003) 269-278.
- [5] Fiedler M., Algebraic connectivity of graphs, *Czechoslovak Math. J.* 23 (1973) 298-305.
- [6] Li J.-S., Zhang D., A new upper bound for eigenvalues of the Laplacian matrix of a graph, *Linear Algebra Appl.* 265 (1997) 93-100.
- [7] Li J.-S., Zhang D., On Laplacian eigenvalues of a graph, *Linear Algebra Appl.* 285 (1998) 305-307.
- [8] Lu M., Zhang L., Tian F., Lower bounds of the Laplacian spectrum of graphs based on diameter, *Linear Algebra Appl.* 420 (2007) 400-406.
- [9] Merris R., A note on Laplacian graph eigenvalues, *Linear Algebra Appl.* 285 (1998) 33-35.
- [10] Merris R., Laplacian matrices of Graphs: a Survey, *Linear Algebra and its Applications*, 1 97, 198(1994) 143-176
- [11] Mohar B., The Laplacian spectrum of graphs, In Graph Theory, *Combinatorics and Applications*, Vol.2 (1998) 871-898.
- [12] Rojo O., Soto R., Rojo H., An always nontrivial upper bound for Laplacian graph eigenvalues, *Linear Algebra Appl.* 312(2000),155-159.
- [13] Wolkowicz H., Styan G.P.H., Bounds for eigenvalues using traces, *Linear Algebra Appl.* 29 (1980) 471-506.
- [14] Maden A.D., Büyükköse Ş., Bounds for Laplacian Graph Eigenvalues, *Mathematical Inequalities and Applications*, Vol 12, Num.3 (2012), 529-536.
- [15] Grone R., Merris R., The Laplacian spectrum of a graph, *SIAM J. Discr. Math.* 7 (1994) 221-229.
- [16] Li J.S., Pan Y.L., A note on the second largest eigenvalue of the Laplacian matrix of a graph, *Linear Multilinear Algebra* 48 (2000) 117-121.