

On the Conversion of Convex Functions to Certain within the Unit Disk

ISSN: 2651-544X

<http://dergipark.gov.tr/cpost>

Hasan Şahin^{1,*}, İsmet Yıldız¹, Umran Menek¹

¹ Department of Mathematics, Faculty of Science and Arts, Düzce University, Düzce, Turkey, ORCID:0000-0002-5227-5300

* Corresponding Author E-mail: hasansahin13@gmail.com

Abstract: A function $g(z)$ is said to be univalent in a domain D if it provides a one-to-one mapping onto its image, $g(D)$. Geometrically, this means that the representation of the image domain can be visualized as a suitable set of points in the complex plane. We are mainly interested in univalent functions that are also regular (analytic, holomorphic) in U . Without loss of generality we assume D to be unit disk $U = \{z : |z| < 1\}$. One of the most important events in the history of complex analysis is Riemann's mapping theorem, that any simply connected domain in the complex plane \mathbb{C} which is not the whole complex plane, can be mapped by any analytic function univalently on the unit disk U . The investigation of analytic functions which are univalent in a simply connected region with more than one boundary point can be confined to the investigation of analytic functions which are univalent in U . The theory of univalent functions owes the modern development the amazing Riemann mapping theorem. In 1916, Bieberbach proved that for every $g(z) = z + \sum_{n=2}^{\infty} a_n z^n$ in class S , $|a_2| \leq 2$ with equality only for the rotation of Koebe function $k(z) = \frac{z}{(1-z)^2}$. We give an example of this univalent function with negative coefficients of order $\frac{1}{4}$ and we try to explain $B_{\frac{1}{4}}(1, \frac{\pi}{3}, -1)$ with convex functions.

Keywords: Class s , Convex functions, Univalent functions.

1 Introduction

A indicates the class of the functions of form $g(z)$

$$g(z) = z + a_2 z^2 + a_3 z^3 + a_4 z^4 + a_5 z^5 + a_6 z^6 + a_7 z^7 + a_8 z^8 + \dots$$

that are analytic and univalent in the open unit disk $U = \{z : |z| < 1\}$.

Let $A(n)$ show the A subclass of form's functions

$$g(z) = z + a_2 z^2 + a_3 z^3 + a_4 z^4 + a_5 z^5 + a_6 z^6 + a_7 z^7 + a_8 z^8 + \dots \quad (a_k \geq 0).$$

Let $T(n)$ denote the subclass of $A(n)$ consisting of functions which are univalent in U . Further a function in $T(n)$ is said to be starlike of order $\frac{1}{4}$ if and only if satisfies

$$\Re\left(\frac{zg'(z)}{g(z)}\right) > \frac{1}{4} \quad (z \in U),$$

and such a subclass of $A(n)$ consisting of all the starlike functions of order $\frac{1}{4}$ is denote by $T_{\frac{1}{4}}(n)$. Also, $g(z) \in T(n)$ is said to be convex of order $\frac{1}{4}$ if and only if satisfies

$$\Re\left\{1 + \frac{zg''(z)}{g'(z)}\right\} > \frac{1}{4} \quad (z \in U),$$

and the subclass by $C_{\frac{1}{4}}(n)[1][2][3][6]$.

For $n = 1$, these notations are usually used as $T_{\frac{1}{4}}(1) = T^*(\frac{1}{4})$ this form with starlike function and show us we have this form for convex functions with $C_{\frac{1}{4}}(1) = C^*(\frac{1}{4})[5]$.

Theorem 1. [1] A function $g(z)$ in $A(n)$ is in $T_{\frac{1}{4}}(n)$ if and only if

$$\sum_{k=n+1}^{\infty} (k - \frac{1}{4}) a_k \leq 1 - \frac{1}{4} = \frac{3}{4}.$$

Theorem 2. [1] A function $g(z)$ in $A(n)$ is in $C_{\frac{1}{4}}(n)$ if and only if

$$\sum_{k=n+1}^{\infty} k (k - \frac{1}{4}) a_k \leq 1 - \frac{1}{4} = \frac{3}{4}.$$

We introduced subclass $A(n, \theta)$ of A , and the subclass $T_{\frac{1}{4}}^*(n, \theta)$ and $C_{\frac{1}{4}}(n, \theta)$ of $A(n, \theta)$ in the we define the subclass with this way. Let $A(n, \theta)$ denote the subclass of A consisting of function of the form

$$g(z) = z - \sum_{k=n+1}^{\infty} e^{i(k-1)\theta} a_k z^n \quad (a_k \geq 0, \quad n \in \mathbb{N}) [4].$$

We note that $A(n, \theta) = A(n)$, that is $A(n, \theta)$ is the subclass of analytic functions with negative coefficients. We denote by $T_{\frac{1}{4}}^*(n, \theta)$ starlike functions and $C_{\frac{1}{4}}(n, \theta)$ the subclass of $A(n, \theta)$ of convex functions of order $\frac{1}{4}$ in U .

Theorem 3. A function $g(z)$ in $A(n, \theta)$ is in $T_{\frac{1}{4}}^*(n, \theta)$ if and only if

$$\sum_{k=n+1}^{\infty} (k - \frac{1}{4}) a_k \leq 1 - \frac{1}{4} = \frac{3}{4} [4].$$

Theorem 4. A function $g(z)$ in $A(n, \theta)$ is in $C_{\frac{1}{4}}(n, \theta)$ if and only if

$$\sum_{k=n+1}^{\infty} k (k - \frac{1}{4}) a_k \leq 1 - \frac{1}{4} = \frac{3}{4} [4].$$

Theorem 5. If $g(z)$ is in $C_{\frac{1}{4}}(n, \theta)$, then

$$|z| - \frac{3}{(n+1)(4n+3)} |z|^{n+1} \leq |g(z)| \leq |z| + \frac{3}{(n+1)(4n+3)} |z|^{n+1}.$$

The right hand equality holds for the function

$$g(z) = z - e^{in\theta} \frac{3}{(n+1)(4n+3)} z^{n+1} \quad \left(z = r e^{-i(\theta + \frac{\pi}{n})}, \quad r < 1 \right)$$

and the left hand equality holds for the function

$$g(z) = z - e^{in\theta} \frac{3}{(n+1)(4n+3)} z^{n+1} \quad \left(z = r e^{-i\theta}, \quad r < 1 \right) [4].$$

Theorem 6. (Main theorem) If $g(z) \in B_{\frac{1}{4}}(1, \frac{\pi}{3}, -1)$, then we have

$$g(z) = z - \frac{3+3i\sqrt{3}}{49} z^2 + \frac{3-3i\sqrt{3}}{49} z^3 + \frac{6}{49} z^4 - \frac{3+3i\sqrt{3}}{49} z^5 - \frac{3-3i\sqrt{3}}{49} z^6 - \dots$$

Proof:

Let $B_{\frac{1}{4}}(n, \theta, h)$ denote the subclass of $A(n, \theta)$ consisting of functions of the form

$$g(z) = z - \sum_{k=n+1}^{\infty} e^{i(k-1)\theta} a_k z^n \quad (h \geq -n),$$

where

$$b_{k,h} = \frac{(1 - \frac{1}{4})^2}{(2 - 1 - \frac{1}{4})(2 + 1 + 1 - \frac{1}{4})(2 - \frac{1}{4})2} = \frac{\frac{9}{16}}{\frac{3}{4} \cdot \frac{7}{4} \cdot \frac{7}{4} \cdot 2} = \frac{6}{49}.$$

If we put in place at that $b_{k,h} = \frac{6}{49}$

$$\begin{aligned} g(z) &= z - \sum_{k=n+1}^{\infty} e^{i(k-1)\theta} \frac{6}{49} z^n \quad (h \geq n, \quad n \in \mathbb{N}, \quad n \geq 1). \\ g(z) &= z - \frac{6e^{i\frac{\pi}{3}}}{49} z^2 - \frac{6e^{2i\frac{\pi}{3}}}{49} z^3 - \frac{6e^{3i\frac{\pi}{3}}}{49} z^4 - \frac{6e^{4i\frac{\pi}{3}}}{49} z^5 - \frac{6e^{5i\frac{\pi}{3}}}{49} z^6 - \dots \\ &= z - \frac{6cis\frac{\pi}{3}}{49} z^2 - \frac{6cis\frac{2\pi}{3}}{49} z^3 - \frac{6cis\pi}{49} z^4 - \frac{6cis\frac{4\pi}{3}}{49} z^5 - \frac{6cis\frac{5\pi}{3}}{49} z^6 - \dots \\ &= z - \frac{2\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)}{49} z^2 - \frac{2\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)}{49} z^3 - \frac{-2}{49} z^4 - \frac{2\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)}{49} z^5 - \dots \\ &= z - \frac{3+3i\sqrt{3}}{49} z^2 + \frac{3-3i\sqrt{3}}{49} z^3 + \frac{6}{49} z^4 + \frac{3+3i\sqrt{3}}{49} z^5 - \frac{3-3i\sqrt{3}}{49} z^6 + \dots \end{aligned}$$

We have proved the desired answer and we show that $g(z) \in B_{\frac{1}{4}}(1, \frac{\pi}{3}, -1)$ and $B_{\frac{1}{4}}(1, \frac{\pi}{3}, -1) \in C_{\frac{1}{4}}(1, \frac{\pi}{3})$ so $g(z)$ is convex function. \square

2 References

- [1] S. K. Chatterjea, On starlike functions, *J. Pure Math.*, **1**(1981), 23-26.
- [2] V. S. Kiryakova, M. Saigo, S. Owa, Distortion and characterization theorems for starlike and convex functions related to generalized fractional calculus, *Publ. Res. Inst. Math. Sci.*, **1012**(1997), 25-46.
- [3] T. Sekine, On new generalized classes of analytic functions with negative coefficients, *Report Res. Inst. Sci. Tec. Nihon Univ.*, **35**(1987), 1-26.
- [4] T. Sekine, S. Owa, New problems of coefficients inequalities, *Publ. Res. Inst. Math. Sci.*, **1012**(1997), 164-176.
- [5] H. Silverman, Univalent functions with negative coefficients, *Proc. Amer. Math. Soc.*, **51**(1975), 109-116.
- [6] H. M. Srivasta, S. Owa, S. K. Chatterjea, A note on certainleclass of starlike functions, *Rend. Sem. Mat. Univ. Padova*, **77**(1987), 115-124.