

EXISTENCE OF ENTIRE RADIALY SYMMETRIC SOLUTIONS FOR A QUASILINEAR SYSTEM WITH d -EQUATIONS

Dragos-Patru Covei*

Received 13:05:2010 : Accepted 24:09:2010

Abstract

The studies developed within this article will be focused on achieving results related to the existence and qualitative properties of entire radially symmetric solutions for a Schrodinger problem of type $\Delta_p u_i + h_i(r) |\nabla u_i|^{p-1} = a_i(r) f_i(u_{i+1})$ for $i = \overline{1, d-1}$ and $\Delta_p u_d + h_d(r) |\nabla u_d|^{p-1} = a_d(r) f_d(u_1)$ on \mathbb{R}^N , where $p > 1$, $d \geq 2$, h_i and a_i are nonnegative radial continuous functions and f_i are nonnegative increasing continuous functions on $[0, \infty)$.

Keywords: Large solutions, Bounded solutions, Radially symmetric solutions.

2000 AMS Classification: 35J57, 35J92.

1. Introduction

In this article we generalize existence results for systems such as

$$(1.1) \quad \begin{cases} \Delta_p u_1 + h_1(r) |\nabla u_1|^{p-1} = a_1(r) f_1(u_2) & \text{in } \mathbb{R}^N, \\ \dots\dots\dots \\ \Delta_p u_i + h_i(r) |\nabla u_i|^{p-1} = a_i(r) f_i(u_{i+1}) & \text{in } \mathbb{R}^N, \\ \dots\dots\dots \\ \Delta_p u_d + h_d(r) |\nabla u_d|^{p-1} = a_d(r) f_d(u_1) & \text{in } \mathbb{R}^N, \end{cases}$$

where $r := |x| \geq 0$, $N \geq 3$, $d \geq 2$ is an integer, $i = \overline{1, d-1}$, Δ_p is the p -Laplacian operator defined by

$$\Delta_p u := \operatorname{div}(|\nabla u|^{p-2} \nabla u), \quad 1 < p < \infty,$$

$h_{j=\overline{1,d}}, a_{j=\overline{1,d}} : [0, \infty) \rightarrow [0, \infty)$ are radial continuous functions and $f_{j=\overline{1,d}}$ satisfy the following hypotheses

*Constantin Brancusi University of Tg-Jiu, Calea Eroilor, No 30, Tg-Jiu, Gorj, Romania;
 West University of Timisoara, Bld. Parvan, No. 4, 300223, Timisoara, Timis, Romania.
 E-mail: coveid@yahoo.com

(C1) $f_{j=\overline{1,d}} : [0, \infty) \rightarrow [0, \infty)$ are continuous;

(C2) $f_{j=\overline{1,d}}$ are increasing on $[0, \infty)$.

Answers to questions related to the existence of radially symmetric solutions for problem like (1.1) have been gradually achieved by several authors over many years (see [1]-[13] and references therein).

Before we start describing our results let us mention that the problem of the existence of solutions to (1.1) has received an increased interest with Alan Lair's recent paper [6]. In [6] the author considered problem (1.1) under hypotheses $p = d = 2$, $h_{i=\overline{1,2}} = 0$, $a_1(x) f_1(u_2(x)) := a_1(|x|) u_2^\alpha(|x|)$ ($\alpha \in (0, 1)$) and $a_2(x) f_2(u_1(x)) := a_2(|x|) u_1^\beta(|x|)$ ($\beta \in (0, 1)$).

The main result achieved by Alan Lair may be summarized as follows:

1.1. Proposition. *Problem (1.1) has an explosive radial symmetric solution on \mathbb{R}^N if and only if the continuous radially symmetric functions $a_{i=\overline{1,2}} : [0, \infty) \rightarrow [0, \infty)$ simultaneously meet the following conditions*

$$\int_0^\infty t a_1(t) \left(t^{2-N} \int_0^t s^{N-3} \int_0^s \tau a_2(\tau) d\tau ds \right)^\alpha dt = \infty$$

$$\int_0^\infty t a_2(t) \left(t^{2-N} \int_0^t s^{N-3} \int_0^s \tau a_1(\tau) d\tau ds \right)^\beta dt = \infty.$$

Moreover, the author proposes the following problem:

"It remains unknown whether an analogous result holds for the system

$$\begin{cases} \Delta u_1(|x|) = a_1(|x|) f_1(u_2(|x|)) \text{ for } x \in \mathbb{R}^N, \\ \Delta u_2(|x|) = a_2(|x|) f_2(u_1(|x|)) \text{ for } x \in \mathbb{R}^N, \end{cases}$$

where f_1 and f_2 meet, for example

$$(1.2) \quad \int_1^\infty \left[\int_0^s f_i(t) dt \right]^{-1/2} ds = \infty, \quad i = \overline{1,2},$$

or satisfy the stronger condition

$$(1.3) \quad \int_1^\infty [f_i(t)]^{-1} dt = \infty, \quad i = \overline{1,2}."$$

When $h_{j=\overline{1,d}} = 0$ and $p = 2$ we mention that in the paper [3] the author established the existence results for the problems (1.1) under conditions of type (1.2).

The starting point for the analysis and research of the more general problems (1.1) in order to discover the novelty which is in fact the scientific task of the article, is generated by the proposed problem in [6]. In this sense, we establish results related to the existence and various qualitative properties of solutions for problems (1.1) under conditions of the (1.3) type.

The principal difficulty in the treatment of (1.1) is due to the nonlinear gradient term combined with multiple equations of the system. The solving methods that we will use for the problem are based on results of nonlinear analysis, the above cited articles as well as other techniques that will be discovered during the research.

Throughout this paper we use the notations

$$j = \overline{1,d} \quad H_j(r) := r^{N-1} e^{\int_0^r h_j(t) dt}$$

$$A_j(\infty) := \lim_{r \rightarrow \infty} A_j(r), \quad A_j(r) = \int_0^r \left(\frac{1}{H_j(t)} \int_0^t H_j(s) a_j(s) ds \right)^{1/(p-1)} dt$$

and

$$F(\infty) := \lim_{r \rightarrow \infty} F(r), F(r) = \int_a^r \left(\sum_{j=1}^d f_j(s) \right)^{-1/(p-1)} ds; \quad r \geq a > 0.$$

We see that

$$F'(r) = \left(\sum_{j=1}^d f_j(r) \right)^{-1/(p-1)} > 0 \text{ for all } r > a$$

and F has the inverse function F^{-1} on $[a, \infty)$.

Our main results will be stated in what follows.

1.2. Theorem. *Assume that (C1)–(C2) hold and that*

(C3) $F(\infty) = \infty$.

Then the system (1.1) possesses at least one positive radial solution (u_1, \dots, u_d) . If, in addition, $A_{j=\overline{1,d}}(\infty) < \infty$, the positive radial solution (u_1, \dots, u_d) is bounded. Moreover, when $A_{j=\overline{1,d}}(\infty) = \infty$, the positive solution (u_1, \dots, u_d) is an entire large solution, i.e.

$$\lim_{r \rightarrow \infty} u_1(r) = \dots = \lim_{r \rightarrow \infty} u_d(r) = \infty.$$

1.3. Theorem. *Assume that (C1)–(C2) hold and that*

(C4) $F(\infty) < \infty$;

(C5) $A_{j=\overline{1,d}}(\infty) < \infty$;

(C6) *there exists $\beta > \frac{a}{d}$ such that*

$$\sum_{j=1}^d A_j(\infty) < F(\infty) - F(d\beta).$$

Then, the system (1.1) possesses at least one positive bounded radial solution (u_1, \dots, u_d) satisfying

$$\beta + f_j^{1/(p-1)}(\beta) A_j(r) \leq u_j(r) \leq F^{-1} \left(F(d\beta) + \sum_{j=1}^d A_j(r) \right), \quad j = \overline{1, d}.$$

2. Proof of the theorems

2.1. Proof of Theorem 1.2. We note that radial solutions of (1.1) are positive solutions (u_1, \dots, u_d) of the integral equations

$$\begin{cases} u_1(r) = \beta + \int_0^r \left(\frac{1}{H_1(t)} \int_0^t H_1(s) a_1(s) f_1(u_2(s)) ds \right)^{1/(p-1)} dt, \\ \dots\dots \\ u_i(r) = \beta + \int_0^r \left(\frac{1}{H_i(t)} \int_0^t H_i(s) a_i(s) f_i(u_{i+1}(s)) ds \right)^{1/(p-1)} dt, \\ \dots\dots \\ u_d(r) = \beta + \int_0^r \left(\frac{1}{H_d(t)} \int_0^t H_d(s) a_d(s) f_d(u_1(s)) ds \right)^{1/(p-1)} dt, \end{cases}$$

where β may be any non-negative number.

Define sequences $\{u_j^k\}_{j=\overline{1,d}}^{k \geq 1}$ on $[0, \infty)$ by

$$\begin{cases} u_1^0 = \dots = u_d^0 = \beta, & r \geq 0, \\ u_1^k(r) = \beta + \int_0^r \left(\frac{1}{H_1(t)} \int_0^t H_1(s) a_1(s) f_1(u_2^{k-1}(s)) ds \right)^{1/(p-1)} dt, & \\ \dots\dots\dots & \\ u_i^k(r) = \beta + \int_0^r \left(\frac{1}{H_i(t)} \int_0^t H_i(s) a_i(s) f_i(u_{i+1}^{k-1}(s)) ds \right)^{1/(p-1)} dt, & i = \overline{1, d-1} \\ \dots\dots\dots & \\ u_d^k(r) = \beta + \int_0^r \left(\frac{1}{H_d(t)} \int_0^t H_d(s) a_d(s) f_d(u_1^k(s)) ds \right)^{1/(p-1)} dt, & d \geq 2 \end{cases}$$

We remark that, for all $r \geq 0$, $j = \overline{1,d}$ and $k \in N$,

$$u_j^k(r) \geq \beta.$$

We claim that $\{u_j^k\}_{j=\overline{1,d}}^{k \geq 1}$ are increasing sequences on $[0, \infty)$. Since

$$u_j^0 = \beta \leq u_j^1, \quad j = \overline{1,d},$$

it follows that

$$f_j(u_{j+1}^0) \leq f_j(u_{j+1}^1), \quad j = \overline{1, d-1},$$

and so

$$\begin{aligned} u_j^1(r) &= \beta + \int_0^r \left(\frac{1}{H_j(t)} \int_0^t H_j(s) a_j(s) f_j(u_{j+1}^0(s)) ds \right)^{1/(p-1)} dt \\ &\leq \beta + \int_0^r \left(\frac{1}{H_j(t)} \int_0^t H_j(s) a_j(s) f_j(u_{j+1}^1(s)) ds \right)^{1/(p-1)} dt \\ &= u_j^2(r), \quad j = \overline{1, d-1}. \end{aligned}$$

From the above relation we obtain

$$\begin{aligned} u_d^1(r) &= \beta + \int_0^r \left(\frac{1}{H_d(t)} \int_0^t H_d(s) a_d(s) f_d(u_1^1(s)) ds \right)^{1/(p-1)} dt \\ &\leq \beta + \int_0^r \left(\frac{1}{H_d(t)} \int_0^t H_d(s) a_d(s) f_d(u_1^2(s)) ds \right)^{1/(p-1)} dt \\ &= u_d^2(r). \end{aligned}$$

Consequently,

$$u_j^2 \leq u_j^3, \quad j = \overline{1, d-1},$$

which yield

$$u_d^2 \leq u_d^3.$$

Continuing this line of reasoning, we obtain that $\{u_j^k\}_{j=\overline{1,d}}^{k \geq 1}$ are increasing sequences.

By conditions (C1) and (C2) we obtain

$$\begin{aligned}
 (u_1^k(r))' &= \left(\frac{1}{H_1(r)} \int_0^r H_1(s) a_1(s) f_1(u_2^{k-1}(s)) ds \right)^{1/(p-1)} \\
 &\leq f_1^{1/(p-1)}(u_2^k(r)) A_1'(r) \\
 &\leq \left(\sum_{j=1}^d f_j \left(\sum_{j=1}^d u_j^k(r) \right) \right)^{1/(p-1)} A_1'(r), \\
 &\dots\dots\dots \\
 (u_i^k(r))' &= \left(\frac{1}{H_i(r)} \int_0^r H_i(s) a_i(s) f_i(u_{i+1}^{k-1}(s)) ds \right)^{1/(p-1)} \\
 (2.1) \quad &\leq f_i^{1/(p-1)}(u_{i+1}^k(r)) A_i'(r) \\
 &\leq \left(\sum_{j=1}^d f_j \left(\sum_{j=1}^d u_j^k(r) \right) \right)^{1/(p-1)} A_i'(r), \quad i = \overline{1, d-1} \\
 &\dots\dots\dots \\
 (u_d^k(r))' &= \left(\frac{1}{H_d(r)} \int_0^r H_d(s) a_d(s) f_d(u_1^k(s)) ds \right)^{1/(p-1)} \\
 &\leq f_d^{1/(p-1)}(u_1^k(r)) A_d'(r) \\
 &\leq \left(\sum_{j=1}^d f_j \left(\sum_{j=1}^d u_j^k(r) \right) \right)^{1/(p-1)} A_d'(r), \quad d \geq 2.
 \end{aligned}$$

Summing up gives

$$\left(\sum_{j=1}^d f_j \left(\sum_{j=1}^d u_j^k(t) \right) \right)^{-1/(p-1)} \cdot \left(\sum_{j=1}^d u_j^k(t) \right)' \leq \sum_{j=1}^d A_j'(r).$$

Integrating the above equation between 0 and r , we have

$$\int_0^r \left(\sum_{j=1}^d f_j \left(\sum_{j=1}^d u_j^k(t) \right) \right)^{-1/(p-1)} \cdot \left(\sum_{j=1}^d u_j^k(t) \right)' dt \leq \sum_{j=1}^d A_j(r) \text{ for each } r > 0,$$

which is equivalent to

$$\int_0^r F' \left(\sum_{j=1}^d u_j^k(t) \right) dt \leq \sum_{j=1}^d A_j(r) \text{ for each } r > 0.$$

Consequently,

$$(2.2) \quad F \left(\sum_{j=1}^d u_j^k(r) \right) - F(d\beta) \leq \sum_{j=1}^d A_j(r)$$

for all $r \geq 0$.

Since F^{-1} is increasing on $[0, \infty)$, it follows that

$$(2.3) \quad \sum_{j=1}^d u_j^k(r) \leq F^{-1} \left(F(d\beta) + \sum_{j=1}^d A_j(r) \right)$$

for all $r \geq 0$.

Since (C3) holds, we can see that

$$(2.4) \quad F^{-1}(\infty) = \infty.$$

It follows that the sequences $\{u_j^k\}_{j=1, \overline{d}}^{k \geq 1}$ are bounded and increasing on $[0, c]$ for $c > 0$.

Thus,

$$(u_1^k, \dots, u_d^k) \text{ converges to } (u_1, \dots, u_d) \text{ on } [0, c]^d.$$

Consequently, (u_1, \dots, u_d) is the positive entire radial solution of system (1.1). Moreover, when

$$A_{j=1, \overline{d}}(\infty) < \infty,$$

we see by (2.3) that

$$\sum_{j=1}^d u_j(r) \leq F^{-1} \left(F(d\beta) + \sum_{j=1}^d A_j(\infty) \right) \text{ for all } r \geq 0.$$

When

$$A_j(\infty) = \infty \text{ for } j = \overline{1, d}$$

then by (C2) and the monotonicity of $\{u_j^k\}_{j=1, \overline{d}}^{k \geq 1}$ it follows that

$$u_j(r) \geq \beta + f_j^{1/(p-1)}(\beta) A_j(r), \text{ for all } r \geq 0 \text{ and } j = \overline{1, d}.$$

Thus

$$\lim_{r \rightarrow \infty} u_1(r) = \dots = \lim_{r \rightarrow \infty} u_d(r) = \infty,$$

ending the proof of the Theorem. □

2.2. Proof of Theorem 1.3. We proceed as in the proof of the Theorem 1.2. Then, by (2.2) we see that

$$(2.5) \quad F \left(\sum_{j=1}^d u_j^k(r) \right) \leq F(d\beta) + \sum_{j=1}^d A_j(\infty) < F(\infty) < \infty.$$

Since F^{-1} is strictly increasing on $[0, \infty)$, we have

$$(2.6) \quad \sum_{j=1}^d u_j^k(r) \leq F^{-1} \left(F(d\beta) + \sum_{j=1}^d A_j(\infty) \right) < \infty \text{ for all } r \geq 0.$$

Therefore, since the sequence $\{u_j^k(r)\}_{j=1, \overline{d}}$ is monotone it converges to some function $\{u_j(r)\}_{j=1, \overline{d}}$ on \mathbb{R}^N that in fact is a solution to (1.1). This concludes the proof. □

2.1. Remark. If (C1), (C2), (C3) are satisfied then

$$\int_a^\infty \frac{ds}{f_j^{1/(p-1)}(s)} = \infty \text{ for all } j = \overline{1, d}.$$

Acknowledgements

This article is dedicated to my daughter Daria, my son Rareş and to the anonymous reviewers for their support in my work. The work is supported by CNCSIS-UEFISCSU, project number PN II- IDEI 1080/2008 from West University of Timisoara.

References

- [1] Bandle, C. and Giarrusso, E. *Boundary blowup for semilinear elliptic equations with nonlinear gradient terms*, Advances in Differential Equations **1** (1), 133–150, 1996.
- [2] Covei, D.P. *Large and entire large solution for a quasilinear problem*, Nonlinear Analysis: Theory, Methods & Applications **70** (4), 1738–1745, 2009.
- [3] Covei, D.P. *Radial and nonradial solutions for a semilinear elliptic system of Schrödinger type*, Funkcialaj Ekvacioj-Serio Internacia, accepted manuscript.
- [4] Li, H., Zhang, P. and Zhang, Z. *A remark on the existence of entire positive solutions for a class of semilinear elliptic systems*, Journal of Mathematical Analysis and Applications (2009), doi:10.1016/j.jmaa.2009.10.036.
- [5] Lair, A. V. and Wood, A. W. *Existence of entire large positive solutions of semilinear elliptic systems*, Journal of Differential Equations **164**, 380–394, 2000.
- [6] Lair, A. V. *A necessary and sufficient condition for the existence of large solutions to sublinear elliptic systems*, Journal of Mathematical Analysis and Applications **365** (1), 103–108, 2010.
- [7] Keller, J.B. *On solution of $\Delta u = f(u)$* , Communications on Pure and Applied Mathematics **10**, 503–510, 1957.
- [8] Osserman, R. *On the inequality $\Delta u \geq f(u)$* , Pacific Journal of Mathematics **7**, 1641–1647, 1957.
- [9] Smith, D.N. *Existence of large solutions to semilinear elliptic equations with multiple terms*, Master’s Thesis, Air Force Inst of Tech Wright-Patterson Afb OH School of Engineering and Management, 41 Pages, Sep 2006.
- [10] Teramoto, T. *Existence and nonexistence of positive radial entire solutions of second order quasilinear elliptic systems*, Hiroshima Math. J. **30**, 437–461, 2000.
- [11] Toumi, F. *Existence of blowup solutions for nonlinear problems with a gradient term*, International Journal of Mathematics and Mathematical Sciences Volume 2006 (2006), Article ID 80605, 11 pages.
- [12] Yang, Z. *Existence of entire explosive positive radial solutions for a class of quasilinear elliptic systems*, Journal of Mathematical Analysis and Applications **288**, 768–783, 2003.
- [13] Ye, D. and Zhou, F. *Existence and nonexistence of entire large solutions for some semilinear elliptic equations*, Journal of Partial Differential Equations Year **3**, 253–262, 2008.