

Ostrowski and Trapezoid Type Inequalities for the Generalized k - g -Fractional Integrals of Functions with Bounded Variation

Silvestru Sever Dragomir^{1*}

Abstract

In this paper we establish some Ostrowski and trapezoid type inequalities for the k - g -fractional integrals of functions of bounded variation. Applications for mid-point and trapezoid inequalities are provided as well. Some examples for a general exponential fractional integral are also given.

Keywords: Functions of bounded variation, Generalized Riemann-Liouville fractional integrals, Hadamard fractional integrals, Ostrowski type inequalities, Trapezoid inequalities

2010 AMS: 26D15, 26D10, 26D07, 26A33

¹ Mathematics, College of Engineering & Science Victoria University, Melbourne City, MC 8001, Australia, ORCID: 0000-0003-2902-6805

*Corresponding author: sever.dragomir@vu.edu.au

Received: 02 October 2019, Accepted: 21 November 2019, Available online: 29 December 2019

1. Introduction

Assume that the kernel k is defined either on $(0, \infty)$ or on $[0, \infty)$ with complex values and integrable on any finite subinterval. We define the function $K : [0, \infty) \rightarrow \mathbb{C}$ by

$$K(t) := \begin{cases} \int_0^t k(s) ds & \text{if } 0 < t, \\ 0 & \text{if } t = 0. \end{cases}$$

As a simple example, if $k(t) = t^{\alpha-1}$ then for $\alpha \in (0, 1)$ the function k is defined on $(0, \infty)$ and $K(t) := \frac{1}{\alpha} t^\alpha$ for $t \in [0, \infty)$. If $\alpha \geq 1$, then k is defined on $[0, \infty)$ and $K(t) := \frac{1}{\alpha} t^\alpha$ for $t \in [0, \infty)$.

Let g be a strictly increasing function on (a, b) , having a continuous derivative g' on (a, b) . For the Lebesgue integrable function $f : (a, b) \rightarrow \mathbb{C}$, we define the k - g -left-sided fractional integral of f by

$$S_{k,g,a+} f(x) = \int_a^x k(g(x) - g(t)) g'(t) f(t) dt, \quad x \in (a, b) \quad (1.1)$$

and the k - g -right-sided fractional integral of f by

$$S_{k,g,b-} f(x) = \int_x^b k(g(t) - g(x)) g'(t) f(t) dt, \quad x \in [a, b). \quad (1.2)$$

If we take $k(t) = \frac{1}{\Gamma(\alpha)}t^{\alpha-1}$, where Γ is the *Gamma function*, then

$$\begin{aligned} S_{k,g,a+}f(x) &= \frac{1}{\Gamma(\alpha)} \int_a^x [g(x) - g(t)]^{\alpha-1} g'(t) f(t) dt \\ &=: I_{a+,g}^\alpha f(x), \quad a < x \leq b \end{aligned}$$

and

$$\begin{aligned} S_{k,g,b-}f(x) &= \frac{1}{\Gamma(\alpha)} \int_x^b [g(t) - g(x)]^{\alpha-1} g'(t) f(t) dt \\ &=: I_{b-,g}^\alpha f(x), \quad a \leq x < b, \end{aligned} \tag{1.3}$$

which are the *generalized left- and right-sided Riemann-Liouville fractional integrals* of a function f with respect to another function g on $[a, b]$ as defined in [1, p. 100]

For $g(t) = t$ in (1.3) we have the classical *Riemann-Liouville fractional integrals* while for the logarithmic function $g(t) = \ln t$ we have the *Hadamard fractional integrals* [1, p. 111]

$$H_{a+}^\alpha f(x) := \frac{1}{\Gamma(\alpha)} \int_a^x \left[\ln \left(\frac{x}{t} \right) \right]^{\alpha-1} \frac{f(t) dt}{t}, \quad 0 \leq a < x \leq b$$

and

$$H_{b-}^\alpha f(x) := \frac{1}{\Gamma(\alpha)} \int_x^b \left[\ln \left(\frac{t}{x} \right) \right]^{\alpha-1} \frac{f(t) dt}{t}, \quad 0 \leq a < x < b.$$

One can consider the function $g(t) = -t^{-1}$ and define the "*Harmonic fractional integrals*" by

$$R_{a+}^\alpha f(x) := \frac{x^{1-\alpha}}{\Gamma(\alpha)} \int_a^x \frac{f(t) dt}{(x-t)^{1-\alpha} t^{\alpha+1}}, \quad 0 \leq a < x \leq b$$

and

$$R_{b-}^\alpha f(x) := \frac{x^{1-\alpha}}{\Gamma(\alpha)} \int_x^b \frac{f(t) dt}{(t-x)^{1-\alpha} t^{\alpha+1}}, \quad 0 \leq a < x < b.$$

Also, for $g(t) = \exp(\beta t)$, $\beta > 0$, we can consider the " *β -Exponential fractional integrals*"

$$E_{a+,\beta}^\alpha f(x) := \frac{\beta}{\Gamma(\alpha)} \int_a^x [\exp(\beta x) - \exp(\beta t)]^{\alpha-1} \exp(\beta t) f(t) dt,$$

for $a < x \leq b$ and

$$E_{b-,\beta}^\alpha f(x) := \frac{\beta}{\Gamma(\alpha)} \int_x^b [\exp(\beta t) - \exp(\beta x)]^{\alpha-1} \exp(\beta t) f(t) dt,$$

for $a \leq x < b$.

If we take $g(t) = t$ in (1.1) and (1.2), then we can consider the following *k-fractional integrals*

$$S_{k,a+}f(x) = \int_a^x k(x-t) f(t) dt, \quad x \in (a, b) \tag{1.4}$$

and

$$S_{k,b-}f(x) = \int_x^b k(t-x) f(t) dt, \quad x \in [a, b). \tag{1.5}$$

In [2], Raina studied a class of functions defined formally by

$$\mathcal{F}_{\rho,\lambda}^\sigma(x) := \sum_{k=0}^{\infty} \frac{\sigma(k)}{\Gamma(\rho k + \lambda)} x^k, \quad |x| < R, \quad \text{with } R > 0 \tag{1.6}$$

for $\rho, \lambda > 0$ where the coefficients $\sigma(k)$ generate a bounded sequence of positive real numbers. With the help of (1.6), Raina defined the following left-sided fractional integral operator

$$\mathcal{I}_{\rho, \lambda, a+; w}^{\sigma} f(x) := \int_a^x (x-t)^{\lambda-1} \mathcal{F}_{\rho, \lambda}^{\sigma}(w(x-t)^{\rho}) f(t) dt, \quad x > a \quad (1.7)$$

where $\rho, \lambda > 0, w \in \mathbb{R}$ and f is such that the integral on the right side exists.

In [3], the right-sided fractional operator was also introduced as

$$\mathcal{I}_{\rho, \lambda, b-; w}^{\sigma} f(x) := \int_x^b (t-x)^{\lambda-1} \mathcal{F}_{\rho, \lambda}^{\sigma}(w(t-x)^{\rho}) f(t) dt, \quad x < b \quad (1.8)$$

where $\rho, \lambda > 0, w \in \mathbb{R}$ and f is such that the integral on the right side exists. Several Ostrowski type inequalities were also established.

We observe that for $k(t) = t^{\lambda-1} \mathcal{F}_{\rho, \lambda}^{\sigma}(wt^{\rho})$ we re-obtain the definitions of (1.7) and (1.8) from (1.4) and (1.5).

In [4], Kirane and Torebek introduced the following *exponential fractional integrals*

$$\mathcal{T}_{a+}^{\alpha} f(x) := \frac{1}{\alpha} \int_a^x \exp\left\{-\frac{1-\alpha}{\alpha}(x-t)\right\} f(t) dt, \quad x > a \quad (1.9)$$

and

$$\mathcal{T}_{b-}^{\alpha} f(x) := \frac{1}{\alpha} \int_x^b \exp\left\{-\frac{1-\alpha}{\alpha}(t-x)\right\} f(t) dt, \quad x < b \quad (1.10)$$

where $\alpha \in (0, 1)$.

We observe that for $k(t) = \frac{1}{\alpha} \exp(-\frac{1-\alpha}{\alpha}t)$, $t \in \mathbb{R}$ we re-obtain the definitions of (1.9) and (1.10) from (1.4) and (1.5).

Let g be a strictly increasing function on (a, b) , having a continuous derivative g' on (a, b) . We can define the more general exponential fractional integrals

$$\mathcal{T}_{g, a+}^{\alpha} f(x) := \frac{1}{\alpha} \int_a^x \exp\left\{-\frac{1-\alpha}{\alpha}(g(x)-g(t))\right\} g'(t) f(t) dt, \quad x > a$$

and

$$\mathcal{T}_{g, b-}^{\alpha} f(x) := \frac{1}{\alpha} \int_x^b \exp\left\{-\frac{1-\alpha}{\alpha}(g(t)-g(x))\right\} g'(t) f(t) dt, \quad x < b$$

where $\alpha \in (0, 1)$.

Let g be a strictly increasing function on (a, b) , having a continuous derivative g' on (a, b) . Assume that $\alpha > 0$. We can also define the *logarithmic fractional integrals*

$$\mathcal{L}_{g, a+}^{\alpha} f(x) := \int_a^x (g(x)-g(t))^{\alpha-1} \ln(g(x)-g(t)) g'(t) f(t) dt,$$

for $0 < a < x \leq b$ and

$$\mathcal{L}_{g, b-}^{\alpha} f(x) := \int_x^b (g(t)-g(x))^{\alpha-1} \ln(g(t)-g(x)) g'(t) f(t) dt,$$

for $0 < a \leq x < b$, where $\alpha > 0$. These are obtained from (1.4) and (1.5) for the kernel $k(t) = t^{\alpha-1} \ln t, t > 0$.

For $\alpha = 1$ we get

$$\mathcal{L}_{g, a+} f(x) := \int_a^x \ln(g(x)-g(t)) g'(t) f(t) dt, \quad 0 < a < x \leq b$$

and

$$\mathcal{L}_{g, b-} f(x) := \int_x^b \ln(g(t)-g(x)) g'(t) f(t) dt, \quad 0 < a \leq x < b.$$

For $g(t) = t$, we have the simple forms

$$\mathcal{L}_{a+}^{\alpha} f(x) := \int_a^x (x-t)^{\alpha-1} \ln(x-t) f(t) dt, \quad 0 < a < x \leq b,$$

$$\mathcal{L}_{b-}^{\alpha} f(x) := \int_x^b (t-x)^{\alpha-1} \ln(t-x) f(t) dt, \quad 0 < a \leq x < b,$$

$$\mathcal{L}_{a+} f(x) := \int_a^x \ln(x-t) f(t) dt, \quad 0 < a < x \leq b$$

and

$$\mathcal{L}_{b-} f(x) := \int_x^b \ln(t-x) f(t) dt, \quad 0 < a \leq x < b.$$

In the recent paper [5] we obtained the following Ostrowski and trapezoid type inequalities for the *generalized left- and right-sided Riemann-Liouville fractional integrals* of a function f with respect to another function g on $[a, b]$.

Theorem 1.1. *Let $f : [a, b] \rightarrow \mathbb{C}$ be a function of bounded variation on $[a, b]$. Also let g be a strictly increasing function on (a, b) , having a continuous derivative g' on (a, b) . Then we have*

$$\begin{aligned} & \left| I_{x-,g}^{\alpha} f(a) + I_{x+,g}^{\alpha} f(b) - \frac{1}{\Gamma(\alpha+1)} [(g(x) - g(a))^{\alpha} + (g(b) - g(x))^{\alpha}] f(x) \right| \\ & \leq \frac{1}{\Gamma(\alpha)} \left[\int_a^x (g(t) - g(a))^{\alpha-1} g'(t) \mathcal{V}_t^x(f) dt + \int_x^b (g(b) - g(t))^{\alpha-1} g'(t) \mathcal{V}_x^t(f) dt \right] \\ & \leq \frac{1}{\Gamma(\alpha+1)} \left[(g(x) - g(a))^{\alpha} \mathcal{V}_a^x(f) + (g(b) - g(x))^{\alpha} \mathcal{V}_x^b(f) \right] \\ & \leq \frac{1}{\Gamma(\alpha+1)} \left\{ \begin{array}{l} \left[\frac{1}{2} (g(b) - g(a)) + \left| g(x) - \frac{g(a)+g(b)}{2} \right| \right]^{\alpha} \mathcal{V}_a^b(f); \\ ((g(x) - g(a))^{\alpha p} + (g(b) - g(x))^{\alpha p})^{1/p} \left((\mathcal{V}_a^x(f))^q + (\mathcal{V}_x^b(f))^q \right)^{1/q} \\ \text{with } p, q > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \left[\frac{1}{2} \mathcal{V}_a^b(f) + \frac{1}{2} \left| \mathcal{V}_a^x(f) - \mathcal{V}_x^b(f) \right| \right] ((g(x) - g(a))^{\alpha} + (g(b) - g(x))^{\alpha}) \end{array} \right. \end{aligned}$$

and

$$\begin{aligned} & \left| I_{a+,g}^{\alpha} f(x) + I_{b-,g}^{\alpha} f(x) - \frac{1}{\Gamma(\alpha+1)} [(g(x) - g(a))^{\alpha} f(a) + (g(b) - g(x))^{\alpha} f(b)] \right| \\ & \leq \frac{1}{\Gamma(\alpha)} \left[\int_a^x (g(x) - g(t))^{\alpha-1} g'(t) \mathcal{V}_a^t(f) dt + \int_x^b (g(t) - g(x))^{\alpha-1} g'(t) \mathcal{V}_t^b(f) dt \right] \\ & \leq \frac{1}{\Gamma(\alpha+1)} \left[(g(x) - g(a))^{\alpha} \mathcal{V}_a^x(f) + (g(b) - g(x))^{\alpha} \mathcal{V}_x^b(f) \right] \\ & \leq \frac{1}{\Gamma(\alpha+1)} \left\{ \begin{array}{l} \left[\frac{1}{2} (g(b) - g(a)) + \left| g(x) - \frac{g(a)+g(b)}{2} \right| \right]^{\alpha} \mathcal{V}_a^b(f); \\ ((g(x) - g(a))^{\alpha p} + (g(b) - g(x))^{\alpha p})^{1/p} \left((\mathcal{V}_a^x(f))^q + (\mathcal{V}_x^b(f))^q \right)^{1/q} \\ \text{with } p, q > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \left[\frac{1}{2} \mathcal{V}_a^b(f) + \frac{1}{2} \left| \mathcal{V}_a^x(f) - \mathcal{V}_x^b(f) \right| \right] ((g(x) - g(a))^{\alpha} + (g(b) - g(x))^{\alpha}) \end{array} \right. \end{aligned}$$

for any $x \in (a, b)$.

For applications to the classical Riemann-Liouville fractional integrals, Hadamard fractional integrals and Harmonic fractional integrals see [5].

For several Ostrowski type inequalities for Riemann-Liouville fractional integrals see [6]-[21], [22]-[32] and the references therein.

Motivated by the above results, in this paper we establish some Ostrowski and trapezoid type inequalities for the k - g -fractional integrals of functions of bounded variation. Applications for mid-point and trapezoid inequalities are provided as well. Some examples for a general exponential fractional integral are also given.

2. Some identities for the operator $S_{k,g,a+,b-}$

For k and g as at the beginning of Introduction, we consider the mixed operator

$$\begin{aligned} S_{k,g,a+,b-}f(x) &:= \frac{1}{2} [S_{k,g,a+}f(x) + S_{k,g,b-}f(x)] \\ &= \frac{1}{2} \left[\int_a^x k(g(x) - g(t))g'(t)f(t)dt + \int_x^b k(g(t) - g(x))g'(t)f(t)dt \right] \end{aligned}$$

for the Lebesgue integrable function $f : (a, b) \rightarrow \mathbb{C}$ and $x \in (a, b)$.

The following two parameters representation for the operator $S_{k,g,a+,b-}$ holds:

Lemma 2.1. *With the above assumptions for k , g and f we have*

$$\begin{aligned} S_{k,g,a+,b-}f(x) &= \frac{1}{2} [\gamma K(g(b) - g(x)) + \lambda K(g(x) - g(a))] \\ &\quad + \frac{1}{2} \int_a^x k(g(x) - g(t))g'(t)[f(t) - \lambda]dt \\ &\quad + \frac{1}{2} \int_x^b k(g(t) - g(x))g'(t)[f(t) - \gamma]dt \end{aligned} \tag{2.1}$$

for any $\lambda, \gamma \in \mathbb{C}$.

Proof. We have, by taking the derivative over t and using the chain rule, that

$$[K(g(x) - g(t))] = K'(g(x) - g(t))(g(x) - g(t))' = -k(g(x) - g(t))g'(t)$$

for $t \in (a, x)$ and

$$[K(g(t) - g(x))] = K'(g(t) - g(x))(g(t) - g(x))' = k(g(t) - g(x))g'(t)$$

for $t \in (x, b)$.

Therefore, for any $\lambda, \gamma \in \mathbb{C}$ we have

$$\begin{aligned} &\int_a^x k(g(x) - g(t))g'(t)[f(t) - \lambda]dt \\ &= \int_a^x k(g(x) - g(t))g'(t)f(t)dt - \lambda \int_a^x k(g(x) - g(t))g'(t)dt \\ &= S_{k,g,a+}f(x) + \lambda \int_a^x [K(g(x) - g(t))] = S_{k,g,a+}f(x) + \lambda [K(g(x) - g(t))]_a^x \\ &= S_{k,g,a+}f(x) + \lambda [K(g(x) - g(t))]_a^x = S_{k,g,a+}f(x) - \lambda K(g(x) - g(a)) \end{aligned} \tag{2.2}$$

and

$$\begin{aligned} &\int_x^b k(g(t) - g(x))g'(t)[f(t) - \gamma]dt \\ &= \int_x^b k(g(t) - g(x))g'(t)f(t)dt - \gamma \int_x^b k(g(t) - g(x))g'(t)dt \\ &= S_{k,g,b-}f(x) - \gamma \int_x^b [K(g(t) - g(x))] = S_{k,g,b-}f(x) - \gamma [K(g(t) - g(x))]_x^b \\ &= S_{k,g,b-}f(x) - \gamma [K(g(t) - g(x))]_x^b = S_{k,g,b-}f(x) - \gamma K(g(b) - g(x)) \end{aligned} \tag{2.3}$$

for $x \in (a, b)$.

If we add the equalities (2.2) and (2.3) and divide by 2 then we get the desired result (2.1). □

Corollary 2.2. *With the above assumptions for k , g and f we have the Ostrowski type identity*

$$\begin{aligned}
 S_{k,g,a+,b-}f(x) &= \frac{1}{2} [K(g(b) - g(x)) + K(g(x) - g(a))] f(x) \\
 &+ \frac{1}{2} \int_a^x k(g(x) - g(t)) g'(t) [f(t) - f(x)] dt \\
 &+ \frac{1}{2} \int_x^b k(g(t) - g(x)) g'(t) [f(t) - f(x)] dt
 \end{aligned} \tag{2.4}$$

and the trapezoid type identity

$$\begin{aligned}
 S_{k,g,a+,b-}f(x) &= \frac{1}{2} [K(g(b) - g(x)) f(b) + K(g(x) - g(a)) f(a)] \\
 &+ \frac{1}{2} \int_a^x k(g(x) - g(t)) g'(t) [f(t) - f(a)] dt \\
 &+ \frac{1}{2} \int_x^b k(g(t) - g(x)) g'(t) [f(t) - f(b)] dt
 \end{aligned} \tag{2.5}$$

for any $x \in (a, b)$.

For $x = \frac{a+b}{2}$ we can consider

$$\begin{aligned}
 M_{k,g,a+,b-}f &:= S_{k,g,a+,b-}f\left(\frac{a+b}{2}\right) \\
 &= \frac{1}{2} \int_a^{\frac{a+b}{2}} k\left(g\left(\frac{a+b}{2}\right) - g(t)\right) g'(t) f(t) dt \\
 &+ \frac{1}{2} \int_{\frac{a+b}{2}}^b k\left(g(t) - g\left(\frac{a+b}{2}\right)\right) g'(t) f(t) dt.
 \end{aligned}$$

By (2.4) we have the representation

$$\begin{aligned}
 M_{k,g,a+,b-}f &= \frac{1}{2} \left[K\left(g(b) - g\left(\frac{a+b}{2}\right)\right) + K\left(g\left(\frac{a+b}{2}\right) - g(a)\right) \right] f\left(\frac{a+b}{2}\right) \\
 &+ \frac{1}{2} \int_a^{\frac{a+b}{2}} k\left(g\left(\frac{a+b}{2}\right) - g(t)\right) g'(t) \left[f(t) - f\left(\frac{a+b}{2}\right) \right] dt \\
 &+ \frac{1}{2} \int_{\frac{a+b}{2}}^b k\left(g(t) - g\left(\frac{a+b}{2}\right)\right) g'(t) \left[f(t) - f\left(\frac{a+b}{2}\right) \right] dt
 \end{aligned}$$

and (2.5) we have

$$\begin{aligned}
 M_{k,g,a+,b-}f &= \frac{1}{2} \left[K\left(g(b) - g\left(\frac{a+b}{2}\right)\right) f(b) + K\left(g\left(\frac{a+b}{2}\right) - g(a)\right) f(a) \right] \\
 &+ \frac{1}{2} \int_a^{\frac{a+b}{2}} k\left(g\left(\frac{a+b}{2}\right) - g(t)\right) g'(t) [f(t) - f(a)] dt \\
 &+ \frac{1}{2} \int_{\frac{a+b}{2}}^b k\left(g(t) - g\left(\frac{a+b}{2}\right)\right) g'(t) [f(t) - f(b)] dt.
 \end{aligned}$$

If g is a function which maps an interval I of the real line to the real numbers, and is both continuous and injective then we can define the g -mean of two numbers $a, b \in I$ as

$$M_g(a, b) := g^{-1}\left(\frac{g(a) + g(b)}{2}\right).$$

If $I = \mathbb{R}$ and $g(t) = t$ is the identity function, then $M_g(a, b) = A(a, b) := \frac{a+b}{2}$, the arithmetic mean. If $I = (0, \infty)$ and $g(t) = \ln t$, then $M_g(a, b) = G(a, b) := \sqrt{ab}$, the geometric mean. If $I = (0, \infty)$ and $g(t) = \frac{1}{t}$, then $M_g(a, b) = H(a, b) := \frac{2ab}{a+b}$, the harmonic mean. If $I = (0, \infty)$ and $g(t) = t^p$, $p \neq 0$, then $M_g(a, b) = M_p(a, b) := \left(\frac{a^p+b^p}{2}\right)^{1/p}$, the power mean with exponent p . Finally, if $I = \mathbb{R}$ and $g(t) = \exp t$, then

$$M_g(a, b) = LME(a, b) := \ln \left(\frac{\exp a + \exp b}{2} \right),$$

the LogMeanExp function.

Using the g -mean of two numbers we can introduce

$$\begin{aligned} P_{k,g,a+,b-}f &:= S_{k,g,a+,b-}f(M_g(a, b)) \\ &= \frac{1}{2} \int_a^{M_g(a,b)} k \left(\frac{g(a)+g(b)}{2} - g(t) \right) g'(t) f(t) dt \\ &\quad + \frac{1}{2} \int_{M_g(a,b)}^b k \left(g(t) - \frac{g(a)+g(b)}{2} \right) g'(t) f(t) dt. \end{aligned}$$

Using (2.4) and (2.5) we have the representations

$$\begin{aligned} P_{k,g,a+,b-}f &= K \left(\frac{g(b)-g(a)}{2} \right) f(M_g(a, b)) \\ &\quad + \frac{1}{2} \int_a^{M_g(a,b)} k \left(\frac{g(a)+g(b)}{2} - g(t) \right) g'(t) [f(t) - f(M_g(a, b))] dt \\ &\quad + \frac{1}{2} \int_{M_g(a,b)}^b k \left(g(t) - \frac{g(a)+g(b)}{2} \right) g'(t) [f(t) - f(M_g(a, b))] dt \end{aligned}$$

and

$$\begin{aligned} P_{k,g,a+,b-}f &= K \left(\frac{g(b)-g(a)}{2} \right) \frac{f(b)+f(a)}{2} \\ &\quad + \frac{1}{2} \int_a^{M_g(a,b)} k \left(\frac{g(a)+g(b)}{2} - g(t) \right) g'(t) [f(t) - f(a)] dt \\ &\quad + \frac{1}{2} \int_{M_g(a,b)}^b k \left(g(t) - \frac{g(a)+g(b)}{2} \right) g'(t) [f(t) - f(b)] dt. \end{aligned}$$

3. Some identities for the dual operator $\check{S}_{k,g,a+,b-}$

Observe that

$$S_{k,g,x+}f(b) = \int_x^b k(g(b) - g(t)) g'(t) f(t) dt, \quad x \in [a, b]$$

and

$$S_{k,g,x-}f(a) = \int_a^x k(g(t) - g(a)) g'(t) f(t) dt, \quad x \in (a, b].$$

Define also the mixed operator

$$\begin{aligned} \check{S}_{k,g,a+,b-}f(x) &:= \frac{1}{2} [S_{k,g,x+}f(b) + S_{k,g,x-}f(a)] \\ &= \frac{1}{2} \left[\int_x^b k(g(b) - g(t)) g'(t) f(t) dt + \int_a^x k(g(t) - g(a)) g'(t) f(t) dt \right] \end{aligned}$$

for any $x \in (a, b)$.

Lemma 3.1. *With the above assumptions for k , g and f we have*

$$\begin{aligned} \check{S}_{k,g,a+,b-}f(x) &= \frac{1}{2} [\lambda K(g(b) - g(x)) + \gamma K(g(x) - g(a))] \\ &+ \frac{1}{2} \int_a^x k(g(t) - g(a)) g'(t) [f(t) - \gamma] dt \\ &+ \frac{1}{2} \int_x^b k(g(b) - g(t)) g'(t) [f(t) - \lambda] dt \end{aligned} \tag{3.1}$$

for any $\lambda, \gamma \in \mathbb{C}$.

Proof. We have, by taking the derivative over t and using the chain rule, that

$$[K(g(b) - g(t))] = K'(g(b) - g(t)) (g(b) - g(t))' = -k(g(b) - g(t)) g'(t)$$

for $t \in (x, b)$ and

$$[K(g(t) - g(a))] = K'(g(t) - g(a)) (g(t) - g(a))' = k(g(t) - g(a)) g'(t)$$

for $t \in (a, x)$.

For any $\lambda, \gamma \in \mathbb{C}$ we have

$$\begin{aligned} &\int_x^b k(g(b) - g(t)) g'(t) [f(t) - \lambda] dt \\ &= \int_x^b k(g(b) - g(t)) g'(t) f(t) dt - \lambda \int_x^b k(g(b) - g(t)) g'(t) dt \\ &= S_{k,g,x+}f(b) + \lambda \int_x^b [K(g(b) - g(t))] dt \\ &= S_{k,g,x+}f(b) - \lambda K(g(b) - g(x)) \end{aligned} \tag{3.2}$$

and

$$\begin{aligned} &\int_a^x k(g(t) - g(a)) g'(t) [f(t) - \gamma] dt \\ &= \int_a^x k(g(t) - g(a)) g'(t) f(t) dt - \gamma \int_a^x k(g(t) - g(a)) g'(t) dt \\ &= \int_a^x k(g(t) - g(a)) g'(t) f(t) dt - \gamma \int_a^x [K(g(t) - g(a))] dt \\ &= \int_a^x k(g(t) - g(a)) g'(t) f(t) dt - \gamma K(g(x) - g(a)) \end{aligned} \tag{3.3}$$

for $x \in (a, b)$.

If we add the equalities (3.2) and (3.3) and divide by 2 then we get the desired result (3.1). □

Corollary 3.2. *With the assumptions of Lemma 3.1 we have the Ostrowski type identity*

$$\begin{aligned} \check{S}_{k,g,a+,b-}f(x) &= \frac{1}{2} [K(g(b) - g(x)) + K(g(x) - g(a))] f(x) \\ &+ \frac{1}{2} \int_a^x k(g(t) - g(a)) g'(t) [f(t) - f(x)] dt \\ &+ \frac{1}{2} \int_x^b k(g(b) - g(t)) g'(t) [f(t) - f(x)] dt \end{aligned} \tag{3.4}$$

and the trapezoid identity

$$\begin{aligned} \check{S}_{k,g,a+,b-}f(x) &= \frac{1}{2} [K(g(b) - g(x)) f(b) + K(g(x) - g(a)) f(a)] \\ &+ \frac{1}{2} \int_a^x k(g(t) - g(a)) g'(t) [f(t) - f(a)] dt \\ &+ \frac{1}{2} \int_x^b k(g(b) - g(t)) g'(t) [f(t) - f(b)] dt \end{aligned} \tag{3.5}$$

for $x \in (a, b)$.

For $x = \frac{a+b}{2}$ we can consider

$$\begin{aligned} \check{M}_{k,g,a+,b-f} &:= \check{S}_{k,g,a+,b-f} \left(\frac{a+b}{2} \right) \\ &= \frac{1}{2} \int_{\frac{a+b}{2}}^b k(g(b) - g(t)) g'(t) f(t) dt \\ &\quad + \frac{1}{2} \int_a^{\frac{a+b}{2}} k(g(t) - g(a)) g'(t) f(t) dt. \end{aligned}$$

Using the equalities (3.4) and (3.5), we have

$$\begin{aligned} \check{M}_{k,g,a+,b-f} &= \frac{1}{2} \left[K \left(g(b) - g \left(\frac{a+b}{2} \right) \right) + K \left(g \left(\frac{a+b}{2} \right) - g(a) \right) \right] f \left(\frac{a+b}{2} \right) \\ &\quad + \frac{1}{2} \int_a^{\frac{a+b}{2}} k(g(t) - g(a)) g'(t) \left[f(t) - f \left(\frac{a+b}{2} \right) \right] dt \\ &\quad + \frac{1}{2} \int_{\frac{a+b}{2}}^b k(g(b) - g(t)) g'(t) \left[f(t) - f \left(\frac{a+b}{2} \right) \right] dt \end{aligned}$$

and

$$\begin{aligned} \check{M}_{k,g,a+,b-f} &= \frac{1}{2} \left[K \left(g(b) - g \left(\frac{a+b}{2} \right) \right) f(b) + K \left(g \left(\frac{a+b}{2} \right) - g(a) \right) f(a) \right] \\ &\quad + \frac{1}{2} \int_a^{\frac{a+b}{2}} k(g(t) - g(a)) g'(t) [f(t) - f(a)] dt \\ &\quad + \frac{1}{2} \int_{\frac{a+b}{2}}^b k(g(b) - g(t)) g'(t) [f(t) - f(b)] dt. \end{aligned}$$

Using the g -mean of two numbers we can introduce

$$\begin{aligned} \check{P}_{k,g,a+,b-f} &:= \check{S}_{k,g,a+,b-f} (M_g(a, b)) \\ &= \frac{1}{2} \int_{M_g(a,b)}^b k(g(b) - g(t)) g'(t) f(t) dt \\ &\quad + \frac{1}{2} \int_a^{M_g(a,b)} k(g(t) - g(a)) g'(t) f(t) dt. \end{aligned}$$

Using the equalities (3.4) and (3.5), we have

$$\begin{aligned} \check{P}_{k,g,a+,b-f} &= K \left(\frac{g(b) - g(a)}{2} \right) f(M_g(a, b)) \\ &\quad + \frac{1}{2} \int_a^{M_g(a,b)} k(g(t) - g(a)) g'(t) [f(t) - f(M_g(a, b))] dt \\ &\quad + \frac{1}{2} \int_{M_g(a,b)}^b k(g(b) - g(t)) g'(t) [f(t) - f(M_g(a, b))] dt \end{aligned}$$

and

$$\begin{aligned} \check{P}_{k,g,a+,b-f} &= K \left(\frac{g(b) - g(a)}{2} \right) \frac{f(b) + f(a)}{2} \\ &\quad + \frac{1}{2} \int_a^{M_g(a,b)} k(g(t) - g(a)) g'(t) [f(t) - f(a)] dt \\ &\quad + \frac{1}{2} \int_{M_g(a,b)}^b k(g(b) - g(t)) g'(t) [f(t) - f(b)] dt. \end{aligned}$$

4. Trapezoid functional $T_{k,g,a+,b-}$

We can also introduce the functional

$$\begin{aligned} T_{k,g,a+,b-f} &:= \frac{1}{2} [S_{k,g,a+}f(b) + S_{k,g,b-}f(a)] \\ &= \frac{1}{2} \int_a^b [k(g(b) - g(t)) + k(g(t) - g(a))] g'(t) f(t) dt. \end{aligned}$$

We have:

Lemma 4.1. *With the assumption of Lemma 2.1, we have*

$$\begin{aligned} T_{k,g,a+,b-f} &= K(g(b) - g(a)) \delta \\ &\quad + \frac{1}{2} \int_a^b [k(g(b) - g(t)) + k(g(t) - g(a))] g'(t) [f(t) - \delta] dt \end{aligned} \tag{4.1}$$

for any $\delta \in \mathbb{C}$.

Proof. Observe that

$$\begin{aligned} &\int_a^b [k(g(b) - g(t)) + k(g(t) - g(a))] g'(t) dt \\ &= \int_a^b k(g(b) - g(t)) g'(t) dt + \int_a^b k(g(t) - g(a)) g'(t) dt \\ &= - \int_a^b [K(g(b) - g(t))] dt + \int_a^b [K(g(t) - g(a))] dt \\ &= -K(g(b) - g(t)) \Big|_a^b + K(g(t) - g(a)) \Big|_a^b \\ &= K(g(b) - g(a)) + K(g(b) - g(a)) = 2K(g(b) - g(a)). \end{aligned}$$

Therefore

$$\begin{aligned} &\frac{1}{2} \int_a^b [k(g(b) - g(t)) + k(g(t) - g(a))] g'(t) [f(t) - \delta] dt \\ &= \frac{1}{2} \int_a^b [k(g(b) - g(t)) + k(g(t) - g(a))] g'(t) f(t) dt \\ &\quad - \frac{1}{2} \delta \int_a^b [k(g(b) - g(t)) + k(g(t) - g(a))] g'(t) dt \\ &= T_{k,g,a+,b-f} - \delta K(g(b) - g(a)), \end{aligned}$$

which proves the desired equality (4.1). □

Corollary 4.2. *With the assumptions of Lemma 4.1 we have the Ostrowski type identity*

$$\begin{aligned} T_{k,g,a+,b-f} &= K(g(b) - g(a)) f(x) \\ &\quad + \frac{1}{2} \int_a^b [k(g(b) - g(t)) + k(g(t) - g(a))] g'(t) [f(t) - f(x)] dt \end{aligned} \tag{4.2}$$

for any $x \in [a, b]$ and the trapezoid identity

$$\begin{aligned} T_{k,g,a+,b-f} &= K(g(b) - g(a)) \frac{f(a) + f(b)}{2} \\ &\quad + \frac{1}{2} \int_a^b [k(g(b) - g(t)) + k(g(t) - g(a))] g'(t) \left[f(t) - \frac{f(a) + f(b)}{2} \right] dt. \end{aligned} \tag{4.3}$$

We observe that for $x = \frac{a+b}{2}$ we obtain from (4.2) that

$$\begin{aligned} T_{k,g,a+,b-}f &= K(g(b) - g(a))f\left(\frac{a+b}{2}\right) \\ &+ \frac{1}{2} \int_a^b [k(g(b) - g(t)) + k(g(t) - g(a))]g'(t) \left[f(t) - f\left(\frac{a+b}{2}\right) \right] dt. \end{aligned}$$

5. Inequalities for functions of bounded variation

We considered the cumulative function $K : [0, \infty) \rightarrow \mathbb{C}$ by

$$K(t) := \begin{cases} \int_0^t k(s) ds & \text{if } 0 < t, \\ 0 & \text{if } t = 0. \end{cases}$$

We also define the function $\mathbf{K} : [0, \infty) \rightarrow [0, \infty)$ by

$$\mathbf{K}(t) := \begin{cases} \int_0^t |k(s)| ds & \text{if } 0 < t, \\ 0 & \text{if } t = 0. \end{cases}$$

We observe that if k takes nonnegative values on $(0, \infty)$, as it does in some of the examples in Introduction, then $\mathbf{K}(t) = K(t)$ for $t \in [0, \infty)$.

Theorem 5.1. *Assume that the kernel k is defined either on $(0, \infty)$ or on $[0, \infty)$ with complex values and integrable on any finite subinterval. Let $f : [a, b] \rightarrow \mathbb{C}$ be a function of bounded variation on $[a, b]$ and g be a strictly increasing function on (a, b) , having a continuous derivative g' on (a, b) . Then we have the Ostrowski type inequality*

$$\begin{aligned} & \left| S_{k,g,a+,b-}f(x) - \frac{1}{2} [K(g(b) - g(x)) + K(g(x) - g(a))]f(x) \right| \\ & \leq \frac{1}{2} \left[\int_x^b |k(g(t) - g(x))| \bigvee_x^t(f) g'(t) dt + \int_a^x |k(g(x) - g(t))| \bigvee_t^x(f) g'(t) dt \right] \\ & \leq \frac{1}{2} \left[\mathbf{K}(g(b) - g(x)) \bigvee_x^b(f) + \mathbf{K}(g(x) - g(a)) \bigvee_a^x(f) \right] \\ & \leq \frac{1}{2} \begin{cases} \max \{ \mathbf{K}(g(b) - g(x)), \mathbf{K}(g(x) - g(a)) \} \bigvee_a^b(f); \\ [\mathbf{K}^p(g(b) - g(x)) + \mathbf{K}^p(g(x) - g(a))]^{1/p} \left((\bigvee_a^x(f))^q + (\bigvee_x^b(f))^q \right)^{1/q} \\ \text{with } p, q > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ [\mathbf{K}(g(b) - g(x)) + \mathbf{K}(g(x) - g(a))] \left[\frac{1}{2} \bigvee_a^b(f) + \frac{1}{2} \left| \bigvee_a^x(f) - \bigvee_x^b(f) \right| \right] \end{cases} \quad (5.1) \end{aligned}$$

and the trapezoid type inequality

$$\begin{aligned}
 & \left| S_{k,g,a+,b-} f(x) - \frac{1}{2} [K(g(b) - g(x))f(b) + K(g(x) - g(a))f(a)] \right| \\
 & \leq \frac{1}{2} \left[\int_a^x |k(g(x) - g(t))| \bigvee_a^t(f) g'(t) dt + \int_x^b |k(g(t) - g(x))| \bigvee_t^b(f) g'(t) dt \right] \\
 & \leq \frac{1}{2} \left[\mathbf{K}(g(b) - g(x)) \bigvee_x^b(f) + \mathbf{K}(g(x) - g(a)) \bigvee_a^x(f) \right] \\
 & \leq \frac{1}{2} \begin{cases} \max \{ \mathbf{K}(g(b) - g(x)), \mathbf{K}(g(x) - g(a)) \} \bigvee_a^b(f); \\ [\mathbf{K}^p(g(b) - g(x)) + \mathbf{K}^p(g(x) - g(a))]^{1/p} \\ \times \left((\bigvee_a^x(f))^q + (\bigvee_x^b(f))^q \right)^{1/q} \\ \text{with } p, q > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ [\mathbf{K}(g(b) - g(x)) + \mathbf{K}(g(x) - g(a))] \\ \times \left[\frac{1}{2} \bigvee_a^b(f) + \frac{1}{2} \left| \bigvee_a^x(f) - \bigvee_x^b(f) \right| \right] \end{cases} \tag{5.2}
 \end{aligned}$$

for any $x \in (a, b)$.

Proof. Using the equality (2.4) we have

$$\begin{aligned}
 & \left| S_{k,g,a+,b-} f(x) - \frac{1}{2} [K(g(b) - g(x)) + K(g(x) - g(a))] f(x) \right| \\
 & \leq \frac{1}{2} \left| \int_a^x k(g(x) - g(t)) g'(t) [f(t) - f(x)] dt \right| \\
 & + \frac{1}{2} \left| \int_x^b k(g(t) - g(x)) g'(t) [f(t) - f(x)] dt \right| \\
 & \leq \frac{1}{2} \int_a^x |k(g(x) - g(t)) g'(t) [f(t) - f(x)]| dt \\
 & + \frac{1}{2} \int_x^b |k(g(t) - g(x)) g'(t) [f(t) - f(x)]| dt \\
 & = \frac{1}{2} \int_a^x |k(g(x) - g(t))| |f(x) - f(t)| g'(t) dt \\
 & + \frac{1}{2} \int_x^b |k(g(t) - g(x))| |f(t) - f(x)| g'(t) dt \\
 & =: B(x)
 \end{aligned}$$

for $x \in (a, b)$.

Since f is of bounded variation, then

$$|f(x) - f(t)| \leq \bigvee_t^x(f) \leq \bigvee_a^x(f) \text{ for } a < t \leq x \leq b$$

and

$$|f(t) - f(x)| \leq \bigvee_x^t(f) \leq \bigvee_x^b(f) \text{ for } a \leq x \leq t < b.$$

Therefore

$$\begin{aligned} B(x) &\leq \frac{1}{2} \int_a^x |k(g(x) - g(t))| \bigvee_t^x(f) g'(t) dt \\ &\quad + \frac{1}{2} \int_x^b |k(g(t) - g(x))| \bigvee_x^t(f) g'(t) dt \\ &\leq \frac{1}{2} \bigvee_a^x(f) \int_a^x |k(g(x) - g(t))| g'(t) dt \\ &\quad + \frac{1}{2} \bigvee_x^b(f) \int_x^b |k(g(t) - g(x))| g'(t) dt \\ &=: C(x) \end{aligned}$$

for $x \in (a, b)$.

We have, by taking the derivative over t and using the chain rule, that

$$[\mathbf{K}(g(x) - g(t))] = \mathbf{K}'(g(x) - g(t)) (g(x) - g(t))' = -|k(g(x) - g(t))| g'(t)$$

for $t \in (a, x)$ and

$$[\mathbf{K}(g(t) - g(x))] = \mathbf{K}'(g(t) - g(x)) (g(t) - g(x))' = |k(g(t) - g(x))| g'(t)$$

for $t \in (x, b)$.

Then

$$\int_a^x |k(g(x) - g(t))| g'(t) dt = - \int_a^x [\mathbf{K}(g(x) - g(t))] dt = \mathbf{K}(g(x) - g(a))$$

and

$$\int_x^b |k(g(t) - g(x))| g'(t) dt = \int_x^b [\mathbf{K}(g(t) - g(x))] dt = \mathbf{K}(g(b) - g(x))$$

giving that

$$C(x) = \frac{1}{2} \left[\mathbf{K}(g(b) - g(x)) \bigvee_x^b(f) + \mathbf{K}(g(x) - g(a)) \bigvee_a^x(f) \right],$$

for $x \in (a, b)$, which proves the first and the second inequality in (5.1).

The last part of (4.1) is obvious by making use of the elementary Hölder type inequalities for positive real numbers $c, d, m, n \geq 0$

$$mc + nd \leq \begin{cases} \max\{m, n\}(c + d); \\ (m^p + n^p)^{1/p} (c^q + d^q)^{1/q} \text{ with } p, q > 1, \frac{1}{p} + \frac{1}{q} = 1. \end{cases}$$

Further, by the identity (2.5) we have, as above,

$$\begin{aligned}
 & \left| S_{k,g,a+,b-} f(x) - \frac{1}{2} [K(g(b) - g(x))f(b) + K(g(x) - g(a))f(a)] \right| \\
 & \leq \frac{1}{2} \int_a^x |k(g(x) - g(t))| |f(t) - f(a)| g'(t) dt \\
 & \quad + \frac{1}{2} \int_x^b |k(g(t) - g(x))| |f(t) - f(b)| g'(t) dt \\
 & \leq \frac{1}{2} \int_a^x |k(g(x) - g(t))| \bigvee_a^t (f) g'(t) dt \\
 & \quad + \frac{1}{2} \int_x^b |k(g(t) - g(x))| \bigvee_t^b (f) g'(t) dt \\
 & \leq \frac{1}{2} \bigvee_a^x (f) \int_a^x |k(g(x) - g(t))| g'(t) dt \\
 & \quad + \frac{1}{2} \bigvee_x^b (f) \int_x^b |k(g(t) - g(x))| g'(t) dt \\
 & = \frac{1}{2} \mathbf{K}(g(x) - g(a)) \bigvee_a^x (f) + \frac{1}{2} \mathbf{K}(g(b) - g(x)) \bigvee_x^b (f),
 \end{aligned}$$

which proves (5.2). □

The following particular case for the functional

$$\begin{aligned}
 P_{k,g,a+,b-} f & := S_{k,g,a+,b-} f(M_g(a,b)) \\
 & = \frac{1}{2} \int_a^{M_g(a,b)} k\left(\frac{g(b) + g(a)}{2} - g(t)\right) g'(t) f(t) dt \\
 & \quad + \frac{1}{2} \int_{M_g(a,b)}^b k\left(g(t) - \frac{g(b) + g(a)}{2}\right) g'(t) f(t) dt.
 \end{aligned}$$

is of interest:

Corollary 5.2. *With the assumptions of Theorem 5.1 we have*

$$\begin{aligned}
 \left| P_{k,g,a+,b-} f - K\left(\frac{g(b) - g(a)}{2}\right) f(M_g(a,b)) \right| & \leq \frac{1}{2} \int_{M_g(a,b)}^b \left| k\left(g(t) - \frac{g(b) + g(a)}{2}\right) \right| \bigvee_{M_g(a,b)}^t (f) g'(t) dt \\
 & \quad + \frac{1}{2} \int_a^{M_g(a,b)} \left| k\left(\frac{g(b) + g(a)}{2} - g(t)\right) \right| \bigvee_t^{M_g(a,b)} (f) g'(t) dt \\
 & \leq \frac{1}{2} \mathbf{K}\left(\frac{g(b) - g(a)}{2}\right) \bigvee_b^b (f)
 \end{aligned} \tag{5.3}$$

and

$$\begin{aligned}
 \left| P_{k,g,a+,b-} f - K\left(\frac{g(b) - g(a)}{2}\right) \frac{f(b) + f(a)}{2} \right| & \leq \frac{1}{2} \int_a^{M_g(a,b)} \left| k\left(\frac{g(b) + g(a)}{2} - g(t)\right) \right| \bigvee_a^t (f) g'(t) dt \\
 & \quad + \frac{1}{2} \int_{M_g(a,b)}^b \left| k\left(g(t) - \frac{g(b) + g(a)}{2}\right) \right| \bigvee_t^b (f) g'(t) dt \\
 & \leq \frac{1}{2} \mathbf{K}\left(\frac{g(b) - g(a)}{2}\right) \bigvee_b^b (f).
 \end{aligned} \tag{5.4}$$

We have:

Theorem 5.3. *With the assumptions of Theorem 5.1 we have the Ostrowski type inequality*

$$\begin{aligned}
 & \left| \check{S}_{k,g,a+,b-} f(x) - \frac{1}{2} [\mathbf{K}(g(b) - g(x)) + \mathbf{K}(g(x) - g(a))] f(x) \right| \\
 & \leq \frac{1}{2} \int_a^x |k(g(t) - g(a))| \bigvee_a^t(f) g'(t) dt + \frac{1}{2} \int_x^b |k(g(b) - g(t))| \bigvee_x^t(f) g'(t) dt \\
 & \leq \frac{1}{2} \left[\mathbf{K}(g(b) - g(x)) \bigvee_x^b(f) + \mathbf{K}(g(x) - g(a)) \bigvee_a^x(f) \right] \\
 & \leq \frac{1}{2} \begin{cases} \max \{ \mathbf{K}(g(b) - g(x)), \mathbf{K}(g(x) - g(a)) \} \mathcal{V}_a^b(f); \\ [\mathbf{K}^p(g(b) - g(x)) + \mathbf{K}^p(g(x) - g(a))]^{1/p} \left((\mathcal{V}_a^x(f))^q + (\mathcal{V}_x^b(f))^q \right)^{1/q} \\ \text{with } p, q > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ [\mathbf{K}(g(b) - g(x)) + \mathbf{K}(g(x) - g(a))] \left[\frac{1}{2} \mathcal{V}_a^b(f) + \frac{1}{2} \left| \mathcal{V}_a^x(f) - \mathcal{V}_x^b(f) \right| \right] \end{cases} \quad (5.5)
 \end{aligned}$$

and the trapezoid inequality

$$\begin{aligned}
 & \left| \check{S}_{k,g,a+,b-} f(x) - \frac{1}{2} [\mathbf{K}(g(b) - g(x)) f(b) + \mathbf{K}(g(x) - g(a)) f(a)] \right| \\
 & \leq \frac{1}{2} \int_a^x |k(g(t) - g(a))| \bigvee_a^t(f) g'(t) dt + \frac{1}{2} \int_x^b |k(g(b) - g(t))| \bigvee_x^t(f) g'(t) dt \\
 & \leq \frac{1}{2} \left[\mathbf{K}(g(b) - g(x)) \bigvee_x^b(f) + \mathbf{K}(g(x) - g(a)) \bigvee_a^x(f) \right] \\
 & \leq \frac{1}{2} \begin{cases} \max \{ \mathbf{K}(g(b) - g(x)), \mathbf{K}(g(x) - g(a)) \} \mathcal{V}_a^b(f); \\ [\mathbf{K}^p(g(b) - g(x)) + \mathbf{K}^p(g(x) - g(a))]^{1/p} \left((\mathcal{V}_a^x(f))^q + (\mathcal{V}_x^b(f))^q \right)^{1/q} \\ \text{with } p, q > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ [\mathbf{K}(g(b) - g(x)) + \mathbf{K}(g(x) - g(a))] \left[\frac{1}{2} \mathcal{V}_a^b(f) + \frac{1}{2} \left| \mathcal{V}_a^x(f) - \mathcal{V}_x^b(f) \right| \right] \end{cases} \quad (5.6)
 \end{aligned}$$

for any $x \in (a, b)$.

Proof. Using the identity (3.4) we have

$$\begin{aligned}
 & \left| \check{S}_{k,g,a+,b-} f(x) - \frac{1}{2} [K(g(b) - g(x)) + K(g(x) - g(a))] f(x) \right| \\
 & \leq \frac{1}{2} \int_a^x |k(g(t) - g(a))| |f(t) - f(x)| g'(t) dt \\
 & \quad + \frac{1}{2} \int_x^b |k(g(b) - g(t))| |f(t) - f(x)| g'(t) dt \\
 & \leq \frac{1}{2} \int_a^x |k(g(t) - g(a))| \overset{x}{\underset{t}{V}}(f) g'(t) dt \\
 & \quad + \frac{1}{2} \int_x^b |k(g(b) - g(t))| \overset{t}{\underset{x}{V}}(f) g'(t) dt \\
 & \leq \frac{1}{2} \overset{x}{\underset{a}{V}}(f) \int_a^x |k(g(t) - g(a))| g'(t) dt \\
 & \quad + \frac{1}{2} \overset{b}{\underset{x}{V}}(f) \int_x^b |k(g(b) - g(t))| g'(t) dt \\
 & = \frac{1}{2} \left[\mathbf{K}(g(x) - g(a)) \overset{x}{\underset{a}{V}}(f) + \mathbf{K}(g(b) - g(x)) \overset{b}{\underset{x}{V}}(f) \right],
 \end{aligned}$$

for any $x \in (a, b)$, which proves (5.5).

By the identity (3.5) we have

$$\begin{aligned}
 & \left| \check{S}_{k,g,a+,b-} f(x) - \frac{1}{2} [K(g(b) - g(x)) f(b) + K(g(x) - g(a)) f(a)] \right| \\
 & \leq \frac{1}{2} \int_a^x |k(g(t) - g(a))| |f(t) - f(a)| g'(t) dt \\
 & \quad + \frac{1}{2} \int_x^b |k(g(b) - g(t))| |f(b) - f(t)| g'(t) dt \\
 & \leq \frac{1}{2} \int_a^x |k(g(t) - g(a))| \overset{t}{\underset{a}{V}}(f) g'(t) dt \\
 & \quad + \frac{1}{2} \int_x^b |k(g(b) - g(t))| \overset{b}{\underset{t}{V}}(f) g'(t) dt \\
 & \leq \frac{1}{2} \overset{x}{\underset{a}{V}}(f) \int_a^x |k(g(t) - g(a))| g'(t) dt \\
 & \quad + \frac{1}{2} \overset{b}{\underset{x}{V}}(f) \int_x^b |k(g(b) - g(t))| g'(t) dt \\
 & = \frac{1}{2} \left[\mathbf{K}(g(x) - g(a)) \overset{x}{\underset{a}{V}}(f) + \mathbf{K}(g(b) - g(x)) \overset{b}{\underset{x}{V}}(f) \right]
 \end{aligned}$$

for any $x \in (a, b)$, which proves (5.6). □

Also, we have the particular inequalities for

$$\begin{aligned}
 \check{P}_{k,g,a+,b-} f & := \check{S}_{k,g,a+,b-} f(M_g(a, b)) \\
 & = \frac{1}{2} \int_{M_g(a,b)}^b k(g(b) - g(t)) g'(t) f(t) dt \\
 & \quad + \frac{1}{2} \int_a^{M_g(a,b)} k(g(t) - g(a)) g'(t) f(t) dt.
 \end{aligned}$$

Corollary 5.4. *With the assumptions of Theorem 5.1 we have*

$$\begin{aligned} \left| \check{P}_{k,g,a+,b-}f - K \left(\frac{g(b) - g(a)}{2} \right) \frac{f(b) + f(a)}{2} \right| &\leq \frac{1}{2} \int_a^{M_g(a,b)} |k(g(t) - g(a))| \bigvee_t^{M_g(a,b)} (f) g'(t) dt \\ &\quad + \frac{1}{2} \int_{M_g(a,b)}^b |k(g(b) - g(t))| \bigvee_{M_g(a,b)}^t (f) g'(t) dt \\ &\leq \frac{1}{2} \mathbf{K} \left(\frac{g(b) - g(a)}{2} \right) \bigvee_b^b (f) \end{aligned}$$

and

$$\begin{aligned} \left| \check{P}_{k,g,a+,b-}f - K \left(\frac{g(b) - g(a)}{2} \right) \frac{f(b) + f(a)}{2} \right| &\leq \frac{1}{2} \int_a^{M_g(a,b)} |k(g(t) - g(a))| \bigvee_a^t (f) g'(t) dt \\ &\quad + \frac{1}{2} \int_{M_g(a,b)}^b |k(g(b) - g(t))| \bigvee_t^b (f) g'(t) dt \\ &\leq \frac{1}{2} \mathbf{K} \left(\frac{g(b) - g(a)}{2} \right) \bigvee_b^b (f). \end{aligned}$$

Finally, we have the following result for the trapezoid functional

$$\begin{aligned} T_{k,g,a+,b-}f &:= \frac{1}{2} [S_{k,g,a+}f(b) + S_{k,g,b-}f(a)] \\ &= \frac{1}{2} \int_a^b [k(g(b) - g(t)) + k(g(t) - g(a))] g'(t) f(t) dt. \end{aligned}$$

Theorem 5.5. *With the assumptions of Theorem 5.1 we have the trapezoid type inequality*

$$\left| T_{k,g,a+,b-}f - K(g(b) - g(a)) \frac{f(a) + f(b)}{2} \right| \leq \frac{1}{2} \mathbf{K}(g(b) - g(a)) \bigvee_a^b (f). \quad (5.7)$$

Proof. From the identity (4.3) we have

$$\begin{aligned} &\left| T_{k,g,a+,b-}f - K(g(b) - g(a)) \frac{f(a) + f(b)}{2} \right| \\ &\leq \frac{1}{2} \int_a^b |k(g(b) - g(t)) + k(g(t) - g(a))| \left| f(t) - \frac{f(a) + f(b)}{2} \right| g'(t) dt \\ &\leq \frac{1}{2} \int_a^b [|k(g(b) - g(t))| + |k(g(t) - g(a))|] \left| f(t) - \frac{f(a) + f(b)}{2} \right| g'(t) dt \\ &=: D. \end{aligned}$$

Since $f : [a, b] \rightarrow \mathbb{C}$ is of bounded variation, then for any $t \in [a, b]$ we have

$$\begin{aligned} \left| f(t) - \frac{f(a) + f(b)}{2} \right| &= \left| \frac{f(t) - f(a) + f(t) - f(b)}{2} \right| \\ &\leq \frac{1}{2} [|f(t) - f(a)| + |f(b) - f(t)|] \leq \frac{1}{2} \bigvee_a^b (f). \end{aligned}$$

Therefore

$$\begin{aligned} D &\leq \frac{1}{4} \bigvee_a^b (f) \int_a^b [|k(g(b) - g(t))| + |k(g(t) - g(a))|] g'(t) dt \\ &= \frac{1}{4} \bigvee_a^b (f) [\mathbf{K}(g(b) - g(a)) + \mathbf{K}(g(b) - g(a))] = \frac{1}{2} \mathbf{K}(g(b) - g(a)) \bigvee_a^b (f), \end{aligned}$$

which proves the desired result (5.7). □

6. Example for an exponential kernel

The above inequalities may be written for all the particular fractional integrals introduced in the introduction.

If we take, for instance $k(t) = \frac{1}{\Gamma(\alpha)} t^{\alpha-1}$, where Γ is the *Gamma function*, then we recapture the results for the *generalized left- and right-sided Riemann-Liouville fractional integrals* of a function f with respect to another function g on $[a, b]$ as outlined in [5].

For $\alpha, \beta \in \mathbb{R}$ we consider the kernel $k(t) := \exp[(\alpha + \beta i)t]$, $t \in \mathbb{R}$. We have

$$K(t) = \frac{\exp[(\alpha + \beta i)t] - 1}{(\alpha + \beta i)}, \text{ if } t \in \mathbb{R}$$

for $\alpha, \beta \neq 0$.

Also, we have

$$|k(s)| := |\exp[(\alpha + \beta i)s]| = \exp(\alpha s) \text{ for } s \in \mathbb{R}$$

and

$$\mathbf{K}(t) = \int_0^t \exp(\alpha s) ds = \frac{\exp(\alpha t) - 1}{\alpha} \text{ if } 0 < t,$$

for $\alpha \neq 0$.

Let $f : [a, b] \rightarrow \mathbb{C}$ be a function of bounded variation on $[a, b]$ and g be a strictly increasing function on (a, b) , having a continuous derivative g' on (a, b) . We have

$$\begin{aligned} \mathcal{E}_{g, a+, b-}^{\alpha + \beta i} f(x) &= \frac{1}{2} \int_a^x \exp[(\alpha + \beta i)(g(x) - g(t))] g'(t) f(t) dt \\ &\quad + \frac{1}{2} \int_x^b \exp[(\alpha + \beta i)(g(t) - g(x))] g'(t) f(t) dt \end{aligned}$$

for $x \in (a, b)$.

If $g = \ln h$ where $h : [a, b] \rightarrow (0, \infty)$ is a strictly increasing function on (a, b) , having a continuous derivative h' on (a, b) , then we can consider the following operator as well

$$\begin{aligned} &\mathbf{K}_{h, a+, b-}^{\alpha + \beta i} f(x) \\ &:= \mathcal{E}_{\ln h, a+, b-}^{\alpha + \beta i} f(x) \\ &= \frac{1}{2} \left[\int_a^x \left(\frac{h(x)}{h(t)} \right)^{\alpha + \beta i} \frac{h'(t)}{h(t)} f(t) dt + \int_x^b \left(\frac{h(t)}{h(x)} \right)^{\alpha + \beta i} \frac{h'(t)}{h(t)} f(t) dt \right], \end{aligned}$$

for $x \in (a, b)$.

By using the inequality (5.1) we have for $x \in (a, b)$ that

$$\begin{aligned}
 & \left| \mathcal{E}_{g,a+,b-}^{\alpha+\beta i} f(x) - \frac{1}{2} \left[\frac{\exp[(\alpha+\beta i)(g(b)-g(x))] - 1 + \exp[(\alpha+\beta i)(g(x)-g(a))] - 1}{(\alpha+\beta i)} \right] f(x) \right| \\
 & \leq \frac{1}{2} \left[\int_x^b \exp(\alpha(g(t)-g(x))) g'(t) \mathcal{V}_x^t(f) dt + \int_a^x \exp(\alpha(g(x)-g(t))) g'(t) \mathcal{V}_t^x(f) dt \right] \\
 & \leq \frac{1}{2} \left[\frac{\exp(\alpha(g(b)-g(x))) - 1}{\alpha} \mathcal{V}_x^b(f) + \frac{\exp(\alpha(g(x)-g(a))) - 1}{\alpha} \mathcal{V}_a^x(f) \right] \\
 & \leq \frac{1}{2} \begin{cases} \max \left\{ \frac{\exp(\alpha(g(b)-g(x))) - 1}{\alpha}, \frac{\exp(\alpha(g(x)-g(a))) - 1}{\alpha} \right\} \mathcal{V}_a^b(f); \\ \left[\left(\frac{\exp(\alpha(g(b)-g(x))) - 1}{\alpha} \right)^p + \left(\frac{\exp(\alpha(g(x)-g(a))) - 1}{\alpha} \right)^p \right]^{1/p} \left((\mathcal{V}_a^x(f))^q + (\mathcal{V}_x^b(f))^q \right)^{1/q} \\ \text{with } p, q > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \left[\frac{\exp(\alpha(g(b)-g(x))) - 1 + \exp(\alpha(g(x)-g(a))) - 1}{\alpha} \right] \left[\frac{1}{2} \mathcal{V}_a^b(f) + \frac{1}{2} \left| \mathcal{V}_a^x(f) - \mathcal{V}_x^b(f) \right| \right] \end{cases}
 \end{aligned}$$

for $\alpha, \beta \in \mathbb{R}$ with $\alpha \neq 0$.

By using the inequality (5.2) we also have for $x \in (a, b)$ that

$$\begin{aligned}
 & \left| \mathcal{E}_{g,a+,b-}^{\alpha+\beta i} f(x) - \frac{1}{2} \left[\frac{(\exp[(\alpha+\beta i)(g(b)-g(x)] - 1) f(b) + (\exp[(\alpha+\beta i)(g(x)-g(a)] - 1) f(a))}{(\alpha+\beta i)} \right] \right| \\
 & \leq \frac{1}{2} \left[\int_a^x \exp(\alpha(g(t)-g(x))) g'(t) \mathcal{V}_a^t(f) dt + \int_x^b \exp(\alpha(g(x)-g(t))) g'(t) \mathcal{V}_t^b(f) dt \right] \\
 & \leq \frac{1}{2} \left[\frac{\exp(\alpha(g(b)-g(x))) - 1}{\alpha} \mathcal{V}_x^b(f) + \frac{\exp(\alpha(g(x)-g(a))) - 1}{\alpha} \mathcal{V}_a^x(f) \right] \\
 & \leq \frac{1}{2} \begin{cases} \max \left\{ \frac{\exp(\alpha(g(b)-g(x))) - 1}{\alpha}, \frac{\exp(\alpha(g(x)-g(a))) - 1}{\alpha} \right\} \mathcal{V}_a^b(f); \\ \left[\left(\frac{\exp(\alpha(g(b)-g(x))) - 1}{\alpha} \right)^p + \left(\frac{\exp(\alpha(g(x)-g(a))) - 1}{\alpha} \right)^p \right]^{1/p} \\ \times \left((\mathcal{V}_a^x(f))^q + (\mathcal{V}_x^b(f))^q \right)^{1/q} \\ \text{with } p, q > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \left[\frac{\exp(\alpha(g(b)-g(x))) - 1 + \exp(\alpha(g(x)-g(a))) - 1}{\alpha} \right] \\ \times \left[\frac{1}{2} \mathcal{V}_a^b(f) + \frac{1}{2} \left| \mathcal{V}_a^x(f) - \mathcal{V}_x^b(f) \right| \right] \end{cases}
 \end{aligned}$$

for $\alpha, \beta \in \mathbb{R}$ with $\alpha \neq 0$.

If we denote

$$\begin{aligned}
 \overline{\mathcal{E}}_{g,a+,b-}^{\alpha+\beta i} f & := \mathcal{E}_{g,a+,b-}^{\alpha+\beta i} f(M_g(a, b)) \\
 & = \frac{1}{2} \int_a^x \exp \left[(\alpha + \beta i) \left(\frac{g(b) + g(a)}{2} - g(t) \right) \right] g'(t) f(t) dt \\
 & \quad + \frac{1}{2} \int_x^b \exp \left[(\alpha + \beta i) \left(g(t) - \frac{g(b) + g(a)}{2} \right) \right] g'(t) f(t) dt
 \end{aligned}$$

then by (5.3) and (5.4) we have the simpler results

$$\begin{aligned} \left| \bar{\mathcal{E}}_{g,a+,b-}^{\alpha+\beta i} f - \frac{\exp\left[(\alpha+\beta i)\frac{g(b)-g(a)}{2}\right]-1}{(\alpha+\beta i)} f(M_g(a,b)) \right| &\leq \frac{1}{2} \int_{M_g(a,b)}^b \exp\left(\alpha\left(g(t) - \frac{g(b)+g(a)}{2}\right)\right) g'(t) \bigvee_{M_g(a,b)}^t(f) dt \\ &\quad + \frac{1}{2} \int_a^{M_g(a,b)} \exp\left(\alpha\left(\frac{g(b)+g(a)}{2} - g(t)\right)\right) g'(t) \bigvee_t^{M_g(a,b)}(f) dt \\ &\leq \frac{1}{2} \frac{\exp\left(\alpha\left(\frac{g(b)-g(a)}{2}\right)\right)-1}{\alpha} \bigvee_b^b(f) \end{aligned} \quad (6.1)$$

and

$$\begin{aligned} \left| \bar{\mathcal{E}}_{g,a+,b-}^{\alpha+\beta i} f - \frac{\exp\left[(\alpha+\beta i)\frac{g(b)-g(a)}{2}\right]-1}{(\alpha+\beta i)} \frac{f(b)+f(a)}{2} \right| &\leq \frac{1}{2} \int_a^{M_g(a,b)} \exp\left(\alpha\left(g(t) - \frac{g(b)+g(a)}{2}\right)\right) g'(t) \bigvee_a^t(f) dt \\ &\quad + \frac{1}{2} \int_{M_g(a,b)}^b \exp\left(\alpha\left(\frac{g(b)+g(a)}{2} - g(t)\right)\right) g'(t) \bigvee_t^b(f) dt \\ &\leq \frac{1}{2} \frac{\exp\left(\alpha\left(\frac{g(b)-g(a)}{2}\right)\right)-1}{\alpha} \bigvee_b^b(f). \end{aligned} \quad (6.2)$$

In particular, if we take in (6.1) and (6.2) $g = \ln t$, $t \in [a, b] \subset (0, \infty)$, then by using the notation $G(\gamma, \delta) := \sqrt{\gamma\delta}$ for the geometric mean of the positive real numbers $\gamma, \delta > 0$ we have

$$\begin{aligned} \left| \bar{\mathcal{K}}_{a+,b-}^{\alpha+\beta i} f - \frac{\left(\frac{b}{a}\right)^{\alpha+\beta i} - 1}{(\alpha+\beta i)} f(G(a,b)) \right| &\leq \frac{1}{2} \int_{G(a,b)}^b \left(\frac{t}{G(a,b)}\right)^\alpha \frac{1}{t} \bigvee_{G(a,b)}^t(f) dt + \frac{1}{2} \int_a^{G(a,b)} \left(\frac{G(a,b)}{t}\right)^\alpha \frac{1}{t} \bigvee_t^{G(a,b)}(f) dt \\ &\leq \frac{1}{2} \frac{\left(\frac{b}{a}\right)^\alpha - 1}{\alpha} \bigvee_b^b(f) \end{aligned}$$

and

$$\begin{aligned} \left| \bar{\mathcal{K}}_{a+,b-}^{\alpha+\beta i} f - \frac{\left(\frac{b}{a}\right)^{\alpha+\beta i} - 1}{(\alpha+\beta i)} \frac{f(b)+f(a)}{2} \right| &\leq \frac{1}{2} \int_{G(a,b)}^b \left(\frac{G(a,b)}{t}\right)^\alpha \frac{1}{t} \bigvee_t^b(f) dt + \frac{1}{2} \int_a^{G(a,b)} \left(\frac{t}{G(a,b)}\right)^\alpha \frac{1}{t} \bigvee_a^t(f) dt \\ &\leq \frac{1}{2} \frac{\left(\frac{b}{a}\right)^\alpha - 1}{\alpha} \bigvee_b^b(f), \end{aligned}$$

where

$$\bar{\mathcal{K}}_{a+,b-}^{\alpha+\beta i} f := \frac{1}{2} \int_{G(a,b)}^b \left(\frac{t}{G(a,b)}\right)^{\alpha+\beta i} \frac{1}{t} f(t) dt + \frac{1}{2} \int_a^{G(a,b)} \left(\frac{G(a,b)}{t}\right)^{\alpha+\beta i} \frac{1}{t} f(t) dt.$$

References

- [1] A. Kilbas, H. M. Srivastava, J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematics Studies, Elsevier Science B.V., Amsterdam, 2006.
- [2] R. K. Raina, *On generalized Wright's hypergeometric functions and fractional calculus operators*, East Asian Math. J., **21** (2) (2005), 191-203.
- [3] R. P. Agarwal, M.-J. Luo, R. K. Raina, *On Ostrowski type inequalities*, Fasc. Math., **56** (2016), 5-27.
- [4] M. Kirane, B. T. Torebek, *Hermite-Hadamard, Hermite-Hadamard-Fejer, Dragomir-Agarwal and Pachpatte type inequalities for convex functions via fractional integrals*, Preprint arXiv:1701.00092.

- [5] S. S. Dragomir, *Further Ostrowski and trapezoid type inequalities for the generalized Riemann-Liouville fractional integrals of functions with bounded variation*, RGMIA Res. Rep. Coll., **20** (2017), Art. 84.
- [6] A. Aglič Aljinović, *Montgomery identity and Ostrowski type inequalities for Riemann-Liouville fractional integral*, J. Math., **2014**, Art. ID 503195, 6 pp.
- [7] T. M. Apostol, *Mathematical Analysis*, Second Edition, Addison-Wesley Publishing Company, 1975.
- [8] A. O. Akdemir, *Inequalities of Ostrowski's type for m - and (α, m) -logarithmically convex functions via Riemann-Liouville fractional integrals*, J. Comput. Anal. Appl., **16**(2) (2014), 375–383.
- [9] G. A. Anastassiou, *Fractional representation formulae under initial conditions and fractional Ostrowski type inequalities*, Demonstr. Math., **48**(3) (2015), 357–378.
- [10] G. A. Anastassiou, *The reduction method in fractional calculus and fractional Ostrowski type inequalities*, Indian J. Math., **56**(3) (2014), 333-357.
- [11] H. Budak, M. Z. Sarikaya, E. Set, *Generalized Ostrowski type inequalities for functions whose local fractional derivatives are generalized s -convex in the second sense*, J. Appl. Math. Comput. Mech., **15**(4) (2016), 11–21.
- [12] P. Cerone, S. S. Dragomir, *Midpoint-type rules from an inequalities point of view. Handbook of analytic-computational methods in applied mathematics*, 135–200, Chapman & Hall/CRC, Boca Raton, FL, 2000.
- [13] S. S. Dragomir, *The Ostrowski's integral inequality for Lipschitzian mappings and applications*, Comput. Math. Appl., **38**(11-12) (1999), 33-37.
- [14] S. S. Dragomir, *The Ostrowski integral inequality for mappings of bounded variation*, Bull. Austral. Math. Soc., **60**(3) (1999), 495-508.
- [15] S. S. Dragomir, *On the midpoint quadrature formula for mappings with bounded variation and applications*, Kragujevac J. Math., **22** (2000), 13–19.
- [16] S. S. Dragomir, *On the Ostrowski's integral inequality for mappings with bounded variation and applications*, Math. Ineq. Appl. **4** (2001), No. 1, 59-66. Preprint: RGMIA Res. Rep. Coll., **2** (1999), Art. 7.
- [17] S. S. Dragomir, *Refinements of the generalised trapezoid and Ostrowski inequalities for functions of bounded variation*, Arch. Math., **91**(5) (2008), 450–460.
- [18] S. S. Dragomir, *Refinements of the Ostrowski inequality in terms of the cumulative variation and applications*, Analysis (Berlin) **34** (2014), No. 2, 223–240. Preprint: RGMIA Res. Rep. Coll., **16** (2013), Art. 29.
- [19] S. S. Dragomir, *Ostrowski type inequalities for Lebesgue integral: a survey of recent results*, Australian J. Math. Anal. Appl., **14**(1) (2017), 1-287.
- [20] S. S. Dragomir, *Ostrowski type inequalities for Riemann-Liouville fractional integrals of bounded variation, Hölder and Lipschitzian functions*, Preprint RGMIA Res. Rep. Coll., **20** (2017), Art. 48.
- [21] S. S. Dragomir, *Ostrowski type inequalities for generalized Riemann-Liouville fractional integrals of functions with bounded variation*, RGMIA Res. Rep. Coll., **20** (2017), Art. 58.
- [22] A. Guezane-Lakoud, F. Aissaoui, *New fractional inequalities of Ostrowski type*, Transylv. J. Math. Mech., **5**(2) (2013), 103–106
- [23] A. Kashuri, R. Liko, *Ostrowski type fractional integral inequalities for generalized (s, m, φ) -preinvex functions*, Aust. J. Math. Anal. Appl., **13**(1) (2016), Art. 16, 11 pp.
- [24] M. A. Noor, K. I. Noor, S. Iftikhar, *Fractional Ostrowski inequalities for harmonic h -preinvex functions*, Facta Univ. Ser. Math. Inform., **31**(2) (2016), 417–445.
- [25] M. Z. Sarikaya, H. Filiz, *Note on the Ostrowski type inequalities for fractional integrals*, Vietnam J. Math., **42**(2) (2014), 187–190.
- [26] M. Z. Sarikaya, H. Budak, *Generalized Ostrowski type inequalities for local fractional integrals*, Proc. Amer. Math. Soc., **145**(4) (2017), 1527–1538.
- [27] E. Set, *New inequalities of Ostrowski type for mappings whose derivatives are s -convex in the second sense via fractional integrals*, Comput. Math. Appl., **63**(7) (2012), 1147-1154.
- [28] M. Tunç, *On new inequalities for h -convex functions via Riemann-Liouville fractional integration*, Filomat, **27**(4) (2013), 559–565.

- [29] M. Tunç, *Ostrowski type inequalities for m - and (α, m) -geometrically convex functions via Riemann-Louville fractional integrals*, Afr. Mat., **27**(5-6) (2016), 841–850.
- [30] H. Yildirim, Z. Kirtay, *Ostrowski inequality for generalized fractional integral and related inequalities*, Malaya J. Mat., **2**(3) (2014), 322-329.
- [31] C. Yildiz, E. Özdemir, Z. S. Muhamet, *New generalizations of Ostrowski-like type inequalities for fractional integrals*, Kyungpook Math. J., **56**(1) (2016), 161–172.
- [32] H. Yue, *Ostrowski inequality for fractional integrals and related fractional inequalities*, Transylv. J. Math. Mech., **5**(1) (2013), 85–89.