

REDUCTION OF THE GENERALIZED POPOV'S DIFFERENTIAL EQUATION TO BIRKHOFF MATRIX FORM

Ivo B. Risteski Kostadin G. Trenčevski
Valéry C. Covachev*

Dedicated to Academician Blagoj S. Popov on his 75th anniversary

Abstract

In this paper the reduction of the generalized Popov's linear differential equation with one n -tuple regular singularity to the Birkhoff canonical matrix form is given.

1 Introduction

In [1, p. 32] B. S. Popov determined the necessary and sufficient condition for reducibility of the differential equation

$$x^2 y'' + (\alpha x + \beta) x y' + (\lambda x^2 + Bx + C)y = 0$$

to a system of differential equations.

In the present paper we will show that the differential equation of B. S. Popov in a generalized form of n -th order can be reduced to a Birkhoff canonical matrix form, using the previous algebraic method given in [2].

*Partially supported by the Bulgarian Ministry of Education and Science under Grant MM 706.

1991 Mathematics Subject Classification: *Primary 34A30, Secondary 15A18.*

Key words and phrases: *Reduction, Generalized Popov's differential equation, Birkhoff matrix form.*

2 Preliminaries

Consider the generalized Popov's differential equation

$$(1) \quad x^n y^{(n)} = \sum_{i=1}^n \left(\sum_{j=0}^i a_{ij} x^j \right) x^{n-i} y^{(n-i)}$$

For the equation (1) the characteristic constants ρ_i ($1 \leq i \leq n$) of the regular singularity at the coordinate origin $x = 0$ are given as roots of the equation

$$I(\rho) \equiv [\rho]_n \cdot \sum_{i=1}^n a_{i0} [\rho]_{n-i} = 0,$$

where

$$[\rho]_k = \rho(\rho-1)\cdots(\rho-k+1), \quad [\rho]_0 = 1,$$

such that $\rho_i \not\equiv \rho_j \pmod{1}$ ($i \neq j; 1 \leq i, j \leq n$). The last equation is derived when we look for a solution of (1) behaving as x^ρ near $x = 0$. The constants ρ_i ($1 \leq i \leq n$) enter the formulation of our main result. For the sake of completeness we note that the principal characteristic constants λ_i ($1 \leq i \leq n$) of the irregular singularity of rank 1 at infinity $x = \infty$ are given as roots of the equation

$$J(\lambda) \equiv \lambda^n - \sum_{i=1}^n a_{ii} \lambda^{n-i} = 0.$$

It is derived when looking for a solution of (1) behaving as $e^{\lambda x}$ times an arbitrary power of x at infinity.

3 Main result

Now we will prove the following result.

Theorem 3.1 *The equation (1) is reduced to the Birkhoff matrix form*

$$(2) \quad xI \frac{dU}{dx} = Q(x)U \equiv (Q_0 + Q_1 x)U.$$

where

$$(3) \quad Q(x) = \begin{bmatrix} q_{11}(x) & x & 0 & \cdots & 0 & 0 \\ q_{21}(x) & q_{22}(x) & x & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ q_{n-1,1}(x) & q_{n-1,2}(x) & q_{n-1,3}(x) & \cdots & q_{n-1,n-1}(x) & x \\ q_{n1}(x) & q_{n2}(x) & q_{n3}(x) & \cdots & q_{n,n-1}(x) & q_{nn}(x) \end{bmatrix}$$

and

$$(4) \quad q_{ij}(x) = q_{ij}^0 = \text{const} \quad (j \leq i \neq n), \quad q_{nj}(x) = q_{nj}^0 + a_{n+1-j,n+1-j}x \quad (j \leq n)$$

(in particular, $q_{ii}^0 = \rho_i - i + 1$, $1 \leq i \leq n$) by a linear transformation with polynomials in x^{-1} as its coefficients.

Proof. Let us denote

$$(5) \quad y_i(x) = y^{(i)}.$$

After a differentiation this yields

$$(6) \quad y_i' = y_{i+1} \quad (0 \leq i \leq n-1).$$

If we multiply the equation (1) by x^{-n} , then it takes on the form

$$(7) \quad y_n = \sum_{i=1}^n F_i(x)y_{n-i}.$$

where

$$(8) \quad F_i(x) = \sum_{j=0}^i a_{ij}x^{j-i} \quad (1 \leq i \leq n).$$

Let us denote $\mathcal{D} \equiv \frac{d}{dx}$. Now according to (6) and (7) we obtain

$$\mathcal{D}Y \equiv \mathcal{D} \begin{bmatrix} y_0 \\ y_1 \\ \vdots \\ \vdots \\ y_{n-2} \\ y_{n-1} \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ \vdots & & & & & \\ \vdots & & & & & \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ F_n(x) & F_{n-1}(x) & F_{n-2}(x) & \cdots & F_2(x) & F_1(x) \end{bmatrix} \begin{bmatrix} y_0 \\ y_1 \\ \vdots \\ \vdots \\ y_{n-2} \\ y_{n-1} \end{bmatrix} \\ \cong L(x)Y.$$

i.e.

$$(9) \quad \mathcal{D}Y = L(x)Y.$$

By the linear transformation of Turrittin [3]

$$(10) \quad U = C'(x)Y,$$

where

$$C'(x) = \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 & 0 \\ c_{21}(x) & 1 & 0 & \cdots & 0 & 0 \\ c_{31}(x) & c_{32}(x) & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ c_{n1}(x) & c_{n2}(x) & c_{n3}(x) & \cdots & c_{n,n-1}(x) & 1 \end{bmatrix}$$

we obtain the equality

$$(11) \quad x\mathcal{D}U = x[\mathcal{D}C'(x) + C'(x)L(x)]C'^{-1}(x)U = Q(x)U,$$

and hence it follows that

$$(12) \quad x[\mathcal{D}C'(x) + C'(x)L(x)] = Q(x)C'(x).$$

From (12) $c_{ij}(x)$ ($i > j$) can be uniquely determined so that all entries of the matrix $Q(x)$ are polynomials of the required form (4).

If we denote

$$C'(x)L(x) = [p_{ij}(x)]_{n \times n} \quad \text{and} \quad Q(x)C'(x) = [l_{ij}(x)]_{n \times n},$$

then we have

$$(13_1) \quad p_{ij}(x) = c_{i,j-1}(x) \quad (1 \leq i \leq n-1).$$

$$(13_2) \quad p_{nj}(x) = c_{n,j-1}(x) + F_{n-j+1}(x)$$

and

$$(14) \quad t_{ij}(x) = q_{ij}(x) + \sum_{\nu=j+1}^i q_{i\nu}(x)c_{\nu j}(x) + xc_{i+1,j}(x),$$

where $c_{ii}(x) = 1$, $c_{ij}(x) = 0$ and $q_{ij}(x) = 0$ ($j > i$), $c_{i0}(x) = 0$ ($1 \leq i < n$), $c_{n+1,j} = 0$.

Then (12) can be written as

$$(15) \quad x[\mathcal{D}c_{ij}(x) + p_{ij}(x)] = t_{ij}(x) \quad (j < i + 2).$$

These relations are identically satisfied for the entries above the diagonal because $p_{i,i+1}(x) = 1$ and $t_{i,i+1} = x$.

For the diagonal entries (15) becomes

$$xp_{ii}(x) = t_{ii}(x),$$

which implies that

$$(16) \quad xc_{i,i-1}(x) = q_{ii}(x) + xc_{i+1,i}(x) \quad (1 \leq i \leq n-1),$$

$$(17) \quad x[c_{n,n-1}(x) + F_1(x)] = q_{nn}(x),$$

and for the j -th subdiagonal entries we obtain

$$(18) \quad \begin{aligned} & x[\mathcal{D}c_{i,i-j}(x) + c_{i,i-j-1}(x)] \\ &= q_{i,i-j}(x) + \sum_{\nu=0}^{j-1} q_{i,i-\nu}(x)c_{i-\nu,i-j}(x) + xc_{i+1,i-j}(x) \quad (2 \leq i \leq n-1). \end{aligned}$$

$$(19) \quad \begin{aligned} & x[\mathcal{D}c_{n,n-j}(x) + c_{n,n-j-1}(x) + F_{j+1}(x)] \\ &= q_{n,n-j}(x) + \sum_{\nu=0}^{j-1} q_{n,n-\nu}(x)c_{n-\nu,n-j}(x) \quad (1 \leq j \leq n-1). \end{aligned}$$

From the equalities (16)–(19) the subdiagonal entries of $C(x)$ can be determined as polynomials of the same degree with respect to x^{-1} . Now we will substitute

$$(20) \quad c_{i,i-j}(x) = \sum_{m=1}^j c_{i,i-j}^m x^{-m} \quad (2 \leq i \leq n; \quad 1 \leq j \leq i-1),$$

where $c_{i,i-j}^m$ ($1 \leq m \leq j$) are constants (we assume $c_{i,i-j}^m = 0$ for $m > j$). We recall the representation

$$(21) \quad F_i(x) = \sum_{m=0}^i a_{i,j-m} x^{-m} \quad (1 < i < n),$$

and we write

$$(22) \quad q_{ij}(x) = q_{ij}^0 + q_{ij}^1 x.$$

Substituting (20), (21) and (22) into (18) and (19), we obtain

$$(23_j) \quad -j c_{i,i-j}^j + c_{i,i-j-1}^{j+1} = q_{ii}^0 c_{i,i-j}^j + c_{i+1,i-j}^{j+1},$$

$$(23_m) \quad -m c_{i,i-j}^m + c_{i,i-j-1}^{m+1} \\ = \sum_{\nu=0}^{j-m} q_{i,i-\nu}^0 c_{i,i-j-\nu}^m + \sum_{\nu=0}^{j-m-1} q_{i,i-\nu}^1 c_{i,i-j-\nu}^{m+1} + c_{i+1,i-j}^{m+1} \quad (1 \leq m \leq j-1),$$

$$(23_0) \quad c_{i,i-j-1}^1 = q_{i,i-j}(x) + \sum_{\nu=0}^{j-1} q_{i,i-\nu}^1 c_{i,i-j-\nu}^1 + c_{i+1,i-j}^1.$$

$$(24_j) \quad -j c_{n,n-j}^j + c_{n,n-j-1}^{j+1} + a_{j+1,0} = q_{nn}^0 c_{n,n-j}^j,$$

$$(24_m) \quad -m c_{n,n-j}^m + c_{n,n-j-1}^{m+1} + a_{j+1,j-m} \\ = \sum_{\nu=0}^{j-m} q_{n,n-\nu}^0 c_{n,n-j-\nu}^m + \sum_{\nu=0}^{j-m-1} q_{n,n-\nu}^1 c_{n,n-j-\nu}^{m+1} \quad (1 \leq m \leq j-1),$$

$$(24) \quad c_{n,n-j}^1 + a_{j+1,j} + a_{j+1,j+1}x = q_{n,n-j}(x) + \sum_{\nu=0}^{j-1} q_{n,n-\nu}^1 c_{n,n-j}^1.$$

From (24) it just follows that

$$(25) \quad q_{n,n-j}(x) = a_{j+1,j+1}x + q_{n,n-j}^0 \quad (0 \leq j \leq n-1).$$

Now we are able to determine all coefficients c_{ij}^m and the polynomials $q_{ij}(x)$. We shall need the following

Lemma 3.2 *Let $\xi_0^1, \xi_0^2, \dots, \xi_0^n$ be constants satisfying the equality*

$$(26) \quad [\rho]_n + \xi_0^1[\rho]_{n-1} + \dots + \xi_0^n = \prod_{\nu=1}^n (\rho - \rho_\nu).$$

Denote

$$f(\rho) = \begin{vmatrix} \mu_0^1 + \rho - (n-2) & -1 & 0 & \dots & 0 \\ \mu_0^2 & \rho - (n-3) & -1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mu_0^{n-2} & 0 & 0 & \dots & -1 \\ \mu_0^{n-1} & 0 & 0 & \dots & \rho \end{vmatrix}$$

where μ_0^j ($1 \leq j \leq n-1$) are some constants. Let the unknown variable ζ satisfy the following equalities

$$(27) \quad \begin{aligned} \xi_0^1 + [\zeta - (n-1)] &= \mu_0^1, \\ \xi_0^j + [\zeta - (n-j)]\mu_0^{j-1} &= \mu_0^j \quad (2 \leq j \leq n-1), \\ \zeta\mu_0^{n-1} + \xi_0^n &= 0. \end{aligned}$$

If ζ takes one of the values ρ_ν , say $\zeta = \rho_n$, then

$$(28) \quad f(\rho) = [\rho]_{n-1} + \mu_0^1[\rho]_{n-2} + \dots + \mu_0^{n-1} = \prod_{\nu=1}^{n-1} (\rho - \rho_\nu).$$

Proof. Since

$$\begin{aligned} & \begin{vmatrix} \mu_0^1 + y_1 & -1 & 0 & \cdots & 0 \\ \mu_0^2 & y_2 & -1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mu_0^{n-2} & 0 & 0 & \cdots & -1 \\ \mu_0^{n-1} & 0 & 0 & \cdots & y_{n-1} \end{vmatrix} \\ &= y_1 y_2 \cdots y_{n-1} + \mu_0^1 y_2 y_3 \cdots y_{n-1} + \mu_0^2 y_3 y_4 \cdots y_{n-1} + \cdots + \mu_0^{n-1}. \end{aligned}$$

then

$$f(\rho) = [\rho]_{n-1} + \mu_0^1 [\rho]_{n-2} + \cdots + \mu_0^{n-1}$$

and

$$\begin{aligned} & (\rho - \rho_n) f(\rho) = \\ & [\rho]_n - (\rho_n - a + 1) [\rho]_{n-1} + \mu_0^1 [\rho]_{n-1} - (\rho_n - a + 2) \mu_0^1 [\rho]_{n-2} + \cdots + \mu_0^{n-1} [\rho]_1 - \rho_n \mu_0^{n-1} \\ &= [\rho]_n + \xi_0^1 [\rho]_{n-1} + \cdots + \xi_0^n = \prod_{\nu=1}^n (\rho - \rho_\nu) = (\rho - \rho_n) \prod_{\nu=1}^{n-1} (\rho - \rho_\nu). \quad \square \end{aligned}$$

Now we will start by the determination of $c_{n,n-j}^j$ ($1 \leq j \leq n-1$) and $q_{nn}(x)$. From (17) and (25) we obtain

$$(29) \quad c_{n,n-1}^1 + a_{10} = q_{nn}^0.$$

If we substitute

$$c_{n,n-j}^j = \mu_0^j \quad (1 \leq j \leq n-1),$$

then from (29) and (24_j) it follows that

$$(30) \quad \mu_0^1 = q_{nn}^0 - a_{10}.$$

$$(31) \quad \mu_0^{j+1} = (q_{nn}^0 + j) \mu_0^j - a_{j+1,0} \quad (1 \leq j \leq n-1),$$

where $\mu_0^0 = 0$. Applying Lemma 3.2 to (30) and (31), it follows that $q_{nn}^0 + n - 1$ is one of the roots ρ_i ($1 \leq i \leq n$) of the equation

$$[\rho]_n - \sum_{i=1}^n a_{ni} [\rho]_{n-i} = 0.$$

By the substitution

$$(32) \quad q_{nn}^0 + n - 1 = \rho_n$$

the coefficients μ_0^j ($1 \leq j \leq n - 1$) can be determined. In fact, according to the equation (28) the determinant

$$\begin{vmatrix} \mu_0^1 + \rho_n - (n - 2) & -1 & 0 & \cdots & 0 \\ \mu_0^2 & \rho_n - (n - 3) & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mu_0^{n-2} & 0 & 0 & \cdots & -1 \\ \mu_0^{n-1} & 0 & 0 & \cdots & \rho_n \end{vmatrix} \\ = [\rho_n]_{n-1} + \mu_0^1[\rho_n]_{n-2} + \cdots + \mu_0^{n-1} = \prod_{\nu=1}^{n-1} (\rho_n - \rho_\nu)$$

is different from zero, which means that the coefficients $c_{n,n-j}^j$ and q_{nn}^0 can be uniquely determined.

Further the coefficients $c_{i,i-j}^j$ ($1 \leq j < i - 1$) and $q_{ii}(x)$ will be successively determined. From (16) and (23_j), it follows that

$$(33_1) \quad c_{i,i-1}^1 = q_{ii} + c_{i+1,i}^1.$$

$$(33_2) \quad j c_{i,i-j}^j + c_{i,i-j-1}^{j+1} = q_{ii}^0 c_{i,i-j}^j + c_{i+1,i-j}^{j+1} \quad (1 \leq j \leq i - 1).$$

Further we apply the mathematical induction. Let

$$c_{i+1,i-j}^{j+1}(x) = \xi_0^{j+1} \quad (0 \leq j < i - 1),$$

and let the constants ξ_0^j ($1 \leq j \leq i$) satisfy the equation

$$(3_4) \quad [\rho]_i + \xi_0^1[\rho]_{i-1} + \cdots + \xi_0^i = \prod_{\nu=1}^i (\rho - \rho_\nu).$$

This is so for $i = n - 1$ by virtue of Lemma 3.2.

By the changes

$$c_{i,i-j}^j = \mu_0^j \quad (1 \leq j \leq i - 1)$$

we obtain

$$(35) \quad \mu_0^1 = q_{ii}^0 + \xi_0^1,$$

$$(36) \quad \mu_0^{j+1} = (q_{ii}^0 + j)\mu_0^j + \xi_0^{j+1} \quad (1 \leq j \leq i-1),$$

where $\mu_0^0 = 0$. Applying the Lemma to the equations (35) and (36), it follows that

$$(37) \quad q_{ii}^0 + i - 1 = \rho_i$$

and

$$(38) \quad [\rho]_{i-1} + \mu_0^1[\rho]_{i-2} + \cdots + \mu_0^{i-1} = \prod_{\nu=1}^{i-1} (\rho - \rho_\nu).$$

Again, assuming that (32) and (37) hold, $c_{i,i-j}^j$ ($1 \leq j \leq i-1$) and $q_{ii}(x)$ can be determined. Thus by mathematical induction we proved that for each i ($1 \leq i \leq n$) $c_{i,i-j}^j$ ($1 \leq j \leq i-1$) and $q_{ii}(x)$ can be determined. We also obtain that $q_{ii}(x)$ has the form

$$q_{ii}(x) = \rho_i - i + 1 + q_{ii}^1 x \quad (1 \leq i \leq n), \quad ($$

where $q_{ii}^1 = 0$ ($i \neq n$) and $q_{nn}^1 = a_{11}$.

Finally, we will prove that the sets

$$\{q_{n,n-j}(x), c_{n-j,n-m}^m; \quad 1 \leq m \leq n-j-1\}$$

and

$$\{q_{i,i-j}(x), c_{i,i-j-m}^m; \quad 1 \leq m \leq i-j-1\} \quad (n-1 \geq i \geq 1)$$

can be determined successively for the values n and i ($1 \leq i \leq n-1$). It will be proved by induction on j . Let the sets $\{q_{n,n-\nu}, c_{n-\nu,n-m}^m\}$ and $\{q_{i,i-\nu}, c_{i-\nu,n-m}^m\}$ ($1 \leq i \leq n-1$) be known for $0 \leq \nu \leq j-1$. Then from (21_m) we obtain

$$(39) \quad -m c_{n,n-j-m}^m + c_{n,n-j-m-1}^{m+1} + a_{j+m+1,j} \\ = \sum_{\nu=0}^j q_{n,n-\nu}^0 \nu c_{n-\nu,n-j-m}^m + \sum_{\nu=0}^{j-1} q_{n,n-\nu}^1 \nu c_{n-\nu,n-j-m}^{m+1} \quad (1 \leq m \leq n-j-1).$$

By the change

$$c_{n-j, n-j-m}^j = \xi_0^m \quad (1 \leq m \leq n-j-1)$$

and using (34) and (38), we obtain

$$(40) \quad [\rho]_{n-j-1} + \xi_0^1 [\rho]_{n-j-2} + \cdots + \xi_0^{n-j-1} = \prod_{i=1}^{n-j-1} (\rho - \rho_i).$$

If we substitute

$$c_{n, n-j}^m = \mu_0^m \quad (1 \leq m \leq n-j-1),$$

then from (24₀) there follow the equalities

$$(41) \quad q_{n, n-j}^1 = a_{j+1, j+1}.$$

$$(42) \quad q_{n, n-j}^0 = \mu_0^1 + \text{known terms.}$$

and from (39) we obtain

$$(43) \quad -m\mu_0^m + \mu_0^{m+1} = q_{mm}^0 \mu_0^m + q_{n, n-j}^0 \xi_0^m + \cdots \quad (1 \leq m \leq n-j-1).$$

The equation (43) can be represented in the following form

$$\mu_0^{m+1} = [\rho_n - j - (n-j-m-1)]\mu_0^m + q_{n, n-j}^0 \xi_0^m + \cdots$$

and hence we obtain the determinant

$$\begin{vmatrix} \xi_0^1 + \rho_n - j - (n-j-2) & -1 & 0 & \cdots & 0 \\ \xi_0^2 & \rho_n - j - (n-j-3) & -1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \xi_0^{n-j-2} & 0 & 0 & \cdots & -1 \\ \xi_0^{n-j-1} & 0 & 0 & \cdots & \rho_n - j \end{vmatrix} \\ = [\rho_n - j]_{n-j-1} + \xi_0^1 [\rho_n - j]_{n-j-2} + \cdots + \xi_0^{n-j-1} \\ = \prod_{i=1}^{n-j-1} (\rho_n - j - \rho_i) \neq 0.$$

Thus, the set $\{q_{n, n-j}, c_{n, n-j-m}^m\}$ is uniquely determined. In order to prove that the sets $\{q_{i, i-j}, c_{i, i-j-m}^m\}$ ($1 \leq i \leq n-1$) can be uniquely determined, we again apply mathematical induction. The arguments are similar to those above, so we omit the details. \square

References

- [1] B. S. Popov. *Formation des Critériums de Réductibilité des Equations Différentielles Linéaires Ayant des Formes Données à l'Avance*. Annuaire Fac. Phil. Univ. Skopje **5** (1952), 1-68.
- [2] I. B. Risteski. *A Simple Reduction of the Poincaré Differential Equation to Cauchy Matrix Form*. Missouri J. Math. Sci. **11** (1999), 4-9.
- [3] H. L. Turrittin. *Reduction of Ordinary Differential Equation to Birkhoff Canonical Form*. Trans. Amer. Math. Soc. **107** (1963), 485-507.

Bratfordska 2/3-9,

91000 Skopje, MACEDONIA

Institute of Mathematics,

St. Cyril and Methodius University,

P.O.Box 162, 91000 Skopje, MACEDONIA.

Email: kostatre@imom.pmf.ukim.edu.mk

Institute of Mathematics,

Bulgarian Academy of Sciences,

8 Acad. G. Bonchev Str.,

1113 Sofia, BULGARIA.

Email: rotp@math.bas.bg