

## Mistletoe in the treatment of malignant melanoma

### *Malign melanomun tedavisinde ökse otunun yeri*

Esin Sakallı Çetin<sup>1</sup>, Pınar Aslan Koşar<sup>2</sup>, Nurten Özçelik<sup>2</sup>

#### ABSTRACT

Malignant melanoma is a malignant neoplasia drives from melanocytes. Malignant melanoma, the most causing death, is seen in the third place at skin cancer. Malignant melanoma shows intrinsic resistance to chemotherapeutic agents and variability in the course of the disease which are distinct features separating from other solid tumors. These features prevent the development and standardization of non-surgical treatment models of malignant melanoma. Although there is a large number of chemotherapeutic agents used in the treatment of metastatic malignant melanoma, it hasn't been demonstrated the survival advantage of adjuvant treatment with chemotherapeutic agents. Because of the different clinical course of malignant melanoma, the disease is thought to be closely associated with immune system. Therefore, immunomodulatory therapy models were developed. Mistletoe stimulates the immune system by increasing the number and activity of dendritic cells, thus it has been shown to effect on tumor growth and metastasis of malignant melanoma patient. Outlined in this review are the recent developments in the understanding the role of mistletoe as a complementary therapy for malignant melanoma. *J Clin Exp Invest* 2014; 5 (1): 145-152

**Key words:** Malignant melanoma, mistletoe, *Viscum album*

#### INTRODUCTION

Malignant melanoma is the malignant transformation of melanocytes, the pigment producing cells found in the skin, eye, inner ear, and leptomeninges, of which the skin is the most common site for melanoma development [1]. Superficial spreading, nodular, and acral lentiginous, of which nodular melanomas have the worst prognosis are three histologic types of malignant melanoma [2]. Malignant melanoma is the most serious type of skin cancer with the fastest increasing incidence and survival of patients with distant metastases is generally less

#### ÖZET

Melanoma melanositlerden köken alan malign tümördür. Deri kanserleri içinde 3. sıklıkta görülen melanoma en fazla ölüme neden olan kanser tipidir. Malign melanomu, diğer solid tümörlerden ayıran belirgin özellikleri, kemoterapötik ajanlara gösterdiği intrinsic direnç ve hastalık seyrinin değişkenliğidir. Bu özellikler, malign melanomda cerrahi dışı tedavi modellerinin geliştirilmesini ve standart hale getirilmesini engellemektedir. Metastatik malign melanom tedavisinde kullanılan çok sayıda kemoterapötik ajan olmasına rağmen bu ajanlar ile yapılan adjuvan tedavi çalışmalarında sağ kalım avantajı gösterilememiştir. Malign melanomun farklı klinik seyir göstermesi nedeniyle, hastalığın immun sistem ile yakından ilişkili olabileceği düşünülmektedir. Bu nedenle immunomodulator tedavi modelleri geliştirilmiştir. Ökse otunun dendritik hücrelerin sayısını ve aktivitesini artırmak suretiyle immün sistemi uyardığı ve böylece malign melanomlu hastada tümör büyümesi ve metastazı üzerine etkili olduğu gösterilmiştir. Derlemede malign melanomun komplementer tedavisinde ökseotunun rolünün anlaşılmasına yönelik güncel gelişmeler özetlenmiştir.

**Anahtar kelimeler:** Malign melanom, ökse otu, *viscum album*

than one year [3]. The treatment of early-stage malignant melanoma is possible with wide surgical excision and regional lymph node curettage [4]. Despite intensive research, no curative treatment exists for metastatic malignant melanoma [5]. Conventional chemotherapy or combination of radio and chemotherapy have yet no broadly successful therapies and have not led to any considerable prolongation of survival [6]. As scientific medicine have been disappointingly ineffective to offer for melanoma patients with the threat of metastatic disease, the most of the patients turn to supplementary therapy such as aqueous mistletoe (*Viscum album*, VA) extracts

<sup>1</sup> Department of Medical Biology, School of Medicine, Muğla Sıtkı Koçman University, 48100, Muğla, Turkey

<sup>2</sup> Department of Medical Biology, School of Medicine, Süleyman Demirel University, 32260, Isparta, Turkey

**Correspondence:** Esin Sakallı Çetin,

Department of Medical Biology, Muğla Sıtkı Koçman University Medical School, Muğla, Turkey Email: esinsakalli@yahoo.com

Received: 02.10.2013, Accepted: 24.11.2013

Copyright © JCEI / Journal of Clinical and Experimental Investigations 2014, All rights reserved

[7]. The mechanism underlying the anti-tumoral activity of mistletoe preparations has been poorly understood. Moreover, the proposed mechanisms include induction of apoptosis of tumor cells and lymphocytes, inhibition of angiogenesis and stimulation of the cellular compartment of the immune system, raising the number and the activity of natural killer (NK) cells, dendritic cells (DCs) and granulocytes [8-11]. This review investigates the role of the mistletoe preparations on the reduction of melanoma growth and number of metastases in experimental models [8,12-17] with the enhancement of DC infiltration and apoptosis induction in the melanoma cells. In addition, a case report presents in literature, emphasized on a complete remission of malignant melanoma with mistletoe treatment [18]. These findings suggest that mistletoe brings new clinical perspectives as a complementary therapy for malignant melanoma.

### Epidemiology

Malignant melanoma is an aggressive skin disease with high incidence mortality. According to a World Health Organization estimate, there are 132,000 new cases of melanoma per year worldwide [19]. The American Cancer Society (ACS) estimates 68,130 new cases of melanoma in the United States in 2010 with 8,700 deaths, mostly male deaths, constituting a serious public health issue [20]. Malignant melanoma has become a cancer with a major socioeconomic impact because of a high mortality rate of metastatic disease and a relatively high incidence among adolescents and young adults [21]. In addition to its incidence and propensity to affect young adults, melanoma is a major health problem with high metastatic potential toward the skin, lung, brain and the gastro-intestinal tract, aggressive clinical behavior and notable resistance to currently available chemotherapeutic and immunological treatments [22]. The higher incidence of melanoma is also closely associated with the some host factors important risk factors for melanoma such as skin type, presence of dysplastic nevi, degree of pigmentation, susceptibility to ultraviolet radiation and episodes of sunburn, immunosuppression, certain melanoma susceptibility genes and family history of the disease [23]. The molecular mechanisms underlying the melanoma development remain unclear, even though there is advances in the knowledge of risk factors for melanoma [22].

### The treatment of malignant melanoma

The therapeutic management of malignant melanoma is quite challenging issue. It requires care-

ful clinical examination, skin biopsies and precise immuno-histological analysis. After tumor staging, medical or surgical intervention is made. Survival has been found to be strongly correlated with thickness and ulceration of primary tumour [24]. For patients with thicker than 4 mm melanoma, 5-year survival rate is approximately 30-50% (17). Therefore, early diagnosis is the most critical step in the disease management. For metastatic disease the prognosis is poor, with a 5-year life expectancy of <10% and a median survival of 6-8,5 months. Chemotherapy for advanced disease remains unsatisfactory [25]. Since melanoma is one of the most resistant malignancy to medical therapy among the solid tumours, therefore early diagnosis and surgical removal of the primary tumor is virtually the only curative approach currently available [26]. In 2011, for advanced malignant melanoma treatment, the Food and Drug Administration (FDA) and the European Medicines Agency (EMA) have approved new drugs, ipilimumab and vemurafenib [27]. However, time is needed to conclude whether these drugs can be replaced with dacarbazine (DTIC), a drug used for over 30 years in the therapy of metastatic melanoma, even if the response rate was only 10-15%, so the number of living is less than 6 years [26]. When all survival rates were evaluated, it was found that there was no difference between the single use or combination therapies, including CVD (cisplatin, vinblastin, DTIC) and PVB (cisplatin, vinblastin, bleomisin). The response rates of these combination therapies at phase II studies is 20-40%, the tumor response rate is <5% and <2% for log-term. As a result, the adjuvant treatment with chemotherapeutic agents seems to be not effective for survival [28].

Because malignant melanoma shows a different clinical course, it is thought to be closely associated with the immune system. Therefore, immunomodulatory therapy models, interferon alpha (INF- $\alpha$ ) and interleukin-2 (IL-2), were developed [29]. The respond rate of these therapies is 10-20 % and long time remission is seen in only 3-5 % of the cases. Also the findings of several randomized controlled trials (RTCs) conducted on the use of INF- $\alpha$  are conflicting in terms of therapeutic efficiency [30-32]. Most importantly, so far there has not been demonstrated any overall survival (OS) benefit, even after adjustment for quality of life incorporating patient's values for the toxic effects of INF- $\alpha$  treatment and melanoma recurrence [33]. Therefore, routine use of INF- $\alpha$  accompanied by clinically relevant toxic effects and represents a substantial economic burden for the health-care system is fostering a continues debate among oncologists [34-36].

## Dendritic cell-based immunotherapy of malignant melanoma

Dendritic cells (DC) are also an important tool for tumor-antigen-specific immunotherapy of cancer because of their critical role in mounting a specific immune response where their intratumoral and peritumoral density as well as their functional status are correlated with clinical staging of the disease and patient's survival [24-37]. DC originates from hematopoietic stem cells within the bone marrow. Under physiological conditions both myeloid and lymphoid precursors differentiate into immature dendritic cells. In peripheral tissues, upon the uptake of antigen, receiving danger signals or in the context of inflammation, DC undergo maturation process and migrate to the secondary lymphoid organs [37]. This maturation process is characterized by increased surface expressing of antigen presenting surface molecules like major histocompatibility complexes (MHC class I interacts with CD8+ cells whereas MHC class II interacts with CD4+ cells) and co-stimulatory molecules such as CD-54, CD-58, CD-83, CD-80 and CD-86 and secrete several pro-inflammatory cytokines (IL-2, IL-4, IL-6, IL-12, TNF- $\alpha$  etc.) [19,38]. (Figure 1) As a result of this maturation process, DC are well equipped to activate naive T cells; CD 8+ T cells differentiate into cytotoxic T cells (CTL) while CD+4 T cells differentiate into T helper-1 (Th1) or T helper-2 cells (Th2) which interact with macrophages and B cells, respectively, thus providing a link between the adaptive and innate immune system in the secondary lymphoid organs [19,37]. A Th1 immune response and also CTL activation are the goal of utilizing DC as a cancer vaccine in order to eliminate tumor [19]. DC are not just regulators of the immune response, but also maintenance of the immune tolerance in the thymus and secondary lymphoid organs. DC present self antigen (Ag) to develop immune cell and induce deletion of autoreactive T cells result in self tolerance depending on the status of the immune system: steady state versus infection. This situation very important for DC-based anti-tumor immunotherapies to control DC differentiation to prevent undesirable effects of vaccination such as tolerance induction by tolerogenic DC [29]. Moreover, immune evasion of tumor cells by down-regulation of surface or intracellular molecules are also limiting factors for the efficiency of the DC-based vaccination in tumor patients. The other limiting factors are the secretion of soluble immunosuppressive cytokines by tumor cells that convert immature into tolerogenic DC and presence of naturally occurring, antigen-specific regulatory T cells. However, DC-based vaccination

led to regression of individual metastases but no complete remission [24].

## Mistletoe and malignant melanoma

In studies using mistletoe preparations, regression of tumor metastases and complete remission was observed in experimental models with increasing the number and activity of DC (8,12-17). Mistletoe preparations, the most common form of complementary/alternative cancer therapy, are used often the protocols of adjuvant treatment with standard chemotherapy or radiotherapy. This combined treatment increased the cancer patient's quality of life with stimulating immune system [39].

Mistletoe is a semiparasitic plant of the *Loranthaceae* family that grows wild on deciduous hardwood trees like the apple, oak, ash, and elm (var *album*), or on coniferous trees like pine and fir (*Pinus* and *Abies* varieties) [40]. Mistletoe has been used in traditional medicine since ancient times, especially used by the Druids and ancient Greeks, and it appears in legend and folklore as a panacea [41]. Extracts from mistletoe have been used, as a sedative, vasodilator, diuretic, analgesic, cardiogenic, anti-spasmodic, therapeutically against various diseases including cancer, atherosclerosis, hypertension, dizziness, chorea, hysteria, periarthritis, spondylitis and arthritis [42-44].

Mistletoe was firstly used as an anticancer agent in 1917 by Steiner and Wegman, founders of anthroposophic medicine. Moreover, it has become the most commonly used form of adjuvant cancer therapies in Europe from 1970s. Particularly in Germany, Austria and Switzerland, mistletoe preparations are most frequently used in the treatment of cancer patient [45,46]. Commercially available extracts, prepared from *Viscum album* L., are marketed under a variety of brand names, including Iscador (Iscar), Eurixor, Helixor, Isorel (Vysorel), Iscucin, Plenisol (Lektinol) and Abnova-viscum [46,47]. Commercial preparations are widely differ with regard to their chemical compositions that depends on the species of the host tree (apple, elm, oak, pine, poplar and spruce) and the time of year harvested, lectin content the main active ingredient, and methods of preparations, including alcoholic-aqueous extraction and fermentation of aqueous extract with lactic acid bacteria [48].

The fermented aqueous extracts of mistletoe are biologically and biochemically standardized. One of these preparations is Iscador, which is marketed as Iscador M (from apple trees) contains 250 ng total lectins/ml, Iscador Qu (from oak trees) contains 375 ng total lectins/ml whereas Iscador P (from pine

trees) contains only trace amount of lectins [12]. The antiproliferative effects of Iscador M special, Iscador Qu special and Iscador P were investigated in a panel of 16 human tumor cell lines comprising nine different tumor types, central nervous system, gastric, non-small cell lung, mammary, prostate, renal, uterine cancer cell lines as well as hematological malignancies and melanomas. While Iscador M special and Iscador Qu special showed antitumor activity with a more than 70% growth inhibition in the mammary cell line, Iscador P showed no proliferative activity [12]. Eurixor, an unfermented aqueous extract of mistletoe, harvested from poplar trees, is standardized to contain a specific amount of ML-I. Helixor, unfermented aqueous extract of mistletoe is marketed as Helixor A (from spruce trees), Helixor M (from apple trees) and Helixor P (from pine trees) [49].

Mistletoe gained attention as a potential anti-cancer agent because of their immunomodulatory and cytotoxic properties [9,39,50-53]. Mistletoe extracts stimulate the immune system non-specifically, by increasing the number and activity of T lymphocytes, B lymphocytes, dendritic cells, natural-killer (NK) cells, neutrophils and activating phagocytic activity of granulocytes with subsequent release of cytokines such as tumor necrosis factor alpha (TNF- $\alpha$ ), interferon gamma (IFN- $\gamma$ ), granulocyte macrophage colony-stimulating factor (GM-CSF), interleukin-1 (IL-1), IL-2, IL-4, IL-5, IL-6, IL-8, IL-10, IL-12. All of these immune mechanisms, if stimulated, can induce tumor cell lysis [50-53].

From the various components found in most mistletoe extracts, lectins, alkaloids, viscotoxins and polysaccharides, several enzymes, peptides (such as viscumamide), amino acids, thiols, amines, cyclitols, lipids, phytosterols, triterpenes, flavonoids, phenylpropanes and minerals, the most active compounds in cytotoxicity and immunomodulatory effects are lectins and viscotoxins [54-56]. The three mistletoe lectins (MLs) ML-I, -II, -III are the main therapeutic components of mistletoe extracts [57]. The antitumor effect of lectins is thought to be induction of the death of tumor cells via binding of B-chain carrying the carbohydrate-binding site to the target cell surface, which enables the protein to enter the cell, and inhibition of protein synthesis by A-chain removing an adenine residue from the 28S RNA of the 60S subunit of the ribosome due to its ribosome-inactivating properties (Rip type II) [58,59].

Of the three MLs, ML-I is the widely investigated in the immunomodulation and anticancer studies [53,60,61]. Malignant melanoma cells represent an ideal target for ML-I cytotoxic therapy because pri-


mary malignant melanomas and their metastases express particularly high numbers of ML-I binding sites [62,63]. A highly significant antiproliferative effect of ML-I via the induction of apoptosis on malignant melanoma cells was demonstrated in vitro experiments [14,64]. One of these processes was observed by Thies A. et al. (2005) that all three MLs inhibited melanoma cell proliferation in a dose-dependent manner starting at very low ML concentrations (0.001-100 ng/ml) with ML-I being the most cytotoxic lectin on ultra sensitive cell line MV3 at  $1 \times 10^{-13}$  ng ML-I/ml [13]. In addition their obtained in vitro results, they (Thies et al. 2008) established a clinically relevant human melanoma xenograft scid mouse model to analyze the effect of ML-I on melanoma growth and spread [15]. Because of the expressing high number of ML-I binding sites and the being ultra sensitive to ML-I cytotoxicity in vitro (Thies et al. 2005), the human melanoma cell line MV3 was used to model a targeted therapy in vivo to analyse apoptosis rates, the number of infiltrating DCs and vascular counts in primary melanomas (PM) and their spontaneous lung metastases (LM) [15]. Purified ML-I administered at low-dose (30 ng ML-I per kg) daily for 19 days reduced both tumor weight by 35% and the number of LMs by 56 % compared to control mice. They demonstrated a significant direct cytotoxic effect of ML-I on malignant melanoma cells in the PTs and LMs by assessing apoptotic rates of the malignant melanoma cells. ML-I increased apoptosis rates in the melanoma cells of PTs at all doses, however, apoptosis rates didn't increase in parallel to increased ML-I concentrations as one would have expected from their in vitro data (Thies et al, 2005) [15]. The direct cytotoxic effect of ML-I seems to play only a minor role in the antitumorigenic effect in vivo because no significant reduction in tumor weight was noted in the higher ML-I doses, all of these results demonstrate the importance of in vivo models [33]. The number of tumour-infiltrating DCs was examined to evaluate the importance of immunomodulatory effects of ML-I on melanoma growth and spread. Low-dose lectin-I treatment significantly raised the total number of DCs and also protected them against apoptosis in PT [15].

Clinical studies with mistletoe lectins demonstrated that mistletoe preparations stimulate the cytokine secretion and monocyte function, the precursors of DCs [65]. The previous *in vitro* studies by Stern et al. indicated that aqueous mistletoe extract induced the maturation of DCs with an increased expression of co-stimulatory molecules CD80, CD86 as well as antigen presenting molecules HLA class I and II [66]. Like these studies, Elluru et al. demonstrated

that mistletoe QuSpez induced maturation and activation of DCs presented with increased expression of co-stimulatory molecules CD40, CD80 and CD86 and secretion of inflammatory cytokines IL-6 and IL-8 [8]. Duong Van Huyen et al. demonstrated that antitumoral effects of mistletoe extract (Iscador Qu FrF) are mediated by IL-12 dependent pathway, investigated on B16F1 melanoma implanted mice. In IL-12-deficient strain of mice, the inhibition of melanoma growth were abrogated by mistletoe [16]. Antitumor properties of IL-12 include an enhancement of Th1, CD8+CTL cells (19) and NK-cell functions [67]. Preliminary results indicated that mistletoe extract treatment induces an increase in the NK cells activity [8,68]. Furthermore, DCs educated by mistletoe Qu Spez stimulated CD4+T cells and activated melanoma antigen Melan-A/MART-I-specific M77-80 CD8+T cells as suggested by the increased levels of secretion of TNF- $\alpha$  and IFN- $\gamma$  in a allogenic mixed lymphocyte reaction [8]. In addition, an increase in CD4 + T and CD8 + T cells were reported by Gardin (2008) in immunological tests carried out before and after mistletoe treatment in four cancer patients, hodgkin disease, breast cancer and mul-

iple myeloma who received seven subcutaneous doses of mistletoe 20 mg, twice weekly [69].

In contrast to the effects of ML-I on primary tumours (PTs), there was not a significant induction of apoptosis in the melanoma cells in the lung metastases (LMs) at any ML-I concentration [15]. The significant reduction of LMs at low-dose ML-I (30 ng ML-I per kg) seemed to be the consequence of successful reduction of the PT mass due to ML-I treatment because correlation analyses revealed a highly significant positive correlation between the weight of the PTs and the number of corresponding LMs in all groups [15]. The reduction of the number of LMs but not of their size underlines the considerably stronger cytotoxic effect on circulating tumour cells, which are more prone to apoptosis induction than cells within the PT [70], while in established metastases cytotoxic effects of ML-I seem to play a minor role [15]. The stimulation of the immune system plays the prominent role in its antimelanoma effect via induction of maturation and activation of tumor infiltrating DCs in the PTs of all treatment groups since DC express high number of ML-I-binding sites [15,17].



**Figure 1.** The maturation process of dendritic cells, co-stimulatory molecules and pro-inflammatory cytokines

**A CASE REPORT**

**The complete remission of malignant melanoma with mistletoe treatment**

Clinical benefit from adjuvant treatment with a standardized mistletoe extract in patient with malignant melanoma was also demonstrated by Kirsch (2007)

[18]. The complete tumor remission was achieved with twice-weekly subcutaneous administration of Iscador®M (Weleda AG, CH-Arlesheim, Switzerland). A 68 years-old male patient with one malignant melanoma at the upper part of the right arm first diagnosed in 1992. In November 1999, another melanoma was surgically removed at the patient's right

shoulder, which the histologic examination revealed nodular melanoma, stage IIA (pT3, pN0, M0). After discovery of the second melanoma and surgical resection, treatment with standardized mistletoe extract (Iscador, M; Weleda AG, CH-Arlesheim, Switzerland) was initiated in November 1999 by the patient's general practitioner. During the course of the mistletoe therapy, axillary removal of 8 lymph nodes became necessary, 3 of which proved to be metastatic. In September 2001, first signs of a defined solitary liver metastasis with a maximum diameter of 2 cm in an area next to segments IV and V were detected during an abdominal ultrasound examination. Moreover, this finding was also confirmed by further sonographic examinations. The solitary liver metastasis was not resected, nor was classical antitumor treatment (chemotherapy or radiotherapy) initiated. The patient continued subcutaneous treatment with Iscador M after dose adaptation to 2 mg twice weekly (0.2 mL of a 10-mg vial) from October 2001 until November 2005. By June 2002, complete remission of the liver metastasis was diagnosed by liver ultrasound examination. No further metastases were discovered in May 2006. The use of low-dose Iscador as the sole postoperative modality for the adjuvant treatment of metastatic melanoma was extremely effective and very well tolerated in this patient. It achieved complete response and absence of all complaints [18]. These findings are in accordance with *in vivo* studies showing that mistletoe treatment had no negative influence on the vitality, behaviour and physiological responses, appearance, or food and water habits of the animals at any dosage [15].

In conclusion, despite improvements in chemotherapy and immunotherapy, therapeutic quest of the malignant melanoma continues. The early stage malignant melanoma is curable with surgical resection. However, prognosis is poor for patients with advanced melanoma. Malignant melanoma shows intrinsic resistance to chemotherapeutic agents and variability in the course of the disease which are distinct features separating from other solid tumors. These features prevent the development and standardization of non-surgical treatment models of malignant melanoma. These circumstances call for new treatment modalities for melanoma patients; one potential treatment strategy under evaluation is mistletoe, a natural immunomodulator, based treatment. Because of apoptosis induction in the melanoma cells and simultaneously enhancement the number of DCs, mistletoe treatment could have important impact in future melanoma therapy.

## REFERENCES

- Pópulo H, Soares P, Lopes JM. Insights into melanoma: targeting the mTOR pathway for therapeutics. *Expert Opin Ther Targets* 2012;16:689-705.
- Kaplan MA, Küçüköner M, İnal A, Işıkdoğan A, Uraççı Z. Ovarian malignant melanoma presenting with hypercalcemia and bone marrow infiltration: a case report and review of the literature. *J Clin Exp Invest* 2012;3:96-98.
- Goldszmid RS, İdoyaga J, Bravo AI, et al. Dendritic cells charged with apoptotic tumor cells induce long-lived protective CD4+ and CD8+ T cell immunity against B16 melanoma. *J Immunol* 2003;171:5940-5947.
- Ho VC, Sober AJ. Therapy for cutaneous melanoma. *J Am Acad Dermatol* 1990;22:159-177.
- Eigentler TK, Caroli UM, Radny P, et al. Palliative therapy of disseminated malignant melanoma: a systematic review of 41 randomised clinical trials. *Lancet Oncol* 2003;4: 748-759.
- Edler L. Mistel in der Krebstherapie. *Dtsch Arztebl* 2004;101:44-49.
- Steuer-Vogt MK, Bonkowsky A, Ambrosch P, et al. The effect of an adjuvant mistletoe treatment programme in resected head and neck cancer patients: a randomised controlled clinical trial. *Eur J Cancer* 2000;37:23-31.
- Elluru SR, Duong van Huyen JP, Delignat S, et al. Induction of maturation and activation of human dendritic cells: A mechanism underlying the beneficial effect of *Viscum album* as complimentary therapy in cancer. *BMC Cancer* 2008;8:161.
- Bussing A, Schietzel M. Apoptosis-inducing properties of *Viscum album* L. extracts from different host trees, correlate with their content of toxic mistletoe lectins. *Anticancer Res* 1999;19:23-28.
- Stein GM, Pfuller U, Schietzel M, Bussing A. Intracellular expression of IL-4 and inhibition of IFN-gamma by extracts from European mistletoe is related to induction of apoptosis. *Anticancer Res* 2000; 20:2987-2994.
- Elluru S, Van Huyen JP, Delignat S, Prost F, Bayry J, Kazatchkine MD. Molecular mechanisms underlying the immunomodulatory effects of mistletoe (*Viscum album* L.) extracts Iscador. *Arzneimittelforschung* 2006;56:461-466.
- Maier G, Fiebig HH. Absence of tumor growth stimulation in a panel of 16 human tumor cell lines by mistletoe extracts *in vitro*. *Anticancer Drugs* 2002;13:373-379.
- Thies A, Nugel D, Pfuller U, et al. Influence of mistletoe lectins and cytokines induced by them on cell proliferation of human melanoma cells *in vitro*. *Toxicology* 2005;207:105-116.
- Freudlsperger C, Thies A, Pfuller U, Schumacher U. The Proteasome Inhibitor Bortezomib Augments Anti-proliferative Effects of Mistletoe Lectin-I and the

- PPAR- $\gamma$  Agonist Rosiglitazone in Human Melanoma Cells. *Anticancer Res.* 2007;27:207-214
15. Thies A, Dautel P, Meyer A, et al. Low-dose mistletoe lectin-I reduces melanoma growth and spread in a scid mouse xenograft model. *Brit J Cancer* 2008;98:106 - 112.
  16. Duong Van Huyen JP, Delignat S, Bayry J, et al. Interleukin-12 is associated with the in vivo anti-tumor effect of mistletoe extracts in B16 mouse melanoma. *Cancer Lett* 2006;243:32-37.
  17. Thies A, Mauer S, Fodstad O, Schumacher U. Clinically proven markers of metastasis predict metastatic spread of human melanoma cells engrafted in scid mice. *Br J Cancer* 2007b;96: 609-616.
  18. Kirsch A. Successful Treatment of Metastatic Malignant Melanoma with *Viscum album* Extract (Iscador® M). *J Altern Complement Med* 2007;13:443-446.
  19. Engell-Noerregaard L, Hansen TH, Andersen MH, et al. Review of clinical studies on dendritic cell-based vaccination of patients with malignant melanoma: assessment of correlation between clinical response and vaccine parameters. *Cancer Immunol Immunother* 2009;58:1-14.
  20. Jemal A, Siegel R, Xu J, Ward E. Cancer statistics, 2010. *Cancer J Clin* 2010;60:277-300.
  21. Tsao H, Atkins MB, Sober AJ. Management of cutaneous melanoma. *N Engl J Med* 2004;351:998-1012.
  22. Li W, Sanki A, Karim RZ, et al. The role of cell cycle regulatory proteins in the pathogenesis of melanoma. *Pathology* 2006;38:287-301.
  23. Markovic SN, Erickson LA, Rao RD, et al. Malignant melanoma in the 21st century, part 1: epidemiology, risk factors, screening, prevention, and diagnosis. *Mayo Clin Proc* 2007;82:364-380.
  24. El Marsafy S, Bagot M, Bensussan A, and Mauviel A. Dendritic cells in the skin—potential use for melanoma treatment. *Pigment Cell Melanoma Res* 2008;22:30-41.
  25. Bajetta E, Del Vecchio M, Bernard-Marty C, et al. “Metastatic melanoma: chemotherapy,” *Seminars in Oncology* 2002;29:427-445.
  26. Mocellin S, Pasquali S, Rossi CR, Nitti D. Interferon alpha adjuvant therapy in patients with high-risk melanoma: a systematic review and meta-analysis. *J Natl Cancer Inst* 2010;102:493-501.
  27. Graziani G, Tentori L, Navarra P. Ipilimumab: a novel immunostimulatory monoclonal antibody for the treatment of cancer. *Pharmacol Res* 2012;65:9-22.
  28. Karabulut B, Sezgin VC, Göksel G, et al. The Efficacy and Tolerability of Intermediate High Dose Interferon Alpha 2B Treatment as an Adjuvant Therapy of High Risk Malignant Melanoma. *The Turk J Hematol and Oncol* 2005;15:6-14.
  29. Kadison AS, Morton DL. Immunotherapy of malignant melanoma. *Surg Clin North Am.* 2003;83:343-70.
  30. Eggermont AM, Suci S, Santinami M, et al. Adjuvant therapy with pegylated interferon alfa-2b versus observation alone in resected stage III melanoma: final results of EORTC 18991, a randomised phase III trial. *Lancet* 2008;372:117-126.
  31. Hauschild A, Weichenthal M, Balda BR, et al. Prospective randomized trial of interferon alfa-2b and interleukin-2 as adjuvant treatment for resected intermediate- and high-risk primary melanoma without clinically detectable node metastasis. *J Clin Oncol* 2003;21:2883-2888.
  32. Pectasides D, Dafni U, Bafaloukos D, et al. Randomized phase III study of 1 month versus 1 year of adjuvant high-dose interferon alfa-2b in patients with resected high-risk melanoma. *J Clin Oncol* 2009;27:939-944.
  33. Kilbridge KL, Cole BF, Kirkwood JM, et al. Quality-of-life-adjusted survival analysis of high-dose adjuvant interferon alpha-2b for high-risk melanoma patients using intergroup clinical trial data. *J Clin Oncol* 2002;20:1311-1318.
  34. Eggermont AM, Voit C. Management of melanoma: a European perspective. *Surg Oncol Clin N Am* 2008;17:635-648.
  35. Ascierto PA, Kirkwood JM. Adjuvant therapy of melanoma with interferon: lessons of the past decade. *J Transl Med* 2008;6:62.
  36. Dummer R, Hauschild A, Jost L. Cutaneous malignant melanoma: ESMO clinical recommendations for diagnosis, treatment and follow-up. *Ann Oncol* 2008;19:86-88.
  37. Tuettenberg A, Schmitt E, Knop J, Jonuleit H. Dendritic Cell-Based Immunotherapy of Malignant Melanoma: Success and Limitations. *JDDG* 2007;5:190-196.
  38. Satthaporn S, Eremin O. Dendritic cells (I): Biological functions. *J R Coll Surg Edinb* 2001;46:9-19.
  39. Eggenschwieler J, von Balthazar L, Stritt B, et al. Mistletoe lectin is not the only cytotoxic component in fermented preparations of *Viscum album* from white fir (*Abies pectinata*). *BMC Complement and Altern Med* 2007;7:14
  40. Legnani W. Integrative Cancer Therapies. *Anticancer Res* 2008;28:1893-7
  41. Becker H: Botany of European mistletoe (*Viscum album* L.). *Oncology* 1986;43:2-7.
  42. Lee CH, Kim JK, Kim HY, et al. Immunomodulating effects of Korean mistletoe lectin in vitro and in vivo. *Int Immunopharmacol* 2009;9:1555-1561.
  43. Deliorman D, Çalış İ, Ergun F, et al. “Studies on the vascular effects of the fractions and phenolic compounds isolated from *Viscum album* ssp. *album*”. *J of Ethnopharmacology* 2000;72:323-329.
  44. Yesilada E, Deliorman D, Ergun F, et al. Effects of the Turkish subspecies of *Viscum album* on macrophage-derived cytokines. *J Ethnopharmacol* 1998;61:195-200.
  45. Maier G, Fiebig HH. Absence of tumor growth stimulation in a panel of 16 human tumor cell lines by mistletoe extracts in vitro. *Anticancer Drugs* 2002;13:373-379.

46. Sakallı Çetin E, Özcelik N. Apoptotic mechanism of mistletoe (*viscum album*) extract used in the treatment of cancer: review. *J Med Sci* 2007;27:533-539.
47. Horneber MA, Bueschel G, Huber R, et al. Mistletoe therapy in oncology. *Cochrane Database Syst Rev* 2008;16:CD003297.
48. Ribéreau-Gayon G, Jung ML, Di Scala D, Beck JP. Comparison of the effects of fermented and unfermented mistletoe preparations on cultured tumor cells. *Oncology* 1986;43:35-41.
49. Stauder H, Kreuser ED. Mistletoe extracts standardised in terms of mistletoe lectins (ML I) in oncology: current state of clinical research. *Onkologie* 2002;25:374-380.
50. Hajto T, Hostanska K, Frei K, et al. Increased secretion of tumor necrosis factor- $\alpha$  interleukin-1, and interleukin-6 by human mononuclear cells exposed to the  $\beta$ -galactoside-specific lectin from clinically applied mistletoe extract. *Cancer Res* 1990;50:3322-3326.
51. Hajto T, Hostanska K, Fischer J, Saller R. Immunomodulatory effects of *Viscum album* agglutinin-I on natural immunity. *Anticancer Drugs* 1997;8:43-46.
52. Schink M. Mistletoe therapy for human cancer: The role of the natural killer cells. *Anticancer Drugs* 1997;8:47-51.
53. Pryme IF, Bardocz S, Pusztaï A, Ewen SWB. Suppression of growth of tumour cell lines in vitro and in tumours in vivo by mistletoe lectins. *Histol Histopathol* 2006;21:285-299.
54. Orhan DD, Küpeli E, Yesilada E, Ergun F. Anti-inflammatory and antinociceptive activity of flavonoids isolated from *Viscum album* ssp. *album*. *Z Naturforsch C*. 2006;61:26-30.
55. Jager S, Winkler K, Pfuller U, Scheffler A. Solubility studies of oleanolic acid and betulinic acid in aqueous solutions and plant extracts of *Viscum album* L. *Planta Med* 2007;73:157-162.
56. Bussing A, ed. Mistletoe. The Genus *Viscum*. Amsterdam, the Netherlands: Hardwood Academic; 2000.
57. Bussing A, Suzart K, Bergmann J, et al. Induction of apoptosis in human lymphocytes treated with *Viscum album* L. is mediated by the mistletoe lectins. *Cancer Lett* 1996;99:59-72.
58. Endo Y, Tsurugi K, Franz H. The site of action of the A-chain of mistletoe lectin I on eukaryotic ribosomes. *FEBS Lett* 1988;231:378-380.
59. Stirpe F, Barbieri L, Battelli MG, et al. Ribosome inactivating proteins from plants: present status and future prospects. *Biotechnology* 1992;10:405-412.
60. Barbieri L, Battelli MG, Stirpe F. Ribosome-inactivating proteins from plants. *Biochem Biophys Acta* 1993;1154:237-282.
61. Beuth J. Clinical relevance of immunoactive mistletoe lectin-I. *Anticancer Drugs* 1997;8:53-55.
62. Thies A, Pfuller U, Schachner M, et al. Binding of mistletoe lectins to cutaneous malignant melanoma: implications for prognosis and therapy. *Anticancer Res* 2001;21:2883-2888.
63. Thies A, Berlin A, Brunner G, et al. Glycoconjugate profiling of primary melanoma and its sentinel node and distant metastases: implications for diagnosis and pathophysiology of metastases. *Cancer Lett* 2007a;248: 68-80.
64. Thies A, Nugel D, Pfuller U, et al. Influence of mistletoe lectins and cytokines induced by them on cell proliferation of human melanoma cells in vitro. *Toxicology* 2005;207:105-116.
65. Heinzerling L, Von Baehr V, Liebenthal C, et al. Immunologic effector mechanisms of a standardized mistletoe extract on the function of human monocytes and lymphocytes in vitro, ex vivo, and in vivo. *J Clin Immunol* 2006; 26:347-359.
66. Stein GM, Bussing A, Schietzel M. Activation of dendritic cells by an aqueous mistletoe extract and mistletoe lectin-3 in vitro. *Anticancer Res* 2002;22:267-274.
67. Trinchieri G. Interleukin-12 and the regulation of innate resistance and adaptive immunity. *Nat Rev Immunol* 2003;3:133-146.
68. Hajto T, Hostanska K, Gabius HJ. Modulatory potency of the beta-galactoside-specific lectin from mistletoe extract (Iscador) on the host defense system in vivo in rabbits and patients. *Cancer Res* 1989;49:4803-4808.
69. Gardin N.E. Immunological Response to Mistletoe (*Viscum album* L) in Cancer Patients: A Four-Case Series. *Phytother Res* 2008;23:407-411.
70. Weber C, Mengs U, Schwarz T, et al. Effects of a standardized mistletoe preparation on metastatic B16 melanoma colonization in murine lungs. *Drug Res* 1998;48:497-502.