

REVIEW ARTICLE

West Nile virus infections in South-eastern Europe and in the Eastern Mediterranean area

Giovanni Rezza

Department of Infectious, Parasitic and Immunomediated Diseases Istituto Superiore di Sanità, Roma, Italy

ABSTRACT

The West Nile virus (WNV) was identified in Uganda in 1937. This virus appeared in the Mediterranean area in the early 1950s, causing an outbreak in Egypt. Since then, WNV caused a series of outbreaks, in human and/or horses, especially in North-Africa and in the Middle East. In the last few decades, the lineage 1 of West Nile started to cause sporadic cases and outbreaks of West Nile fever and neuroinvasive disease in Southern and Eastern Europe, and a large outbreak was reported in Romania in 1997; then, sporadic cases continued to be reported in central and southern Europe for years. Since 2010, West Nile has reemerged in Europe and in the Mediterranean area. Studies of molecular epidemiology show a co-circulation of WNV lineage 1 and lineage 2 in several European countries. In particular, WNV lineage 2 appears now to be established in South-eastern Europe and in the Balkans. This suggests that WNV is continuously reintroduced in this geographical area. Specifically, the two lineages of WNV are dispersed throughout Europe after arriving through bird migration, following the western and the eastern routes. The latter route is likely to be involved in the emergence and reemergence of WNV infection in South-eastern Europe, where this viral infection represent an important public health challenge. *J Microbiol Infect Dis 2014; Special Issue 1: S10-S16*

Key words: West Nile virus infections, South-eastern Europe, Eastern Mediterranean area, Emerging Infectious Diseases

Güney-batı Avrupa ve Doğu Akdeniz Bölgesinde Batı Nil Virüsü Enfeksiyonları

ÖZET

Batı Nil virus (WNV) etkeni ilk olarak Uganda'da 1937 yılında identifiye edilmiştir. Bu virus 1950 yıllarının başında Mısır'da yaptığı salgınla birlikte Akdeniz bölgesinde kendini göstermiştir. O tarihten itibaren WNV özellikle Kuzey Afrika ve Orta-Doğu bölgesinde insan ve/veya atlarda birtakım salgınlara sebep olmuştur. Son bir kaç dekad içerisinde ise WNV-lineage 1 Güney Doğu ve Doğu Avrupa'da Batı Nil Virüs ateşi ve nöroinvazif hastalık şeklinde bir takım sporadik vakalar ve salgınlara sebep olmaya başlamıştır. Romanya'da 1997 yılında büyük bir salgın yaşanmış olup orta ve güney Avrupa'da yıllardır rapor edilmeye devam etmiştir. Son olarak 2010 yılından itibaren, Batı Nil Virüs enfeksiyonları Avrupa ve Akdeniz ülkelerinde yeniden önem kazanmıştır. Moleküler epidemiyoloji çalışmaları birçok Avrupa ülkesinde "WNV lineage 1 ve lineage 2'nin varlığını göstermiştir. Özellikle, WNV lineage 2'nin Güney doğu Avrupa ve Balkan ülkelerinde günümüzde dolaşmakta olduğu gözükmektedir. Bu WNV'nin sürekli olarak bu coğrafik bölgelerde yeniden önem kazandığını düşündürmektedir. Spesifik olarak, iki WNV türünün batı ve doğu yollarını takiben kuş göçleriyle birlikte tüm Avrupa'ya dağılmaktadır. Doğu hattı boyunca gerçekleşen kuş göçlerinin Güney doğu Avrupa'da önemli bir halk sağlığı problemi haline gelen WNV enfeksiyonlarının ortaya çıkması ve yeniden önem kazanmasında muhtemelen rol oynadığı değerlendirilmektedir.

Anahtar kelimeler: Batı Nil virus enfeksiyonu, Güney-doğu Avrupa, Doğu Akdeniz Bölgesi, Yeniden önem kazanan enfeksiyon hastalıkları

BACKGROUND

The West Nile virus (WNV) is a mosquito-borne flavivirus belonging to the Japanese Encephalitis complex. The virus was firstly identified in 1937 in a febrile patient in northern Uganda.¹

Birds (i.e., waterfowl, passeriformes and columbiformes, plus many synanthropic birds) are the natural reservoir of the WNV, which is transmitted by mosquitoes of the *Culex* genus. The virus is maintained in a sylvatic cycle, probably by transmission

Correspondence: Giovanni Rezza, Department of Infectious, Parasitic and Immunomediated Diseases, Istituto Superiore di Sanità, Viale Regina Elena 299 - Area Code 00161 Rome, Italy Email: giovanni.rezza@iss.it

Received: 01 October 2013, Accepted: 19 January 2014

Copyright © Journal of Microbiology and Infectious Diseases 2014, All rights reserved

through *Culex modestus* mosquitoes, and in a urban cycle, through *Culex pipiens*, a highly ornithophilic species, which is abundant in urban areas and is a major epizootic vector among birds.^{2,3} Horses and humans are only accidental dead-end hosts.

The genome of this single-strand RNA virus encodes for a single polyprotein which is then processed by viral and cellular proteases, in order to produce three structural (capsid, C; membrane, M; and envelope, E) and seven non-structural proteins (NS1, NS2A, NS2B, NS3, NS4A, NS4B and NS5).⁴

From a phylogenetic point of view, two lineages of WNV have so far been described (3). The lineage 1 is widespread and further segregates into different subclades (1a-c): WNV-1a, which includes strains from all over the world, the Australian Kunjin (1b), and some Indian strains (1c) 5, whereas lineage 2 is mainly present in sub-Saharan Africa and has only recently been detected also in Europe. Other rare lineages have been identified (lineage 3 represented by a single isolate from Czech Republic, lineage 4 in Southern Russia, and lineage 5 in India, plus other putative lineages).⁶

WNV infection is often asymptomatic, causing a febrile syndrome in about 20% of the cases; West Nile fever (WNF) may be accompanied by a roseo-

lar or maculopapular skin rash in up to half of the patients. In less than 1% of the cases WNV infection may cause a neuroinvasive disease (WNND), with a variety of symptoms such as high fever, severe headache, stiff neck, disorientation or confusion, stupor or coma, tremors or muscle jerking, lack of coordination, convulsions, pain, partial paralysis or sudden weakness.^{2,7,8} Age-specific attack rates of WNND and fatality/case ratios tend to increase with age as a consequence of increased host susceptibility.³

WNV is widely distributed over the world. Beyond its original ecological niche in sub-Saharan Africa, WNV has been identified in Northern Africa, Europe, Asia, Australia⁹, and, since the summer of 1999, in North-America.^{10,11}

WNV distribution in South-Eastern Europe and in the Mediterranean basin

In the Mediterranean area, WNV was first identified in Egypt in the early 1950s, and then caused a series of outbreaks, in human and/or horses, especially in North-Africa and in the Middle East.⁸ Figure 1 shows the place and year of emergence of the main WNV outbreaks occurred in the Mediterranean basin and in the Balkans.




Figure 1. Main WNV outbreaks reported in the Mediterranean area: place of occurrence and year of identification.

In Europe, specific antibodies against WNV were detected for the first time in 2 Albanians in 1958.¹² In 1963, WNV was isolated from humans and mosquitoes in the Rhone Delta (13), and from humans and *Hyalomma marginatum* in the Volga Delta.^{14,15} Since the 1960s, WNV outbreaks occurred in several European countries, such as

Southern France, Southern Russia, Spain, Romania, Belarus, and Czech Republic.⁸

However, WNV was only sporadically detected in humans until 1996, when a major outbreak characterized by a high fatality rate (~10%) occurred in Romania.^{3,9} Since then, the virus has been isolated from horses, humans, and mosquitoes, in a number

of Eastern and Western European countries.^{16,17} In particular, with the exception of an outbreak among horses in Camargue, Southern France, in the year 2000¹⁸, and few cases detected in Spain¹⁹ and north-Africa²⁰, WNV activity in Europe has been mostly concentrated in Italy and in south-eastern and eastern Europe (i.e., sporadic cases have occurred in Hungary every year, whereas large outbreaks have been reported in Russia).

Hereby we concentrate the attention on the Balkan and bordering countries (i.e., Italy, Romania and Turkey) and on the Eastern Mediterranean area.

ROMANIA

Between mid-July and mid October 1996, 393 cases of WNND (352 cases) and WNF were reported in Romania. The outbreak was confined to Bucharest and to the lower Danube valley, with higher rates in the capital city. At the end of the outbreak, a seroprevalence of anti-WNV antibodies of 4.1% was estimated. *Culex pipiens* mosquito was the most likely vector of WNV.³ Case-control studies identified "mosquitoes" in the home and "flooded apartment building basements" as risk factors for WNV infection.²¹

Sporadic cases of WNND continued to be identified in consecutive years after the 1996 epidemic,^{22,23} until a large outbreak of 57 cases occurred in 2010.²⁴

ITALY

In Italy, WNV was firstly identified, in the summer of 1998; overall, 14 horses residing in the Tuscany region developed neurological symptoms²⁵; no human cases occurred during the outbreak, but sero-epidemiological surveys identified 4 asymptomatic cases among individuals working with horses.²⁶ The virus re-emerged in 2008 in the North-East of the country; since then, sporadic cases and/or clusters of West Nile neuroinvasive disease (WNND) among humans and horses continued to occur every year during the summertime.²⁷⁻³⁰ Up to now, human cases of WNF or WNND have been reported in 4 regions of the North (Veneto, Friuli Venetia Giulia, Emilia-Romagna, and Lombardy), in 2 regions of the South (Puglia and Basilicata), and in the Sardinia island.

Greece and other Balkan countries

During 2010, Greece experienced the second largest outbreak of WNV infection since the one which

occurred in Romania in the summer of 1996. Overall, 262 cases were reported, mostly in the Central Macedonia region.^{31,32} In 2011 and 2012, 69 and 161 cases were identified, also from areas that had not been affected before. The reoccurrence of human cases in two consecutive years suggests that WNV is established in the country.^{32,33} As expected, WNV has reappeared in the summer season of 2013, causing 81 cases by September.²⁶

Other Balkan countries have reported WNV cases over the last few years. In Albania, 1 to 2 cases were reported in 2010-2011, whereas few sporadic cases were reported in FYRM, Kosovo, Montenegro and Bulgaria in the last three years. In Serbia WNV was firstly identified in 2012, when 69 cases of disease were notified, and a new outbreak is occurring in 2013, with 238 cases identified up to the end of September.³³

TURKEY

Cases of WNND and/or WNF were identified between 2010 and 2012 in several areas of the country, such as eastern Thrace³⁴ and the city of Ankara³⁵; moreover, WNV-RNA was found in the CSF of humans and in the buffy coat of horses in Central Anatolia.³⁶ Human cases of WNV infection were also identified in the Mesopotamia Region in 2009, Syria border of Turkey.³⁷ Overall, surveillance data show that few cases were reported in 2010 compared with 2011, when a total of 47 human cases of WN disease was reported in the country.^{33,38} Sero-surveys conducted in 2009, using the plaque reduction neutralization test (PRNT), found a prevalence of 0.56% and 0.80% among blood donors in Anatolia³⁹ and in Hacettepe, respectively.⁴⁰ However, none of 729 healthy blood donors from Ankara tested by PCR resulted positive for WNV sequences.⁴¹ Although data on human disease suggest a first, unexpected, epidemic event, with a peak of cases in 2010, followed by a rapid decline in the following year, studies conducted in the first half of the 2000s show the presence of neutralizing antibodies to WNV in a wide range of mammals and in up to 20% of humans, suggesting previous virus circulation.⁴²

MIDDLE EAST

Israel is apparently the most affected country in the Middle East, with over 100 cases reported in 2010 and recurrence of outbreaks in the following years. Human cases of WNF have been reported in Israel since the early 1950s, when several outbreaks

occurred; large outbreaks were also reported in 1980.⁴³ In the district of Tel Aviv, cases of WNN and WNF were notified every year in the period 2005-2010; incidence peaks observed in 2005 and 2010 were attributed to either the emergence of new virus variants or to extremely hot summer.⁴⁴ Sporadic cases of WNV infection have been also reported in the Occupied Palestinian Territory, whereas no human cases have been identified in other Eastern Mediterranean countries such as Syria, Lebanon or Egypt.^{33,45}

Molecular epidemiology and phylogeography of WNV

WNV lineage 1, clade 1a, has been largely predominant in Europe and in the Mediterranean basin for a long time.

WNV strains belonging to lineage 1 have been identified over the years in Israel.⁴⁴ The large

outbreak occurred in Romania in 1996 was also caused by WNV lineage 1.⁴⁶ Moreover, lineage 1 strains were detected also in Hungary⁴⁷ and were responsible of sporadic cases and outbreaks occurred among humans and animals in northeastern Italy between 2008 and 2012.⁴⁸⁻⁵¹ Finally, this lineage was isolated in Turkey, specifically in Thrace, in 2012³⁴ and in central Anatolia.³⁶

The spatiotemporal distribution of WNV lineage 1 suggests that the virus was introduced from Africa to Europe by birds migrating through different routes (see Figure 2). In fact, microorganisms infecting birds tend to be dispersed along migration routes. WNV lineage 1 originated somewhere in Africa and was exported northward, mainly through the eastern route connecting Israel with central Europe and Russia. Another sub-Saharan strain spread to Morocco and the Maghreb, then entered Europe through Spain and possibly Italy, following the "Western route".




Figure 2. Bird migration routes and dispersion ways of WNV lineage 1 and 2.

WNV lineage 1a includes two different subclades: A and B. Subclade A, predominant in the Western Mediterranean area, has been detected also in Eastern Europe (i.e. Rumania and Volgograd, Russia), whereas the subclade B includes several strains which have been circulating in Israel (from where it was introduced into the United States of America) and in Eastern Europe.⁵² Since the eastern-European isolates are included in at least two different clades, it is likely that the virus was reintroduced several times in Eastern Europe, probably following different bird migration routes.

The lineage 2 of WNV, which was probably introduced in Europe at the beginning of the years 2000s, is now predominant in several areas of the

continent, in particular in Eastern and south-eastern Europe.

WNV Lineage 2 was isolated for the first time outside Sub-Saharan Africa in 2004, in a goshawk, in Hungary.⁴⁷ Interestingly, a lineage 1 strain had been isolated in a flock of geese in the same area in 2003, suggesting co-circulation of two different lineages in central Europe.⁴⁷ Since then, viral infections due to WNV lineage 2 have been identified in humans affected by WNF or WNN, in horses, birds, and mosquitoes. Large outbreaks of WNV disease in humans were identified in Russia, in the area of Volgograd, in 2007 and then in 2010-2012.⁵³ Surprisingly, WNV lineage 2, similar to the one isolated in Volgograd, was detected in 2010 in Roma-

nia, where all previous outbreaks had been due to lineage 1.²⁴ In Greece, large human outbreaks due to WNV lineage 2 occurred since 2010; viral sequences were obtained from *Culex pipiens* mosquitoes trapped in the town of Athanasios, west of Thessaloniki, on June 2011.⁵⁴ The enzootic circulation of WNV lineage 2 was confirmed by the detection of this strain among wild birds and chickens.^{55,56}

In Italy, where the lineage 1 of WNV was responsible for all cases up to 2011, lineage 2 was firstly detected in two patient, one from central-eastern Italy and one from north-eastern Sardinia.^{57,58} Unexpectedly, almost all cases of WNND and WNF identified in 2013 were caused by the lineage 2.⁵⁹

Long-distance migratory birds overwintering in central Africa are likely having introduced WNV lineage 2 in Eastern Europe, starting from the wetlands of Hungary at the turn of the century. This virus lineage probably arrived following the eastern route, expanding its geographic range of activity within Europe, to Austria, Italy, and in particular to Greece, where large outbreaks occurred since 2010. To this regard, short-distance migratory birds are likely to play a role in virus migration within Europe.⁶

Phylogenetic analysis of eastern European strains suggests the introduction of a slightly different strain of WNV-2 to the Volgograd region of Russia at the beginning of the years 2000s; this appears consistent with an alternative route, which overlaps with the eastern flyway till Israel, where it turns towards the Black Sea and the Russian steppes.⁶

Implications for South-eastern Europe

In summary, the two main lineages of WNV are continuously reemerging in south-eastern Europe and in the Mediterranean basin, as shown in Figure 3. Viral reintroduction, especially of WNV lineage 2, appears to occur more frequently through the eastern route of bird migration; as a consequence, south-eastern Europe appears at high risk for WNV outbreaks. However, the impact of WNV disease appears to vary largely among different countries and within the country in the same European Region. Whether there are ecological factors which may explain such differences, including the influence of climatic changes, needs to be further investigated. Recently, the lineage 2 of the WNV has become predominant in European countries, and it seems to be the unique cause of ongoing outbreaks in south-eastern Europe, with particular regards to Greece and other Balkan countries (see Table 1). In other areas, co-circulation of two different lineages appears to occur. Whether the seasonal reoccurrence of sporadic cases and outbreaks of WNV may be due to continuous reintroduction of the virus or to the establishment of an enzootic cycle, due to virus overwintering in natural hosts (i.e., non-migrant or short distance migrant birds) or in infected mosquitoes, needs to be better defined. In fact, even though the co-circulation of different WNV lineages and the detection of viral diversity within the same lineage support the hypothesis of viral reintroduction, the detection of similar viral strains across the years suggests that virus overwintering in birds and/or in mosquitoes may also play a role.

Table 1. Characteristics of WNV outbreaks in South-eastern Europe and in the Eastern Mediterranean area

Country	Year of activity	Lineage	Max N. human cases (Source. ECDC)
Albania	2011	2	2(2011)
Bosnia-Herzegovina	2013	-	1 (2013)
Bulgaria	2012	-	2 (2012)
Croatia	2012-2013	-	14 (2013)
Macedonia (FYRM)	2011-2013	-	6 (2012)
Greece	2010-2013	2	262 (2010)
Montenegro	2013	-	2 (2013)
Kosovo	2012	-	4 (2012)
Romania	1996-1997	1	393 (1996)
	2010-2013	2	57 (2010)
Serbia	2012-2013	2	238 (2013)
Turkey	2010,2 011	1	47 (2010)
Israel	1951/1980		N/A
	1998-2013	1	435 (2000), 114 (2010)
Palestine (OPT)	2012	-	2 (2012)

FYRM: Former Yugoslavian Republic of Macedonia, OPT: Occupied Palestinian Territory




Figure 3. WNV outbreaks: date of notification of the first case and virus lineage involved in the outbreak/human cases.

In conclusion, Mediterranean and Middle-East countries, along with countries located in the Balkan inlands, have been dealing with recurrent seasonal outbreaks of WNV in the last few years. This urges a coordinated and appropriate public health response. Preparedness plans aimed at improving surveillance, diagnosis, treatment and control of WNV is an essential tool for reducing the burden of infection and disease in this geographic area.

REFERENCES

- Smithburn KC, Hughes TP, Burke AW, Paul JH. A neurotropic virus isolated from the blood of a native of Uganda. *Am J Trop Med* 1940;20:471-492.
- Campbell GL, Marfin AA, Lanciotti RS, Gubler DJ. West Nile virus. *Lancet* 2002;2:519-529.
- Tsai TF, Popovici F, Cernescu C, et al. West Nile encephalitis epidemic in southeastern Romania. *Lancet* 1998;352:767-771.
- Brinton, M.A., 2002. The molecular biology of West Nile Virus: a new invader of the western hemisphere. *Annu Rev Microbiol* 56:371-402.
- Kramer L.D, Styer L.M, Ebel G.D, 2008. A global perspective on the epidemiology of West Nile virus. *Annu Rev Entomol* 53:61-81.
- Ciccozzi M, Peletto S, Cella E, et al. Epidemiological history and phylogeography of West Nile virus lineage 2. *Infect Genet Evol* 2013;17:46-50.
- Granwehr BP, LillibrIDGE KM, Higgs S, et al. West Nile virus: where are we now? *Lancet* 2004;4:547-555.
- Hubalek Z, Halouzka J. West Nile fever – a reemerging mosquito-borne viral disease in Europe. *Emerg Infect Dis* 1999;5:643-650.
- Zeller H.G, Schuffenecker I, 2004. West Nile virus: an overview of its spread in Europe and the Mediterranean basin in contrast to its spread in the Americas. *Eur J Clin Microbiol Infect Dis* 23,147-156.
- Hayes C.G, 2001. West Nile virus: Uganda, 1937, to New York City, 1999. *Ann N Y Acad Sci* 951:25-37.
- Lanciotti RS, Roehrig JT, Deubel V, et al. Origin of the West Nile virus responsible for an outbreak of encephalitis in the northeastern United States. *Science* 1999;286:2333-2337.
- Bardos V, Adamcova J, Dedei S, et al. Neutralizing antibodies against some neurotropic viruses determined in human sera in Albania. *Journal of Hygiene, Epidemiology, Microbiology and Immunology (Prague)* 1959;3:277-282.
- Hannoun C, Panthier R, Mouchet J, Eouzan JP. Isolement en France du virus West Nile a partir de maladies et du vecteur *Culex molestus* Ficalbi. *Compte Rendu de l'Academie de Sciences* 1964 ; D259:4170-4172.
- Chumakov MP, Belyaeva AP, Butenko AM. Isolation and study of an original virus from *Hyalomma plumbeum plumbeum* ticks and from the blood of a febrile patient in the Astrakhan region (in Russian). *Materialy XI Nauchnoi Sessii Instituta Poliomielitita I Virusnykh Encefalitov (Moskva)* 1964:5-7.
- Butenko AM, Chumakov MP, Stolbov DN. Serological and virological examinations in a natural focus of West Nile fever in the Astrakhan region (in Russian). *Voprosy Medicinskoi Virusologii* 1967;1:208-211.
- Calistri P, Giovannini A, Hubalek Z, et al. Clinical and neuropathological features of West Nile virus equine encephalomyelitis in Italy. *Equine Vet J* 2000;32:31-35.
- Murgue B, Murri S, Triki H, et al. 2001. West Nile in the Mediterranean basin: 1950-2000. *Ann N Y Acad Sci* 951,117-126.
- Murgue B, Murri S, Zientara S, et al. West Nile outbreak in horses in Southern France, 2000: the return after 35 years. *Emerg Infect Dis* 2001;7:62-696.
- Garcia-Bocanegra I, Jarn-Tellez JA, Napp S, et al. West Nile fever outbreak in horses and humans, Spain, 2010. *Emerg Infect Dis* 2011;2397-2398.
- Feki I, Marrakchi C, Ben Hmida M, et al. Epidemic West Nile virus encephalitis in Tunisia. *Neuroepidemiol* 2005;24:1-7.
- Han LL, Popovici F, Alexander JP, Jr, et al. Risk factors for West Nile virus infection and meningoencephalitis, Romania, 1996. *J Infect Dis* 1999;179:230-233.
- Cernescu C, Nedelcu N-I, Tardei G, et al. Continued transmission of West Nile virus to humans in Southeastern Romania, 1997-1998. *J Infect Dis* 2000;181:710-712.
- Popovici F, Sarbu A, Nicolae O, et al. West Nile fever in a patient in Romania, August 2008: case report. *Euro Surveill* 2008;13.Doi:pil: 18989.
- Sirbu A, Ceianu CS, Panculescu-Gatej RI, et al. Outbreak of West Nile virus infection in humans, Romania, July to October 2010. *Euro Surveill* 2011;16. Doi:pil:19762.
- Autorino GL, Battisti A, Deubel V, et al. West Nile virus epidemic in horses, Tuscany region, Italy. *Emerg Infect Dis* 2002; 8:1372-1378.
- Rezza G. Chikungunya and West Nile virus outbreaks: what is happening in north-eastern Italy? *Eur J Publ Health* 2009;19:236-237.

27. Rossini G, Cavrini F, Pierro A, et al. First human case of West Nile virus neuroinvasive infection in Italy, September 2008-case report. *EuroSurv* 2008;13:1-2.
28. Barzon L, Pacenti M, Cusinato R, et al. Human cases of West Nile Virus infection in north-eastern Italy, 15 June to 15 November 2010. *Euro Surveill*. 2011 Aug 18;16.
29. Rizzo C, Vescio F, Declich S, et al. West Nile virus transmission with human cases in Italy, August–September 2009. *Euro Surveill*. 2009;14. Doi:pii: 19353.
30. Rizzo C, Salcuni P, Nicoletti L, et al. Epidemiological surveillance of West Nile neuro invasive diseases in Italy, 2008 to 2011. *Eurosurv* 2012;17. Doi:pii: 20171.
31. Papa A, Danis K, Baka A, et al. Ongoing outbreak of West Nile virus infections in humans in Greece, July-August 2010. *Euro Surveill* 2010;15. Doi:pii: 19644.
32. Danis K, Papa A, Papanikolaou E, et al. Ongoing outbreak of West Nile virus infection in humans, Greece, July to August 2011. *Euro Surveill* 2011;16:pii=19951.
33. ECDC. West nile fever maps, 29/09/2013. <http://www.ecdc.europa.eu/en/Pages/home.aspx>.
34. Erdem H, Ergunay K, Yilmaz A, et al. Emergence and co-infections of West Nile virus and Toscana virus in Eastern Thrace, Turkey. *Clin Microbiol Infect* 2013; Jun 25. doi: 10.1111/1469-0691.12310 [Epub ahead of print].
35. Ergunay K, Sayiner AA, Litzba N, et al. Multicentre evaluation of central nervous system infections due to Flavi and Phleboviruses in Turkey. *J Infect* 2012 Oct;65:343-349. Doi: 10.1016/j.jinf.2012.05.010. Epub 2012 Jun 13.
36. Ozkul A, Ergunay K, Koysuren A, et al. Concurrent occurrence of human and equine West Nile virus infections in Central Anatolia, Turkey: the first evidence for circulation of lineage 1 viruses. *Int J Infect Dis* 2013; Jul; 17: e546-51. Doi: 10.1016/j.ijid.2013.02.005. Epub 2013 Mar 19.
37. Karakoc ZC, Tuzuner BM, Ergonul O, et al. West Nile virus infection in the Mesopotamia Region, Syria border of Turkey. *Vector Borne Zoonotic Dis* 2013 Jun 29 [Epub ahead of print].
38. Kalaycioglu H, Korukluoglu G, Ozkul A, et al. Emergence of West Nile virus infections in human in Turkey, 2010 to 2011. *Euro Surveill* 2012 May 24;17. Doi:pii: 20182
39. Ergunay K, Saygan MB, Aydogan S, et al. West Nile virus seroprevalence in blood donors from Central Anatolia, Turkey. *Vector Borne Zoonotic Dis* 2010 Oct;10:771-775. Doi: 10.1089/vbz.2009.0130. Epub 2009 Dec 20.
40. Ayturan S, Aydogan S, Ergunay K, et al. Investigation of West Nile virus seroprevalence in Hacettepe University Hospital blood donors and confirmation of the positive results by plaque reduction neutralization test. *Mikrobiyol Bul* 2011;45:113-124.
41. Sahiner F, Avci IY, Bedir O, et al. Investigation of West Nile virus RNA in blood donors by real-time RT-PCR. *Mikrobiyol Bul* 2012;46:464-469.
42. Ozkul A, Yildirim Y, Pinar D, et al. Serological evidence of West Nile virus (WNV) in mammalian species in Turkey. *Epidemiol Infect* 2006;134:826-829.
43. Weinberger M, Pitlik SD, Gandacu D, et al. West Nile fever outbreak, Israel, 2000 : epidemiologic aspects. *Emerg Infect Dis* 2001;7:686-691.
44. Kopel E, Amitai Z, Bin H, et al. Surveillance of West Nile virus disease, Tel Aviv district, Israel, 2005 to 2010. *Euro Surveill* 2011;16:pii=19894.
45. EpiSouth. West Nile virus circulation in the EpiSouth countries and in neighbouring areas. Update 1st July 2012. http://www.episouthnetwork.org/sites/default/files/outputs/note_west_nile_episouth_2010_2011_july2012.pdf
46. Savage HM, Caianu C, Nicolescu G, et al. Entomologic and avian investigations of an epidemic of West Nile fever in Romania in 1996, with serologic and molecular characterization of a virus isolate from mosquitoes. *Am J Trop Med Hyg* 1999;61:600-611.
47. Bakonyi T, Ivanics E, Erdelyi K, et al. Lineage 1 and 2 strains of encephalitic West Nile virus, Central Europe. *Emerg Infect Dis* 2006;12:618-623.
48. Savini G, Monaco F, Calistri P, Lelli R. Phylogenetic analysis of West Nile virus isolated in Italy in 2008. *Euro Surveill* 2008;13. Doi:pii: 19048.
49. Barzon L, Franchin E, Squarzon L, et al. Genome sequence analysis of the first human West Nile virus isolated in Italy in 2009. *Euro Surveill* 2009;14. doi:pii: 19384
50. Rossini G, Carletti F, Bordin L, et al. Phylogenetic analysis of West Nile virus isolates, Italy, 2008-2009. *Emerg Infect Dis* 2011;17:903-906.
51. Barzon L, Pacenti M, Franchin E, et al. Clinical and virological findings in the ongoing outbreak of West Nile virus Livorno strain in northern Italy, July to September 2012. *Euro Surveill* 2012;17:20260.
52. Zehender G, Ebranati E, Bernini F, et al. Phylogeography and epidemiological history of West Nile virus genotype 1° in Europe and the Mediterranean basin. *Infect Genet Evol* 2011;11:646-653.
53. Platonov AE, Fedorova MV, Shopenskaia TA, et al. Genotyping of West Nile virus strains circulating in southern Russia as an epidemiological investigation method: principles and results. *Zh Microbiol Epidemiol Immunobiol* 2011;2:29-37 (in Russian).
54. A. Papa, T. Bakonyi, K. Xanthopoulou, ET AL. Genetic characterization of West Nile virus lineage 2 Greece, 2010. *Emerg Infect Dis* 17:2011, pp. 920–922
55. Chaskopoulou A, Dovas CI, Chaintoutis SC, ET AL. Evidence of enzootic circulation of West Nile virus (Nea Santa-Greece-2010, lineage 2), Greece, May to July 2011. *Euro Surveill* 2011;16 pii=19933.
56. Valiakos G, Touloudi A, Iacovakis C, et al. Molecular detection and phylogenetic analysis of West Nile virus lineage 2 in sedentary wild birds (Eurasian magpie), Greece, 2010. *Euro Surveill* 2011; 16(18) pii=19862.
57. Bagnarelli P, Marinelli K, Trotta D, et al. Human case of autochthonous West Nile virus lineage 2 infection in Italy, September 2011. *Euro Surveill*. 2011:16.
58. Magurano F, Remoli ME, Baggieri M, et al. Circulation of West Nile virus 1 and 2 during an outbreak in Italy. *Clin Microbiol Infect* 2012;18: E545-E547.
59. Barzon L, Pacenti M, Franchin E, et al. Whole genome sequencing and phylogenetic analysis of West Nile virus lineage 1 and lineage 2 from human cases of infection, Italy, August 2013. *Euro Surveill* 2013.