



Terrestrial Vertebrate Fauna of Beytepe Campus, Hacettepe University

Hacettepe Üniversitesi Beytepe Yerleşkesinin Karasal Omurgalı Faunası

Mehmet Kürşat Şahin^{1*}, Burak Akbaba², Şafak Bulut³ and Zafer Ayaş²

¹Department of Biology, Kamil Ozdag Faculty of Science, Karamanoglu Mehmetbey University, Karaman, Turkey.

²Department of Biology, Faculty of Science, Hacettepe University, Ankara, Turkey.

³Department of Molecular Biology, Faculty of Science and Literature, Hitit University, Corum, Turkey.

ABSTRACT

Beytepe Campus (Hacettepe University) is one of the few isolated areas with large scale for wildlife within Ankara metropolis (Turkey). However, there is no checklist regarding to the terrestrial vertebrate (amphibian, reptile, avian and mammalian) fauna of this campus has ever been published so far. Therefore, field studies and literature surveys were conducted towards Beytepe Campus from 2005 to 2019. According to the results, a total of 3 amphibians, 11 reptilians, 93 avian and 14 mammalian species were identified in the study area. Due to increasing urbanization trends in this metropolitan city, this study will provide us a useful guide for further surveys and conservation activities in similar areas.

Key Words

Ankara, urbanization, amphibians, reptiles, birds, mammals.

Öz

Beytepe Yerleşkesi (Hacettepe Üniversitesi), Ankara'daki (Türkiye) kentsel yaşamda yaban hayatı için birkaç büyük izole alandan birisidir. Bununla birlikte, şimdiye kadar bu yerleşkenin karasal omurgalı (amfibi, sürüngen, kuş ve memeli) faunasına ilişkin genel bir tür listesi bulunmamaktadır. Bu amaçla 2005 – 2019 yılları arasında Beytepe Yerleşkesi'nde yapılan çalışmalar sonucunda toplam 3 amfibi, 11 sürüngen, 93 kuş ve 14 memeli türü tespit edilmiştir. Kentselleşmenin büyükşehirlerde giderek artan trendi nedeniyle, bu çalışma bize benzer alanlarda yapılacak araştırmalar ve koruma faaliyetleri için faydalı bir rehber olacaktır.

Anahtar Kelimeler

Ankara, kentselleşme, amfibiler, sürüngenler, kuşlar, memeliler.

Article History: Received: Apr 18, 2020; Revised: Jun 26, 2020; Accepted: Sep 22, 2020; Available Online: Nov 22, 2020.

DOI: <https://doi.org/10.15671/hjbc.722876>

Correspondence to: M.K.Şahin, Department of Biology, Karamanoglu Mehmetbey University, Karaman, Turkey.

E-Mail: mkursatsahin@kmu.edu.tr

INTRODUCTION

Urbanization is a key component of land-use changes due to extremely modified and intricate landscapes urban regions, within which green or open areas are seen as valuable for human welfare as well as wildlife [1,2]. The biological processes of dispersal interact with the landscape structure in influencing the distribution of populations of species present [3]. During the last decade, many researches, related to urban biodiversity, embed this issue under conservation perspective [4–7]. Urbanization phenomenon has been an important component of land use and land cover change, and its significance will certainly continue to increase in Turkey's metropolitans, like many others in the world [8]. In addition to that, while focusing on urbanization in Turkey, it cannot be neglected that rapid inhabitant increase, immigration from rural areas to urban places, and industrial progress after 1960s' has an important role to change the dynamics for people and natural habitats. The other major case for the land and biological resources around the cities is that they have tended to be consumed. Therefore, some restoration areas, like city parks, school campuses, which were built on such areas, need to be evaluated under the terms of biodiversity [9].

Here, we would like to introduce the actual terrestrial vertebrate fauna of Hacettepe University Beytepe Campus as an official record for a guideline about forthcoming studies on human – wildlife interactions in urban zones.

MATERIALS and METHODS

Study area

The field studies have been carried out from 2005 to 2019 to identify each species of terrestrial vertebrate class (amphibians, reptiles, birds and mammals) in Beytepe Campus, Ankara, which is transformed from rural to urban area year after year. Beytepe Campus comprises about 1500 ha and this region is about 918 m asl (Figure 1). It was established in 1967 on the western side of Ankara. There are two artificial wetlands in (Beytepe pond) and around Beytepe Campus (Ministry of Agriculture and Forest Work Campus pond) that contribute to maintain water sources for related terrestrial vertebrate groups. In addition to these wetlands, the study area covers coniferous forests, steppe vegetation, urbanized areas with the constructions of educational buildings

and dormitories. Average annual precipitation of 413 mm falls, and relative humidity is 45.61 ± 18.36 in recent years. The average annual temperature is 10.1°C [10,11].

Methodologies for detecting species

Amphibia surveys were carried out by visual search [12]. The survey has been done between March and October irregularly once a month in studied years. However, we focused on spring and early summer months particularly for reproduction activities. During the daytime, visual searches were made by simply walking around the pond edge and the surrounding of terrestrial habitat to look for amphibians and clusters of their spawn in the water. And also amphibians under refugia such as logs and large stones were also searched.

General observations on the reptile fauna of the region were irregularly made by visual encounter surveys from spring to fall seasons while intensive activity of specimens was clearly seen [13]. Surveys were conducted both daytime and nighttime. Flashlights were used at night patrols. All habitats including water bodies, under rocks, logs and decaying vegetation, and woodlands and bushes up to 5 m were thoroughly searched for the presence of reptiles. All collected specimens were examined carefully and recorded prior to being released back to their original point of capture.

Both amphibian and reptile species were identified by nation-wide guides [14] and websites which provide the latest articles with identification of new species about herpetofauna [15,16]. However, the species former common synonym names were also given in related fauna lists (Table 1,2).

Bird watching was done from 6:30 to 10:00 irregularly per month when weather conditions were suitable. Bird species were identified by naked eyes or professional binocular and documented immediately after observation. Several local, regional and international avian guidebooks were used in the identification process of bird species [17,18]. For nocturnal birds, such as owls, night surveys were carried out.

Mammal surveys were undertaken during different seasons of the year (2005–2019) and the report was compiled. Occasional field surveys were conducted to cover several habitats, like forests, steppe vegetation in Beytepe Campus and surroundings. The most favorable time for observation was between March and July. All



Figure 1. Study area: Beytepe Campus and near environment.

mammal sightings and signs were recorded and identified according to the trustful guide [19]. Active ground searches were undertaken throughout the study site. The survey was principally opportunistic and based on observations by using different techniques. Noninvasive methods such as capturing with camera traps, counting scats, footprints and other remains were used to determine medium and large size mammals like foxes. In the detection of small mammals, Sherman live capture traps were used. Individuals were released after identification of the species. Ultrasonic bat sound recorders (Pettersson D500X), BatSound and BatExplorer computer software were used to identify bat species.

RESULTS and DISCUSSION

The climate regime in the study area is affected by steppe conditions, and the plantation of coniferous trees might have an influence on local climatic conditions. That might provide suitable conditions for several terrestrial vertebrate groups. The results from each class given respectively in below:

A total of 3 amphibian and 11 reptile species were recorded in different areas of the campus (Table 1, 2). According to the recent publishing data, number of amphibian species have been reported from Ankara and entire Turkey are 9 and 33 respectively [20,21] so far. The situation of reptiles in province and country based is 30 and 132 respectively. Therefore, the records from a limited area (Beytepe Campus) is remarkable, compared to the entire country [20,21]. Although herpetile species studies are undervalued in urbanized areas of developing countries, like Turkey, there are serious issues that should be focused immediately and efficiently. Because Turkey is about to be in a crisis from a biodiversity perspective [22]. Here, we recommend some crucial take-action plans to monitor amphibian habitat sustainability. For example, focusing on *Pelophylax ridibundus* population level might contribute to assessing aquatic life dynamics in an urban area. On the other hand, there are several studies which are still going on to understand the ecosystem services of reptiles; such as *Ablepharus kitaibelii* habitat preferences, *Emys orbicularis-Trachemys scripta* competition in terms of these action plans. Therefore, this checklist is beneficial for continuous and further studies to understand the urban wildlife in Beytepe Campus scale.

Table 1. The list of amphibians of Beytepe Campus.

ORDO	Scientific Name	Former Synonym	Common name	IUCN	BERN	CITES
	<i>Bufotes variabilis</i>	<i>Pseudepidalea variabilis</i>	Varying toad	DD	APP-III	-
ANURA	<i>Pelophylax ridibundus</i>	<i>Rana ridibunda</i>	Eurasian marsh frog	LC	APP-III	-
	<i>Hyla orientalis</i>	<i>Hyla arborea orientalis</i>	Oriental tree frog	LC	APP-II	-

Table 2. The list of reptiles of Beytepe Campus.

ORDO	Scientific Name	Former Synonym	Common name	IUCN	BERN	CITES
TESTUDINES	<i>Testudo graeca</i>		Spur-thighed tortoise	VU	APP-II	APP-II
	<i>Emys orbicularis</i>		European pond turtle	NT	APP-II	-
	<i>Trachemys scripta</i>		Yellow-bellied slider turtle	LC	APP-II	-
SQUAMATA	<i>Mediodactylus danilewskii</i>	<i>Cyrtopodion kotschy</i>	Thin-toed gecko	LC	APP-II	-
	<i>Lacerta diplochondrodes</i>	<i>Lacerta trilineata ssp. diplochondrodes</i>	Green lizard	-	-	
	<i>Ophisops elegans</i>		Snake-eyed lizard	LC	APP-II	
	<i>Ablepharus kitaibellii</i>		Juniper skink	LC	APP-II	
	<i>Xerotyphlops vermicularis</i>	<i>Typhlops vermicularis</i>	Eurasian blind snake	LC	APP-III	
	<i>Natrix natrix</i>		Grass snake	LC	APP-II	
	<i>Elaphe sauromates</i>	<i>Elaphe quatuorlineata sauromates</i>	Eastern four-lined ratsnake	LC	APP-II	
	<i>Dolichophis caspius</i>	<i>Coluber caspius</i>	Large Whip Snake	LC	APP-III	

Table 3. The list of birds of Beytepe Campus.

ORDO	Scientific Name	Common name	IUCN	BERN	CITES	STATUS*	
PODICIPEDIFORMES	<i>Tachybaptus ruficollis</i>	Little grebe	LC	APP-II	-	R	
	<i>Podiceps cristatus</i>	Great crested grebe	LC	APP-III	-	R	
	<i>Podiceps nigricollis</i>	Black-necked grebe	LC	APP-II	-	W	
PELECANIFORMES	<i>Phalacrocorax carbo</i>	Great cormorant	LC	APP-III	-	R	
	<i>Ixobrychus minutus</i>	Little bittern	LC	APP-II	-	R	
	<i>Nycticorax nycticorax</i>	Black-crowned night heron	LC	APP-II	-	R	
	<i>Ardeola ralloides</i>	Squacco heron	LC	APP-II	-	W	
	<i>Casmerodius albus</i>	Great white egret	LC	APP-II	-	W	
	<i>Egretta garzetta</i>	Little egret	LC	APP-II	-	R	
	<i>Ardea cinerea</i>	Grey heron	LC	APP-III	-	W	
	<i>Ardea purpurea</i>	Purple heron	LC	APP-II	-	W	
	CICONIIFORMES	<i>Ciconia nigra</i>	Black stork	LC	APP-II	APP-II	M
		<i>Ciconia ciconia</i>	White stork	LC	APP-II	-	M
ANSERIFORMES	<i>Alopochen aegyptiacus</i>	Egyptian goose	LC	APP-III	-	V	
	<i>Tadorna ferruginea</i>	Ruddy shelduck	LC	APP-II	-	R	
	<i>Anas crecca</i>	Common teal	LC	APP-III	-	W	
	<i>Anas platyrhynchos</i>	Mallard	LC	APP-III	-	W	
	<i>Anas querquedula</i>	Garganey	LC	APP-III	-	W	
	<i>Aythya ferina</i>	Common pochard	LC	APP-II	-	W	

Table 3. The list of birds of Beytepe Campus. (Continue)

ORDO	Scientific Name	Common name	IUCN	BERN	CITES	STATUS*
ACCIPITRIFORMES	<i>Pernis apivorus</i>	European honey buzzard	LC	APP-III	APP-II	M
	<i>Milvus migrans</i>	Black kite	LC	APP-III	APP-II	M
	<i>Neophron percnopterus</i>	Egyptian vulture	EN	APP-III	APP-II	M
	<i>Circaetus gallicus</i>	Short-toed snake eagle	LC	APP-III	APP-II	M
	<i>Circus aeruginosus</i>	Western marsh harrier	LC	APP-III	APP-II	R
	<i>Circus cyaneus</i>	Northern harrier	LC	APP-III	APP-II	W
	<i>Accipiter gentilis</i>	Northern goshawk	LC	APP-III	APP-II	M
	<i>Accipiter nisus</i>	Sparrowhawk	LC	APP-III	APP-II	R
	<i>Buteo buteo</i>	Buzzard	LC	APP-III	APP-II	R
	<i>Buteo rufinus</i>	Long-legged buzzard	LC	APP-III	APP-II	R
FALCONIFORMES	<i>Falco tinnunculus</i>	Common kestrel	LC	APP-II	APP-II	R
	<i>Falco peregrinus</i>	Peregrine falcon	LC	APP-II	APP-I	R
GALLIFORMES	<i>Alectoris chukar</i>	Chukar partridge	LC	APP-III	-	R
	<i>Perdix perdix</i>	Grey partridge	LC	APP-III	-	R
GRUIFORMES	<i>Gallinula chloropus</i>	Moorhen	LC	APP-III	-	R
	<i>Fulica atra</i>	Coot	LC	APP-III	-	W
CHARADRIIFORMES	<i>Charadrius dubius</i>	Little ringed plover	LC	APP-II	-	W
	<i>Actitis hypoleucos</i>	Common sandpiper	LC	APP-II	-	R
COLUMBIFORMES	<i>Columba livia</i>	Rock pigeon	LC	APP-III	-	R
	<i>Columba palumbus</i>	Woodpigeon	LC	-	-	W
	<i>Streptopelia decaocto</i>	Collared dove	LC	APP-III	-	R

Table 3. The list of birds of Beytepe Campus. (Continue)

ORDO	Scientific Name	Common name	IUCN	BERN	CITES	STATUS*	
STRIGIFORMES	<i>Athene noctua</i>	Little owl	LC	APP-II	APP-II	R	
	<i>Asio otus</i>	Long-eared owl	LC	APP-II	-	R	
CORACIIFORMES	<i>Alcedo atthis</i>	Kingfisher	LC	APP-II	-	R	
	<i>Merops apiaster</i>	Bee-eater	LC	APP-III	-	M	
BUCEROTIFORMES	<i>Upupa epops</i>	Hoope	LC	APP-II	-	N	
PICIFORMES	<i>Dendrocopos syriacus</i>	Syrian woodpecker	LC	APP-II	-	R	
	<i>Hirundo rustica</i>	Barn swallow	LC	APP-II	-	N	
	<i>Delichon urbicum</i>	House martin	LC	APP-II	-	N	
	<i>Anthus campestris</i>	Twany pipit	LC	APP-II	-	W	
	<i>Anthus trivialis</i>	Tree pipit	LC	APP-II	-	M	
	<i>Anthus pratensis</i>	Meadow pipit	LC	APP-II	-	W	
	<i>Motacilla flava</i>	Western yellow wagtail	LC	APP-II	-	W	
	<i>Motacilla cinerea</i>	Gray wagtail	LC	APP-II	-	W	
	<i>Motacilla alba</i>	White wagtail	LC	APP-II	-	R	
	PASSERIFORMES	<i>Erithacus rubecula</i>	European robin	LC	APP-II	-	R
		<i>Luscinia megarynchos</i>	Nightingale	LC	APP-II	-	N
		<i>Phoenicurus ochruros</i>	Black redstart	LC	APP-II	-	W
		<i>Phoenicurus phoenicurus</i>	Redstart	LC	APP-II	-	M
		<i>Saxicola rubetra</i>	Whinchat	LC	APP-II	-	N
		<i>Oenanthe isabellina</i>	Isabelline wheather	LC	APP-II	-	N
<i>Turdus merula</i>		Blackbird	LC	APP-II	-	R	
<i>Cettia cetti</i>		Cetti's warbler	LC	APP-II	-	R	

Table 3. The list of birds of Beytepe Campus. (Continue)

ORDO	Scientific Name	Common name	IUCN	BERN	CITES	STATUS*
PASSERIFORMES	<i>Hippolais pallida</i>	Olivaceous warbler	LC	APP-II	-	N
	<i>Phylloscopus collybita</i>	Chiffchaff	LC	APP-II	-	R
	<i>Muscicapa striata</i>	Spotted flycatcher	LC	APP-II	-	N
	<i>Ficedula parva</i>	Red-breasted flycatcher	LC	APP-II	-	M
	<i>Aegithalos caudatus</i>	Long-tailed tit	LC	APP-II	-	W
	<i>Parus ater</i>	Coal tit	LC	APP-II	-	R
	<i>Parus caeruleus</i>	Blue tit	LC	APP-II	-	R
	<i>Parus major</i>	Great tit	LC	APP-II	-	R
	<i>Certhia brachydactyla</i>	Short-toed treecreeper	LC	APP-II	-	R
	<i>Lanius collurio</i>	Red-backed shrike	LC	APP-III	-	N
	<i>Lanius minor</i>	Lesser gray shrike	LC	APP-III	-	M
	<i>Lanius nubicus</i>	Masked shrike	LC	APP-III	-	M
	<i>Garrulus glandarius</i>	Jay	LC	-	-	R
	<i>Pica pica</i>	Magpie	LC	-	-	R
	<i>Corvus monedula</i>	Jackdaw	LC	-	-	R
	<i>Corvus frugilegus</i>	Rook	LC	-	-	W
	<i>Corvus cornix</i>	Carrion crow	-	-	-	R
	<i>Corvus corax</i>	Raven	LC	APP-III	-	R
	<i>Sturnus vulgaris</i>	Starling	LC	-	-	R
	<i>Passer domesticus</i>	House sparrow	LC	-	-	R
<i>Passer montanus</i>	Tree sparrow	LC	APP-III	-	R	
<i>Fringilla coelebs</i>	Chaffinch	LC	APP-III	-	R	

Table 3. The list of birds of Beytepe Campus. (Continue)

ORDO	Scientific Name	Common name	IUCN	BERN	CITES	STATUS*
	<i>Carduelis chloris</i>	Greenfinch	LC	APP-II	-	W
	<i>Carduelis carduelis</i>	Goldfinch	LC	APP-II	-	R
	<i>Carduelis cannabina</i>	Linnet	LC	APP-II	-	W
	<i>Coccothraustes coccothraustes</i>	Hawfinch	LC	APP-II	-	W
PASSERIFORMES	<i>Emberiza cirtrinella</i>	Yellow hummer bunting	LC	APP-II	-	W
	<i>Emberiza hortulana</i>	Ortolan bunting	LC	APP-III	-	W
	<i>Emberiza schoeniclus</i>	Reed bunting	LC	APP-II	-	R
	<i>Emberiza melanocephala</i>	Black-headed bunting	LC	APP-II	-	N
	<i>Miliaria calandra</i>	Corn bunting	LC	APP-III	-	N

*Status N: nesting, V: vagrant, W: wintering, M: migratory/stopover

The results pointed out that a total number of 93 bird species, belonging to 15 orders. Passeriformes is the largest order and comprised 47 (50%) of the recorded bird species. The rest 47 species (50%) represented the non-passerines, which form the other bird orders (Table 3). According to recent reports, while 324 species were recorded in Ankara, the species number increased up to 478 (364 of them are resident) in Turkey [20,23]. Therefore, avifauna records from Beytepe Campus are remarkable and that's why this area can be identified as an isolated refuge within this metropolis.

The mammalian diversity recorded in the campus till 2019 is represented by 5 orders and 14 species (Table 4). Compared to Turkey (165-168 species) [24], Beytepe Campus and its near environment host 14 species that consist of 8.48% of the entire country's fauna. While 51 mammal species were recorded from Ankara Province, the records from Beytepe Campus shows us that, this

isolated refuge does not only provide suitable habitats for other vertebrate groups, but also hosts several mammal species. Most species belong to the Rodentia and Carnivora order followed by Chiroptera.

It is clear that human values, perceptions, and limited city budgets often cause many dilemmas for urban conservation biology. In addition to that, sometimes it is difficult for the general public to comprehend why this biotic diversity should be protected. People directly affect urban vegetation, and as a result urban habitat quality and quantity, due to habitats, defined by vegetation [25]. The rest of the biological community is mostly

Table 4. The list of mammals of Beytepe Campus.

ORDO	Scientific Name	Common name	IUCN	BERN	CITES
EULIPOTYPHILA	<i>Erinaceus concolor</i>	Hedgehog	LC	-	-
	<i>Crocidura suaveolens</i>	Lesser shrew	LC	APP-II	-
CHIROPTERA	<i>Hypsigo savii</i>	Savi's pipistrelle	LC	-	-
	<i>Pipistrellus pipistrellus</i>	Common pipistrelle	LC	-	-
LAGOMORPHA	<i>Lepus europaeus</i>	European hare	LC	-	-
RODENTIA	<i>Microtus guentheri</i>	Mediterranean vole	LC	-	-
	<i>Apodemus witherbyi</i>	Steppe field mouse	LC	-	-
	<i>Apodemus flavicollis</i>	Yellow-necked field mouse	LC	-	-
	<i>Mus macedonicus</i>	Macedonian mouse	LC	-	-
	<i>Mus domesticus</i>	House mouse	LC	-	-
	<i>Rattus rattus</i>	Black rat	LC	-	-
CARNIVORA	<i>Vulpes vulpes</i>	Red fox	LC	-	APP-III
	<i>Mustela nivalis</i>	Least weasel	LC	APP-III	-
	<i>Martes foina</i>	Stone marten	LC	APP-III	APP-III

determined by this vegetation template and interactions with other species and its environment [25]. Moreover, fragmentation via constructions is also inherent in cities and might be one of the causes of biodiversity loss. Except for a few exceptional examples, terrestrial vertebrate species feel the pressure of urbanization in their life cycles via several factors, such as habitat loss or fragmentation, destruction to their nests, distanced by ecological corridors, and etc [26-28]. However, campus areas maintain relatively better sites to focus on many kinds of anthropogenic risks on terrestrial or aquatic wildlife sustainability. As a result, Beytepe Campus hosts 8.64% of Turkish herpetofauna, 19.45% of avifauna and 8.48% of mammal fauna elements. We can conc-

lude that isolated areas such as campuses are essential to maintain urban biodiversity [26].

Here, conservation status of recorded species were organized according to the IUCN, BERN, and CITES criteria in related tables (Table 1-4) [29-31]. Species that represent the class were given in Appendix:

Acknowledgments: This study was supported by Hacettepe University Scientific Research Projects Coordination Unit (012 D10 601 001-46).

APPENDIX: Representative species from Beytepe Campus fauna



a *Pelophylax ridibundus*



b *Hyla orientalis*



c *Testudo graeca*



d *Ophisops elegans*



e *Phylloscopus collybita*



f *Asio otus*



g *Crocidura suaveolens*



h *Vulpes vulpes*

Basimil

10-10-2015 00:45:24

References

1. S.T. Pickett, M.L. Cadenasso, J.M. Grove, C.H. Nilon, R.V. Pouyat, W.C. Zipperer, R. Costanza, Urban ecological systems: linking terrestrial ecological, physical, and socioeconomic components of metropolitan areas, *Annual Review of Ecology and Systematics*. 32 (2001) 127–157.
2. S.T. Pickett, M.L. Cadenasso, J.M. Grove, Resilient cities: meaning, models, and metaphor for integrating the ecological, socio-economic, and planning realms, *Landscape and Urban Planning*. 69 (2004) 369–384.
3. J. Niemelä, Ecology and urban planning, *Biodiversity & Conservation*. 8 (1999) 119–131.
4. D.C. Dearborn, S. Kark, Motivations for conserving urban biodiversity, *Conservation Biology*. 24 (2010) 432–440.
5. M.A. Goddard, A.J. Dougill, T.G. Benton, Scaling up from gardens: biodiversity conservation in urban environments, *Trends in Ecology & Evolution*. 25 (2010) 90–98.
6. B. Güneralp, K.C. Seto, Futures of global urban expansion: uncertainties and implications for biodiversity conservation, *Environmental Research Letters*. 8 (2013) 014025.
7. M.F. Aronson, C.A. Lepczyk, K.L. Evans, M.A. Goddard, S.B. Lerman, J.S. MacIvor, C.H. Nilon, T. Vargo, Biodiversity in the city: key challenges for urban green space management, *Frontiers in Ecology and the Environment*. 15 (2017) 189–196.
8. X. Zhang, M.A. Friedl, C.B. Schaaf, A.H. Strahler, A. Schneider, The footprint of urban climates on vegetation phenology, *Geophysical Research Letters*. 31 (2004).
9. H. Sevgili, A. Karataş, O. Candan, Biodiversity in Urban environments of Ordu City and Nearby Areas: Mammals, Birds, Reptiles and Amphibians, *Hacettepe Journal of Biology and Chemistry*. 44 (2016) 47–63.
10. P. Faramarzi, Determination of Chemical Composition in Pm2. 5 Samples Collected in Urban and Semi-Urban Two Stations in Ankara, Hacettepe University, Graduate School of Science and Engineering, 2018.
11. Climate:Ankara, Ankara_Climate. (n.d.). <https://tr.climate-data.org/asya/tuerkiye/ankara/ankara-172/> (accessed April 10, 2020).
12. R. Heyer, M.A. Donnelly, M. Foster, R. Mcdiarmid, Measuring and monitoring biological diversity: standard methods for amphibians, Smithsonian Institution, 2014.
13. N.J. Scott, M.L. Crump, B.L. Zimmerman, R.G. Jaeger, R.F. Inger, P.S. Corn, B.D. Woodward, C.K. Dodd, D.E. Scott, H.B. Shaffer, Standard techniques for inventory and monitoring, Measuring and Monitoring Biological Diversity. Standard Methods for Amphibians/Heyer, W. Ronald. (1994).
14. İ. Baran, Ç. Ilgaz, A. Avcı, Y. Kumlutaş, K. Olgun, Türkiye Amfibi ve Sürüngenleri [The Amphibians and Reptiles of Turkey]; Ankara Tübitak Popüler Bilim Kitapları no: 207, Semih Publishing House.(in Turkish), 2013.
15. AmphibiaWeb, (n.d.). <https://amphibiaweb.org/> (accessed February 10, 2020).
16. The Reptile Database, (n.d.). <http://www.reptile-database.org/> (accessed June 10, 2019).
17. C.J.O. Harrison, field guide to the nests, eggs and nestlings of European birds, Collins, 1975.
18. H. Heinzel, R. Fitter, J. Parslow, A. Boyla, Türkiye'nin ve Avrupa'nın kuşları: Kuzey Afrika ve Ortadoğu dahil, Doğal Hayatı Koruma Derneği, 2002 (in Turkish).
19. P. Bang, P. Dahlström, G. Vevers, Collins guide to animal tracks and signs: The tracks and signs of British and European mammals and birds, Collins, 1974.
20. Ankara İli Tüm Yüzölçüm Alanı için Karasal ve İç su Ekosistemleri Biyolojik Çeşitlilik Envanteri ve İzleme Projesi Sonuç Raporu, Tarım Ve Orman Bakanlığı, 2014 (in Turkish).
21. B. Akman, M. Çakmak, M.Z. Yildiz, On the Herpetofauna of the Central Anatolian Province of Kırkkale (Turkey), *Acta Biologica Turcica*. 33 (2020) 70–79.
22. Ç.H. Şekercioğlu, S. Anderson, E. Akçay, R. Bilgin, Ö.E. Can, G. Semiz, Ç. Tavşanoğlu, M.B. Yokeş, A. Soyumert, K. Ipekdal, Turkey's globally important biodiversity in crisis, *Biological Conservation*. 144 (2011) 2752–2769.
23. G. Kirwan, B. Demirci, H. Welch, K. Boyla, M. Özen, P. Castell, T. Marlow, The birds of Turkey, Bloomsbury Publishing, 2010.
24. A. Karataş, A. Budak, The mammals (Mammalia) of Yamanlar Mountain (İzmir and Manisa provinces) and their ectoparasites, *Acta Biologica Turcica*. 32 (2019) 53–60.
25. J. Beninde, M. Veith, A. Hochkirch, Biodiversity in cities needs space: a meta-analysis of factors determining intra-urban biodiversity variation, *Ecology Letters*. 18 (2015) 581–592.
26. M.L. McKinney, Urbanization, Biodiversity, and Conservation The impacts of urbanization on native species are poorly studied, but educating a highly urbanized human population about these impacts can greatly improve species conservation in all ecosystems, *Bioscience*. 52 (2002) 883–890.
27. A.J. Hamer, M.J. McDonnell, The response of herpetofauna to urbanization: inferring patterns of persistence from wildlife databases, *Austral Ecology*. 35 (2010) 568–580.
28. K. Szlavecz, P. Warren, S. Pickett, Biodiversity on the urban landscape, in: *Human Population*, Springer, 2011: pp. 75–101.
29. D.S. Favre, International trade in endangered species: a guide to CITES [Convention on International Trade in Endangered Species], Brill, 1989.
30. P. Genovesi, C. Shine, European strategy on invasive alien species: Convention on the Conservation of European Wildlife and Habitats (Bern Convention), Council of Europe, 2004.
31. S. IUCN, IUCN red list of threatened species, Version. 2017 (2017) 3.