



ORIGINAL ARTICLE

Clin Exp Ocul Trauma Infect. 2020; 2(3):56-60

Outcomes of Early Pars Plana Vitrectomy for Acute Post Operative Endophthalmitis with or without Silicone Oil**● Hussain Ahmad Khaqan¹, ● Usman Imtiaz¹, ● Hasnain Muhammad Buksh¹, ● Hafiz Ateeq Ur Rehman¹, ● Raheela Naz¹**

1 Ameer Ud Din Medical College PGMI, Lahore General Hospital, Lahore, Pakistan

Abstract

Purpose: To evaluate the anatomical and functional outcomes of pars plana vitrectomy (PPV) in acute post operative endophthalmitis with or without endotamponade.

Material and Methods: Quasi experimental study was conducted at Lahore General hospital, Lahore. One hundred ninety patients of acute post-surgical endophthalmitis were included in the study. Patients were randomized into two groups after no clinical improvement was seen post primary vitreous tap and intravitreal vancomycin and ceftazidime. In group 1 patients underwent PPV with endotamponade (silicone oil) were included while in group 2 patients underwent PPV without endotamponade were included. Study was divided in two phases. In first phase 30 patients underwent PPV without endotamponade and 30 patients with endotamponade. Considering the results of phase 1, rest of the 130 patients underwent PPV with endotamponade in phase 2. Removal of silicone oil in all patients was done at 12 weeks.

Results: In first phase of study 23 (76.66%) patients in Group 2 showed retinal detachment within four weeks of follow up, while no patient (0%) in Group 1 showed retinal detachment within four weeks of follow up. Later 6 (7.31%) patients in group 1 showed retinal detachment within four weeks of silicone oil removal. In second phase all 130 patients showed no

retinal detachment after undergoing PPV with endotamponade as in group 1. Overall 160 patients underwent PPV with endotamponade including first and second phase and only 6 patients got retinal detachment.

Conclusion: Early PPV with endotamponade should be preferred to PPV without endotamponade in cases of acute post-operative endophthalmitis due to statistically significant improvement in anatomical and functional outcomes.

Introduction

Endophthalmitis is one of the most devastating vision threatening intraocular inflammation.(1) There are two main routes for inoculation of this condition: Exogenous and endogenous. Exogenous endophthalmitis can occur post-operative, post-traumatic, or post-intravitreal injections due to ocular contamination by infective agents from the external environment.(1) Endogenous endophthalmitis is less common and is caused by spread of microorganisms through blood from different parts. Endophthalmitis causes severe anatomical and functional damage of intraocular structures leading to marked visual deterioration.(2) Acute post-operative endophthalmitis usually occurs within 5-6 days after an intra-ocular surgery. Most of the cases occur after cataract surgery.(3,4) After cataract surgery the incidence of acute-postoperative endophthalmitis ranges from 0.03% to 0.2% in different pub-

lications.(5-12) Ocular surgeries other than cataract surgery such as; penetrating keratoplasty (5,13,14) scleral buckling (15) and glaucoma drainage device implantation (16) show less incidence of acute postoperative endophthalmitis as compared with post cataract surgery. There are many treatment options for this sight threatening condition including intravitreal antibiotics, parsplana vitrectomy (PPV) and adjunctive systemic antibiotics. Endophthalmitis vitrectomy study (EVS) provides us the guidelines for the treatment of endophthalmitis with respect to vision at presentation. PPV is generally recommended in patients presenting with light perception (LP) vision while in patients presenting with visual acuity of better than LP intravitreal antibiotics is recommended. (17) PPV can be performed with and without endotamponade (Silicon oil). A study was conducted to evaluate the efficacy of PPV with endotamponade (silicon oil) and they found silicon oil having intrinsic bactericidal properties.(18) Another study was conducted which showed silicone oil a beneficial adjunct to vitrectomy in the treatment of endophthalmitis. In PPV with silicone oil endotamponade all the patients were found to have better visual outcomes (19). As endophthalmitis causes diffuse tissue necrosis and post-operative retinal detachment so endotamponade plays an important role in securing visual and anatomical outcomes (19). A study was done to compare the post PPV outcomes with and without endotamponade in the treatment of endophthalmitis. There was markedly increased incidence of postoperative retinal detachment in PPV without endotamponade (20) . This study was done to evaluate and quantify the effect of endotamponade, in preventing post PPVretinal detachment, done for endophthalmitis.

Material And Methods

A total of 190 subjects with acute post-operative endophthalmitis were enrolled in this study on the basis of EVS recommendations. This was a quasi experimental study conducted at Lahore General Hospital, Lahore Pakistan, from 2011 to 2019. Sample size was calculated by WHO standard formula with 95% confidence interval. Written and informed consent was taken from all participants. Approval of the Ethical Committee of Lahore General Hospital, Lahore was obtained. A detailed history and evaluation of all the participants was done systematically to identify any risk factors causing endophthalmitis. All participants were randomly divided in two equal groups. In first phase 30 patients of Group 1 underwent PPV with silicon oil and 30 patients of Group 2 underwent PPV only. Second phase started 4 weeks after first phase and rest of all 130 patients underwent PPV with endotamponade (table 2). 23 G PPV with and without silicon oil was done and patients were evaluated at first day, first week, first month, third months and sixth months. Silicon oil was removed 2 months

after surgery. On every follow up visual acuity, IOP and fundus examination were recorded. Statistical package SPSS version 15.0 was used for data analysis.

Results

Total 190 patients with acute post-operative endophthalmitis were enrolled in this study. 111 (58.42%) were male and 79 (41.57%) were female. Mean age of participants was 48 years. 23 (76.66%) patients who underwent PPV alone in Group 2 presented with retinal detachment during first four weeks follow up (table 2). In first phase of study 23 (76.66%) patients out of 30 who underwent PPV only (Group 2) showed retinal detachment within first four weeks of follow up, while among 30 patients of Group 1 who underwent PPV with endotamponade, no patient showed retinal detachment in first four weeks postoperatively . Later 6 (7.31%) patients in group 1 showed retinal detachment within four weeks of silicone oil removal (figure 1). In second phase rest of 130 patients underwent PPV with endotamponade. No patient showed retinal detachment at first 4 weeks follow up (figure 2). Patients with retinal detachment underwent redo surgery for PPV with endotamponade of silicone. Silicon oil was removed after 2 months in all patients except 6 patients. In those six patients silicon oil was removed in first month and RD was seen in all those patients. 149 (92.68%) participants showed improved vision (6/36-6/60) in Group 1 and 7 (23.33%) participants in Group 2 showed improved vision (6/36-6/60).




Figure 1: Retinal detachment ratio in two groups.

In every case vitreous and aqueous tap was done and sent for culture, antibiotics sensitivity and gram/giemsa staining. Vitreous examination provided more positive results as compared to aqueous sample examination (92% in vitreous tap; 78% aqueous tap). Staphylococcus aureus was isolated in most of the cases (41.5%), followed by Streptococcus pneumo-

niae (20.5%), and Pseudomonas aeruginosa (25%).




Figure 2: After removal of oil ratio of retinal detachment in second phase.

Discussion

Considerable differences were observed between two groups in our study. At fourth post-operative week and sixth post-operative month after surgery, Group 1 (who underwent PPV

Table 1: Patient distribution in the study phases.

First Phase		
	Group 1	Group 2
N	30	30
RD	6	23
%	20%	76.67%
Second Phase		
	Group 1	Group 2
N	130	0
RD	0	0
%	0%	0%

Table 2: Results of the two groups.

Combined	Group 1	Group 2
Total Patients	160	30
RD	6	23
%	3.75%	76.67%

with silicone oil endotamponade had better visual and functional outcomes and less need to re-surgery. The results were in favour of conclusions from previous studies.(19) It shows the significance of endotamponade with silicone oil in PPV for endophthalmitis. Parsplana vitrectomy has improved the anatomical and functional outcomes of endophthalmitis from a success rate of 33% (21) to 40% (22) . Role of surgical management (PPV with endotamponade) in improving visual function has been shown by many studies and plays an important role in securing the useful vision of patients. Success rate of this study 92.68% in Group 1 compared with 30% by another study showed the importance of endotamponade. (23) Improved functional and anatomical outcome of PPV with endotamponade (silicon oil) could be explained as follows: Eradication of microbes by antibiotics is assisted by silicon oil. (18)

A study was published which showed that silicone oil has inhibitory effect on most of the microorganisms including aerobes, facultative aerobes and anaerobes.(18) Postoperative examination and additional laser treatments can be done effectively as silicon oil keeps the media clear. Because of good surface tension, silicon oil pushes the retina against the eye wall, hence giving a good tamponade and sealing the retinal breaks effectively.(19) In severely infected eyes to perform PPV carries hazards. Unexpected damage to the retina can occur due to obscuration of the view because of opaque media.

In endophthalmitis, retina becomes infected, ischemic and fragile and iatrogenic injuries or tractions might be happen during surgery. After surgery there can be necrosis of retina secondary to persistent intraocular inflammation. Postoperative hypotony can result due to ciliary body damage. These issues may cause retinal detachment. (19,24)

Considerable difference in retinal detachment after PPV was seen between groups in our study in eyes with endophthalmitis. 76.66% of participants showed retinal detachment at first post-operative week in Group 2. All these eyes required re-operation with silicone oil endotamponade to obtain better visual outcomes by restoring the anatomical aspects. In the Group 1, there were 6 cases of retinal detachment that occurred later after removing silicon oil. Proliferative vitreo-retinopathy (PVR) plays an important role in the late complications in treating endophthalmitis. In Group 1 all the six cases who got retinal detachment where repaired with silicone oil endotamponade and PVR was the main cause for retinal detachment. (24) Among 160 participants with oil filled eyes of Group 1, silicon oil was removed in all cases at 2 months. Progressive PVR was responsible for recurrent retinal detachment, causing new breaks or tractions emphasizing the need for endotamponade with silicon oil (25) . Retinal breaks and tractions were responsible for recurrent detachments after surgery (26). These recurrent detachments had very poor prognosis and

these eyes became phthisic.

In this study *Staphylococcus aureus* was isolated in most of the cases (41.5%), followed by *Streptococcus pneumoniae* (20.5%), and *Pseudomonas aeruginosa* (25%). In South East Asia region, the most common pathogens were Gram negative rods *Klebsiella* from hepatobiliary infections are the major cause of endophthalmitis in South East Asia region while gram positive cocci i.e. *Staphylococcus* and *Streptococcus* are the leading cause of endophthalmitis in the region of Europe and America. (18)

In the study, we found that endotamponade with silicon oil is an important tool for adequate attachment of retina after PPV. Early PPV for post-surgical endophthalmitis with poor red reflex and vision of light perception show dramatic effects, already shown by Endophthalmitis Vitrectomy Study (17) Early pars plana vitrectomy with endotamponade resulted in improvement in anatomical and functional outcomes.

References

- Mamalis N. Endophthalmitis. *J Cataract Refract Surg.* 2002; 28 (5): 729–730.
- Smith SR, Kroll AJ, Lou PL, Ryan EA. Endogenous bacterial and fungal endophthalmitis. *Int Ophthalmol Clin.* 2007; 47 (2): 173–183.
- Moloney TP, Park J. Microbiological isolates and antibiotic sensitivities in culture-proven endophthalmitis: a 15-year review. *Br J Ophthalmol.* 2014; 98 (11): 1492–1497.
- Verbraeken H. Treatment of postoperative endophthalmitis. *Ophthalmologica.* 1995; 209 (3): 165– 171.
- Wykoff CC, Parrott MB, Flynn HW, Jr, Shi W, Miller D, Alfonso EC. Nosocomial acute-onset postoperative endophthalmitis at a University Teaching Hospital (2002–2009) *Am J Ophthalmol.* 2010; 150 (3): 392.
- Miller JJ, Scott IU, Flynn HW, Jr, Smiddy WE, Newton J, Miller D. Acute-onset endophthalmitis after cataract surgery (2000–2004): incidence, clinical settings, and visual acuity outcomes after treatment. *Am J Ophthalmol.* 2005; 139 (6): 983–987.
- Ravindran RD, Venkatesh R, Chang DF, Sengupta S, Gyatsho J, Talwar B. Incidence of post-cataract endophthalmitis at Aravind Eye Hospital: outcomes of more than 42,000 consecutive cases using standardized sterilization and prophylaxis protocols. *J Cataract Refract Surg.* 2009; 35 (4): 629–636.
- Freeman EE, Roy-Gagnon M-H, Fortin E, Gauthier D, Popescu M, Boisjoly H. Rate of endophthalmitis after cataract surgery in Quebec, Canada, 1996–2005. *Arch Ophthalmol.* 2010; 128 (2): 230–234.
- Moshirfar M, Feiz V, Vitale AT, Wegelin JA, Basavanthappa S, Wolsey DH. Endophthalmitis after uncomplicated cataract surgery with the use of fourth-generation fluoroquinolones: a retrospective observational case series. *Ophthalmology.* 2007; 114 (4): 686–691.
- Jensen MK, Fiscella RG, Moshirfar M, Mooney B. Third- and fourth-generation fluoroquinolones: retrospective comparison of endophthalmitis after cataract surgery performed over 10 years. *J Cataract Refract Surg.* 2008; 34 (9): 1460–1467.
- Friling E, Lundström M, Stenevi U, Montan P. Six-year incidence of endophthalmitis after cataract surgery: Swedish national study. *J Cataract Refract Surg.* 2013; 39 (1): 15–21.
- Keay L, Gower EW, Cassard SD, Tielsch JM, Schein OD. Post-cataract surgery endophthalmitis in the United States: analysis of the complete 2003 to 2004 Medicare database of cataract surgeries. *Ophthalmology.* 2012; 119 (5): 914–922.
- Alharbi SS, Alrajhi A, Alkahtani E. Endophthalmitis following keratoplasty: incidence, microbial profile, visual and structural outcomes. *Ocul Immunol Inflamm.* 2013; 22 (3): 218–223.
- Taban M, Behrens A, Newcomb RL, Nobe MY, McDonnell PJ. Incidence of acute endophthalmitis following penetrating keratoplasty: a systematic review. *Arch Ophthalmol.* 2005; 123 (5): 605–609.
- Tay E, Bainbridge J, da Cruz L. Subretinal abscess after scleral buckling surgery: a rare risk of retinal surgery. *Can J Ophthalmol.* 2007; 42 (1): 141–142.
- Al-Torbak AA, Al-Shahwan S, Al-Jadaan I, AlHommedi A, Edward DP. Endophthalmitis associated with the Ahmed glaucoma valve implant. *Br J Ophthalmol.* 2005; 89 (4): 454–458.
- Endophthalmitis Vitrectomy Study Group Results of the endophthalmitis vitrectomy study: a randomized trial of immediate vitrectomy and of intravenous antibiotics for the treatment of postoperative bacterial endophthalmitis. *Arch Ophthalmol.* 1995; 113 (12): 1479–1496.
- Ozadamar A, Aras C, Ozturk R, Akin E, karacorlu M, Ercikan C. In vitro antimicrobial activity of silicone oil against endophthalmitis-causing agents. *Retina.* 1999; 19 (2): 122–126.

19. Azad R, Ravi K, Talwar D, Rajpal, Kumar N. Pars plana vitrectomy with and without silicone oil tamponade in post-traumatic endophthalmitis. *Graefe's Arch ClinExp Ophthalmol*. 2003; 241 (6): 478–483.

20. Do T, Hon D, Aung T, Hien ND, Cowan CL Jr. Bacterial endogenous endophthalmitis in Vietnam: a randomized controlled trial comparing vitrectomy with silicone oil versus vitrectomy alone. *Clin Ophthalmol*. 2014 Aug 28; 8: 1633-40.

21. Wong JS, Chan TK, Lee HM, et al. Endogenous bacterial endophthalmitis. *Ophthalmology*, 2000; 107 (8): 1483–1491

22. Jackson TL, Eykyn JS, Graham EM, Graham EM, Stanford MR. Endogenous bacterial endophthalmitis: a 17-year prospective study and review of 267 reported cases. *Surv Ophthalmol*. 2003; 48 (4): 403–423.

23. Yoon YH, Lee SU, Sohn JH, Lee SE. Result of early vitrectomy for endogenous *Klebsiella pneumoniae* endophthalmitis. *Retina*. 2003; 23 (3): 366–370.

24. Kuhn F, Gini G. Ten years after... are findings of the Endophthalmitis Vitrectomy Study still relevant today? *Graefe's Arch Clin Exp Ophthalmol*. 2005; 243 (12): 1197–1199.

25. Cowley M, Conway BP, Campochiaro PA, kaiser D, Gaskin H. Clinical risk factors for proliferative vitreo retinopathy. *Arch Ophthalmol*. 1989; 107 (8): 1147–1151

26. Cowley M, Conway BP, Campochiaro PA, kaiser D, Gaskin H. Clinical risk factors for proliferative vitreoretinopathy. *Arch Ophthalmol*. 1989; 107 (8): 1147–1151