

Petrographic and Geochemical Fingerprints of Sub-Volcanic Dykes and their Host Harzburgites from the Ulaş Ultramafics (Sivas, Turkey)

Özgür Bilici^{1*}, Hasan Kolaylı², Ekrem Kalkan³, Tuğba Bilici¹, Necmi Yarbaşı³

¹Ataturk University, Oltu Earth Sciences Faculty, Department of Petroleum and Natural Gas Engineering, 25400 Erzurum, Turkey

²Karadeniz Technical University, Engineering Faculty, Department of Geological Engineering, 25240 Erzurum, Turkey

³Ataturk University, Oltu Earth Sciences Faculty, Department of Geological Engineering, 25400 Erzurum, Turkey

INFORMATION

Article history

Received 18 February 2020

Revised 17 March 2020

Accepted 20 March 2020

Available 30 April 2020

Keywords

Peridotites

Sub-volcanic dyke

Subduction zone

Ulaş ultramafics

Sivas

Contact

*Özgür Bilici

ozgurbilici@atauni.edu.tr

ABSTRACT

In the southern Ulaş region (Sivas, mid-Anatolia), the ophiolitic fragments of the Neotethys oceanic lithosphere are widespread. The study area consists of commonly harzburgite, with minor dunite and chromitite pods and also pyroxenites. The latter are sub-volcanic gabbro and diabase dykes. In this study, the petrographic and geochemical importance of the mantle harzburgites and the sub-volcanic dykes cutting them were evaluated, within these ophiolitic units. The textural properties of the harzburgites were generally anhedral, granular and partially poikilitic. They consisted of olivine, orthopyroxene and minor amount of clinopyroxene and chromian spinel. On the other hand, the textural and mineralogical characteristics of sub-volcanic dykes, consisted of plagioclase, pyroxene, hornblende, opaque minerals and their secondary products, reflected semi-depth characters. The major, trace and rare earth element compositions indicated that the harzburgites were similar to that of both abyssal and supra-subduction (SSZ) peridotites with depleted character. Furthermore, the sub-volcanic dykes that formed the focus of this study in the mantle harzburgites exhibited N-MORB (Normal Mid-Ocean Ridge Basalt) type patterns of rare earth (REE) and trace elements. Especially, negative Nb anomaly showed the presence of a subduction component with the late stage modification of subduction-derived fluids/melts. All results indicated that the dykes characterized as N-MORB and late recorded the characteristics of Ocean Island Basalts (OIB). Consequently, geochemical features of the harzburgites and sub-volcanic dykes that cut them showed that peridotites were affected by two different tectonic setting conditions such as MORB and SSZ.

1. Introduction

Ophiolite suites represent ancient oceanic lithospheric remnants in continental orogenic belts and provide important records on the petrological evolution, metamorphism and tectonic processes of oceanic basins. In these ophiolite suites, sub-volcanic dyke intrusions such as micro-gabbro and diabase and also ultramafic pyroxenite dykes are quite common in mantle peridotites and are known as magmatic rocks produced by partial melting of mantle (Pearce et al., 1981).

In this context, petrological and geochemical studies on the world's best known ophiolites (Troodos, Oman, Anatolia, etc.) show that they offer similar features to the SSZ environment (Pearce et al., 1984; Parlak and Delaloye, 1996; Collins and Robertson, 1998; Dilek et al., 1999; Andrew and Robertson, 2002; Robertson, 2002; Parlak et al., 2006; Parlak et al., 2013; Robertson et al., 2013; Bilici, 2015; Parlak, 2016; Bilici and Kolaylı, 2018).

The Tauride ophiolite belt is represented by fragmented ophiolitic rocks embedded and/or cropped out in north of the Tauride platform (Şengör and Yılmaz, 1981; Parlak et al., 2006). In the Sivas region within this belt, the ophiolitic or related units are overlain by the Tauride carbonate platform. Many studies have been carried out about sub-volcanic dyke intrusions intruded the mantle peridotites of the Tauride ophiolites in Sivas province or in central Anatolia. Especially, the geochemical properties of these dykes display that they formed in a supra-subduction zone environment and point out their origin from an Island Arc Tholeiite (IAT) for Taurid ophiolites (Dilek et al., 1999; Çelik and Delaloye, 2003; Parlak et al., 2006; Parlak, 2016).

These features have been interpreted by previous researchers (Dilek et al., 1999; Shervais, 2001; Parlak et al., 2006) as the explosion of MORB-like or Ocean Island Basalt (OIB) lavas on the previously formed arc-related tholeiitic lavas in the final stage of magmatic activity above a supra-subduction zone environment. In this paper, we presented the whole rock

major, trace and rare earth element geochemistry of the sub-volcanic dykes and their host harzburgites to discussing their

petrogenetic significance and formation conditions within the related tectonic environment.




Fig. 1. Illustration of ophiolite belts and related tectonic map of Turkey with the major sutures and continental blocks of northeastern Mediterranean (modified after MTA (2002) and Okay and Tüysüz (1999)), (b) General geological map of the southern Sivas including the study area (Ulaş district) (modified after Yılmaz et al. (1989))




Fig. 2. Detailed geological map of the study area (Bilici, 2015)

2. Geological Summary

Turkey is located in a very important position in terms of tectonic emplacement and depending on the relative motions of continents have been exposed to intense orogenic activity. As a result of the closure of the Neotethys ocean, which started from the Cretaceous period, many basins were closed by the compressive regime and ophiolite settlements have been thrust onto the continents (Şengör and Yılmaz, 1981; Robertson and Dixon, 1984; Robertson et al., 2013; Parlak, 2016) (Fig. 1a). The remnants of these oceanic basins, which have an important place in understanding the geological evolution of Neotethys, are represented by ophiolites, ophiolite base metamorphics and ophiolitic melange in Anatolia (Dilek et al., 1999; Robertson, 2002). As a remnant basin, Sivas region, which also includes the study area, is located between Taurid and Pontid platforms (Cater et al., 1991; Poisson et al., 1996) and occurred during the closure of the northern branch of Neotethys (Yılmaz and Yılmaz, 2006). The ophiolitic units, previously called the Divriği ophiolite (Yılmaz et al., 2001), bear ultramafics mostly of Cretaceous age transported in soils that are ubiquitous and

stretch over hundreds of kilometres in Turkey, especially in the Sivas region (MTA, 2002).

This region was composed of autochthonous platform carbonates, ophiolitic mélangé, metamorphic sole, ophiolitic rocks, volcano-sedimentary units, granitoid rocks, and sediments (Yılmaz et al., 2001; Yılmaz and Yılmaz, 2004; Parlak, 2016) (Fig. 1b). The mantle peridotites observed in the study area (southeast Ulaş, Sivas) consist of harzburgites containing dunite lenses with chromitites. These widespread harzburgites were cut with different levels of sub-volcanic dykes such as diabase (partially gabbroic and doleritic) (Bilici, 2015; Bilici and Kolaylı, 2018) (Fig. 2).

3. Analytical Method

As a result of field studies, a series of samples were collected for geochemical studies, such as petrological characteristics and the tectonic environment of the peridotites and the sub-volcanic dykes that cut them. All samples were carefully cleaned and then ground in a special ring grinder. For this region, major oxide (ICP-ES method) and trace element

(ICP-MS method) analyses of total nine samples were performed in ACME Laboratories (Canada). The analytical indefiniteness was predicted to be 10% for trace elements with abundances of <10 ppm and approximately 5% for those with abundances of >10 ppm.

4. Results and Discussion

4.1. Field observation and petrography

In the study area located in the southeast of Ulaş district of Sivas province, the most common ultramafic lithology, the harzburgites, were systematically sampled. The harzburgites whose distant appearance were characteristic reddish colours and easily distinguished from other units, were observed as dark brown and green outcrops in some places (Fig. 3). On the other hands, during the field studies, a large number of sub-volcanic dykes were observed, which cut harzburgitic peridotites and lined up with a certain geometry. These dyke

masses were seen as outcrops up to a few meters in diameter which increased in number as they approached the areas where chromitite bodies (Fig. 3).

As a result of the microscopic investigations the harzburgite samples, it was determined that most samples underwent intensely serpentinization. The harzburgites, whose textural properties were generally anhedral, granular and partially poikilitic, also have been seen stockwork and cataclastic textures due to serpentinization (Fig. 4).

The harzburgites generally consisted of olivine (70-80%), orthopyroxene (15-20%) and to a lesser extent (about 1-5%) clinopyroxene and chromian spinel minerals in terms of modal mineral abundances. In addition, there were trace amounts of alteration minerals (serpentine, brucite, chlorite etc.) (Fig. 4).



Fig. 3. Field photos showing the lithological boundaries from the study area




Fig. 4. Photomicrographs of the harzburgite. a and b were taken cross-polarized light and plane polarized light, respectively (ol: olivine; opx: orthopyroxene; spl: chromian spinel; srp: serpentine)




Fig. 5. Photomicrographs of the sub-volcanic dykes. a and b were taken cross-polarized light and plane polarized light, respectively (plg: plagioclase; cpx: orthopyroxene; hrb: hornblende; op: opaque mineral)

All dyke samples collected from the study area were examined in detail under a polarizing microscope. Although the macro samples appear as quite fresh, it was found that they underwent intensely alteration in microscopic observation. Textures were generally observed as intersertal and to a lesser extent doleritic and also poikilitic (Fig. 5). Plagioclases, which were determined pervasively altered, have formed all of the light colored minerals. Altered pyroxenes and opaque minerals were seen, among the plagioclases generally formed as leaning against each other. Partially or fully chloritized and uralitized pyroxenes minerals were determined in the samples at a rate of about 10% in terms of modal composition (Fig. 5).

Moreover, most pyroxene crystals were determined to be augite, altered partially to actinolite and tremolite. Hornblende minerals along with augites, have also participated in a significant amount of modal mineralogy. As a result, it was determined that these dykes were in lithology ranging from micro-gabbro to diabase and even dolerite.

4.2. Geochemistry and tectonic implications

Whole-rock major, trace and rare earth element (REE) analyses were performed on four harzburgite and five sub-volcanic samples and results were given in Table 1. Firstly, LOI values were high more than 10 wt% because of the serpentinization of all harzburgite samples. According to the analysis results, the Al_2O_3 , CaO, Fe_2O_3 , TiO_2 , MnO, Cr_2O_3 contents (% by weight) of the harzburgites ranged between 0.43-1.87, 0.22-0.51, 7.03-8.56, 0-0.05, 0-0.1, 0.15-1.01, respectively. The harzburgite samples have relatively high MgO (average 35.14 wt%), whereas low Al_2O_3 (average 1.87 wt%) and alkalis ($\text{Na}_2\text{O} < 0.07$ wt%; $\text{K}_2\text{O} < 0.04$ wt%) contents.

These elemental abundances of the harzburgites were found to be relatively lower compared to sub-volcanics, and it was observed that these distributions (apart from MgO and Cr_2O_3 (%wt.) contents) showed positive orientations from the harzburgites to the sub-volcanics, according to SiO_2 contents

(Fig. 6a-j). This trend was an expected result, consistent with the observed modal mineralogical compositions. The MgO content in the mantle peridotite represents the consumption or partial melting degree of the mantle. It shows higher degree of the partial melting processes with high MgO contents and low CaO, Al_2O_3 and SiO_2 contents (Coleman, 1977; Hartmann and Wedepohl, 1993). In other words, when evaluated petrologically, incompatible components such as CaO, Al_2O_3 and SiO_2 tend to melt during the melting of a basic magma developing in the upper mantle. As the melting degree increases, mantle peridotites now become richer in MgO and the mantle becomes more depleted.

Here, this process was discussed on the $\text{Al}_2\text{O}_3/\text{SiO}_2$ vs. MgO/SiO_2 diagram (Fig. 7). The bulk compositions of the peridotites tend to run just below the mantle sequence, but in parallel (Hart and Zindler, 1986; Wu et al., 2017). This depletion trend may indicate the MgO loss or SiO_2 addition from the system during serpentinization. While the $\text{Al}_2\text{O}_3/\text{SiO}_2$ ratios are significantly reduced, MgO/SiO_2 ratios increase slightly in orthopyroxene and clinopyroxene during partial melting. As melting progresses, pyroxene, especially clinopyroxene, is depleted very quickly, and the modal content of olivine increases, resulting in an increase in the MgO/SiO_2 ratio. As the degree of partial melting increases, the bulk MgO/SiO_2 ratio increases. It is then understood that partial melting may not be the only reason that the observed bulk MgO/SiO_2 ratio is below the mantle sequence. Here, all of the analysed harzburgite samples have a slightly lower MgO/SiO_2 ratio than the mantle sequence, which means some loss of Mg due to alteration previously suggested by Niu (2004). Moreover, some samples may contain lower concentrations of major and trace elements due to the effect of this alteration or serpentinization.

In this paper, the MgO/SiO_2 and $\text{Al}_2\text{O}_3/\text{SiO}_2$ ratios of the harzburgites were much higher than primitive mantle (PM) despite small differences between samples and they had relatively high mantle depletion or partial melting (Fig. 7). When compared to the abyssal and fore-arc SSZ mantle

peridotites, the compositions of the Ulaş harzburgite samples were similar to that of abyssal peridotites, and two harzburgite samples fall in the fore-arc SSZ peridotites field or near (Fig. 7).

Table 1. Representative major oxide (wt %), trace (ppm) and rare earth (ppm) element composition of the harzburgite and sub-volcanic samples

Lithology	HARZ	HARZ	HARZ	HARZ	SVLC	SVLC	SVLC	SVLC	SVLC
Sample	Y-1	Y-6	AT-12	AT-13	S-8	S-15	S-21	AT-8	AT-9
SiO ₂	37.86	39.92	40.11	39.84	48.50	48.58	47.03	49.98	51.46
Al ₂ O ₃	0.59	0.43	1.18	1.87	15.61	14.91	15.76	15.74	15.47
Fe ₂ O ₃	7.17	7.03	8.54	8.56	10.68	9.28	8.95	9.70	10.66
MgO	37.11	34.27	36.09	33.10	5.80	7.37	8.12	7.15	6.98
CaO	0.22	0.23	0.51	0.34	9.79	9.60	10.91	10.60	9.58
Na ₂ O	0.01	0.00	0.07	0.06	5.11	4.62	3.38	3.54	2.88
K ₂ O	0.00	0.00	0.04	0.02	0.29	0.21	0.87	0.66	0.65
TiO ₂	0.00	0.00	0.05	0.04	1.11	1.00	0.88	1.01	1.03
P ₂ O ₅	0.00	0.00	0.06	0.04	0.07	0.08	0.06	0.12	0.14
MnO	0.10	0.10	0.10	0.07	0.17	0.15	0.15	0.17	0.18
Cr ₂ O ₃	0.31	0.35	0.16	1.01	0.01	0.03	0.04	0.02	0.01
LOI	15.80	16.80	12.55	12.10	2.70	3.90	3.60	2.02	1.04
Total	99.17	99.13	99.46	96.98	99.84	99.73	99.75	100.71	100.08
Ni	1933.00	2031.00	1682.00	2680.00	38.00	75.00	103.00	43.00	36.00
Sc	8.00	7.00	24.00	13.00	35.00	36.00	37.00	48.00	41.00
Ba	2.00	0.00	7.00	4.10	24.00	33.00	93.00	23.00	15.00
Be	0.00	0.00	0.00	0.00	0.00	0.00	1.00	0.00	0.00
Co	95.30	92.20	104.80	113.60	35.80	35.70	38.20	41.00	56.00
Cs	0.00	0.00	0.00	0.00	0.20	0.50	2.50	0.00	0.00
Ga	0.00	0.00	1.60	2.80	15.60	13.70	13.50	12.00	15.00
Hf	0.00	0.00	0.00	0.00	1.90	1.70	1.50	4.00	1.00
Nb	0.00	0.00	0.50	0.20	0.20	0.60	0.40	3.00	0.90
Rb	0.20	0.20	0.30	0.00	3.90	4.60	17.80	4.00	6.20
Sn	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Sr	1.10	2.80	4.70	3.20	103.00	250.20	138.00	34.00	156.00
Ta	0.00	0.00	0.00	0.00	0.10	0.30	0.20	0.20	0.10
Th	0.00	0.00	0.00	0.00	0.50	0.10	0.20	0.10	0.10
U	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
V	38.00	28.00	0.00	0.00	345.00	267.00	266.00	330.00	301.00
W	0.00	0.00	1.10	0.80	0.00	0.00	0.00	0.00	0.00
Zr	0.10	0.00	1.90	1.70	58.10	58.10	48.50	58.00	61.00
Y	0.00	0.00	0.10	0.60	25.20	21.50	20.10	25.00	23.00
La	0.30	0.10	0.50	0.00	2.50	2.30	2.10	3.98	2.32
Ce	0.10	0.00	0.90	0.50	7.20	7.20	5.80	10.36	7.45
Pr	0.00	0.01	0.09	0.06	1.24	1.26	1.08	1.54	1.25
Nd	0.02	0.01	0.00	0.00	6.50	6.80	5.90	7.89	4.01
Sm	0.01	0.00	0.00	0.00	2.38	2.21	2.05	2.55	2.33
Eu	0.01	0.03	0.04	0.03	0.95	0.88	0.83	0.98	0.79
Gd	0.00	0.12	0.07	0.07	3.37	3.25	2.80	3.12	3.39
Tb	0.01	0.00	0.01	0.01	0.66	0.63	0.53	0.55	0.58
Dy	0.00	0.02	0.06	0.09	4.27	3.94	3.37	3.78	4.17
Ho	0.03	0.00	0.02	0.02	0.98	0.78	0.80	0.84	0.98
Er	0.04	0.02	0.05	0.07	2.82	2.49	2.25	2.34	2.68
Tm	0.01	0.00	0.02	0.03	0.42	0.36	0.33	0.35	0.41
Yb	0.06	0.01	0.06	0.09	2.58	2.35	2.12	2.33	2.64
Lu	0.02	0.00	0.01	0.03	0.43	0.37	0.33	0.34	0.39
ΣREE	0.61	0.32	1.83	1.00	36.30	34.82	30.29	40.95	33.39

(HARZ: harzburgite, SVLC: sub-volcanic)




Fig. 6. Whole rock major element variation diagrams of the Ulaş harzburgites and sub-volcanic dykes




Fig. 7. Whole rock major element distribution of harzburgites in MgO/SiO₂ and Al₂O₃/SiO₂ exchange diagram (after Wu et al. (2017))




Fig. 8. REE and trace element distributions in whole rocks of the harzburgites; (a) the REE patterns (b) trace elements spider diagram (chondrite and primitive mantle normalizing values were taken from Sun and McDonough (1989)). Abyssal and fore-arc peridotite fields were taken from Niu et al. (1997) and Parkinson and Pearce (1998), respectively

The harzburgite samples have very low REE compositions, and their ΣREE contents range between 0.324–1.83 (ppm) (Table 1). This indicates that the harzburgites exhibited significant depletion as a result of high and/or moderate degrees of partial melting (Miller et al., 2003). The chondrite normalized REE patterns and primitive mantle normalized multi-element distributions were illustrated in Fig. 8. Although the low REE contents varied not considerably, their distribution patterns were very significant (Fig. 8a). Especially, two of the samples were slightly enriched in light REE. The REE pattern of the Ulaş harzburgites was distinctively different from the REE-depleted pattern of the fore-arc SSZ peridotite with low total REE contents that were quite depleted.

They showed similarity with the abyssal and SSZ peridotites (Fig. 8a). However, this light REE enrichment feature has widely been interpreted as a penetration of subduction-derived melts or fluids (Wu et al., 2017). Here, our data were

not exactly matched with the modelled SSZ or abyssal REE patterns, but they were more compatible with light REE enrichment produced by melt-peridotite interactions beneath a MOR-type setting. Previous researchers suggested that this process originated in the cold thermal boundary layer, through which melts react with an earlier residual mantle, resulting in both light REE and high field strength element (HFSE) enrichments in abyssal peridotites (Niu, 2004). However, this light REE (Fig. 8a) and HFSE (Fig. 8b) enrichments showed by Ulaş harzburgites might have been a result of their interaction with SSZ melts (Dilek and Furnes, 2014). Especially, the primitive mantle-normalized multi-element diagram of the harzburgites exhibited right dipping patterns (Fig. 8b). The large ion-lithophile elements (LILE) enrichments might be due to crustal fluid interaction of the subduction zone. Moreover, high field strength elements exhibit both depletion of Nb, high REE enrichment indicating the depleted mantle source and different degrees of metasomatic processes above a subduction zone.




Fig. 9. Zr/ TiO₂ vs Nb/Y classification diagram for sub-volcanic rocks cutting the harzburgites (Floyd and Winchester 1978)




Fig. 10. Chondrite normalized REE patterns (a) and N-MORB normalized multi-element patterns (b) for the sub-volcanic dykes in the Ulaş ultramafics (normalizing values were taken from Sun and McDonough (1989))




Fig. 11. Tectonic discrimination diagrams for Ulaş sub-volcanic dykes cutting the mantle harzburgites: a) Diagram of Hf/3 vs. Th vs. Ta (suggested by Wood (1980)); b) Diagram of Ta/Yb vs. Th/Yb (suggested by Pearce et al. (1984)). MORB: Mid-Ridge Basalt, WPB: Within Plate Basalts, CAB: Calc-alkaline Basalt, SH: Shoshonite, TH: tholeiite, TR: transition basalt, IAT: Island-Arc Tholeiites




Fig. 12. Tectonic discrimination diagrams for Ulaş sub-volcanic dykes cutting the mantle harzburgites; a) Diagram of Nb/Yb vs. TiO_2/Yb (after Pearce (2008)), b) Diagram of Zr vs. Zr/Y (suggested by Pearce and Norry (1979)). OIB, N-MORB and E-MORB were taken from Sun and McDonough (1989).

Primitive mantle-normalized trace element patterns of the harzburgites showed that they were highly depleted in terms of lithophile elements, with concentrations well below those of abyssal peridotites (Fig. 8b). The sub-volcanic dykes cutting the mantle harzburgites in the Ulaş ultramafic massif showed moderate alteration according to the geochemical results and they had up to 4 wt % LOI values (Table 1). In the Zr/TiO₂ vs. Nb/Y diagram suggested by Floyd and Winchester (1978), these sub-volcanic dykes plotted in the andesite/basalt field with high values of TiO₂ (0.88 to 1.11 wt %), Zr (48.5 to 61 ppm), Nb (0.20 to 3 ppm) and Y (20.10 to 25.20 ppm) (Fig. 9). The chondrite-normalized REE patterns and N-MORB normalized multi-element distributions of the sub-volcanic dykes were illustrated in Fig. 10a-b.

It was observed that there was a similar distribution with N-MORB in REE contents and slightly depletion in terms of incompatible light REEs (Fig. 10a). This is compatible with the REE contents of other diabase dykes in the Taurid belt ophiolites (Parlak and Delaloye, 1996; Çelik and Delaloye, 2003; Parlak et al., 2006). Furthermore, due to these REEs indicated that the Ulaş sub-volcanic dykes were similar as N-MORB, which formed in a MORB-like setting and were mounted by the features of OIB. The N-MORB normalized multi-element distributions of sub-volcanic dykes exhibited enrichment patterns (Fig. 10b). In particular, consumption of Nb in these dykes and their enrichment in large-ion lithophile elements (LILE) such as Sc, Rb, Ba, U and K₂O even offered similarity with the alkali melts changed in the subduction zone. This situation showed similarity with the OIBs suggested by Parlak et al. (2006). The hornblende minerals in the modal mineralogy of the sub-volcanic dykes can also be cited as evidence.

In terms of determining the tectonic formation environments of ophiolites, some elements such as Hf, Ti, Zr, Y, Nb and Sr are distinctive trace elements for volcanic and/or sub-volcanic rocks (Pearce and Cann, 1973; Xu et al., 2019). Here, on the Hf/3-Th-Ta ternary diagram (Fig. 11a), the sub-

volcanic dykes plotted within the MORB area and on the Ta/Yb vs. Th/Yb diagram (Fig. 11b) most of these dyke samples plotted in the N-MORB area except for two samples. One of these two samples plotted in the calc-alkali basalt (CAB) field and other plotted in enriched mid-ocean ridge basalt (E-MORB) field.

In the Nb/Yb vs. TiO₂/Yb diagram, the Ulaş sub-volcanic dykes plotted close to the N-MORB field with low Nb/Yb ratios (Fig. 12a), but one sample fell in the E-MORB field. This situation indicated that the source of the dykes was more depleted than the N-MORB source. On other hand, in the Zr vs. Zr/Y diagram all samples showed similarity in each other and plotted in the intersection field of the MORB and volcanic arc basalt (VAB) fields (Fig. 12b).

5. Concluding Remarks

The Ulaş ultramafic massif as a part of the Divriği ophiolite (Sivas) consists mainly of harzburgite, with minor dunite-chromitite and pyroxenite, suggested the involvement of mantle material. Based on field observations, the results in this study indicated that these harzburgites cutted by sub-volcanic dykes may source from deeper mantle. The composition of the Ulaş harzburgites were similar to that of both abyssal and the SSZ peridotites with depleted character. The Ulaş sub-volcanic dykes in the mantle harzburgites exhibited N-MORB-type patterns of REEs and trace elements. These patterns showed that they were slightly depleted in highly incompatible elements (light REEs) with respect to less incompatible elements (heavy REEs). Especially, pronounced negative Nb anomalies, suggesting the presence of a subduction component. All results indicated that the dykes characterized as N-MORB and late recorded the characteristics of OIBs. The presence of hornblende crystals in these sub-volcanic units showed the late stage modification of subduction-derived fluids. All these features of the studied harzburgites and sub-volcanic dykes suggested that the Ulaş ultramafic section of the Divriği ophiolite might have been formed in two different tectonic settings (MORB and SSZ). These data imply that the ophiolites in the north

of Tauride belt were characteristic of the two stage evolution. First, the development of harzburgitic peridotites with moderate-degree of the partial melting, from the mid-ocean ridges to the subduction zone. The second was the interaction of these peridotites reaching the subduction zone with MORB-like melt, which occurs with different angle and/or breaking of the subducted lithosphere plate.

Acknowledgements

This paper was prepared as a part of the PhD thesis performed by the first author. The authors are grateful to two anonymous referees for their very valuable contributions to improving of the manuscript.

References

- Andrew, T., Robertson, A.H.F., 2002. The Beyflehir-Hoyran-Hadim nappes: genesis and emplacement of Mesozoic marginal and oceanic units of the northern Neotethys in southern Turkey. *Journal of the Geological Society* 159, 529-543.
- Bilici, Ö., 2015. Comparative investigation of Kop (Erzurum-Erzincan-Bayburt), Ulaş (Sivas) and Yeşilova (Burdur) ultramafics and chromitites in terms of mineralogical, petrological and geodynamical aspects. PhD, Karadeniz Technical University, Trabzon, Turkey (in Turkish with English Abstract).
- Bilici, Ö., Kolaylı, H., 2018. Mineral records of the pyroxenites formed within harzburgites (Ulaş, Sivas, Turkey): implications on petrogenesis and tectonic setting. *Turkish Journal of Earth Science* 27, 384-404.
- Cater J.M.L., Hanna, S.S., Ries, A.C., Turner, P., 1991. Tertiary evolution of the Sivas Basin, Central Turkey. *Tectonophysics*, 195, 29-46.
- Coleman, R.G., 1977. Ophiolites, Ancient Oceanic Lithosphere. Berlin: Springer-Verlag, 2-10.
- Collins, A.S., Robertson, A.H.F. 1998. Processes of Late Cretaceous to Late Miocene episodic thrust-sheet translation in the Lycian Taurides, SW Turkey. *Journal of Geological Society* 155, 759-772.
- Çelik, Ö.F., Delaloye, M., 2003. Origin of metamorphic soles and their post-kinematic mafic dyke swarms in the Antalya and Lycian ophiolites, SW Turkey. *Geological Journal* 38 235-256.
- Dilek, Y., Furnes, H., 2014. Ophiolites and their origins. *Elements* 10, 93-100.
- Dilek, Y., Thy, P., Hacker, B., Grundvig, S., 1999. Structure and petrology of Tauride ophiolites and mafic dyke intrusions (Turkey): implications for the Neotethyan ocean. *Geological Society of America Bulletin* 111, 1192-1216.
- Floyd, P.A., Winchester, J.A., 1978. Identification and discrimination of altered and metamorphosed volcanic rocks using immobile elements. *Chemical Geology* 21, 291-306.
- Hart, S.R., Zindler, A., 1986. In search of a bulk-Earth composition. *Chemical Geology* 57, 247-267.
- Hartmann, G., Wedepohl, K.H., 1993. The composition of peridotite tectonics from the Ivrea complex, northern Italy, residues from melt extraction. *Geochimica et Cosmochimica Acta* 57, 1761-1782.
- MTA, 2002. 1:500,000 Scale Geology Map of Turkey. General Directorate of Mineral Research and Exploration, Ankara, Turkey.
- Miller, C., Thöni, M., Frank, W., Schuster, R., Melcher, F., Meisel, T., Zanetti, A., 2003. Geochemistry and tectonomagmatic affinity of the Yungbwa ophiolite, SW Tibet. *Lithos* 66, 155-172.
- Niu, Y., 2004. Bulk-rock major and trace element compositions of abyssal peridotites: implications for mantle melting, melt extraction and post-melting processes beneath mid-ocean ridges. *Journal of Petrology* 45, 2423-2458.
- Niu, Y.L., Langmuir, C.H., Kinzler, R.J., 1997. The origin of abyssal peridotites: A new perspective. *Earth and Planetary Science Letters* 152(1-4), 251-265.
- Okay, I.A., Tüysüz, O., 1999. Tethyan sutures of northern Turkey. *Journal of Geological Society, London Special Publication* 156, 475-515.
- Parkinson, I.J., Pearce, J.A., 1998. Peridotites from the Izu-Bonin-Mariana forearc (ODP Leg 125), evidence for mantle melting, melt-mantle interaction in a suprasubduction zone setting. *Journal of Petrology* 39, 1577-1618.
- Parlak, O., 2016. The Tauride ophiolites of Anatolia (Turkey): a review. *Journal of Earth Science* 27, 901-934.
- Parlak, O., Çolakoğlu, A., Dönmez, C., Sayak, H., Yıdırım, N., Türkel, A., Odabaşı, I., 2013. Geochemistry and tectonic setting of ophiolites along the İzmir-Ankara-Erzincan Suture Zone in northeastern Anatolia. In A.H.F. Robertson, O. Parlak, U.C., Ünlügöç (Eds.), *Geological development of Anatolia and the easternmost Mediterranean*. London: Geological Society, Special Publications 372, 75-106.
- Parlak, O., Delaloye, M., 1996. Geochemistry and timing of post metamorphic dike emplacement in the Mersin ophiolite (southern Turkey): new age constraints from ⁴⁰Ar/³⁹Ar geochronology. *Terra Nova* 8, 585-592.
- Parlak, O., Yılmaz, H., Boztuğ, D., 2006. Origin and tectonic significance of the metamorphic sole and isolated dykes of the Divriği ophiolite (Sivas, Turkey): evidence for slab break-off prior to ophiolite emplacement. *Turkish Journal of Earth Science* 15, 25-45.
- Pearce, J.A., 2008. Geochemical fingerprinting of oceanic basalts with applications to ophiolite classification and the search for Archean oceanic crust. *Lithos* 100, 14-48.
- Pearce, J.A., Alabaster, T., Shelton, A.W., Searle M.P., 1981. The Oman Ophiolite as a Cretaceous Arc-basin Complex, Evidence and Implications. *Philosophical Transactions of the Royal Society a Mathematical Physical and Engineering Sciences* 300, 299-317.
- Pearce, J.A., Cann, J.R., 1973. Tectonic setting of basic volcanic rocks determined using trace element analyses. *Earth and Planetary Science Letters* 19, 290-300.
- Pearce, J.A., Lippard, S.J., Roberts, S., 1984. Characteristics and tectonic significance of suprasubduction zone ophiolites. In: Kokelaar, B.P., Howells, M.F. (eds), *Marginal Basin Geology*. Geological Society, London, Special Publications 16, 77-94.
- Pearce, J.A., Norry, M.J., 1979. Petrogenetic implications of Ti, Zr, Y and Nb variations in volcanic rocks. *Contributions to Mineralogy and Petrology* 69, 33-47.
- Poisson, A., Guezou, J.C., Öztürk, A., İnan, S., Temiz, H., Gürsöy, H., Kavak, K.S., Özden, S., 1996. Tectonic setting and evolution of the Sivas Basin, Central Anatolia, Turkey. *International Geology Review* 38, 838-853.
- Robertson, A.H.F., 2002. Overview of the genesis and emplacement of Mesozoic ophiolites in the eastern Mediterranean Tethyan region. *Lithos* 65, 1-67.
- Robertson, A.H.F., Dixon, J.E., 1984. Introduction: aspects of the geological evolution of the Eastern Mediterranean. In: Dixon, J.E., Robertson, A.H.F. (eds), *The Geological Evolution of the Eastern Mediterranean*. Geological Society, Special Publications 17, 1-74.
- Robertson, A., Parlak, O., Ustaömer, T., Tash, K., İnan, N., Dumitrica, P., Karaoglan, F., 2013. Subduction, ophiolite genesis and collision history of Tethys adjacent to the Eurasian continental margin: new evidence from the Eastern Pontides, Turkey. *Geodinamica Acta* 26, 230-293.
- Şengör, A.M.C., Yılmaz, Y., 1981. Tethyan evolution of Turkey: a plate tectonic approach. *Tectonophysics* 75, 181-241.
- Shervais, J.W. 2001. Birth, death and resurrection: The life cycle of supra-subduction zone ophiolites. *Geochemistry Geophysics Geosystems* 2, 1-45.

- Sun, S.S., McDonough, W.F., 1989. Chemical and isotopic systematics of oceanic basalts: implications for mantle compositions and processes. In: Saunders, A.D., Norry, M.J. (eds.), *Magmatism in the Ocean Basins*. London: Geological Society, Special Publications 42, 313-345.
- Wood, D.A., 1980. The applications of a Th-Hf-Ta diagram to problems of tectonomagmatic classification and to establishing the nature of crustal contamination of basaltic lavas of the British Tertiary volcanic province. *Earth and Planetary Science Letters* 50, 11-30.
- Wu, W., Yang, J., Dilek, Y., Milushi, I., Lian, D., 2017. Multiple episodes of melting, depletion, and enrichment of the Tethyan mantle: Petrogenesis of the peridotites and chromitites in the Jurassic Skenderbeu massif, Mirdita ophiolite, Albania. *Lithosphere* 10(1), 54-78.
- Xu, X., Yang, J., Xiong, F., Guo, G., 2019. Petrology and Geochemistry of the Dangqiong Ophiolite, Western Yarlung-Zangbo Suture Zone, Tibet, China. *Acta Geologica Sinica (English Edition)* 93(2), 344-361.
- Yılmaz, H., Arıkal, T., Yılmaz, A., 2001. Geology of the Güneş ophiolite (Divriği-Sivas). *Proceedings of the 54th Geological Congress of Turkey*, Ankara, Turkey, pp. 54-65 (in Turkish with an abstract in English).
- Yılmaz, A., Sümengen, M., Terlemez, I., Bilgiç, T., 1989. 1:100000 Scale Geology Map of Turkey, Sivas G23. General Directorate of Mineral Research and Exploration, Ankara, Turkey.
- Yılmaz, A., Yılmaz, H., 2006. Characteristic features and structural evolution of a post collisional basin: The Sivas Basin, Central Anatolia, Turkey. *Journal of Asian Earth Science* 27, 164-176.
- Yılmaz, H., Yılmaz, A., 2004. Geology and Structural Evolution of the Divriği (Sivas) Region. *Geological Bulletin of Turkey* 47, 13-45.