

Bowen's Disease: An Uncommon Presentation

Nihal Kılınç*, Özden Zeki Pınar**

*Dicle University Medical Faculty, Department of Pathology, Diyarbakır, Turkey

**Diyarbakır Government Hospital, Department of Dermatology, Diyarbakır, Turkey

Objective: Bowen's disease described by Bowen originally in 1912 is an intraepidermal squamous cell carcinoma. We aimed to discuss a case of Bowen's disease with uncommon presentation in the light of recent literature.

Observation a 65-year-old male presented with a lesion over the skin of the abdomen for 5-6 years duration. The lesion gradually enlarged over the years and was asymptomatic. It had been treated and misdiagnosed by different centers, which had resulted in partial regression; however, it started to enlarge again. Dermatologic examination revealed a plaque, 7cm×7cm in size, on the right side of the umbilicus. It was well defined, and erythematous, pigmented, with firm papules of various sizes at the periphery and center of the lesion. Histopathology showed that the lesion was extending through the whole thickness of the epidermis. The basement membrane zone was intact. Keratinocytes showed a loss of polarity and atypia. A diagnosis of "Bowen's disease" was made, the lesion was surgically excised, and skin grafting was performed.

Conclusion: Bowen's disease should be considered when a skin lesion not responded to treatment or lesion mimics papulosquamous disorders.

Key words: Bowen's Disease, Skin, Dermatopathology

Nadir Yerleşimli Bir Bowen Hastalığı

Amaç: İlk olarak 1912'de Bowen tarafından tanımlanan Bowen hastalığı bir intraepidermal yassı hücreli karsinomdur.

Olgu: 65 yaşında erkek hasta 5-6 yıldan beri karın cildinde bulunan lezyon ile başvurdu. Bu asemptomatik deri lezyonu aylar içinde ilerlemiş. Değişik merkezlerde yanlış tanı alan ve tedavi edilen lezyonda kısmi bir gerileme olmuş ancak tekrar büyümeye başlamış. Dermatolojik muayenesinde; göbeğin sağ tarafında, 7cm x 7cm boyutlarında, plak şeklindeki izlenen lezyonun iyi sınırlı, eritemli, pigmentli ve değişik büyüklükte papüller yapıların lezyonun periferine ve merkezine yerleşmiş olduğu izlendi. Histopatolojik muayenesinde; epidermiste kalınlaşma, bazal tabakanın sağlam olduğu, keratinositlerde polarite kaybı ile atipi izlendi. Bowen hastalığı tanısı alan lezyon cerrahi olarak çıkarıldı ve deri grefti konuldu.

Sonuç: Nadir yerleşimli Bowen hastalığı olgusunu ilgili kaynakların ışığında tartışarak sunmayı amaçladık.

Anahtar Kelimeler: Bowen Hastalığı, Cilt, Dermatopatoloji.


In 1912 and 1915, John T. Bowen described 3 patients with precancerous dermatosis.^{1,2} Darier first called the lesion "Bowen's Disease" in 1914.³ It represents a form of squamous cell carcinoma in situ that clinically presents an asymptomatic, elevated and crusted plaque on the skin.⁴ The general histologic features consist of a full thickness loss of epidermal stratification (windblown appearance), great variability of nuclear size and depth of staining, occurrence of mitoses in all layers of living epidermis, and presence of atypical cells, some with multiple nuclei and dyskeratotic cells.⁵ Bowen's disease usually consists of a solitary lesion. It may be caused by exposure to the sunlight, x-rays or ionizing radiation, and arsenic.⁶

CASE

A 65-year-old man was admitted to Diyarbakır Central Hospital Dermatology Polyclinic with complaints of an asymptomatic, crusted plaque over the skin located on the right side of the umbilicus persisting for 5-6 years. The lesion gradually increased in size over the years and was asymptomatic. Psoriasis, nummular dermatitis, tinea corporis were diagnosed and treated at different centers, which had resulted in partial regression; however, it started to grow

again. There was no preceding history of trauma, cold injury, or similar lesion elsewhere on the body. General examination revealed normal. Dermatological examination revealed an erythematous, yellow-white crusted plaque of about 7x7 cm, with well defined borders, lying over the skin of the right side of the umbilicus. Noteworthy was the presence of keratotic and pigmented papules, and a focal pigmented border. Signs of atrophy, bleeding or lymphadenopathy could not be observed (Figure 1). There were no other similar lesions on other parts of body. Routine laboratory investigations were normal. A differential diagnosis of pagetoid type of basal cell carcinoma, squamous cell carcinoma, Bowen's disease and superficial spreading melanoma were considered.

Figure 1. There is a crusted plaque on the right side of the umbilicus.



Histologic examination of the plaque showed that the epidermis was thickened, and there was hyperkeratosis and acanthosis with irregular elongation of the rete ridges. The border between the epidermis and dermis was sharp. The basal layer was invaded by chronic inflammatory infiltrate (Figure 2). The cells of the stratum malpighii were in complete disorder. Some of them were atypical with hyperchromatic nuclei, and a few others showed individual cell keratinization or dyskeratosis (Figure 3). Histopathologic examination of the plaque revealed Bowen's disease. The lesion was surgically excised, and free skin grafting was performed.

DISCUSSION

The morphology of Bowen's disease mimics other cutaneous disorders like psoriasis, lichen simplex chronicus, and other papulo-squamous disorders.⁷ The disease itself is easily treated and has an excellent prognosis.⁸ If untreated, 3% to 5% of patients may

develop invasive carcinoma.⁹ Histopathological classification of Bowen's disease is uncommon. Darier² classified the lesion into five types (lenticular, discoid, simple hyperkeratotic, invasive carcinoma). In 1980, Strayer and Santa Cruz¹⁰ modified this classification and determined six groups (psoriasiform, atrophic, verrucous-hyperkeratotic, irregular, nesting pattern, and other, including amyloid formation and mucinous or sebaceous metaplasia, and combinations of two or more types).

Figure 2. There is hyperkeratosis, acanthosis and a bandlike inflammatory infiltrate with irregular elongation of the rete ridges. The basement membrane zone is intact (HEX100).





Figure 3. High magnification of figure 2. There are cells within epidermis showing atypical hyperchromatic nuclei and dyskeratosis (HEX200).



In the literature, there are some studies analyzing the age, sex and site distribution of Bowen's disease. In one of the studies, 68% of patients were males. Higher prevalence was reported in elderly in people, over the age of sixty.¹⁰ Reymann et al¹² reported that 82% of 581 patients with Bowen's disease were over the age of 60. Our patient was male and over 60 years of age as well.

Kettler et al¹³ demonstrated the presence of HPV in a lesion of nongenital Bowen's disease. The anatomic distribution of Bowen's disease, similar to that of non-melanoma skin cancers, occurs on sun-exposed skin areas (head, neck, hands); half of all lesions occur on the head (44% to 54%).⁷ Kossard et al¹⁴ found marked female predominance of Bowen's disease located on the cheeks and suggested that superficially placed vellus hair follicles on the cheeks of women might be more susceptible to chronic exposure to sun.

Bowen's disease has also been reported to arise from a scar tissue following trauma. The pathogenesis of malignancy in chronic scars and in traumatized skin is unknown. As scars are poorly vascularized, they are susceptible to ulceration and infection and tissue toxin may act as carcinogens. In cases of repeated damage, repeated proliferative repair can cause carcinoma.¹⁵ In our patient, there was no history of trauma.

Typical Bowen's disease presents a gradually enlarging well demarcated erythematous plaque with an irregular border and surface crusting or scaling. The other peculiar feature is the pigmented border. Bowen's disease has been reported to arise from a pigmented seborrheic keratosis.¹⁶ In the present case there was a scattered pigmented border. In addition there is a pigmented type of Bowen's disease, which should be considered in the differential diagnosis of malignant melanoma.¹⁷

There are some immunohistochemical studies about Bowen's disease in the literature. In a recent study, the expressions of CD40 and FasL in normal skin, squamous cell carcinoma (SCC) and Bowen's disease were investigated immunohistochemically. Tae Jung Jang,¹⁸ suggested that FasL is more strongly expressed in Bowen's disease when compared to SCC, but expression of CD40 was higher in SCC than Bowen's disease.

The paraneoplastic nature of Bowen's disease was first considered when Graham and Helwing in 1959 reported that 80% of the patients had internal malignancy. Several studies initially confirmed this observation.^{1,16} But more recently, carefully

controlled studies have shown that the disease has no clear paraneoplastic association.^{11,19}

The diagnosis of Bowen's disease should be considered and a skin biopsy should be performed when a patient admits with a persistent cutaneous lesion that is of uncertain diagnosis or if an atypical dermatitis does not respond to steroid therapy. Carbon dioxide laser has been used for the treatment of Bowen's disease in one of the most recent studies.²⁰ In our case, the lesion was surgically excised, and free skin grafting was performed.

Finally, Bowen's disease should be considered when a skin lesion does not respond to treatment or lesion mimics papulosquamous disorders.

REFERENCES

1. Bowen JT. Precancerous dermatoses: A study of two cases of chronic typical epithelial proliferation, *Journal of Cutaneous Diseases*, 1912;30:242-55.
2. Bowen JT. Precancerous dermatoses: A sixth case of a type recently described, *Journal of Cutaneous Disease*, 1915;33:787-802.
3. Darier J. La dermatose precanceruse de Bowen-dyskeratose lenticulaire et en disques, *Annal de dermatologie* 1914;5:449-71.
4. Sarveswari KN. Bowen's disease of the palm. *Int J Dermatol* 1998; 37:157-58.
5. Mehregan AH, Hashimoto K. *Pincus' Guide to Dermatohistopathology*. 5th ed. London: Appleton&Lange 1991:507-9.
6. Yerebakan O, Ermis O, Yilmaz E, ve ark. Treatment of arsenical keratosis and Bowen's disease with acitretin. *Int J Dermatol* 2002; 41:84-7.
7. Marshall SF, Ronan SG, Massa MC. Pigmented Bowen's disease arising from pigmented seborrheic keratose. *J Am Acad Dermatol* 1990;23:440-44.
8. Lee M, Wick MM. Bowen's disease. *Clin Dermatol* 1993;11:43-6.
9. Mitsuishi T, Kawashima M, Matsukura T, et. al. Human papillomavirus type 58 in Bowen's disease of the elbow. *Br J Dermatol* 2001;144:384-6.
10. Strayer DS, Santa Cruz DJ. Carcinoma in situ of the skin: A review of histopathology. *J Cutan Pathol* 1980;7:244.
11. Miki Y, Kawatsu T, Matsuda K, et. al. Cutaneous and pulmonary cancers associated with Bowen's disease. *J Am Acad Dermatol*. 1982;6:26-31.
12. Reymann F, Ravnborg L, Schou G, et. al. Bowen's disease and internal malignant diseases, A study of 581 patients. *Arch Dermatol* 1988; 124:677-79.
13. Kettler AH, Rutledge M, Tschien JA, et. al. Detection of human papillomavirus in nongenital Bowen's disease by in situ hybridization. *Arch Dermatol* 1990; 126:777-81.
14. Kovacs A, Yonemoto K, Katsuoka K, Nishiyama S, Harhai I. Bowen's disease: Statistical study of a year period. *J Dermatol* 1996; 23: 267-74.
15. Eedy DJ, Gavin AT. Thirteen-year retrospective study of Bowen's disease in North Ireland. *Br J Dermatol* 1987;68:715-20.
16. Köse O, Safalı M, Taştan MS, ve ark. Genital yerleşimli Bowen's hastalığı. *Türk J Dermatopathol* 2000; 9:118-20.
17. Krishnan R, Lewis A, Orengo IF, et. al. Pigmented Bowen's disease (squamous cell carcinoma in situ): a mimic of malignant melanoma. *Dermatol Surg* 2001; 27:673-4.
18. Tae Jung Jang. Expression of CD40 and Fas Ligand in Bowen's Disease, Squamous cell carcinoma and Basal cell carcinoma *Yonsei Med J* 2002; 43:304-308.
19. Chuang T, Reizner GT. Bowen's disease and internal malignancy, a matched case-control studies. *J Am Acad Dermatol* 1988; 19:47-51.
20. Dave R, Monk B, Mahaffey P. Treatment of Bowen's disease with carbon dioxide laser. *Lasers Surg Med* 2003; 32:335

Corresponding Address:

Yrd. Doç. Dr. Nihal KILINÇ
Dicle Üniversitesi Tıp Fakültesi
Patoloji AD. 21280-Diyarbakır
Tel : 90 412 248 8001/4227 internal
90 412 2488114
Mobile :90 535 3910302
E-Posta :nhl@dicle.edu.tr