



New Desmid Records of Karagöl Lake in Karagöl-Sahara National Park (Şavşat-Artvin/Turkey)

Bülent Akar¹, Bülent Şahin^{2,*}

¹ Gümüşhane University, Kürtün Vocational School, 29830, Kürtün, Gümüşhane, Turkey.

² Karadeniz Technical University, Fatih Education Faculty, Department of Biology, Trabzon, Turkey.

* Corresponding Author: Tel.: +90.533 6338855; Fax: ;
E-mail: bulentsahin61@gmail.com

Received 7 November 2013
Accepted 15 March 2014

Abstract

The benthic algal flora of Karagöl Lake (Şavşat, Artvin) was investigated between October 2008 and September 2010. *Actinotaenium diplosporum* var. *americanum*, *Cosmarium pseudoexiguum*, *Gonatozygon kinahanii*, *Staurastrum boreale* var. *quadriradiatum*, *S. hantzschii* and *Staurodesmus dejectus* var. *apiculatus* were recorded for the first time for the desmid flora of Turkey.

Keywords: Karagöl Lake, new record, desmids, Turkey.

Karagöl-Sahara Milli Parkı'ndaki (Şavşat-Artvin/Türkiye) Karagöl'den Yeni Desmid Kayıtları

Özet

Karagöl'ün (Şavşat, Artvin) bentik alg florası Ekim 2008 ile Eylül 2010 tarihleri arasında incelenmiştir. *Actinotaenium diplosporum* var. *americanum*, *Cosmarium pseudoexiguum*, *Gonatozygon kinahanii*, *Staurastrum boreale* var. *quadriradiatum*, *S. hantzschii* ve *Staurodesmus dejectus* var. *apiculatus* Türkiye'nin desmid florası için ilk kez kaydedilmiştir.

Anahtar Kelimeler: Karagöl, yeni kayıt, desmid, Türkiye.

Introduction

Desmids are typically freshwater and very attractive algae characteristic of acidic and nutrient-poor aquatic ecosystems. The roles of desmids are contributing to the food supply of aquatic animals, primary producers in the aquatic environment and indicators of trophic status and pollution since they have a low tolerance for inorganic salts. Desmids have attracted the attention of microscopists because they exhibit great diversity in their external morphology as well as show remarkably complex cell symmetry making them of great natural beauty and aesthetic appeal (Kadiri, 2002).

Turkey is rather unique among the European countries in that it still contains rich freshwater areas. Ertan and Morkoyunlu (1998) pointed out that Turkey has 906.118 ha of lakes, 18.000 ha of dam lakes and a water network 145.000 km long. A number of lakes, pools and rivers are situated in high mountain areas,

offering good conditions for the development of a rich desmid flora because they are largely ecologically intact and remote from the industrial and agricultural centers of Turkey. In particular, there are a number of high mountain lakes in the Eastern Black Sea region of Turkey.

Among numerous floristic papers from European countries there is relatively little information about Turkish desmids. Due to a shortage of limnologists, a lot of water bodies and difficulties of transport. Nevertheless a few check-lists were published including the results of the studies of freshwater algal flora of Turkey at different times (Aysel, 2005; Gönülol *et al.*, 1996; Şahin, 2005). In addition, new records are given for the freshwater algal flora of Turkey in various dates (Atıcı, 2002; Baykal *et al.*, 2009; Ongun-Sevindik *et al.*, 2010, 2011; Öztürk *et al.*, 1995a, 1995b; Şahin, 1998, 2000, 2002, 2005, 2007a, 2007b, 2009).

The aim of the present study was to make a contribution to the desmids flora of Turkey. Therefore

six new records were described in this paper.

Materials and Methods

Study Area

Karagöl Lake is located between the longitudes of 42°29'10.28" E and latitudes of 41°18'32.47" N in Karagöl-Sahara National Park in Artvin (Şavşat), Turkey (Figure 1). The lake has a surface area of 6 ha and depth of 33 m. It is 1630 m above sea level.

The climate of the region is generally cool and rainy in summer and cold and snowy in winter (annual means temperature 9.8°C, annual precipitation 737.9 mm) (DMİGM, 2001). Karagöl-Sahara National Park is most famous for its high amount of endemic plant and animal species. Terrestrial vegetation is composed of trees, shrubs and herbs, including genera *Abies*, *Astragalus*, *Campanula*, *Fagus*, *Geranium*, *Picea*, *Pinus*, *Ranunculus*, *Trifolium*, *Veronica* and *Vicia* (Eminağaoğlu and Anşin, 2004). *Equisetum ramosissimum*, *Polygonum*

amphibium and *Myriophyllum spicatum* in the littoral zone of the lake are also available. *Tetraogallus caspius*, *Vipera kaznakovi* and *Ursus arctos* are also among the animals found in the Karagöl-Sahara National Park (Anonym, 2002).

In order to examine benthic algal flora of Karagöl Lake, three stations were chosen. The number of sampling stations depended on the size of the lake. The first station is at the north-west of the lake. There are small stones and vascular plants which are *Equisetum ramosissimum*, *Polygonum amphibium* and *Myriophyllum spicatum*. The second station is situated at east of the lake where sediment, stones and vascular plants are located. The third station is at the south-west of the lake. In this station there are only vascular plants. Samples were collected during the snow-free period from October 2008 and September 2010. The samples were taken on a monthly basis. Epipellic samples (from station II) were collected by drawing a glass tube across the surface of the sediment, epilithic (from stations I, II) and epiphytic (from all stations) samples were collected at random (Round, 1953; Sladeckova, 1962). All samples were




Figure 1. Map of the Karagöl Lake.

fixed in 4% formaldehyde. Samples were examined in temporary slides and under a Leica DM 2500 light microscope (magnification x400). Identification of algae were made according to Coesel and Meesters (2007), Dillard (1991), Lenzenweger (1996, 1997), John *et al.* (2003). Taxa were photographed with a camera attached to a Leica DM 2500 light microscope.

At the time of sampling, surface water temperature, pH, conductivity and dissolved oxygen concentration were measured using in situ with portable measuring instruments. On return to the laboratory, concentrations of Ca⁺⁺, Mg⁺⁺, K⁺, Cl⁻, SiO₂, PO₄³⁻-P, NO₂⁻-N, NO₃⁻-N, NH₄⁺-N and total hardness were measured with WTW S12 Model photometer.

Identified taxa were checked with the checklist of Aysel (2005); Gönülol *et al.* (1996) and Şahin (2005) determined as new taxa for Turkish algal flora. Taxonomy of algae was controlled with Guiry and Guiry (2013).

Results

Environmental Conditions

The water temperature of the Karagöl Lake ranged from 1,1 °C (at station II, in March 2009) to 25,4 °C (at station III, in July 2010) (mean 14,4 °C) during the sampling period. pH values varied between 6,87 and 9,22 (mean 7,79). Accordingly, the lake water was slightly alkaline. Dissolved oxygen concentrations were measured between 6,58 mg/L (at station III, in July 2010) and 12,42 mg/L (at station III, in December 2008). Other hydrological characteristics of the lake water are given in Table 1.

Annotated List of Species

A total of 6 new records desmid taxa of 5 genera belonging to 2 families were identified from epipellic, epilithic and epiphytic samples. All genera belonged to placoderm desmids.

Divisio: Charophyta

Classis: Conjugatophyceae

Ordo: Desmidiaceae

Familia: Gonatozygaceae

Genus: *Gonatozygon* de Bary, 1858

G. kinahanii (W.Archer) Rabenhorst, 1868 (Figure 2a) Coesel, F.M. and Meesters, K. (2007) p. 28, pl. 5, figs: 1-2 John, D.M., Whitton, B.A. and Brook, A.J. (2003) p. 529, pl. 128, fig. N Lenzenweger, R. (1996) p. 16, pl. 1, fig: 1

Cells 12,5 µm breadth, 275 µm length. Cell 22 times longer than wide. Cell elongate cylindrical and straight. Apices of the cell truncate; walls smooth; chloroplast an axile ribbon with a series of 10 pyrenoids.

Habitat: Epilithic.

Station: 1

Familia: Desmidiaceae

Genus: *Actinotaenium* (Nägeli) Teiling, 1954

A. diplosporium var. *americanum* (West & West) Teiling, 1954 (Figure 2b)

Coesel, F.M. and Meesters, K. (2007) p. 61, pl. 30, figs: 1-2

Dillard, E.G. (1991) p. 16, pl. 1 fig. 9

Cells 27,4µm breadth, 62,2 µm length.

Cells somewhat ellipsoid. Lateral sides

slightly converging toward the apex. Sinus narrower and sharper marked. Chloroplast asteroid.

Habitat: Epipellic, epilithic, epiphytic.

Stations: 1, 2, 3

Genus: *Cosmarium* Corda ex Ralfs, 1848

C. pseudoexiguum Raciborski, 1885 (Figure 2c)

Coesel, F.M. and Meesters, K. (2007) p.134, pl. 67, figs: 12-13

Cells 8,8 µm breadth, 17 µm length, isthmus 2,9 µm. Cells much longer than broad, in outline rectangular with rounded angles. Sinus deep, linear and closed for the greater part. Cell wall smooth.

Habitat: Epiphytic.

Station: 3

Genus: *Staurastrum* Meyen ex Ralfs, 1848

S. boreale var. *quadri radiatum* Korshikov, 1941 (Figure 2d1,2)

Coesel, F.M. and Meesters, K. (2007) p. 177, pl. 104,

Table 1. Hydrological characteristics of Karagöl Lake

Parameters	Station I			Station II			Station III		
	Min.	Max.	Mean	Min.	Max.	Mean	Min.	Max.	Mean
Temp. (°C)	1.20	24.80	14.42	1.10	23.90	14.30	1.20	25.40	14.66
DO(mg/L)	6.73	11.20	9.10	6.90	12.35	9.11	6.58	12.42	8.98
pH	6.90	9.22	7.78	6.94	9.17	7.77	6.87	8.91	7.82
EC (µS/cm)	57.70	119.90	107.18	66.90	123.60	106.10	73.00	140.20	111.84
TDS(mg/L)	28.00	59.00	52.63	33.00	60.00	52.26	36.00	69.00	54.94
CaCO ₃ (mg/L)	30.00	86.00	61.42	28.00	88.00	57.32	21.00	89.00	57.16
NH ₄ ⁺ -N (mg/L)	0.01	0.32	0.05	0.01	0.07	0.03	0.01	0.13	0.04
NO ₃ ⁻ -N (mg/L)	0.10	0.80	0.31	0.10	1.40	0.42	0.10	1.00	0.30
NO ₂ ⁻ -N (mg/L)	0.01	0.02	0.015	0.01	0.03	0.017	0.01	0.03	0.015
PO ₄ ³⁻ -P (mg/L)	0.01	0.19	0.05	0.01	0.20	0.04	0.01	0.16	0.05
SiO ₂ (mg/L)	0.50	6.70	3.62	0.50	8.10	3.61	0.50	5.90	3.17
Ca ²⁺ (mg/L)	12.00	35.00	24.68	11.00	36.00	22.95	8.00	36.00	22.84




Figure 2. a. *Gonatozygon kinahanii*, b. *Actinotaenium diplosporum* var. *americanum*, c. *Cosmarium pseudoexiguum*, d1,2. *Staurodesmus boreale* var. *quadri radiatum*, e1,2. *S. hantzschii*, f1,2. *Staurodesmus dejectus* var. *apiculatus* (Scale 20 μm)

figs: 13-17

Cells 31,1 μm breadth, 23,7 μm length. Cells broader than long, deeply constricted. Sinus widely open. Semicell cup-shaped, the apical angles produced to form rather short to moderately long, slightly divergent processes tipped by short spines. Cell wall furnished with granules. Semicells in apical view 4 radiate.

Habitat: Epilithic, epiphytic.

Stations: 2, 3.

S. hantzschii Reinsch, 1866 (Figure 2e1,2)

Lenzenweger, R. (1997) p. 93, pl. 41, figs: 9-10

Cells 22,22 μm breadth, 27,40 μm length. Cells 1,2 longer than broad, not including processes; median constriction shallow, sinus open; semicells elliptical in outline, lateral margins convex, with processes short, jagged, concave margins between processes, cell wall smooth; chloroplast with a pyrenoid by

semicell; apical circular, with nine denticulate processes, concave edges between the processes.

Habitat: Epiphytic.

Station: 3

Genus: *Staurodesmus* Teiling, 1948

S. dejectus var. *apiculatus* (Brébisson) Croasdale, 1957 (Figure 2f1,2)

Coesel, F.M. and Meesters, K. (2007) p. 161, pl. 88, figs: 20-25

John, D.M., Whitton, B.A. and Brook, A.J. (2003) p. 579, pl. 141, fig: S

Lenzenweger, R. (1997) p. 26, pl. 22, figs: 8,9,11

Cells 16,2 μm breadth, 17,7 μm length. Cells are almost as broad as long. Cells triradiate, sinus widely open and rounded internally, isthmus slightly elongate, 5,1 μm wide; semicells cup-shaped, apex slightly convex and small, vertically directed spines.

Habitat: Epiphytic.

Station: 3

Discussion

On the basis of the results of the physico-chemical analyses and determined algae (e.g. Ochrophyta, Charophyta, Chlorophyta, Cyanobacteria, Euglenozoa and Myzozoa) it is concluded that the Karagöl Lake has mesotrophic character.

The occurrence of acidophilic desmids, including *Gonatozygon kinahanii*, *Staurastrum hantzschii* and *Staurodesmus dejectus. apiculatus*, in slightly alkaline water is remarkable (Coesel, 2007; John *et al.*, 2003; Lenzenweger 1997).

The same situation was observed in many water bodies in Turkey and Europe (Atıcı, 2002; Baykal *et al.*, 2009; Feher, 2003; Ongun-Sevindik *et al.*, 2010, 2011; Öztürk *et al.*, 1995a, 1995b; Kostkevičienė *et al.*, 2003; Şahin and Akar, 2007, Şahin, 1998, 2000, 2002, 2005, 2007, 2009). In addition, according to the ranges of conductivity, composition and concentrations of ions and biogenic elements, as recorded in this study, these desmids appear to be well-adapted to mesotrophic habitat.

Other desmids (e.g. *Actinotaenium diplosporium* var. *americanum*, *Cosmarium pseudoexiguum* and *Staurastrum boreale. quadriradiatum*) are known as inhabitants of mesotrophic water bodies of Europe (Coesel, 2007).

It is noteworthy that several parameters of physico-chemical analysis such as pH, conductivity, dissolved O₂, total hardness, PO₄³⁻-P, NO₂⁻-N, NO₃⁻-N and NH₄⁺-N, have a low variability in the investigated stations and this may have been a contributing factor to the development of a diverse desmid community.

It is generally acknowledged that *Cosmarium* and *Staurastrum* are the most significant desmids in the northern flora (Getzen, 1985). In this study, these genera are represented with 3 species.

In conclusion, we can say that by increasing the number of studies the number of desmids will rise.

Acknowledgements

We would like to thank to Karadeniz Technical University (KTU) for its financial support (Proje Code: 2007.116.02.4).

References

- Anonymous. 2002. Karadeniz Teknik Üniversitesi inceleme raporu, Orman Bakanlığı Doğu Karadeniz Bölge Müdürlüğü Artvin Milli Parklar ve Av-Yaban Hayatı Başmühendisliği, Artvin.
- Atıcı, T. 2002. Nineteen new records from Sarıyar Dam Reservoir phytoplankton for Turkish freshwater algae. Turkish Journal of Botany, 26: 485-490.
- Aysel, V. 2005. Check-List of the freshwater algae of Turkey. Journal of the Black Sea/ Mediterranean Environment, 11(1):1-124.
- Baykal, T., Akbulut, A., Açıköz, İ., Udoh, A.U., Yıldız, K.

- and Şen, B. 2009. New records for the freshwater algae of Turkey, Turkish Journal of Botany, 33: 141-152.
- Coesel, F.M. and Meesters, K. 2007. Desmids of the Lowlands. KNNV Publishing, Netherlands 351 pp.
- DMİGM, 2001. Ortalama ekstrem ve yağış değerleri bülteni Artvin ili verileri. Devlet Meteoroloji İşleri Genel Müdürlüğü Yayını.
- Dillard, G.E. 1991. Freshwater algae of the Southeastern United States Part 4. Chlorophyceae Zygnematales: Desmidiaceae (Section 2). J. Cramer, Berlin-Stuttgart, 205 pp.
- Eminağaoğlu, Ö. and Anşin, R. 2004. Flora of the Karagöl-Sahara National Park (Artvin) and its environs. Turkish Journal of Botany, 28: 557-590.
- Ertan, Ö.O. and Morkoyunlu, A. 1998. The algal flora of Aksu Stream (Isparta/Turkey). Turkish Journal of Botany, 22: 239-255.
- Feher, G. 2003. The desmid flora of some alkaline lakes and wetlands in Southern Hungary. Biologia, Bratislava, 58(4):671-683.
- Getzen, M.V. 1985. Algae in ecosystems of the extreme North (on the example of Bolshezemlskaja tundra). Nauka, Leningrad, 165 pp.
- Gönülol, A., Öztürk, M. and Öztürk, M. 1996. A check-list of the freshwater algae of Turkey. Ondokuz Mayıs Üniversitesi Fen Edebiyat Fakültesi Fen Dergisi, 7(1): 8-46.
- Guiry, M.D. and Guiry, G.M. 2013. AlgaeBase. – Worldwide electronic publication. National University of Ireland, Galway. <http://www.algaebase.org> (28 April 2013).
- John, D.M., Whitton, B.A. and Brook, A.J. 2003. The freshwater algal flora of the British Isles, an identification guide to freshwater and terrestrial algae. Cambridge University Pres., Cambridge, 702 pp.
- Kadiri, M.O. 2002. A Checklist of desmids in Nigeria. Global Journal of Pure and Applied Sciences. 8(2):223-237. <http://dx.doi.org/10.4314/gjpas.v8i2.16036>
- Kostkevičienė, J., Briskaite, R., Bakunaite, J. and Jakimavičiute, I. 2003. Desmids (Chlorophyta, Desmidiales) from Lithuania. Biologia, Bratislava, 58(4):685-695.
- Lenzenweger, R. 1996. Desmidiaceen flora von Österreich Teil 1. J.Cramer, Berlin-Stuttgart, 162 pp.
- Lenzenweger, R. 1997. Desmidiaceen flora von Österreich Teil 2. J.Cramer, Berlin-Stuttgart, 216 pp.
- Ongun-Sevindik, T., Çelik, K. and Gönülol, A. 2010. Twentyfour new records from Çaygören Reservoir phytoplankton for Turkish freshwater algae. Turkish Journal of Botany, 31(4): 249-259.
- Ongun-Sevindik, T., Çelik, K. and Gönülol, A. 2011. Twenty new records for Turkish freshwater algal flora from Çaygören and İkizcetepeler Reservoirs (Balıkesir, Turkey). Turkish Journal of Fisheries and Aquatic Sciences, 11: 399-406. http://dx.doi.org/10.4194/1303-2712-v11_3_09
- Öztürk, M., Gönülol, A. and Öztürk, M. 1995a. Türkiye alg florası için yeni bir kayıt: *Pleurotaenium trabecula* (Ehr.) ex Nägeli (Desmidiaceae), O.M.Ü. Fen Dergisi, 6(1): 212-218.
- Öztürk, M., Gezerler-Şipal, U., Güner, H., Gönülol, A. and Aysel, V. 1995b. *Closterium kuetzingii* Bréb. var. *kuetzingii* (Conjugatophyceae, Desmidiales), A new record for the algal flora of Turkey. E.Ü. Su Ürünleri Dergisi, 12(1-2): 145-149.

- Round, F.E. 1953. An investigation of two benthic algal communities in Malharm Tarn, Yorkshire. *J. Ecol.*, 41: 97-174. doi.org/10.2307/2257108
- Sladeckova, A. 1962. Limnological investigation methods for periphyton Aufwuchs) Community. *Botanical Review*, 28: 86-350. <http://dx.doi.org/10.1007/BF02860817>
- Şahin, B. 1998. Some new records of desmids from Turkey. *Pak. J. Bot.*, 30(1): 7-13.
- Şahin, B. 2000. Some new desmids records for freshwater algal flora of Turkey. *Flora Mediterranea*, 10: 223-226.
- Şahin, B. 2002. Contribution to the desmid flora of Turkey. *Algological Studies*, 107: 39-48.
- Şahin, B. 2005. A preliminary checklist of desmids of Turkey. *Cryptogamie, Algol.*, 26(4): 399-415.
- Şahin, B. 2007. Two new records for the freshwater algae of Turkey. *Turkish Journal of Botany*.31: 153-156.
- Şahin, B. and Akar, B. 2007. The desmid flora of some high mountain lakes of the Turkish Eastern Black Sea region. *Pak. J. Bot.*: 39(5): 1817-1832.
- Şahin, B. 2009. Contribution to the desmid flora of Turkey. *Turkish Journal of Botany*, 33: 457-460.