

COMMUNICATIONS

DE LA FACULTÉ DES SCIENCES
DE L'UNIVERSITÉ D'ANKARA

Série A₁: Mathématiques

TOME : 30

ANNEE : 1981

A LA MEMOIRE D'ATATÜRK AU CENTENAIRE DE SA NAISSANCE



On The Degree Of Approximation Of Continuous Functions

by

PREM CHANDRA

2

Faculté des Sciences de l'Université d'Ankara
Ankara, Turquie

Communications de la Faculté des Sciences de l'Université d'Ankara

Comité de Redaction de la Série A,
B. Yurtsever H. Hacisalihoğlu, M. Oruç,
Secrétaire de Publication
Ö. Çakar

La Revue "Communications de la Faculté des Sciences de l'Université d'Ankara" est un organe de publication englobant toutes les disciplines scientifiques représentées à la Faculté des Sciences de l'Université d'Ankara.

La Revue, jusqu'à 1975 à l'exception des tomes I, II, III était composé de trois séries

Série A: Mathématiques, Physique et Astronomie,
Série B: Chimie,
Série C: Sciences Naturelles.

A partir de 1975 la Revue comprend sept séries:

Série A₁: Mathématiques,
Série A₂: Physique,
Série A₃: Astronomie,
Série B: Chimie,
Série C₁: Géologie,
Série C₂: Botanique,
Série C₃: Zoologie.

En principe, la Revue est réservée aux mémoires originaux des membres de la Faculté des Sciences de l'Université d'Ankara. Elle accepte cependant, dans la mesure de la place disponible les communications des auteurs étrangers. Les langues Allemande, Anglaise et Française seront acceptées indifféremment. Tout article doit être accompagnés d'un résumé.

Les articles soumis pour publications doivent être remis en trois exemplaires dactylographiés et ne pas dépasser 25 pages des Communications, les dessins et figures portes sur les feuilles séparées devant pouvoir être reproduits sans modifications.

Les auteurs reçoivent 25 extraits sans couverture.

L'Adresse : Dergi Yayın Sekreteri,
Ankara Üniversitesi,
Fen Fakültesi,
Beşevler-Ankara

DEDICATION TO ATATÜRK'S CENTENNIAL

Holding the torch that was lit by Atatürk in the hope of advancing our Country to a modern level of civilization, we celebrate the one hundredth anniversary of his birth. We know that we can only achieve this level in the fields of science and technology that are the wealth of humanity by being productive and creative. As we thus proceed, we are conscious that, in the words of Atatürk, "the truest guide" is knowledge and science.

As members of the Faculty of Science at the University of Ankara, we are making every effort to carry out scientific research, as well as to educate and train technicians, scientists, and graduates at every level. As long as we keep in our minds what Atatürk created for his Country, we can never be satisfied with what we have been able to achieve. Yet, the longing for truth, beauty, and a sense of responsibility toward our fellow human beings that he kindled within us gives us strength to strive for even more basic and meaningful service in the future.

From this year forward, we wish and aspire toward surpassing our past efforts, and with each coming year, to serve in greater measure the field of universal science and our own nation.

On The Degree Of Approximation Of Continuous Functions

by

PREM CHANDRA

(Received on April 25, 1979 and accepted on May 29, 1980)

ABSTRACT

In this paper the author has obtained the degree of approximation of a 2π -periodic function f of the class $\text{Lip } \alpha$, $0 < \alpha \leq 1$, by Eulermeans of its Fourier Series. He has further shown that this degree of approximation is best possible in certain sense.

1. Let f be 2π -periodic and L-integrable over $[-\pi, \pi]$. The Fourier series associated with f at a point x is

$$(1.1) \quad \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx).$$

A function $f \in \text{Lip } \alpha$ ($\alpha > 0$) if

$$(1.2) \quad f(x+h) - f(x) = O(|h|^\alpha).$$

Let $\{s_n\}$ be a sequence of partial sums of the given series

$\sum_{n=0}^{\infty} c_n$, where $s_n = c_0 + c_1 + \dots + c_n$. Then (E, q) ($q > 0$) -means

of $\{s_n\}$ are defined by ([4], p. 180)

$$(1.3) \quad (1+q)^{-n} \sum_{k=0}^n \binom{n}{k} q^{n-k} s_k.$$

The (E, q) means for $q > 0$ are regular ([4], p. 179). Throughout this paper K's will denote positive constants.

2. INTRODUCTION. The following theorem on the degree of approximation of a function f , belonging to the class $\text{Lip } \alpha$ by the (C, δ) - means of its Fourier series is proved by Alexits ([1], p. 301):

Theorem A. If a periodic function f belongs to the class $\text{Lip } \alpha$, then the degree of approximation of (C, δ) -means of its Fourier series for $0 < \alpha \leq \delta \leq 1$ is

$$(2.1) \quad \max_{0 \leq x \leq 2\pi} |f(x) - \sigma_n^\delta(x)| = \begin{cases} O(n^{-\alpha}), & \text{for } 0 < \alpha < \delta \leq 1; \\ O(n^{-\alpha} \log n), & \text{for } 0 < \alpha \leq \delta \leq 1, \end{cases}$$

where $\sigma_n^\delta(x)$ are the (C, δ) -means of the partial sums of (1.1).

By extending the above result, the present author [2] proved the following theorem:

Theorem B. The degree of approximation of a periodic function f with period 2π and belonging to the class $\text{Lip } \alpha$, by (R, p_n) -means of its Fourier series is given by

$$(2.2) \quad \max_{0 \leq x \leq 2\pi} |f(x) - T_n(x)| = \begin{cases} O\left(\frac{p_n}{P_n}\right)^\alpha; & 0 < \alpha < 1 \\ O\left(\frac{p_n}{P_n} \log \frac{P_n}{p_n}\right); & \alpha = 1, \end{cases}$$

where (R, p_n) -means are regular and $0 < p_n \uparrow$ (monotonic increasing) with $n \geq n_0$ and $T_n(x)$ are (R, p_n) -means of the partial sums of (1.1).

It may be observed that the means generating sequences in the above theorems are involved either implicitly or explicitly in the order of approximation. In this paper we obtain the following theorem which provides the degree of approximation free from the means generating sequence:

THEOREM. The degree of approximation of a periodic function f with period 2π and belonging to the class of $\text{Lip } \alpha$, $0 < \alpha \leq 1$, by Euler-means of its Fourier series is given by

$$(2.3) \quad \max_{0 \leq x \leq 2\pi} |f(x) - t_n^q(x)| = O(n^{-\frac{1}{2}\alpha}),$$

where $t_n^q(x)$ are (E, q) ($q > 0$)-means of the partial sums of (1.1). However the inequality (2.3) is best possible in the sense that there exists a positive constant K such that

$$(2.4) \quad \max_{0 \leq x \leq 2\pi} |f(x) - t_n^q(x)| \geq K n^{-\frac{1}{2}\alpha} \quad (n > n_0)$$

3. For the proof of the theorem we shall require the following lemmas:

LEMMA 1. Let $0 < t \leq \pi$. Then

$$(1+q)^{-n} \{1+q^2 + 2q \cos t\}^{\frac{n}{2}} < \exp \{-2nqt^2 / (\pi(1+q))^2\}.$$

PROOF. We have

$$\begin{aligned} (1+q)^{-n}(1+q^2 + 2q \cos t)^{\frac{1}{2}n} &= \left\{ 1 - \left(\frac{2\sqrt{q}}{1+q} \sin \frac{1}{2}t \right)^2 \right\}^{\frac{1}{2}n} \\ &= \left[\cos \left\{ \sin^{-1} \left(\frac{2\sqrt{q}}{1+q} \sin \frac{1}{2}t \right) \right\} \right]^n \\ &< \left\{ \cos \left(\frac{2t\sqrt{q}}{\pi(1+q)} \right) \right\}^n \end{aligned}$$

since

$$\begin{aligned} \sin^{-1} \frac{2\sqrt{q}}{1+q} \sin \frac{1}{2}t &> \frac{2\sqrt{q}}{1+q} \sin \frac{1}{2}t \\ &\geq \frac{2\sqrt{q}}{1+q} \frac{t}{\pi} \quad (0 < t \leq \pi), \end{aligned}$$

where $\left\{ \frac{\sin \theta}{\theta} \right\}$ is decreasing in $0 < \theta < \frac{1}{2}\pi$ and its minimum

value in $(0, \frac{1}{2}\pi)$ is $\frac{2}{\pi}$ so that $\sin \theta \leq \frac{2}{\pi} \theta$.

However, for $0 < t \leq \pi$,

$$0 < \frac{2t\sqrt{q}}{\pi(1+q)} < \frac{1}{2}\pi$$

therefore, we have (see Hardy [4], p. 364)

$$\cos \frac{2t\sqrt{q}}{\pi(1+q)} < \exp \{-2qt^2 / (\pi(1+q))^2\}$$

and hence the proof of Lemma 1 follows.

LEMMA 2. Let $0 < t \leq \frac{4}{\pi} \beta$, where β is strictly less

than $\frac{1}{2} \pi$. Then

$$(1+q)^{-n}(1+q^2+2q \cos t)^{\frac{n}{2}} > \exp \left\{ -nqt^2\pi^2 \sec^2 \beta / (8(1+q)^2) \right\}$$

PROOF. We have

$$\log ((1-y)^{-1}) < y / (1-y) \quad (0 < y < 1).$$

Therefore, on replacing y by $\sin^2 \theta$ ($0 < \theta \leq \beta < \frac{1}{2} \pi$) in the above inequality and simplifying, we get

$$\begin{aligned} \log \sec^2 \theta &< \tan^2 \theta \\ &< \theta^2 \sec^2 \beta \end{aligned}$$

$$\text{or } \cos \theta > \exp \left\{ -\frac{1}{2} \theta^2 \sec^2 \beta \right\}.$$

And, as in Lemma 1,

$$(1+q)^{-n}(1+q^2+2q \cos t)^{\frac{1}{2}n} = (\cos z)^n,$$

where

$$\begin{aligned} z &= \sin^{-1} \left(\frac{2\sqrt{q}}{1+q} \sin \frac{1}{2} t \right) = \sin^{-1} \left(\frac{2}{\pi} \frac{\pi \sqrt{q} \sin \frac{1}{2} t}{(1+q)} \right) \\ &\leq \frac{\pi \sqrt{q}}{1+q} \sin \frac{1}{2} t \\ &\leq \frac{\pi}{2} \frac{q^{\frac{1}{2}}}{1+q} t \\ &\leq \beta \left(0 < t \leq \frac{4}{\pi} \beta \right). \end{aligned}$$

Hence

$$\begin{aligned} \cos z &\geq \cos \left\{ \pi q^{\frac{1}{2}} t / 2 (1+q) \right\} \\ &> \exp \left\{ -\frac{\pi^2}{2} \frac{q t^2 \sec^2 \beta}{4 (1+q)^2} \right\} \end{aligned}$$

from which we may follow the proof of Lemma 2.

4. PROOF OF THE THEOREM. To simplify the proof of the theorem, we use the following notations:

$$\begin{aligned} P(q,t,n) &= (1+q)^{-n}(1+q^2 + 2q \cos t)^{\frac{1}{2}n} \\ Q(q,t) &= \tan^{-1}\{\sin t / (q+\cos t)\} \\ a &= 2q / (\pi(1+q))^2 \\ b &= q\pi^2 / (2(1+q))^2. \end{aligned}$$

We have

$$s_n(x) - f(x) = \frac{1}{\pi} \int_0^\pi \frac{\sin(n+\frac{1}{2})t}{\sin \frac{1}{2}t} \Phi_x(t) dt,$$

where

$$\Phi(t) = \frac{1}{2} \{ f(x+t) + f(x-t) - 2f(x) \}.$$

Then (E, q) - transform of $\{s_n(x) - f(x)\}$ will be given by

$$(1+q)^{-n} \sum_{k=0}^n \binom{n}{k} q^{n-k} \{s_k(x) - f(x)\}$$

which equals to

$$t_n^q(x) - f(x).$$

Therefore

$$t_n^q(x) - f(x) = \frac{1}{\pi(1+q)^n} \int_0^\pi \frac{\Phi_x(t)}{\sin \frac{1}{2}t} \left\{ \sum_{k=0}^n \binom{n}{k} q^{n-k} \sin(k+\frac{1}{2})t \right\} dt,$$

and splitting up the integral \int_0^π into the sub-integrals $\int_0^{\frac{1}{\sqrt{n}}}$ and

$\int_{\frac{1}{\sqrt{n}}}^\pi$ and denoting them respectively by I_1 and I_2 , we obtain that

$$|f(x) - t_n^q(x)| \leq |I_1| + |I_2|.$$

Now

$$\max_{0 \leq x \leq 2\pi} |I_1| \leq \max_{0 \leq x \leq 2\pi} \int_0^{\frac{1}{\sqrt{n}}} \frac{|\Phi_x(t)|}{\sin \frac{1}{2}t} \left\{ (1+q)^{-n} \sum_{k=0}^n \binom{n}{k} q^{n-k} \right\} dt$$

$$\begin{aligned}
 &= \max_{0 \leq x \leq 2\pi} \int_0^{\frac{1}{\sqrt{n}}} \frac{|\Phi_x(t)|}{\sin \frac{1}{2} t} dt \\
 &= O \left\{ \int_0^{\frac{1}{\sqrt{n}}} t^{\alpha-1} dt \right\} \\
 &= O(n^{-\frac{1}{2}\alpha})
 \end{aligned}$$

and, proceeding as in Chandra [3]; (3.2), we obtain that

$$\begin{aligned}
 |I_2| &\leq \int_{\frac{1}{\sqrt{n}}}^{\pi} \frac{|\Phi_x(t)|}{\sin \frac{1}{2} t} |P(q,t,n) \sin \{ \frac{1}{2} t + nQ(q,t) \}| dt \\
 &\leq \int_{\frac{1}{\sqrt{n}}}^{\pi} \frac{|\Phi_x(x)|}{\sin \frac{1}{2} t} P(q,t,n) dt \\
 &= O \left\{ \int_{\frac{1}{\sqrt{n}}}^{\pi} \frac{|\Phi_x(t)|}{\sin \frac{1}{2} t} \exp(-ant^2) dt \right\} \\
 &\quad \text{(by Lemma 1)} \\
 &= O \left\{ \frac{1}{n} \int_{\frac{1}{\sqrt{n}}}^{\pi} \frac{|\Phi_x(t)|}{t \sin \frac{1}{2} t} - \frac{\partial}{\partial t} \{ -\exp(-ant^2) \} dt \right\}
 \end{aligned}$$

Therefore

$$\begin{aligned}
 \max_{0 \leq x \leq 2\pi} |I_2| &= O \left\{ \frac{1}{n} \int_{\frac{1}{\sqrt{n}}}^{\pi} t^{\alpha-2} \left\{ -\frac{\partial}{\partial t} \exp(-ant^2) \right\} dt \right\} \\
 &= O \left\{ n^{-(\alpha-2)/2+1} \int_{\frac{1}{\sqrt{n}}}^{\pi} \left\{ -\frac{\partial}{\partial t} \exp(-nt^2a) \right\} dt \right\} \\
 &= O(n^{-\frac{1}{2}\alpha}).
 \end{aligned}$$

Thus combining the results, as obtained above, we get

$$\max_{0 \leq x \leq 2\pi} |f(x) - t_n^q(x)| = O(n^{-\frac{1}{2}\alpha}) + O(n^{-\frac{1}{2}}) \\ = O(n^{-\frac{1}{2}\alpha}).$$

This completes the proof of (2.3).

To prove (2.4), we suppose that δ is a small number less than $\pi/4$. Then we write

$$\begin{aligned} \pi |f(x) - t_n^q(x)| &= \left| \int_0^\pi \frac{\Phi_x(t)}{\sin \frac{1}{2}t} \left\{ (1+q)^{-n} \sum_{k=0}^n \binom{n}{k} q^{n-k} \sin(k+\frac{1}{2})t \right\} dt \right| \\ &= \left| \left(\int_0^{\frac{1}{n}} + \int_{\frac{1}{n}}^{\delta} + \int_{\delta}^{\pi} \right) \left(\frac{\Phi_x(t)}{\sin \frac{1}{2}t} \right) \left((1+q)^{-n} \sum_{k=0}^n \binom{n}{k} q^{n-k} \right. \right. \\ &\quad \left. \left. \cdot \sin(k+\frac{1}{2})t \right) dt \right| \\ &= |J_1 + J_2 + J_3| \text{ (suppose)} \\ &\geq |J_2| - |J_1| - |J_3|. \end{aligned}$$

Proceeding as in $|I_1|$, we get

$$\max_{0 \leq x \leq 2\pi} |J_1| < K_1 n^{-\alpha}$$

and proceeding as in $|I_2|$, we get

$$\begin{aligned} \max_{0 \leq x \leq 2\pi} |J_3| &= O \left\{ \frac{1}{n} \int_{\delta}^{\pi} t^{\alpha-2} \left\{ -\frac{\partial}{\partial t} \exp(-ant^2) \right\} dt \right\} \\ &= O \left\{ \frac{1}{n} \int_{\delta}^{\pi} \left\{ -\frac{\partial}{\partial t} \exp(-ant^2) \right\} dt \right\} \\ &= O \left(\frac{1}{n} \right), \end{aligned}$$

therefore

$$\max_{0 \leq x \leq 2\pi} |J_3| < K_3/n.$$

Now

$$|J_2| \geq \left| 2 \int_{\frac{1}{n}}^{\delta} \Phi_x(t) t^{-1} P(q, t, n) \sin \left\{ \frac{1}{2}t + nQ(q, t) \right\} dt \right|$$

$$\begin{aligned}
 &= \left| \int_{\frac{1}{n}}^{\delta} \Phi_x(t) \{ \cosec \frac{1}{2} t - 2/t \} P(q,t,n) \right. \\
 &\quad \cdot \sin \left\{ \frac{1}{2} t + nQ(q,t) \right\} dt \left. \right| \\
 &= 2 |J_{2,1}| - |J_{2,2}|, \text{ say.}
 \end{aligned}$$

However

$$\begin{aligned}
 \max_{0 \leq x \leq 2\pi} |J_{2,2}| &= O \left\{ \int_{\frac{1}{n}}^{\delta} t^{1+\alpha} P(q,t,n) dt \right\} \\
 &= O \left\{ \frac{1}{n} \int_{\frac{1}{n}}^{\delta} t^{\alpha} \left\{ -\frac{\partial}{\partial t} \exp(-ant^2) \right\} dt \right\} \\
 &\quad (\text{by Lemma 1}) \\
 &= O \left(\frac{1}{n} \right)
 \end{aligned}$$

so that

$$|J_{2,2}| < K_{2,2}/n.$$

Now, by the hypothesis, there exists a constant $K_4 > 0$ such that

$$-K_4 t^\alpha < \Phi_x(t) < K_4 t^\alpha$$

therefore

$$K_4 t^\alpha < \Phi_x(t) + 2 K_4 t^\alpha$$

and hence

$$\begin{aligned}
 |J_{2,1}| &> \left| \int_{\frac{1}{n}}^{\delta} \frac{\Phi_x(t) + 2K_4 t^\alpha}{t} P(q,t,n) \sin \left(\frac{1}{2} t + nQ(q,t) \right) dt \right| \\
 &\quad - 2 K_4 \left| \int_{\frac{1}{n}}^{\delta} t^{\alpha-1} P(q,t,n) \sin \left(\frac{1}{2} t + nQ(q,t) \right) dt \right| \\
 &= |J_{2,1,1}| - 2 K_4 |J_{2,1,2}|, \text{ say.}
 \end{aligned}$$

However $t^{\alpha-1} P(q,t,n)$ is positive and non-increasing in $\frac{1}{n} \leq t \leq \delta$, therefore by the second mean value theorem

$$J_{2,1,2} = n^{1-\alpha} P(q, \frac{1}{n}, n) \int_{\frac{1}{n}}^{\delta'} \sin \left(\frac{1}{2} t + nQ(q,t) \right) dt$$

$(n^{-1} \leq \delta' \leq \delta)$

$$= O(n^{-\alpha})$$

so that

$$\max_{0 \leq x \leq 2\pi} 2 |K_4| |J_{2,1,2}| < K_{2,1,2} n^{-\alpha}.$$

And, by the first mean value theorem

$$\begin{aligned} |J_{2,1,1}| &= |\sin(\frac{1}{2}t' + nQ(q,t'))| \int_{\frac{1}{n}}^{\delta} \frac{\Phi_x(t) + 2K_4 t^\alpha}{t} P(q,t,n) dt \\ &\geq |\sin(\frac{1}{2}t' + nQ(q,t'))| \int_{\frac{1}{\sqrt{n}}}^{\delta} \frac{\Phi_x(t) + 2K_4 t^\alpha}{t} P(q,t,n) dt \\ &> K_4 |\sin(\frac{1}{2}t' + nQ(q,t'))| \int_{\frac{1}{\sqrt{n}}}^{\delta} t^{\alpha-1} P(q,t,n) dt \\ &> K_4 |\sin(\frac{1}{2}t' + nQ(q,t'))| \int_{\frac{1}{\sqrt{n}}}^{\delta} t^{\alpha-1} \exp(-bnt^2) dt \\ &\quad \text{(by Lemma 2)} \\ &\geq K_{2,1,1} n^{-\frac{1}{2}\alpha}, \end{aligned}$$

where the constant $K_{2,1,1}$ depends upon $\sin(\frac{1}{2}t' + nQ(q,t'))$ and other parameters. However the integral $J_{2,1,1}$ is not zero, therefore the constant $K_{2,1,1}$ is positive.

Now, collecting the results we get

$$\begin{aligned} \max_{0 \leq x \leq 2\pi} \pi |f(x) - t_n^q(x)| &\geq K_{2,1,1} n^{-\frac{1}{2}\alpha} - (K_1 + K_{2,1,2}) n^{-\alpha} \\ &\quad - (K_3 + K_{2,2}) / n \\ &\geq K_{2,1,1} n^{-\frac{1}{2}\alpha} - (K_1 + K_3 + K_{2,2} + K_{2,1,2}) / n \\ &= n^{-\frac{1}{2}\alpha} \left[K_{2,1,1} - (K_1 + K_3 + K_{2,2} + K_{2,1,2}) / n^{1-\frac{\alpha}{2}} \right]. \end{aligned}$$

And, for any given constant K' , such that

$$K_{2,1,1} - K' > 0$$

we can find a positive number $n_0 = n_0(K')$ such that

$$(K_1 + K_3 + K_{2,2} + K_{2,1,2})n^{\frac{1}{2}\alpha-1} \leq K' \text{ for } n > n_0.$$

And, hence writing K for $\frac{1}{\pi}(K_{2,1,1} - K')$, we get

$$\max_{0 \leq x \leq 2\pi} |f(x) - t_n^q(x)| \geq K n^{-\frac{1}{2}\alpha} (n > n_0)$$

This completes the proof of the theorem.

REFERENCES

- 1- **Alexits, G.**, Convergence problems of orthogonal series, *Pergamon Press*, 1961.
- 2- **Chandra, P.**, On the degree of approximation of functions belonging to the Lipschitz class, *Nanta Math.*, 8 (1) (1975), 88-91.
- 3- ———, On Euler summability of Fourier series, *Ranchi University Math. Jour.*, 5 (1974), 53-59.
- 4- **Hardy, G.H.**, Divergent series, Oxford, 1949.

Prix de l'abonnement annuel

Turquie: 15 TL; Etranger: 30 TL.

Prix de ce numéro: 5 TL (pour la vente en Turquie).

Prière de s'adresser pour l'abonnement à: Fen Fakültesi
Dekanlığı Ankara, Turquie.