

# A Note on Horadam Hybrinomials

Can Kızılates

Department of Mathematics, Faculty of Arts and Sciences, Zonguldak Bülent Ecevit University, Zonguldak, Turkey

## Article Info

**Keywords:** Horadam number, Horadam polynomial, Hybrid number  
**2010 AMS:** 11B37, 11B39, 11R52  
**Received:** 9 September 2021  
**Accepted:** 4 December 2021  
**Available online:** 11 February 2021

## Abstract

This paper ensures an extensive survey of the generalization of the various hybrid numbers and hybrid polynomials especially as part of its enhancing importance in the disciplines of mathematics and physics. In this paper, by using the Horadam polynomials, we define the Horadam hybrid polynomials called Horadam hybrinomials. We obtain some special cases and algebraic properties of the Horadam hybrinomials such as recurrence relation, generating function, exponential generating function, Binet formula, summation formulas, Catalan's identity, Cassini's identity and d'Ocagne's identity, respectively. Moreover, we give some applications related to the Horadam hybrinomials in matrices.

## 1. Introduction

Horadam defined the sequence  $w_n = w_n(a, b; p, q)$  by the recurrence relation

$$w_n = pw_{n-1} + qw_{n-2}, \quad n \geq 2$$

with the initial values  $w_0 = a$  and  $w_1 = b$ . For different values  $p, q, a, b \in \mathbb{Z}$ , Horadam sequence turns into several well-known sequences such as Fibonacci, Lucas, Pell and so on. These sequences are studied in many areas such as physics, number theory, algebra, geometry, and combinatorics. For more details, we refer to [1]-[6].

In [7], the Horadam polynomials  $h_n(x) = h_n(x; a, b; p, q)$  are defined by the recurrence relation

$$h_n(x) = pxh_{n-1}(x) + qh_{n-2}(x), \quad n \geq 3 \quad (1.1)$$

with the initial values  $h_1(x) = a$  and  $h_2(x) = bx$ . Let  $\alpha = \frac{px + \sqrt{p^2x^2 + 4q}}{2}$  and  $\beta = \frac{px - \sqrt{p^2x^2 + 4q}}{2}$  be the real roots of the characteristic equation  $t^2 - pxt - q = 0$ . The Binet formula for the polynomial  $h_n(x)$  is given by

$$h_n(x) = A\alpha^{n-1} + B\beta^{n-1}, \quad (1.2)$$

where  $A = \frac{bx - a\beta}{\sqrt{p^2x^2 + 4q}}$  and  $B = \frac{a\alpha - bx}{\sqrt{p^2x^2 + 4q}}$ .

The generating function of the Horadam polynomials is

$$\sum_{n=0}^{\infty} h_n(x)t^n = \frac{a + xt(b - ap)}{1 - pxt - qt^2}. \quad (1.3)$$

Hybrid numbers were studied by Ozdemir in [8], extensively. A hybrid number is defined as

$$\mathbb{K} = \{a + bi + c\varepsilon + d\mathbf{h} : a, b, c, d \in \mathbb{R}, \mathbf{i}^2 = -1, \varepsilon^2 = 0, \mathbf{h}^2 = 1, \mathbf{ih} = \mathbf{hi} = \varepsilon + \mathbf{i}\}.$$

Addition and subtraction of hybrid numbers are done by adding and subtracting corresponding terms. Two hybrid numbers are equal if all their components are equal, one by one.

Using the equalities  $\mathbf{i}^2 = -1$ ,  $\varepsilon^2 = 0$ ,  $\mathbf{h}^2 = 0$ ,  $\mathbf{ih} = -\mathbf{hi} = \varepsilon + \mathbf{i}$ , the multiplication table of the basis of hybrid numbers is as follows:

**Table 1:** Multiplication table for  $\mathbb{K}$

.	<b>1</b>	<b>i</b>	$\varepsilon$	<b>h</b>
<b>1</b>	<b>1</b>	<b>i</b>	$\varepsilon$	<b>h</b>
<b>i</b>	<b>i</b>	<b>-1</b>	<b>1-h</b>	$\varepsilon + \mathbf{i}$
$\varepsilon$	$\varepsilon$	<b>h+1</b>	0	$-\varepsilon$
<b>h</b>	<b>h</b>	$-\varepsilon - \mathbf{i}$	$\varepsilon$	<b>1</b>

Recently, many researchers have studied related to hybrid numbers. For example, in [9] Szynal-Liana and Wloch considered the Fibonacci hybrid numbers and obtained some properties of this numbers. In [10, 11] the authors also defined and examined the Jacosthal and Jacosthal–Lucas hybrid numbers and the Pell and Pell–Lucas hybrid numbers respectively. In [12] Szynal-Liana generalized their results and defined the Horadam hybrid numbers. In [13] Kızılateş introduced the another generalization of hybrid numbers and gave miscellaneous properties of these numbers. For more details, we refer to [8]-[23].

We now turn to a recent investigation by Szynal-Liana and Wloch [24], who defined and studied a family of the special polynomials and the special numbers which are related to the Fibonacci hybrid numbers and Lucas hybrid numbers. The Fibonacci hybrid numbers and Lucas hybrid numbers are defined as follows:

$$FH_n(x) = F_n(x) + F_{n+1}(x)\mathbf{i} + F_{n+2}(x)\varepsilon + F_{n+3}(x)\mathbf{h},$$

and

$$LH_n(x) = L_n(x) + L_{n+1}(x)\mathbf{i} + L_{n+2}(x)\varepsilon + L_{n+3}(x)\mathbf{h}.$$

For  $n \geq 2$ , the recurrence relations of the Fibonacci hybrid numbers and the Lucas hybrid numbers are

$$FH_n(x) = xFH_{n-1}(x) + FH_{n-2}(x),$$

and

$$LH_n(x) = xLH_{n-1}(x) + LH_{n-2}(x),$$

with the initial values  $FH_0(x) = \mathbf{i} + x\varepsilon + (x^2 + 1)\mathbf{h}$ ,  $FH_1(x) = 1 + x\mathbf{i} + (x^2 + 1)\varepsilon + (x^3 + 2x)\mathbf{h}$ ,  $LH_0(x) = 2 + x\mathbf{i} + (x^2 + 2)\varepsilon + (x^3 + 3x)\mathbf{h}$  and  $LH_1(x) = x + (x^2 + 2)\mathbf{i} + (x^3 + 3x)\varepsilon + (x^4 + 4x^2 + 2)\mathbf{h}$ , respectively. The Fibonacci hybrid numbers and the Lucas hybrid numbers, namely polynomials, are a generalization of the Fibonacci hybrid and Lucas hybrid numbers.

Motivated by some of the above-mentioned recent papers, we introduce here new polynomials which are called Horadam hybrid numbers. This definition brings about a more general hybrid polynomial sequence by taking components from Horadam polynomials. Thanks to this generalization, we obtain the Fibonacci hybrid numbers  $FH_n(x)$ , the Lucas hybrid numbers  $LH_{n-1}(x)$ , the Pell hybrid numbers  $PH_n(x)$ , the Pell-Lucas hybrid numbers  $QH_{n-1}(x)$ , the Chebyshev hybrid numbers of the first kind  $TH_{n-1}(x)$ , the Chebyshev hybrid numbers of the second kind  $UH_{n-1}(x)$  and the Balancing hybrid numbers  $BH_n(x)$ . We also obtain various results for the Horadam hybrid numbers. Moreover, we give some applications of Horadam hybrid numbers in matrices.

## 2. Horadam hybrid numbers

In this section, we define the Horadam hybrid numbers. Then we give some special cases of Horadam hybrid numbers such as the Fibonacci hybrid numbers, the Fibonacci hybrid numbers, the Lucas hybrid numbers, the Lucas hybrid numbers, the Pell hybrid numbers, the Pell hybrid numbers, the Pell-Lucas hybrid numbers, the Pell-Lucas hybrid numbers, the Chebyshev hybrid numbers of the first kind, the Chebyshev hybrid numbers of the first kind, the Chebyshev hybrid numbers of the second kind, the Chebyshev hybrid numbers of the second kind, the Balancing hybrid numbers and the Balancing hybrid numbers. Finally we obtain some algebraic properties of Horadam hybrid numbers.

**Definition 2.1.** For  $n \geq 1$ , the  $n^{\text{th}}$  Horadam hybrid numbers are defined by

$$\mathbb{H}_n(x) = h_n(x) + h_{n+1}(x)\mathbf{i} + h_{n+2}(x)\varepsilon + h_{n+3}(x)\mathbf{h}. \quad (2.1)$$

Some special cases of Horadam hybrid numbers are as follows:

1. For  $a = b = p = q = 1$ , the Horadam hybrid numbers  $\mathbb{H}_n(x)$  become the Fibonacci hybrid numbers  $FH_n(x)$ ,
2. For  $a = 2$  and  $b = p = q = 1$ , the Horadam hybrid numbers  $\mathbb{H}_n(x)$  become the Lucas hybrid numbers  $LH_{n-1}(x)$ ,

3. For  $a = q = 1$  and  $b = p = 2$ , the Horadam hybrinomials  $\mathbb{H}_n(x)$  become the Pell hybrinomials  $PH_n(x)$ ,
4. For  $a = b = p = 2$  and  $q = 1$ , the Horadam hybrinomials  $\mathbb{H}_n(x)$  become the Pell-Lucas hybrinomials  $QH_{n-1}(x)$ ,
5. For  $a = b = 1$ ,  $p = 2$ , and  $q = -1$ , the Horadam hybrinomials  $\mathbb{H}_n(x)$  become the Chebyshev hybrinomials of the first kind  $TH_{n-1}(x)$ ,
6. For  $a = 1$ ,  $b = p = 2$ , and  $q = -1$ , the Horadam hybrinomials  $\mathbb{H}_n(x)$  become the Chebyshev hybrinomials of the second kind  $UH_{n-1}(x)$ ,
7. For  $a = 1$ ,  $b = p = 6$ , and  $q = -1$ , the Horadam hybrinomials  $\mathbb{H}_n(x)$  become the Balancing hybrinomials  $BH_n(x)$ ,
8. For  $x = 1$ , the Fibonacci hybrinomials  $FH_n(x)$ , reduce to the Fibonacci hybrid numbers  $FH_n$ ,
9. For  $x = 1$ , the Lucas hybrinomials  $LH_{n-1}(x)$ , reduce to the Lucas hybrid numbers  $LH_{n-1}$ ,
10. For  $x = 1$ , the Pell hybrinomials  $PH_n(x)$ , reduce to the Pell hybrid numbers  $PH_n$ ,
11. For  $x = 1$ , the Pell-Lucas hybrinomials  $QH_{n-1}(x)$ , reduce to the Pell-Lucas hybrid numbers  $QH_{n-1}$ ,
12. For  $x = 1$ , the Chebyshev hybrinomials of the first kind  $TH_{n-1}(x)$ , reduce to the Chebyshev hybrid numbers of the first kind  $TH_{n-1}$ ,
13. For  $x = 1$ , the Chebyshev hybrinomials of the second kind  $UH_{n-1}(x)$ , reduce to the Chebyshev hybrid numbers of the second kind  $UH_{n-1}$ ,
14. For  $x = 1$ , the Balancing hybrinomials  $BH_n(x)$ , reduce to the Balancing hybrid numbers  $BH_n$ .

Using (2.1) and (1.1), we obtain that for  $n > 2$ ,

$$\begin{aligned} \mathbb{H}_n(x) &= pxh_{n-1}(x) + qh_{n-2}(x) + (pxh_n(x) + qh_{n-1}(x))\mathbf{i} \\ &\quad + (pxh_{n+1}(x) + qh_n(x))\boldsymbol{\varepsilon} + (pxh_{n+2}(x) + qh_{n+1}(x))\mathbf{h} \\ &= px\mathbb{H}_{n-1}(x) + q\mathbb{H}_{n-2}(x) \end{aligned}$$

and so

$$\mathbb{H}_n(x) = px\mathbb{H}_{n-1}(x) + q\mathbb{H}_{n-2}(x),$$

with the initial values  $\mathbb{H}_1(x) = a + b\mathbf{x}\mathbf{i} + (bp^2x^2 + aq)\boldsymbol{\varepsilon} + (bp^2x^3 + (apq + bq)x)\mathbf{h}$  and  $\mathbb{H}_2(x) = bx + (bp^2x^2 + aq)\mathbf{i} + (bp^2x^3 + (apq + bq)x)\boldsymbol{\varepsilon} + (bp^3x^4 + (ap^2q + 2bpq)x^2 + aq^2)\mathbf{h}$ .

**Theorem 2.2.** *The Binet formula for the Horadam hybrinomial  $\mathbb{H}_n(x)$  is*

$$\mathbb{H}_n(x) = A\alpha^{n-1}\tilde{\alpha} + B\beta^{n-1}\tilde{\beta}, \tag{2.2}$$

where  $\tilde{\alpha} = 1 + \alpha\mathbf{i} + \alpha^2\boldsymbol{\varepsilon} + \alpha^3\mathbf{h}$  and  $\tilde{\beta} = 1 + \beta\mathbf{i} + \beta^2\boldsymbol{\varepsilon} + \beta^3\mathbf{h}$ .

*Proof.* Due to (1.2) and (2.1), we find that

$$\begin{aligned} \mathbb{H}_n(x) &= (A\alpha^{n-1} + B\beta^{n-1}) + (A\alpha^n + B\beta^n)\mathbf{i} + (A\alpha^{n+1} + B\beta^{n+1})\boldsymbol{\varepsilon} + (A\alpha^{n+2} + B\beta^{n+2})\mathbf{h} \\ &= A\alpha^{n-1}(1 + \alpha\mathbf{i} + \alpha^2\boldsymbol{\varepsilon} + \alpha^3\mathbf{h}) + B\beta^{n-1}(1 + \beta\mathbf{i} + \beta^2\boldsymbol{\varepsilon} + \beta^3\mathbf{h}) \\ &= A\alpha^{n-1}\tilde{\alpha} + B\beta^{n-1}\tilde{\beta}. \end{aligned}$$

□

We now give the generating function and exponential generating function for the Horadam hybrinomials.

**Theorem 2.3.** *The generating function for the Horadam hybrinomial  $\mathbb{H}_n(x)$  is*

$$\sum_{n=0}^{\infty} \mathbb{H}_n(x)t^n = \frac{\mathbb{H}_0(x) + (\mathbb{H}_1(x) - px\mathbb{H}_0(x))t}{1 - pxt - qt^2}. \tag{2.3}$$

*Proof.* Suppose that the generating function for the Horadam hybrinomials  $\{\mathbb{H}_n(x)_{n=0}^{\infty}\}$ , has the following formal power series

$$\sum_{n=0}^{\infty} \mathbb{H}_n(x)t^n = \mathbb{H}_0(x) + \mathbb{H}_1(x)t + \dots + \mathbb{H}_k(x)t^k + \dots. \tag{2.4}$$

Hence

$$pxt \sum_{n=0}^{\infty} \mathbb{H}_n(x)t^n = px\mathbb{H}_0(x)t + px\mathbb{H}_1(x)t^2 + \dots + px\mathbb{H}_k(x)t^{k+1} + \dots, \tag{2.5}$$

$$qt^2 \sum_{n=0}^{\infty} \mathbb{H}_n(x)t^n = q\mathbb{H}_0(x)t^2 + q\mathbb{H}_1(x)t^3 + \dots + q\mathbb{H}_k(x)t^{k+2} + \dots. \tag{2.6}$$

From (2.4), (2.5) and (2.6), we find that

$$(1 - pxt - qt^2) \sum_{n=0}^{\infty} \mathbb{H}_n(x)t^n = \mathbb{H}_0(x) + (\mathbb{H}_1(x) - px\mathbb{H}_0(x))t.$$

So

$$\sum_{n=0}^{\infty} \mathbb{H}_n(x)t^n = \frac{\mathbb{H}_0(x) + (\mathbb{H}_1(x) - px\mathbb{H}_0(x))t}{1 - pxt - qt^2}.$$

□

**Corollary 2.4.** ([24, Theorem 2.10]) The generating function for the Fibonacci hybrinomial  $FH_n(x)$  is

$$\sum_{n=0}^{\infty} FH_n(x)t^n = \frac{\mathbf{i} + x\boldsymbol{\varepsilon} + (x^2 + 1)\mathbf{h} + (1 + \boldsymbol{\varepsilon} + x\mathbf{h})t}{1 - xt - t^2}.$$

*Proof.* If we take  $a = b = p = q = 1$  in Equation (2.3), the proof is completed. □

**Corollary 2.5.** ([24, Theorem 2.11]) The generating function for the Lucas hybrinomial  $LH_n(x)$  is

$$\sum_{n=0}^{\infty} LH_n(x)t^n = \frac{LH_0(x) + (LH_1(x) - xLH_0(x))t}{1 - xt - t^2}.$$

*Proof.* If we take  $a = 2$  and  $b = p = q = 1$  in Equation (2.3), the proof is completed. □

**Theorem 2.6.** The exponential generating function for the Horadam hybrinomial  $\mathbb{H}_n(x)$  is

$$\sum_{n=0}^{\infty} \mathbb{H}_n(x) \frac{t^n}{n!} = A\alpha^{-1} \tilde{\alpha} e^{\alpha t} + B\beta^{-1} \tilde{\beta} e^{\beta t}.$$

*Proof.* Using the Equation (2.2), we find that

$$\begin{aligned} \sum_{n=0}^{\infty} \mathbb{H}_n(x) \frac{t^n}{n!} &= \sum_{n=0}^{\infty} (A\alpha^{n-1} \tilde{\alpha} + B\beta^{n-1} \tilde{\beta}) \frac{t^n}{n!} \\ &= \frac{A\tilde{\alpha}}{\alpha} \sum_{n=0}^{\infty} \frac{(\alpha t)^n}{n!} + \frac{B\tilde{\beta}}{\beta} \sum_{n=0}^{\infty} \frac{(\beta t)^n}{n!} \\ &= \frac{A\tilde{\alpha}}{\alpha} e^{\alpha t} + \frac{B\tilde{\beta}}{\beta} e^{\beta t} \\ &= A\alpha^{-1} \tilde{\alpha} e^{\alpha t} + B\beta^{-1} \tilde{\beta} e^{\beta t}. \end{aligned}$$

So the proof is completed. □

We now give the following interesting identities.

**Theorem 2.7.** (Catalan's Identity). For positive integers  $n$  and  $r$ , with  $n \geq r$ , the following identity is true:

$$\mathbb{H}_{n+r}(x)\mathbb{H}_{n-r}(x) - \mathbb{H}_n^2(x) = (-q)^{n-1}AB \left( \tilde{\alpha}\tilde{\beta} \left( \left( \frac{\beta}{\alpha} \right)^r - 1 \right) + \tilde{\beta}\tilde{\alpha} \left( \left( \frac{\alpha}{\beta} \right)^r - 1 \right) \right). \quad (2.7)$$

*Proof.* Using the Equation (2.2), we obtain the LHS of the equality (2.7),

$$\begin{aligned} \mathbb{H}_{n+r}(x)\mathbb{H}_{n-r}(x) - \mathbb{H}_n^2(x) &= \left( A\alpha^{n-r-1} \tilde{\alpha} + B\beta^{n-r-1} \tilde{\beta} \right) \left( A\alpha^{n+r-1} \tilde{\alpha} + B\beta^{n+r-1} \tilde{\beta} \right) \\ &\quad - \left( A\alpha^{n-1} \tilde{\alpha} + B\beta^{n-1} \tilde{\beta} \right)^2 \\ &= AB(\alpha\beta)^{n-1} \alpha^{-r} \beta^r \tilde{\alpha}\tilde{\beta} + BA(\beta\alpha)^{n-1} \beta^{-r} \alpha^r \tilde{\beta}\tilde{\alpha} \\ &\quad - AB(\alpha\beta)^{n-1} \tilde{\alpha}\tilde{\beta} - BA(\beta\alpha)^{n-1} \tilde{\beta}\tilde{\alpha}. \end{aligned}$$

Then, we have

$$\mathbb{H}_{n+r}(x)\mathbb{H}_{n-r}(x) - \mathbb{H}_n^2(x) = (-q)^{n-1}AB \left( \tilde{\alpha}\tilde{\beta} \left( \left( \frac{\beta}{\alpha} \right)^r - 1 \right) + \tilde{\beta}\tilde{\alpha} \left( \left( \frac{\alpha}{\beta} \right)^r - 1 \right) \right).$$

□

**Theorem 2.8.** (Cassini’s Identity). For  $n \geq 1$ , the following equality holds:

$$\mathbb{H}_{n+1}(x)\mathbb{H}_{n-1}(x) - \mathbb{H}_n^2(x) = (-q)^{n-1}AB \left( \tilde{\alpha}\tilde{\beta} \left( \frac{\beta}{\alpha} - 1 \right) + \tilde{\beta}\tilde{\alpha} \left( \frac{\alpha}{\beta} - 1 \right) \right). \tag{2.8}$$

*Proof.* If we take  $r = 1$ , in (2.7), we obtain the assertion of the theorem. □

**Theorem 2.9.** (d’Ocagne’s Identity) Let  $m \geq 0$  and  $n \geq 0$  be integers such that  $m > n + 1$ . Then we have

$$\mathbb{H}_m(x)\mathbb{H}_{n+1}(x) - \mathbb{H}_{m+1}(x)\mathbb{H}_n(x) = \sqrt{\Delta}AB(-q)^{n-1} \left( \beta^{m-n}\tilde{\beta}\tilde{\alpha} - \alpha^{m-n}\tilde{\alpha}\tilde{\beta} \right), \tag{2.9}$$

where  $\Delta = p^2x^2 + 4q$ .

*Proof.* By virtue of Equation (2.2), we get

$$\begin{aligned} \mathbb{H}_m(x)\mathbb{H}_{n+1}(x) - \mathbb{H}_{m+1}(x)\mathbb{H}_n(x) &= \left( A\alpha^{m-1}\tilde{\alpha} + B\beta^{m-1}\tilde{\beta} \right) \left( A\alpha^n\tilde{\alpha} + B\beta^n\tilde{\beta} \right) \\ &\quad - \left( A\alpha^m\tilde{\alpha} + B\beta^m\tilde{\beta} \right) \left( A\alpha^{n-1}\tilde{\alpha} + B\beta^{n-1}\tilde{\beta} \right) \\ &= AB\alpha^{m-1}\beta^n\tilde{\alpha}\tilde{\beta} - AB\alpha^m\beta^{n-1}\tilde{\alpha}\tilde{\beta} \\ &\quad + BA\alpha^n\beta^{m-1}\tilde{\beta}\tilde{\alpha} - BA\alpha^{n-1}\beta^m\tilde{\beta}\tilde{\alpha}. \end{aligned}$$

After some calculations, we can easily see that

$$\mathbb{H}_m(x)\mathbb{H}_{n+1}(x) - \mathbb{H}_{m+1}(x)\mathbb{H}_n(x) = \sqrt{\Delta}AB(-q)^{n-1} \left( \beta^{m-n}\tilde{\beta}\tilde{\alpha} - \alpha^{m-n}\tilde{\alpha}\tilde{\beta} \right).$$

□

If we take  $a = b = p = q = 1$  in (2.7), (2.8) and (2.9), we obtain the Catalan, the Cassini and the d’Ocagne identities for the Fibonacci hybrinomials [24, Theorem 2.4], [24, Corollary 2.6] and [24, Theorem 2.7], respectively. Similarly, if we take  $a = 2$  and  $b = p = q = 1$  in (2.7), (2.8) and (2.9), we obtain the Catalan, the Cassini and the d’Ocagne identities for the Lucas hybrinomials [24, Theorem 2.5], [24, Corollary 2.6] and [24, Theorem 2.9], respectively.

**Theorem 2.10.** Let  $n \geq 2$  be an integer. Then we obtain

$$\sum_{k=1}^{n-1} \mathbb{H}_k(x) = \frac{\mathbb{H}_1(x) - \mathbb{H}_n(x) + q(\mathbb{H}_0(x) - \mathbb{H}_{n-1}(x))}{1 - px - q}. \tag{2.10}$$

*Proof.* By virtue of Equation (2.2), we find that

$$\begin{aligned} \sum_{k=1}^{n-1} \mathbb{H}_k(x) &= \sum_{k=1}^{n-1} \left( A\alpha^{k-1}\tilde{\alpha} + B\beta^{k-1}\tilde{\beta} \right) \\ &= A\tilde{\alpha} \sum_{k=1}^{n-1} \alpha^{k-1} + B\tilde{\beta} \sum_{k=1}^{n-1} \beta^{k-1} \\ &= A\tilde{\alpha} \left( \frac{1 - \alpha^{n-1}}{1 - \alpha} \right) + B\tilde{\beta} \left( \frac{1 - \beta^{n-1}}{1 - \beta} \right) \\ &= \frac{A\tilde{\alpha}(1 - \beta)(1 - \alpha^{n-1}) + B\tilde{\beta}(1 - \alpha)(1 - \beta^{n-1})}{1 - px - q}. \end{aligned}$$

Utilizing the last equation, we have

$$\sum_{k=1}^{n-1} \mathbb{H}_k(x) = \frac{\mathbb{H}_1(x) - \mathbb{H}_n(x) + q(\mathbb{H}_0(x) - \mathbb{H}_{n-1}(x))}{1 - px - q}.$$

□

**Corollary 2.11.** ([24, Theorem 2.13]) Let  $n \geq 2$  be an integer. Then we have

$$\sum_{k=1}^{n-1} FH_k(x) = \frac{FH_n(x) + FH_{n-1}(x) - FH_0(x) - FH_1(x)}{x}.$$

*Proof.* If we take  $a = b = p = q = 1$  in Equation (2.10), the proof is completed. □

**Corollary 2.12.** ([24, Theorem 2.15]) Let  $n \geq 2$  be an integer. Then we have

$$\sum_{k=1}^{n-1} LH_k(x) = \frac{LH_n(x) + LH_{n-1}(x) - LH_0(x) - LH_1(x)}{x}.$$

*Proof.* If we take  $a = 2$  and  $b = p = q = 1$  in Equation (2.10), the proof is completed.  $\square$

**Theorem 2.13.** For  $n \geq 0$ , we have

$$q^n \sum_{i=0}^n \binom{n}{i} \left(\frac{px}{q}\right)^{n-i} \mathbb{H}_{n-i}(x) = \mathbb{H}_{2n}(x). \quad (2.11)$$

*Proof.* Because of the Binet formula of the Horadam hybrinomials, we have the LHS of the equality (2.11),

$$\begin{aligned} & q^n \sum_{i=0}^n \binom{n}{i} (px)^{n-i} q^i (A\alpha^{n-i-1}\tilde{\alpha} + B\beta^{n-i-1}\tilde{\beta}) \\ &= A\tilde{\alpha}\alpha^{-1} \sum_{i=0}^n \binom{n}{i} (px\alpha)^{n-i} q^i + B\tilde{\beta}\beta^{-1} \sum_{i=0}^n \binom{n}{i} (px\beta)^{n-i} q^i \\ &= A\tilde{\alpha}\alpha^{-1} (px\alpha + q)^n + B\tilde{\beta}\beta^{-1} (px\beta + q)^n \\ &= A\tilde{\alpha}\alpha^{2n-1} + B\tilde{\beta}\beta^{2n-1} \\ &= \mathbb{H}_{2n}(x). \end{aligned}$$

Thus the proof is completed.  $\square$

**Corollary 2.14.** For  $n \geq 0$ , we have

$$\sum_{i=0}^n \binom{n}{i} x^{n-i} FH_{n-i}(x) = FH_{2n}(x).$$

*Proof.* If we take  $a = b = p = q = 1$  in Equation (2.11), the proof is completed.  $\square$

**Corollary 2.15.** For  $n \geq 0$ , we have

$$\sum_{i=0}^n \binom{n}{i} (2x)^{n-i} PH_{n-i}(x) = PH_{2n}(x).$$

*Proof.* If we take  $a = q = 1$  and  $b = p = 2$  in Equation (2.11), the proof is completed.  $\square$

**Corollary 2.16.** For  $n \geq 0$ , we have

$$\sum_{i=0}^n (-1)^n \binom{n}{i} (-6x)^{n-i} BH_{n-i}(x) = BH_{2n}(x).$$

*Proof.* If we take  $a = 1$ ,  $b = p = 6$ , and  $q = -1$  in Equation (2.11), the proof is completed.  $\square$

### 3. An application of Horadam hybrinomials in matrices

In this section, we derive the matrix representation of the Horadam hybrinomials. Then we obtain closed formula for the Horadam hybrinomials  $\mathbb{H}_n(x)$ , in terms of tridiagonal determinant (see [26]-[28]).

**Theorem 3.1.** Let  $n \geq 1$  be an integer. The following equality holds:

$$\begin{bmatrix} \mathbb{H}_{n+3}(x) & \mathbb{H}_{n+2}(x) \\ \mathbb{H}_{n+2}(x) & \mathbb{H}_{n+1}(x) \end{bmatrix} = \begin{bmatrix} \mathbb{H}_3(x) & \mathbb{H}_2(x) \\ \mathbb{H}_2(x) & \mathbb{H}_1(x) \end{bmatrix} \begin{bmatrix} px & 1 \\ q & 0 \end{bmatrix}^n. \quad (3.1)$$

*Proof.* For the proof, we use induction method on  $n$ . The equality holds for  $n = 1$ . Now suppose that the equality is true for  $n > 1$ . Then we can verify it for  $n + 1$  as follows:

$$\begin{aligned} \begin{bmatrix} \mathbb{H}_3(x) & \mathbb{H}_2(x) \\ \mathbb{H}_2(x) & \mathbb{H}_1(x) \end{bmatrix} \begin{bmatrix} px & 1 \\ q & 0 \end{bmatrix}^{n+1} &= \begin{bmatrix} \mathbb{H}_3(x) & \mathbb{H}_2(x) \\ \mathbb{H}_2(x) & \mathbb{H}_1(x) \end{bmatrix} \begin{bmatrix} px & 1 \\ q & 0 \end{bmatrix}^n \begin{bmatrix} px & 1 \\ q & 0 \end{bmatrix} \\ &= \begin{bmatrix} \mathbb{H}_{n+3}(x) & \mathbb{H}_{n+2}(x) \\ \mathbb{H}_{n+2}(x) & \mathbb{H}_{n+1}(x) \end{bmatrix} \begin{bmatrix} px & 1 \\ q & 0 \end{bmatrix} \\ &= \begin{bmatrix} \mathbb{H}_{n+4}(x) & \mathbb{H}_{n+3}(x) \\ \mathbb{H}_{n+3}(x) & \mathbb{H}_{n+2}(x) \end{bmatrix}. \end{aligned}$$

So the proof is completed.  $\square$







## 4. Conclusion

In our present research, we have studied Horadam hybrinomials which are defined by dint of the Horadam polynomials. We have obtained some properties of Horadam hybrinomials. Finally in Section 3, with the help of the two different tridiagonal matrix, we have obtained the  $n^{\text{th}}$  term of Horadam hybrinomials. According to the special cases of  $a$ ,  $b$ ,  $p$  and  $q$ , all the results given in Section 2 and Section 3 are applicable to all hybrinomials and hybrid numbers mentioned in this paper. The Horadam hybrinomials that we have defined include previously introduced the Fibonacci hybrinomials  $FH_n(x)$ , the Fibonacci hybrid numbers  $FH_n$ , the Lucas hybrinomials  $LH_{n-1}(x)$ , the Lucas hybrid numbers  $LH_{n-1}$ , the Pell hybrinomials  $PH_n(x)$ , the Pell hybrid numbers  $PH_n$ , the Pell-Lucas hybrinomials  $QH_{n-1}(x)$ , the Pell-Lucas hybrid numbers  $QH_{n-1}$  (see, [24, 25]). From the definition of the Horadam hybrinomials, we also have obtained the Chebyshev hybrinomials of the first kind  $TH_{n-1}(x)$ , the Chebyshev hybrid numbers of the first kind  $TH_{n-1}$ , the Chebyshev hybrinomials of the second kind  $UH_{n-1}(x)$ , the Chebyshev hybrid numbers of the second kind  $UH_{n-1}$ , the Balancing hybrinomials  $BH_n(x)$  and the Balancing hybrid numbers  $BH_n$ .

## Acknowledgements

The author would like to express his sincere thanks to the editor and the anonymous reviewers for their helpful comments and suggestions.

## Funding

There is no funding for this work.

## Availability of data and materials

Not applicable.

## Competing interests

The author declares that he has no competing interests.

## Author's contributions

The author read and approved the final manuscript.

## References

- [1] A. F. Horadam, *Basic properties of a certain generalized sequence of numbers*, Fibonacci Q., **3** (1965), 161-176.
- [2] A. F. Horadam, *Generating functions for powers of a certain generalized sequence of numbers*, Duke Math. J., **32** (1965), 437-446.
- [3] T. Koshy, *Fibonacci and Lucas Numbers with Applications*, Vol. 1, Second edition, Pure and Applied Mathematics (Hoboken), John Wiley & Sons, Inc., Hoboken, NJ, 2018.
- [4] S. Falcon, A. Plaza, *On the Fibonacci k-numbers*, Chaos Solitons Fractals, **32** (5) (2007), 1615-1624.
- [5] E. G. Kocer, N. Tuglu, A. Stakhov, *On the m-extension of the Fibonacci and Lucas p-numbers*, Chaos Solitons Fractals, **40** (4) (2009), 1890-1906.
- [6] C. Kızılateş, *New families of Horadam numbers associated with finite operators and their applications*, Math. Meth. Appl. Sci., **44** (2021), 14371-14381.
- [7] T. Horzum, E. G. Kocer, *On some properties of Horadam polynomials*, Int. Math. Forum., **25** (4) (2009), 1243-1252.
- [8] M. Özdemir, *Introduction to hybrid numbers*, Adv. Appl. Clifford Algebras, **28** (1) (2018).
- [9] A. Szynal-Liana, I. Wloch, *The Fibonacci hybrid numbers*, Utilitas Math., **110** (2019), 3-10.
- [10] A. Szynal-Liana, I. Wloch, *On Jacobsthal and Jacobsthal-Lucas hybrid numbers*, Ann. Math. Sil., **33** (1) (2019), 276-283.
- [11] A. Szynal-Liana, I. Wloch, *On Pell and Pell-Lucas Hybrid Numbers*, Commentat. Math., **58** (2018), 11-17.
- [12] A. Szynal-Liana, *The Horadam hybrid numbers*, Discuss. Math. Gen. Algebra Appl., **38** (1) (2018), 91-98.
- [13] C. Kızılateş, *A new generalization of Fibonacci hybrid and Lucas hybrid numbers*, Chaos Solitons Fractals, **130** (2020), 1-5.
- [14] P. Catarino, *On k-Pell hybrid numbers*, J. Discrete Math. Sci. Cryptography, **22** (1) (2019), 83-89.
- [15] G. Cerda Moreles, *Investigation of generalized hybrid Fibonacci numbers and their properties*, Appl. Math. E-Notes, **21** (2021), 110-118.
- [16] G. Cerda Moreles, *Introduction to third-order Jacobsthal and modified third-order Jacobsthal hybrinomials*, Discuss. Math. Gen. Algebra Appl., **41** (1) (2021), 139-152.
- [17] Y. Alp, E. G. Kocer, *Hybrid Leonardo numbers*, Chaos Solitons Fractals, **150** (2021), 1-5.
- [18] D. Tasci, E. Sevgi, *Some properties between Mersenne, Jacobsthal and Jacobsthal-Lucas hybrid numbers*, Chaos Solitons Fractals, **146** (2021), 1-4.
- [19] E. Tan, NR. Ait-Amrane, *On a new generalization of Fibonacci hybrid numbers*, (2020), arXiv:2006.09727.
- [20] S. Petroudi, M. Pirouz, A. Özkoç, *The Narayana polynomial and Narayana hybrinomial sequences*, Konuralp J. Math., **9** (1) (2021), 90-99.
- [21] A. Özkoç, *A new generalization of Tribonacci hybrinomial*, Bull. Int. Math. Virtual Inst., **11** (3) (2021), 555-568.
- [22] R. Vieira, M. Manguera, F. R. Alves, P. M. M. Cruz Catarino, *Padovan and Perrin Hybrid Number Identities*, Commun. Adv. Math. Sci., **4** (4) (2021), 190-197.
- [23] E. Polath, *A note on ratios of Fibonacci hybrid and Lucas hybrid numbers*, Notes Number Theory Discrete Math., **27** (3) (2021), 73-78.
- [24] A. Szynal-Liana, I. Wloch, *Introduction to Fibonacci and Lucas hybrinomials*, Complex Var. Elliptic Eq., **65** (10) (2020), 1736-1747.
- [25] M. Liana, A. Szynal-Liana, I. Wloch, *On Pell hybrinomials*, Miskolc Math. Notes, **20** (2019), 1051-1062.
- [26] S. Falcón, *On the generating matrices of the k-Fibonacci numbers*, Proyecciones J. Math., **32** (4) (2013), 347-357.
- [27] P. Catarino, *A note on certain matrices with h(x)-Fibonacci quaternion polynomials*, J. Differ. Eq. Appl., **22** (2016), 343-351.
- [28] C. Kızılateş, P. Catarino, N. Tuglu, *On the bicomplex generalized Tribonacci quaternions*, Mathematics, **7** (1) (2019), 80.
- [29] C. Kızılateş, *A Note on Horadam hybrinomials*, (2020), Preprints:2020010116.