



The Effects of GA₃ Treatments and Nutrient Media on *In Vitro* Seed Germination of *Allium tuncelianum* (Kollman), Özhatay, Matthew, Şiraneci

Faika YARALI KARAKAN^{1*} 

¹ Department of Horticulture, Faculty of Agriculture, University Kilis 7 Aralik, 79000, Kilis, Turkey,
E-mail: faikayarali@gmail.com, ORCID ID: [0000-0002-2176-8663](https://orcid.org/0000-0002-2176-8663)

¹ Advanced Technology Application and Research Center, University Kilis 7 Aralik, 79000, Kilis, Turkey
*Corresponding author : faikayarali@gmail.com

<https://doi.org/10.38093/cupmap.1034719>

Received: 09/12/2021
Accepted: 30/12/2021

Abstract

In this study, the effects of GA₃ and nutrient media (BDS and MS) on *in vitro* seed germination of *Allium tuncelianum* (Kollman), N. Özhatay, D. Matthew, Şiraneci, which is one of the endemic plants in Turkey, were investigated. In order to promote seed germination, GA₃ was applied to seeds under sterile conditions for 48 hours at different concentrations (2 mg L⁻¹, 3 mg L⁻¹ and 4 mg L⁻¹). Then the seeds were cultured in BDS and MS medium. The results showed that the germination rate decreased with the increase of the applied concentrations of GA₃. The highest germination rate (17.73%) was obtained from seeds treated with 2 mg L⁻¹ GA₃. It was determined that the effects of nutrient media on the germination of *Allium tuncelianum* seeds were important, and the highest germination values were obtained from the BDS medium.

Key Words: *Allium tuncelianum*, BDS, GA₃, Germination, MS

© CUPMAP. All rights reserved.

1. Introduction

Allium tuncelianum (Kollman), Özhatay, Matthew, Şiraneci, which is locally called as "Tunceli garlic" or "Ovacık garlic", is an endemic species unique to Turkey. It is widely found in especially in Ovacık and Pulümür districts of Tunceli province located at the foot of Munzur Mountain. *Allium tuncelianum* has single cloved, cream-white bulbs and also has the ability to flower and give fertile black seeds different from common garlic (Yanmaz and Ermiş, 2005; Yanmaz et al., 2010; Yarali and Yanmaz, 2016; Kizil et al., 2017; Yarali Karakan, 2019).

It has been reported that consumption of *Allium tuncelianum* stimulates the body's immune system, lowers blood sugar and cholesterol levels, improves blood circulation and thus reduces the risk of heart attack (Agbas et al., 2013; Atila et al., 2017; Yarali Karakan, 2019). In addition, Sehitoglu et al. (2018) stated that *Allium tuncelianum* has a stronger antioxidant, antiradical activity and effective essential omega acid levels than common garlic in terms of fatty acid compositions.

Allium tuncelianum is usually propagated by seeds and cloves. However, there is a problem of germination in the seeds due to dormancy that occurs after the seeds mature.

This situation makes it difficult to propagate the plant by seed (Alper, 2005; Kizil et al., 2017; Yarali Karakan, 2019). Therefore, extensive studies are needed to develop tissue culture techniques for *in vitro* propagation and seed germination of *Allium tuncelianum* (Kizil et al., 2014; Yarali Karakan, 2019). In this study, the effect of GA₃ as a germination stimulant and nutrient media were investigated *in vitro* conditions in order to ensure the propagation of *Allium tuncelianum* by seed.

2. Material and Methods

Allium tuncelianum seeds were kept in sodium hypochlorite solution (containing 5% NaOCl) for 20 minutes to ensure surface

sterilization, and then washed three times for three minutes with distilled water (Kizil et al., 2017). In order to promote germination, different concentrations of GA₃ (2 mg L⁻¹, 3 mg L⁻¹ and 4 mg L⁻¹) were applied to the seeds for 48 hours under sterile conditions. Then the seeds were cultured on BDS (Dunstan and Short, 1977) and MS (Murashige and Skoog, 1962) medium supplemented with 2.0% sucrose at 4 °C (Kizil et al., 2017), with 15 replications from each application and 15 seeds in each replication under *in vitro* conditions (Figure 1). 2 mm rootlets were accepted as germination criterion (Anonymous, 1985; Figure 2).



Figure 1. a) Sterilization of seeds, b) Filter sterilization of GA₃ solutions, c) Seeds treated at different GA₃ concentrations for 48 hours, d) Cultured seeds.



Figure 2. Germinated seeds in BDS medium treated with 2 mg L⁻¹ GA₃

2.1. Statistical Analysis

The experiment was established according to completely randomized design (CRD) with 15 replications. Tukey test was used to identify different groups after analysis of variance. The level of statistical significance was taken as 5% and calculations were made with JMP software package version 5.0.1.

3. Results and Discussion


In this research, in order to reveal the effects of plant growth regulator application, the seeds were treated with gibberellic acid solution at different concentrations (2 mg L⁻¹, 3 mg L⁻¹ and 4 mg L⁻¹ GA₃) prepared under sterile conditions for 48 hours. Then, all seeds were cultured in BDS and MS medium at 4 °C. The effects of different concentrations of GA₃ applications and nutrient media on the germination of *Allium tuncelianum* seeds are shown in Table 1. When Table 1 was examined, it was determined that the seeds applied with 2 mg L⁻¹ GA₃ had the highest germination rate of 17.73%. The germination rate decreased with the increase of the applied dose. Thus, the lowest germination rate was obtained from the application of 4 mg L⁻¹ GA₃ in both nutrient media. It has been reported that storage at 4°C results in hydrolysis of seed starch resulting in mobility of

carbohydrates in *Allium* species. Thus, macro-carbohydrate molecules are converted into sucrose, glucose and fructose, which are utilized during cell metabolism and energy required for the growth of plants. These have significant effects to break seed dormancy of garlic (Kizil et al., 2017). Similarly, it was reported that garlic seeds treated with low temperature at 4°C and gibberellic acid (GA₃) were helpful in seed dormancy break (Arguello et al. 2001). Yanmaz and Ermis (2005) used 0.5, 1.0, 2.0 mg L⁻¹ doses of GA₃ for 24 and 48 hours in their study aiming to solve the germination problem in *Allium tuncelianum* seeds. They stated that GA₃ application for 48 hours was more effective on the germination and also it has been determined that 1.0 mg L⁻¹ and 2.0 mg L⁻¹ doses of GA₃ stimulate germination and increase the germination rate up to 18-20% depending on the low temperature period. In the study, it was determined that the effects of nutrient media on the germination of *Allium tuncelianum* seeds were also important. When BDS and MS media were compared, the highest germination values were obtained from BDS medium (Table 1). On the other hand, in their study to break dormancy in *Allium tuncelianum* seeds, Kizil et al., (2017) reported that it was higher in MS medium.

Table 1. Effects of GA₃ treatments and different nutrient media on germination of *Allium tuncelianum* seeds

GA ₃ treatments	Nutrient media			
	BDS		MS	
	Germinated seeds (number/petri)	Germination rate (%)	Germinated seeds (number/petri)	Germination rate (%)
Control	0,80b	5,33	1,13 ab	7,53
2 mg L ⁻¹	2,66a	17,73	1,33ab	0,86
3 mg L ⁻¹	1,26ab	8,40	1,53ab	10,20
4 mg L ⁻¹	0,40b	2,66	0,46b	3,06

Significant at P = 0.05 level

**Figure 3.** Number of germinated seeds in different GA₃ treatments and nutrient media.

4. Conclusion

The effects of GA₃ treatments and nutrient media on seed germination were found to be significant. The seed germination rate was higher at 2 mg L⁻¹ GA₃ treatment than the other treatments. And also it was determined that the effects of nutrient media on the germination of *Allium tuncelianum* seeds were also important and the highest germination values were obtained from BDS medium. The results obtained from this research will contribute to future research for protection and propagation of *Allium tuncelianum*.

Author Contribution

Faika YARALI KARAKAN designed the experiments and wrote the paper.

Acknowledgements

-

Conflicts of Interest

The author declares no conflict of interest.

References

1. Agbas, B., Karakus, D., Adiguzel, G., Keser, S., Demir, E., (2013). Comparison of total antioxidant properties and dry matter content of Tunceli garlic (*Allium tuncelianum*) and normal garlic

- (*Allium sativum*), *Bilim ve Genclik Dergisi*, 1(2), 50-62. ISSN: 2148- 0273.
2. Alper, A., (2005). Tunceli sarımsağının (*Allium tuncelianum* Kollman, Özhatay, Marhew, Şiraneci) bitkisel özelliklerinin belirlenmesi. Ankara Üniversitesi Fen Bilimleri Enstitüsü, Bahçe Bitkileri Anabilim Dalı, Ankara, p16.
 3. Anonymous, (1985). International Rules for Seed Testing. *Seed Science and Technology*, 13; 421-463.
 4. Arguello, J.A., Falcon, L.R., Seiseddos, L., Milrad, S., Bottini, R., (2001). Morphological changes in garlic (*Allium sativum* L.) microbulblets during dormancy and sprouting as related to peroxidase activity and gibberellin GA₃ content. *Biocell.*, 25: 1-9.
 5. Atila, G., Uslu, H., Erdag, D., Ozkan, O., (2017). Antioxidative and antihyperglycemic effects of *Allium tuncelianum* on streptozotocin induced type I diabetes, *International Congress on Medicinal and Aromatic Plants*, p: 411, May 10-12, 2017, Konya.
 6. Dunstan, D.I., Short, K.C., (1977). Improved growth of tissue cultures of the onion, *Allium cepa*. *Physiol Plant*, 41: 70-72.
 7. Kizil, S., Icgil, D.Y., Khawar, K.M., (2014). Improved in vitro regeneration and propagation of Tunceli garlic (*Allium tuncelianum* L.) from sectioned garlic cloves, leaves and root explants. *Journal of Horticultural Science and Biotechnology*, 89(4): 408-414.
 8. Kizil, S., Sogut, T., Khawar, K.M., (2017). Germination and bulb formation of *Allium tuncelianum* L. under *in vitro* condition, *ICAB 2017: 19th International Conference on Agriculture and Biotechnology*. 13-14 July, Stockholm, Sweden.
 9. Murashige, T., Skoog, F., (1962). A Revised medium for rapid growth and bioassay with tobacco tissue cultures. *Physiologia Plantarum*, 15: 473-497.
 10. Sehitoglu, M.H., Yarali Karakan, F., Kizilkaya, B., Oztopuz, R.O., Gulcin, I., (2018). Investigation of antioxidant properties and bioactive composition of *Allium tuncelianum* ((kollman) ozhatay, matthew & siraneci) and *Allium sativum* L. *Journal of the Institute of Science and Technology*, 8(4): 213-221.
 11. Yanmaz, R., Ermiş, S., (2005). Tunceli sarımsağı (*Allium tuncelianum* Kollman, Özhatay, Matthew, Şiraneci) tohumlarındaki çimlenme probleminin çözülmesi üzerine araştırma/ Researches On Overcoming The Germination Problems Of Tunceli Garlic Seeds (*Allium tuncelianum* . (Kollman), N. Ozhatay, D. Matthew, Siranec . Türkiye 2. Tohumculuk kongresi, Adana, s: 101-106.
 12. Yanmaz, R., Yazar, E., Kantoğlu, Y., and Alper, A., (2010). *In vitro* plant regeneration and bulblet formation of Tunceli garlic (*Allium tuncelianum* (Kollman) Özhatay, Matthew, Siraneci) by shoot and root culture. *Journal of Food, Agriculture & Environment*, Vol: 8(3-4): 572- 576.
 13. Yarali, F., Yanmaz, R., (2016). The effects of media composition on *in vitro* gynogenic embryo induction in *Allium tuncelianum* (Kollman) Ozhatay, Matthew, Siraneci. *International Journal of Agriculture Sciences*, ISSN: 0975-3710 & E-ISSN: 0975-9107, 8(58): 3247-3251.
 14. Yarali Karakan, F., (2019). A review of *in vitro* studies on *Allium tuncelianum* (Kollman) Ozhatay, Matthew, Siraneci, *Scientific Bulletin. Series F. Biotechnologies*, Vol. XXIII, ISSN 2285-1364, CD-ROM ISSN 2285-5521, ISSN Online 2285-1372, ISSN-L 2285-1364, p: 11-16.