

Contribution to the Knowledge of Phylinae (Hemiptera: Heteroptera: Miridae) from Turkey

Gülten YAZICI¹ Erol YILDIRIM²

ÖZ

Türkiye Phylinae (Hemiptera: Heteroptera: Miridae) türlerine katkılar

Bu çalışma çoğunuğu 2006-2014 yıllarında olmak üzere, 1978 ve 2014 yılları arasında Türkiye'nin değişik lokalitelerinden toplanmış olan Miridae familyasının Phylinae altfamilyasına ait materyalden oluşmaktadır. Bu çalışmada Phylinae altfamilyasına ait 20 cinse bağlı 32 tür tespit edilmiştir. Bunlar arasından, *Chlamydatus (Euattus) pullus* (Reuter 1870), *Megalocoleus molliculus* (Fallén 1807), *Oncotylus (Oncotylus) punctipes* Reuter 1875, *Oncotylus (Oncotylus) viridiflavus* (Goeze 1778), *Opisthotaenia (Opisthotaenia) fulvipes* Reuter 1901, *Plagiognathus (Plagiognathus) bipunctatus* Reuter 1883 ve *Plagiognathus chrysanthemi* (Wolff 1804) türleri bol ve yaygın olarak bulunmuştur. Ayrıca, Türkiye'den daha önce bildirilen bazı türleri için yeni lokaliteler ilave edilmiştir.

Anahtar kelimeler: Hemiptera, Heteroptera, Miridae, Phylinae, Türkiye

ABSTRACT

This study is based upon material of subfamily Phylinae from family Miridae collected from different localities of Turkey between 1978 and 2014, mostly 2006-2014. In this study, 32 species from 20 genera of Phylinae species were determined. Among them, *Chlamydatus (Euattus) pullus* (Reuter 1870), *Megalocoleus molliculus* (Fallén 1807), *Oncotylus (Oncotylus) punctipes* Reuter 1875, *Oncotylus (Oncotylus) viridiflavus* (Goeze 1778), *Opisthotaenia (Opisthotaenia) fulvipes* Reuter 1901, *Plagiognathus (Plagiognathus) bipunctatus* Reuter 1883 and *Plagiognathus chrysanthemi* (Wolff 1804) have been found the most abundant and widespread species. In addition, new localities are added for some species previously reported for Turkey.

Keywords: Hemiptera, Heteroptera, Miridae, Phylinae, Turkey.

¹ Plant Protection Central Research Institute, 06172 Yenimahalle, Ankara

² Atatürk University, Faculty of Agriculture, Department of Plant Protection, 25240-Erzurum
Sorumlu Yazar (Corresponding author) e-mail: gulten.yazici@gthb.gov.tr
Alınış (Received): 18.02.2016 , Kabul ediliş (Accepted): 10.07.2016

INTRODUCTION

Miridae (or plant-bugs) is a group of Miroidea, which are represented by nearly 10.040 species, 1507 genera (Cassis et al. 2006). This family comprising eight subfamilies Isometopinae, Psallopinae Cylapinae, Orthotylinae, Bryocorinae, Deraeocorinae, Mirinae and Phylinae (Cassis and Schuh 2012). The subfamily Phylinae is small, rarely medium-sized, usually phytophagous, often with narrow food specialization; less commonly zoophytophagous or predacious, hibernating as eggs (with very rare exception) and the largest subfamily (Lehr 1988). Turkey is biogeographically one of the most interesting countries in the West Palaearctic region. Some faunistic studies on this family in the Turkey have been made by Hoberlandt (1955), Önder (1976), Bingöl (1978), Lodos et al. (1978), Altinayar (1981), Önder et al. (1981), Yayla (1983), Özkan (1984), Karaat (1986), Önder and Lodos (1987), Özbek and Alaoğlu (1987), Çam (1988), Lodos et al. (1989), Kiyak (1990), Önder et al. (1990; 1998), Yıldırım and Özbek (1992), Güçlü et al. (1995 a,b), Çevik (1996), Yaşarakinci and Hincal (1997;2000), Tezcan and Önder (1999; 2003), Yıldırım et al. (1999), Atakan (2000), Beyaz (2000), Yıldırım (2002), Lodos (2003), Kiyak et al. (2004), Çetin and Alaoğlu (2005), Ayyıldız and Atlıhan (2006), Önder et al. (2006).

The aim of this paper is to present new distributional and biological data on Phylinae in Turkey.

MATERIAL AND METHOD

The material of the Phylinae was collected from different localities of Turkey between 1978 and 2014. In addition, in previous years the material collected from different localities of Turkey, and preserved in the Entomology Museum, Erzurum, Turkey (EMET) is also evaluated. Material was obtained by sweeping from meadow and pasture lands containing a variety of flowering plants. Provinces of the collected specimens are given in alphabetical order in the following list. The material is deposited in EMET. The species were identified by Dr. Rauno Linnauori (Finland), Dr. Chérot Frédéric (Belgium) and Prof. Dr. Jacek Gorczyca (Poland).

RESULTS

In this study, 32 species and 20 genera belonging to subfamily Phylinae is recorded from Turkey.

Subfamily Phylinae

Amblytylus nasutus (Kirschbaum 1856)

Material examined: Erzurum: Yakutiye, Dumlubaba, Güngörmez, 2500 m, 1.VII.2010, ♀.

Distribution in Turkey: Edirne (Hoberlandt 1955); Afyonkarahisar, Ankara, Antalya, Çanakkale, Erzincan, Konya, Mersin (Önder 1976; Önder et al. 2006); Bursa, Kocaeli, Sakarya (Önder et al. 1981); İçel (Lodos et al. 2003); İzmir (Tezcan et al. 2010).

Atomoscelis onustus (Fieber 1861)

Material examined: **Erzincan:** 27.VIII.1991, ♀; **Erzurum:** Oltu, 5.VII.1988, ♀, 2 ♂♂, İnanmış, N 40°27'50.1", E 041°42'19.9", 1823 m, 6.VII.2012, ♂; Pasinler, 18.VII.1987, 2 ♀♀, 3 ♂♂; Yakutiye, Atatürk University field, 1850 m, 2.VIII.1988, ♀.

Distribution in Turkey: Ankara, Antalya, Çorum, Eskişehir, Gaziantep, Hatay, Kahramanmaraş, Karabük, Kayseri, Kırıkkale, Kırşehir, Konya, Mersin, Nevşehir, Niğde, Yozgat (Önder 1976; Önder et al. 2006); Bursa, Sakarya (Önder et al. 1981); Diyarbakır (Önder et al. 1995); İçel (Lodos et al. 2003).

Campylomma diversicornis Reuter 1878

Material examined: **Antalya:** 23.VII.1984, ♂, 18.VIII.1984, ♀, ♂; **Erzurum:** Pasinler, 18.VII.1987, 5 ♀♀, 8 ♂♂.

Distribution in Turkey: Adana, İğdır (Hoberlandt 1955); Aydın, Çanakkale, Denizli, Diyarbakır, Elazığ, Mardin, Muğla (Önder 1976); İzmir (Giray 1980; Tezcan et al. 2010); Bursa, Kocaeli, Sakarya (Önder et al. 1981); İçel (Öncüler 1991); Adıyaman (Önder et al. 1995); Manisa (Beyaz 2000); Antalya, Gaziantep, Hatay (Lodos et al. 2003); West, Eastern, Southeast Anatolia Regions (Önder et al. 2006); Kahramanmaraş (Ölmez 2008).

Campylomma nicolasi Puton and Reuter, 1883

Material examined: **Antalya:** 23.VII.1984, 2 ♀♀; **Erzurum:** Yakutiye, Atatürk University field, 1850 m, 7.VIII.1996, 2 ♀♀.

Distribution in Turkey: Adana, Edirne (Hoberlandt 1955); Aydın, Bilecik, Burdur, Hatay, Isparta, Kırklareli, Manisa, Mardin, Muğla, Samsun, Sinop, Tekirdağ (Önder 1976); İzmir (Giray 1980; Yaşarakinci and Hıncal 2000; Tezcan et al. 2010); Bursa, Kocaeli, Sakarya (Önder et al. 1981); Antalya (Öncüler 1991); Diyarbakır, Mardin (Önder et al. 1995); Adana, Ankara, Çankırı, Çorum, Eskişehir, Gaziantep, İçel, Kahramanmaraş, Kastamonu, Kayseri, Kırşehir, Konya, Nevşehir, Niğde (Lodos et al. 2003); Mediterranean, Southeast Anatolia, Aegean, Central Anatolia, Black Sea, Marmara regions (Önder et al. 2006).

Campylomma verbasci (Meyer-Dür 1843)

Material examined: **Bayburt:** Çalıdere, 1700 m, 17.VI.2010, ♂; **Erzincan:** 27.VIII.1991, 5 ♀♀, 4 ♂♂, Mercan, 1381 m, 10.VI.2010, ♂, Yukarı Çakırman, 28.VIII.1991, ♀; **Erzurum:** Aziziye, Başçakmak, N 40°00'5.6", E 040°54'39.3", 1860 m, 20.VII.2011, ♂, 14.VIII.1987, ♀; Hinis, 1742 m, 2.VII.2010, ♀; Köprüköy,

Örentaş, N 40°27'48.6", E 041°39'32.3", 1916 m, 30.VII.2011, ♀, 3 ♂♂; Oltu, 1750 m, 4.VIII.2009, 2 ♂♂, 4.IX.1991, 3 ♀♀, Sarısaç, N 40°32'00.9", E 041°54'27.2", 1421 m, 30.VIII.2012, ♀; Olur, Taşlıköy, 20.VIII.1992, 3 ♀♀, ♂; Pazaryolu, N 40°25'12.8", E 040°46'13.5", 1450 m, 24.VII.2011, ♀, Gülcimen, N 40°24'36.4", E 040°47'28.5", 1663 m, 20.VII.2011, ♀, ♂; Tekman, Güzeldere, N 39°49'17.1", E 042°00'19.8", 1948 m, 16.VII.2011, ♂, Uzundere, Yayla, N 40°27'48.6", E 041°39'32.3", 2005 m, 6.VII.2012, ♂; **Iğdır:** 850 m, 20.VI.2009, ♀; **Kars:** Kağızman, 23.VII.1991, ♂, Sarıkamış, Karakurt, 1500 m, 13.VIII.2009, ♂.

Distribution in Turkey: Adana, Edirne (Hoberlandt 1955); Aydın, Burdur, Çanakkale, Denizli, Edirne, Isparta, İzmir, Mersin (Önder 1976); Bolu, Bursa, Sakarya (Önder et al. 1981); Antalya (Özkan 1984); Diyarbakır (Önder et al. 1995); İçel (Lodos et al. 2003); Balıkesir (Ayyıldız and Atlıhan 2006); Ankara, Aydın, Balıkesir, Burdur, Çanakkale, Çankırı, Denizli, Eskişehir, Gaziantep, Hatay, Isparta, İzmir, Kahramanmaraş, Karaman, Kastamonu, Kayseri, Kırıkkale, Kırşehir, Konya, Mersin, Nevşehir, Niğde, Yozgat, Zonguldak (Önder et al. 2006); Mardin (Matocq and Özgen 2010); Siirt (Matocq et al. 2014).

Chlamydatus (Euattus) pullus (Reuter 1870)

Material examined: Erzincan: Akyazı, 1202 m, 11.VI.2010, ♂; Erzurum: Aşkale, Çayköy, 1600 m, 3.VI.2010, 11 ♂♂, N 39°56'46.6", E 040°48'27.4", 1720 m, 21.VI.2011, 3 ♀♀, 8 ♂♂, Çiftlik, 1620 m, 3.VI.2010, ♂, Kandilli, N 39°54'38.5", E 040°51'19.9", 1710 m, 21.VI.2011, 2 ♀♀, Merdiven, N 39°53'45.2", E 040°48'29.9", 1706 m, 9.VIII.2011, 2 ♀♀; Aziziye, Dadaşkent, 1800 m, 3.VI.2010, 2 ♀♀, 2 ♂♂, Eğerti, N 40°0'6'13.6", E 040°58'44.5", 1906 m, 22.VI.2011, 2 ♂♂, Paşayurdu, 1710 m, 3.VI.2010, 3 ♂♂, Çat, 2200 m, 3.VI.2010, ♂; Horasan, Değirmenli, N 40°0'5'10.7", E 042°07'01.0", 1643 m, 13.VIII.2011, 3 ♀♀, ♂; İspir, Aşağı Özbağ, N 40°28'02.3", E 040°58'25.8", 1177 m, 15.VII.2012, ♀; Narman, 30.VI.1989, ♂, İncedere, 1987 m, 21.VII.2010, ♀, Kamışözü, N 40°20'23.3", E 041°47'29.2", 1835 m, 23.VI.2011, ♀, ♂, Ünlükaya, 1700 m, 8.VI.2012, ♀; Oltu, 12.VII.1987, 2 ♂♂, 22.VII.1987, 2 ♂♂, 23.VIII.1987, ♂, Çamlıbel, N 40°28'34.3", E 041°46'24.1", 1682 m, 31.VII.2011, 2 ♀♀, 2 ♂♂; Olur, Coşkunlar, N 40°44'32.7", E 042°11'11.4", 1054 m, 30.VII.2011, 2 ♀♀, 2 ♂♂, Kaledibi, 1070 m, 4.VIII.2009, ♂; Palandöken, Kümbet, N 39°48'59.7", E 041°4'32.8", 1832 m, 5.VIII.2011, ♀, ♂; Pasınler, 2.VII.1983, 2 ♂♂, 11.VII.1987, ♂, 1.VIII.1983, ♂, Ovaköy, N 39°58'56", E 041°29'38", 1782 m, 31.VII.2014, 3 ♂♂; Şenkaya, Sındırان, N 40°37'14.2", E 042°22'09.3", 1497 m, 10.VII.2011, 2 ♀♀, 5 ♂♂, Timurtaşla, N 40°33'43.8", E 042°16'58.0", 1314 m, 14.VII.2012, ♂; Tortum, N 40°16'38", E 041°33'29.9", 1660 m, 31.VII.2011, ♂, Derekapi, N 40°37'59.5", E 041°42'56.7", 1010 m, 7.VI.2012, ♀, 3 ♂♂, Şenyurt, N 40°26'22.0", E 041°28'39.5", 1265 m, 21.VII.2012, 2 ♀♀; Yakutiye, Atatürk University field, 1850 m, 20.VI.1996, ♀, 23.VI.1980, 2 ♀♀, 12 ♂♂, 29.VII.1987, 4 ♂♂, 14.VIII.1987, ♂, 24.VIII.1987, 3 ♂♂, Dadaşköy, 1806 m, N 39°55'39.5", E 041°15'23.2", 4.VIII.2011, ♀, ♂.

Distribution in Turkey: Bursa, Kocaeli (Hoberlandt 1955); Artvin, Kars, Manisa, Niğde (Önder 1976); Ağrı, Elazığ, Erzincan, Erzurum, Malatya, Muş, Tunceli Van (Bingöl 1978); Kahramanmaraş, Kayseri (Kiyak 1990); Aksaray, Çankırı, Karabük, Nevşehir (Lodos et al. 2003; Önder et al. 2006).

Chlorillus pictus (Fieber 1864)

Material examined: **Erzurum:** Karayazı, Kırgındere, 2215 m, 2.VII.2010, ♀; Oltu, Tutmaç, N 40°26'46.6", E 041°44'06.5", 1756 m, 6.VII.2012, ♂; Şenkaya, Gözebaşı, N 40°24'39.9", E 042°18'11.5", 1794 m, 14.VII.2012, 2 ♀♀; Tortum, Y. Sivri, N 40°20'27.5", E 041°39'01.1", 1962 m, 16.VII.2012, ♀, ♂; Yakutiye, Akdağ, 1820 m, 1.VII.2010, ♀, ♂, Atatürk University field, 1850 m, 25.VI.2009, ♂.

Distribution in Turkey: No locality record (Wagner 1955).

Conostethus roseus (Fallén 1807)

Material examined: **Erzurum:** Aziziye, Paşayurdu, 1710 m, 3.VI.2010, ♀; Çat, 2200 m, 3.VI.2010, ♂.

Distribution in Turkey: Adana (Hoberlandt 1955); Afyonkarahisar, Ağrı, Antalya, Erzurum, Mersin, Uşak (Önder 1976); İçel (Lodos et al. 2003); Ankara, Eskişehir, Karaman, Konya (Önder et al. 2006).

Ephippiocoris lunatus Poppius 1912

Material examined: **Erzurum:** Yakutiye, University field, 1850 m, 16.VII.1996, ♀, ♂; Pasinler, 1.VIII.1983, ♀.

Distribution in Turkey: Iğdır (Hoberlandt 1955).

Eurycolpus aureolus Seidenstucker 1961

Material examined: **Erzurum:** Tekman, Körsu, N 39°34'8.2", E 041°44'30.2", 1901 m, 16.VII.2011, 8 ♀♀, 11 ♂♂.

Distribution in Turkey: Adana, Ankara, Kırşehir, Mersin, Niğde (Önder 1976; Önder et al. 2006); Kahramanmaraş, Kayseri (Kiyak 1990); İçel (Lodos et al. 2003); Diyarbakır (Matocq et al. 2014).

Europiella alpina (Reuter 1875)

Material examined: **Erzurum:** Köprüköy, Eğirmez, N 39°58'09.9", E 041°49'20.7", 1670 m, 13.VIII.2011, 2 ♂♂; Oltu, Toprakkale, N 40°27'55.5", E 041°59'21.1", 1416 m, 23.VI.2011, ♂; Palandöken, Palandöken Mountain, 2300 m, 1.VIII.2010, ♀; Şenkaya, İkizpinar, N 40°36'57.4", E 042°21'30.9", 1589 m, 31.VII.2011, ♀; Yakutiye, Atatürk University field, 1850 m, 25.VI.2009, ♀.

Distribution in Turkey: Ankara (Hoberlandt 1955).

Macrotylus herrichi (Reuter 1873)

Material examined: **Erzurum:** Aziziye, Başçakmak, N 40°00'05.6", E 040°54'39.3", 1860 m, 20.VII.2011, ♂; Çat, N 39°35'42.4", E 040°57'59.6", 1913 m, 23.VII.2011, ♀, Aşağı Çat, N 39°38'29.5", E 041°00'11.7", 2065 m, 9.VIII.2011, ♀; Narman, Toygarlı, N 40°12'42.9", E 041°53'17.8", 2123 m, 19.VII.2011, ♂; Uzundere, Yayla, N 40°27'48.6", E 041°39'32.3", 2005 m, 13.VII.2011, ♂; Yakutiye, Atatürk University field, 1850 m, 7.VIII.1996, ♂.

Distribution in Turkey: Ankara, Kayseri, Konya (Hoberlandt 1955; Önder 1976); Antalya, Gaziantep, Kahramanmaraş (Lodos et al. 2003; Önder et al. 2006).

Megalocoleus molliculus (Fallén 1807)

Material examined: **Bayburt:** 1569 m, 22.VI.2009, ♂; **Erzurum:** Aşkale, 1600 m, 8.VII.2009, ♂, Çayköy, N 39°56'46.6", E 040°48'27.4", 1720 m, 21.VI.2011, 3 ♀♀, 2 ♂♂, Küçükgeçit, N 39°56'43.2", E 040°47'12.1", 1709 m, 23.VII.2011, 2 ♂♂, Küçükova, N 39°47'36.8", E 040°42'53.3", 1896 m, 5.VII.2012, ♂; Aziziye, Dallıkavak Pass, N 40°10'33.55", E 040°58'16.1", 2349 m, 20.VII.2011, ♀, Rizekent, 2070 m, 30.VII.2010, 3 ♂♂; Çat, 1894 m, 29.VI.2009, ♂; Hınıs, 1742 m, 2.VII.2010, ♀; Horasan, Değirmenli, N 40°05'20.7", E 042°06'56.7", 1608 m, 22.VI.2012, 2 ♀♀, ♂, Dönertaş, N 40°13'33.4", E 042°06'58.5", 2009 m, 13.VIII.2012, ♂; Karaçoban, Duman, N 39°18'22", E 041°55'5", 1560 m, 26.VI.2011, 2 ♀♀, Maruf, N 39°18'42", E 041°55'5", 1550 m, 26.VI.2011, 2 ♂♂; Karayazı, Yukarı Söylemez, 1896 m, 2.VII.2010, ♂; Köprüköy, Geyikli, N 39°48'02.1", E 042°04'05.4", 2144 m, 17.VII.2012, ♂; Narman, Demirdağ, N 40°19'23.5", E 041°42'45.5", 2028 m, 16.VII.2012, ♂, İncedere, 1987 m, 21.VII.2010, ♂; Oltu, Çamlıbel, N 40°29'36.1", E 041°45'45.2", 1775 m, 13.VII.2011, ♂, N 40°29'06", E 041°45'47", 1635 m, 14.VII.2014 m, ♂, Özdere, N 40°27'57.7", E 041°44'11.9", 1927 m, 16.VII.2012, ♀, ♂, Tutlu, N 40°36'53.3", E 040°5'39.4", 1240 m, 23.VI.2011, 2 ♀♀, Tutmaç, N 40°26'46.6", E 041°44'06.5", 1756 m, 6.VII.2012, ♂; Olur, Yeşilbağlar, 1000 m, 16.VI.2010, ♂; Palandöken, Taşlıgüney, 2032 m, 26.VI.2009, 2 ♂♂; Pasinler, Yayla, N 40°05'42.7", E 041°44'03.5", 1990 m, 17.VII.2011, 2 ♀♀; Şenkaya, Başaklı, N 40°29'19.2", E 041°48'18.6", 1586 m, 13.VII.2011, 2 ♀♀, İlkizpinar, N 40°36'57.4", E 042°21'30.9", 1589 m, 31.VII.2011, ♀; Tekman, Çiçekdağı, N 39°34'40.8", E 041°44'26.8", 1846 m, 26.VI.2011, ♂, Geyikli, N 39°47'48.5", E 042°4'31.4", 2185 m, 16.VII.2011, ♂; Yakutiye, Atatürk University field, 1850 m, 3.VII.1996, ♂, 10.VII.1996, ♂, 16.VII.1996, ♂, 14.VIII.1998, ♂, Dumlubaba, Güngörmez, 2500 m, 1.VII.2010, ♂, Güzelyayla, 2000 m, 19.VIII.2009, ♂.

Distribution in Turkey: Ankara (Hoberlandt 1955); Balıkesir, Bilecik, Isparta, Kars, Kırklareli, Kütahya, Manisa, Muğla, Tekirdağ, (Önder 1976); Bolu, Bursa, İstanbul, Kocaeli, Sakarya (Önder et al. 1981); Şanlıurfa (Önder et al. 1995); Çankırı, Karaman, Kastamonu, Kayseri, Kırıkkale, Kırşehir, Konya, Nevşehir (Lodos et al. 2003; Önder et al. 2006); İzmir (Tezcan et al. 2010); Diyarbakır (Matocq et al. 2014).

***Monosynamma bohemanni* (Fallén 1829)**

Material examined: Erzurum: Pasinler, 10.VI.1983, ♀.

Distribution in Turkey: Balıkesir, Kırklareli (Önder 1976; Önder et al. 2006); Diyarbakır (Önder et al. 1995); Konya (Lodos et al. 2003).

***Nanopsallus carduellus* (Horvath 1888)**

Material examined: Erzurum: Çat, N 39°35'42.4", E 040°57'59.6", 1913 m, 23.VII.2011, 5 ♀♀, 2 ♂♂, Yukarı Çat, N 39°39'46.3", E 041°00'56.2", 2162 m, 23.VII.2011, 2 ♀♀; Oltu, İnanmış, N 40°27'50.1", E 041°42'19.9", 1823 m, 6.VII.2012, ♀; Yakutiye, Atatürk University field, 1850 m, 7.VIII.1996, ♂.

Distribution in Turkey: Ankara, Mersin (Önder 1976); Bursa, İstanbul (Önder et al. 1981); Diyarbakır (Önder et al. 1995); İçel (Lodos et al. 2003); Antalya, Çorum, Gaziantep, Kahramanmaraş, Karaman, Kırşehir (Önder et al. 2006); İzmir (Tezcan et al. 2010); Mardin, Siirt (Matocq et al. 2014).

***Oncotylus (Cylindromelus) setulosus* (Herrick-Schaeffer 1837)**

Material examined: Erzurum: Aziziye, Rizekent, 2070 m, 30.VII.2010, ♀, 2 ♂♂, Yoncalık, 1900 m, 16.VIII.2014, ♀, ♂; Çat, Yaylasuyu, N 39°43'31.4", E 040°58'42.5", 2208 m, 18.VII.2012, 2 ♂♂; Horasan, Dönertaş, N 40°13'33.4", E 042°06'58.5", 2009 m, 13.VIII.2012, ♀, 2 ♂♂; Narman, İncedere, 1987 m, 21.VII.2010, ♀, Samikale, N 40°21'57.5", E 041°55'20.6", 1541 m, 30.VII.2011, ♀; Oltu, Duralar, N 40°28'31.5", E 041°58'42.4", 1421 m, 30.VII.2011, ♀; Palandöken, 2400 m, 1.VIII.2010, ♀, Nenehatun, 1975 m, 29.VII.2010, ♀, Tepeköy, N 39°50'47.5", E 041°10'30.9", 1873 m, 5.VIII.2011, 6 ♀♀, 4 ♂♂; Pasinler, Ovaköy, N 39°58'56", E 041°29'38", 1782 m, 31.VII.2014, ♀; Pazaryolu, Gülcimene, N 40°24'36.4", E 040°47'28.5", 1663 m, 20.VII.2011, 2 ♂♂; Tortum, 1573 m, 21.VII.2010, 2 ♀♀, 3 ♂♂; Yakutiye, Atatürk University field, 1850 m, 22.VII.1997, 2 ♂♂; Konya: Selçuk University field, 18.VII.1997, 2 ♀♀, 23.VII.1997, ♀.

Distribution in Turkey: Ankara (Hoberlandt 1955; Kiyak and Akar 2010); Elazığ, Isparta, Kars (Önder 1976); Sakarya (Önder et al. 1981); Diyarbakır (Önder et al. 1995); Eskişehir (Lodos et al. 2003); Nevşehir (Kiyak et al. 2004); Adana, Bartın, Çankırı, Çorum, Gaziantep, Kahramanmaraş, Karabük, Karaman, Kastamonu, Kayseri, Konya (Önder et al. 2006); Diyarbakır (Matocq et al. 2014).

***Oncotylus (Oncotylus) pyrethri* (Becker 1864)**

Material examined: Erzurum: Oltu, Demirtaş, N 40°24'58.6", E 041°44'14.4", 1888 m, 6.VII.2012, 2 ♂♂, Özdere, 1927 m, 16.VII.2012, ♂, Yarbaşı, 1784 m, 6.VII.2012, ♂; Yakutiye, Atatürk University field, 1850 m, 20.VII.1997, 2 ♂♂.

Distribution in Turkey: Ankara, Isparta (Önder 1976; Önder et al. 2006).

Oncotylus (Oncotylus) punctipes Reuter 1875

Material examined: **Erzurum:** Aşkale, Çayköy, N 39°56'44.2", E 040°48'18.5", 1700 m, 19.VI.2012, ♀, N 39°56'46.6", E 040°48'27.4", 1720 m, 21.VI.2011, ♀, Kandilli, N 39°54'38.5", E 040°51'19.9", 1710 m, 21.VI.2011, ♂, Küçükgeçit, N 39°56'03.1", E 040°43'32.8", 1660 m, 19.VI.2012, ♀, ♂, Küçükova, N 39°47'36.8", E 040°42'53.3", 1896 m, 5.VII.2012, ♀, Yeniköy, N 39°52'50.3", E 040°42'10.0", 1905 m, 19.VI.2012, 2 ♂♂, N 39°51'28.8", E 040°41'24.1", 1935 m, 5.VII.2012, ♀; Aziziye, Başçakmak, N 40°00'09.8", E 040°54'48.1", 1863 m, 24.VI.2012, ♀, Eğerti, N 40°0.6'13.6", E 040°58'44.5", 1906 m, 22.VI.2011, ♂, Rizekent, 2070 m, 30.VII.2010, 3 ♀♀, Toprakkale, 2157 m, 30.VII.2010, ♀; Çat, Çatköy, 2098 m, 29.VI.2009, ♂, Yaylasuyu Pass, 2322 m, 29.VI.2009, ♂; Hinis, N 39°52'33", E 041°16'46", 1955 m, 26.VI.2011, ♀; Horasan, Muratbağı, N 40°09'12.2", E 042°03'02.7", 1852 m, 22.VI.2012, ♀, ♂, Yeşildere, N 40°07'04.9", E 042°05'29.5", 1698 m, 22.VI.2012, ♀; Karaçoban, Maruf, N 39°18'42", E 041°55'5", 1550 m, 26.VI.2011, 3 ♀♀, ♂; Karayazı, Karasu, N 39°42'51.1", E 042°00'00.5", 2186 m, 16.VII.2011, ♂, Kırgındere, 2215 m, 2.VII.2010, ♂; Köprüköy, Eyüpler, N 39°47'0.5", E 041°46'55.4", 1745 m, 26.VI.2011, 2 ♂♂, Güzelhisar, 1930 m, 2.VII.2010, ♀, Marifet, N 39°50'03.1", E 041°47'38.2", 1685 m, 26.VI.2011, ♀; Narman, Demirbağ, 2028 m, 16.VII.2012, ♀, ♂, Göllü, N 40°13'52.1", E 041°52'06.0", 1842 m, 30.VI.2012, ♀; Oltu, Tutmaç, N 40°26'46.6", E 041°44'06.5", 1756 m, 6.VII.2012, ♀; Palandöken, Taşlıgüney, 2032 m, 26.VI.2009, ♂; Şenkaya, Gözebaşı, N 40°24'39.9", E 042°18'11.5", 1794 m, 14.VII.2012, ♀; Yakutiye, Dumlubaba, Güngörmez, 2200 m, 19.VIII.2009, 8 ♀♀, Güzelyayla, 2200 m, 19.VIII.2009, ♀, Karagöbek, N 40°09'28.9", E 041°25'58.5", 1961 m, 16.VII.2012, ♀.

Distribution in Turkey: Ağrı, Erzurum, Kütahya (Önder 1976; Önder et al. 2006).

Oncotylus (Oncotylus) viridiflavus (Goeze 1778)

Material examined: **Bayburt:** 1550 m, 12.VIII.2009, 3 ♀♀, Örence, 2400 m, 12.VIII.2009, ♀; **Erzincan:** Mercan, 1381 m, 10.VI.2010, ♀; **Erzurum:** Aşkale, Abdalcık, 1756 m, 1.VIII.2010, 3 ♀♀, Büyükleçit, N 39°57'32.3", E 040°46'17.1", 1730 m, 18.VII.2012, ♀, ♂, Çayköy, N 39°58'35.9", E 040°55'49.8", 1876 m, 23.VII.2011, 2 ♂♂, 1691 m, 1.VIII.2010, ♀, Demirkiran, N 39°56'30.2", E 040°41'18.8", 1699 m, 11.VIII.2012, ♂, Gökçebük, N 39°56'55.8", E 040°48'22.6", 1699 m, 23.VII.2011, ♀, ♂, Kandilli, 1707 m, 1.VIII.2010, ♀; Aziziye, Ağzıaçık Pass, 2300 m, 30.VII.2010, 3 ♀♀, 4 ♂♂, Başçakmak, N 40°00'05.6", E 040°54'39.3", 1860 m, 20.VII.2011, 3 ♂♂, Dallıkavak Pass, 2349 m, 30.VII.2010, 3 ♀♀, 2 ♂♂, Eskipolat, 1857 m, 30.VII.2010, 2 ♀♀, ♂, Toprakkale, 2157 m, 30.VII.2010, 5 ♀♀, 3 ♂♂; Çat, N 39°35'42.4", E 040°57'59.6", 1913 m, 23.VII.2011, 2 ♀♀, 1898 m, 6.VIII.2010, 2 ♀♀, Aşağı Çat, 1950 m, 6.VIII.2010, ♀, Değirmenli, N 39°38'51.1", E 040°47'38.6", 1955 m, 11.VIII.2012, 3 ♀♀, Güzelyurt, N 39°46'4.7", E 041°2'42.6", 2099 m, 5.VIII.2011, ♀, ♂, Sarıkaya, N 39°38'15.1", E 040°52'37.7", 1798 m, 11.VIII.2012, ♀, Tuzlataşı, 1910 m, 6.VIII.2010, ♀, Yaylasuyu, N 39°43'31.4", E

040°58'42.5", 2208 m, 18.VII.2012, ♂; Horasan, 1588 m, 2.VII.2010, 3 ♂♂, Aşağı Bademözü, N 40°07'07.7", E 042°10'49.1", 1665 m, 14.VII.2012, ♂, Dalbaşı, N 40°3'2.4", E 042°7'48.2", 1601 m, 17.VII.2011, 4 ♀♀, Kırkgözeler, 1599 m, 29.VII.2010, 3 ♀♀, Kırkdikme, 1601 m, 29.VII.2010, 6 ♀♀, 2 ♂♂, Yörükatlı, 1854 m, 29.VII.2010, ♀, ♂; İspir, Madenköprübaşı, 7.VII.1996, ♀, ♂, N 40°25'58.4", E 040°50'11.3", 1307 m, 20.VII.2011, ♀; Karaçoban, Bozyar, N 39°16'24.9", E 042°00'35.5", 1615 m, 16.VII.2011, ♀, 3 ♂♂; Karayazı, Yukarı Söylemez, N 39°36'46.8", E 041°50'24.3", 1905 m, 17.VII.2012, ♂; Köprüköy, Güzelhisar, 1930 m, 2.VII.2010, ♀, 3 ♂♂; Narman, 1565 m, 21.VII.2010, 2 ♀♀, 1428 m, 4.VIII.2009, 2 ♀♀, 2 ♂♂, Çimenli, N 40°8'47", E 041°53'22.5", 2311 m, 19.VII.2011, ♀, Dikmetaş, 2065 m, 17.VII.2010, ♀, 3 ♂♂, Kışlaköy, N 40°19'20.7", E 042°02'08.3", 1892 m, 13.VIII.2012, ♀; Oltu, 2.VII.1987, ♂, 1750 m, 4.VIII.2009, 4 ♀♀; Olur, Kaledibi, 1074 m, 17.VII.2010, ♀, 1057 m, 4.VIII.2009, 3 ♀♀, 2 ♂♂; Palandöken, Tepeköy, N 39°50'47.5", E 041°10'30.9", 1873 m, 5.VIII.2011, 3 ♀♀, 2 ♂♂; Pasinler, 1600 m, 25.VII.2009, 2 ♀♀, 2 ♂♂, Acı, N 40°02'56.0", E 041°35'21.7", 1905 m, 22.VII.2012, ♂, Çögənder, 1737 m, 29.VII.2010, 5 ♀♀, N 39°59'15.6", E 041°31'07.6", 1768 m, 30.VII.2011, 2 ♂♂, Demirdöven, 1727 m, 29.VII.2010, 2 ♀♀, Korucuk, 1792 m, 29.VII.2010, 2 ♀♀, 2 ♂♂, Ovaköy, N 39°58'57.7", E 041°29'40.8", 1803 m, 30.VII.2011, 2 ♀♀, N 39°58'56", E 041°29'38", 1782 m, 31.VII.2014, ♀, ♂, Serçeboğazı, N 40°00'39.4", E 041°38'23.9", 1725 m, 12.VIII.2012, ♀, Üğümü, N 40°00'39.9", E 041°43'57.0", 1724 m, 17.VII.2011, 2 ♂♂; Şenkaya, 1815 m, 31.VII.2010, ♀, Hoş, N 40°38'50.0", E 042°19'26.0", 1193 m, 31.VII.2011, ♀, Paşalı, N 40°40'08.7", E 042°12'18.4", 1106 m, 19.VII.2011, 2 ♀♀, Sarıyar, N 40°37'57.4", E 042°12'14.9", 1249 m, 13.VII.2011, ♀, ♂, Sındırın, N 40°37'20", E 042°21'34", 1409 m, 14.VII.2014, 2 ♂♂, 1491 m, 31.VII.2010, ♀; Tekman, Körsu, N 39°34'8.2", E 041°44'30.2", 1901 m, 16.VII.2011, ♂; Tortum, 1573 m, 21.VII.2010, 2 ♀♀, 2 ♂♂; Yakutiye, Atatürk University field, 1850 m, 11.VII.2008, ♂, 15.VII.1998, ♀, 23.VII.1986, ♂, 30.VII.1987, ♂♂, 30.VII.2008, ♂, N 39°48'59.7", E 041°04'32.8", 1880 m, 5.VIII.2011, 2 ♀♀, 14.VIII.1987, ♂, 1850 m, 17.VIII.1998, ♀, Güzelyayla, 2000 m, 19.VIII.2009, ♂; **Iğdır:** 850 m, 20.VI.2009, 4 ♂♂, Bayraktar, 850 m, 23.VI.2010, ♀, Aralık, 830 m, 23.VI.2010, ♀, Karakoyunlu, Alican, 850 m, 23.VI.2010, 3 ♀♀, 2 ♂♂, Tuzluca, 900 m, 18.VI.2009, ♂; **Kars:** Sarıkamış, Akkurt, 1650 m, 13.VIII.2009, ♀, Karakurt, 1470 m, 29.VII.2010, 2 ♀♀.

Distribution in Turkey: Bilecik, Burdur, Çanakkale, Denizli, Edirne, İzmir (Önder 1976); Bursa, Sakarya (Önder et al. 1981); Erzurum (Özbek and Alaoglu 1987); Kahramanmaraş, Kayseri (Kiyak 1990); Diyarbakır, Siirt, Şanlıurfa (Önder et al. 1995); Adana, Aksaray, Ankara, Antalya, Bolu, Çankırı, Çorum, Eskeşehir, İçel, Karaman, Kastamonu, Kırıkkale, Konya, Nevşehir, Niğde, Sinop (Lodos et al. 2003); Aegean, Marmara, Central Anatolia and Southeast Anatolia Regions (Önder et al. 2006).

Opisthotaenia (Opisthotaenia) fulvipes Reuter 1901

Material examined: **Artvin:** Ardanuç, Ferhatlı, 573 m, 15.VI.2010, 2 ♀♀, ♂; **Bayburt:** 1650 m, 17.VI.2010, 4 ♀♀, ♂, Aydıntepe, Arpalı, 1550 m, 17.VI.2009, ♀, Çalıdere, 1700 m, 17.VI.2010, ♀, ♂, Demirözü, 1650 m, 17.VI.2010, ♂, Kopdağı Pass, 2344 m, 12.VIII.2009, 4 ♀♀, 3 ♂♂; **Erzurum:** Aşkale, 1600 m, 8.VII.2009, ♀, Çayköy, 1691 m, 1.VIII.2010, ♀, Kandilli, N 39°54'15.3", E 040°50'40.7", 1754 m, 19.VI.2012, ♀, Yaylayolu, N 39°46'53.2", E 040°49'51.4", 2108 m, 5.VII.2012, ♀; Aziziye, Rizekent, N 40°09'56.2", E 040°59'53.1", 2088 m, 24.VI.2012, ♀, 2070 m, 30.VII.2010, ♀, Yoncalık, 1900 m, 16.VIII.2014, 5 ♀♀, ♂; Hınıs, N 39°52'33", E 041°16'46", 1955 m, 26.VI.2011, ♀, 1742 m, 2.VII.2010, 6 ♀♀, ♂, Akören, N 39°29'08.6", E 041°42'02.3", 1939 m, 17.VII.2012, ♂; Horasan, 1536 m, 22.VI.2014, ♀, N 40°00'53", E 041°59'06", 1588 m, 22.VII.2014, ♀, Muratbağı, N 40°09'12.2", E 042°03'02.7", 1852 m, 22.VI.2012, ♀; Karayazı, Kırgındere, 2215 m, 2.VII.2010, ♀, 3 ♂♂; Köprüköy, Eğirmez, N 39°58'09.9", E 041°49'20.7", 1670 m, 13.VIII.2011, 3 ♀♀, 2 ♂♂, Eyüpler, N 39°47'05.", E 041°46'55.4", 1745 m, 26.VI.2011, ♀, 3 ♂♂, İlicasu, N 40°5'39.6", E 041°51'57.1", 2111 m, 30.VII.2011, 3 ♀♀, Örengtaş, N 40°04'15.1", E 041°50'40.7", 1916 m, 30.VII.2011, ♀; Narman, 1900 m, 24.VII.2009, ♀, Dikmetaş, 2665 m, 17.VII.2010, ♀, İncedere, 1987 m, 21.VII.2010, ♀, 2 ♂♂, Kamışözü, N 40°20'23.3", E 041°47'29.2", 1835 m, 23.VI.2011, 2 ♂♂; Oltu, Başbağlar, N 40°27'56.1", E 041°41'37.4", 1868 m, 6.VII.2012, 3 ♀♀, 3 ♂♂, Çamlıbel, 1700 m, 14.VII.1996, ♀, Demirtaş, N 40°24'58.6", E 041°44'14.4", 1888 m, 6.VII.2012, 3 ♀♀, 2 ♂♂, Özdere, N 40°27'57.7", E 041°44'11.9", 1927 m, 16.VII.2012, 2 ♂♂; Palandöken, Taşlıgüney, 2032 m, 26.VI.2009, 2 ♀♀, N 39°48'41.2", E 041°07'10.0", 1875 m, 5.VII.2012, ♀; Pasinler, Korucuk, 1650 m, 7.VII.2009, ♀, Yayla, N 40°05'42.7", E 041°44'03.5", 1990 m, 17.VII.2011, 2 ♀♀; Pazaryolu, Çiftepinar, N 40°22'05.3", E 041°43'07.4", 1742 m, 15.VII.2012, ♀; Tekman, Güzeldere, N 39°49'17.1", E 042°00'19.8", 1948 m, 16.VII.2011, ♀, ♂, Körsu, 1940 m, 2.VII.2010, ♂; Tortum, N 40°16'37.4", E 041°33'30", 1661 m, 23.VI.2011, 5 ♀♀, 3 ♂♂, Uzundere, Yayla, N 40°27'48.6", E 041°39'32.3", 2005 m, 6.VII.2012, ♀; Yakutiye, Atatürk University field, N 39°48'59.7", E 041°04'32.8", 1880 m, 20.VI.2012, ♂, 1850 m, 24.VI.2009, 7 ♀♀, 4 ♂♂, 25.VI.2009, 2 ♀♀, ♂, 2.VII.2008, ♀, 7.VII.2008, ♀, 8.VII.1986, ♀, 8.VII.1987, 2 ♂♂, 9.VII.1986, 4 ♀♀, 9.VII.1996, ♂, 10.VII.1996, ♂, 13.VII.2009, 5 ♀♀, 20.VII.2009, ♀, ♂, N 39°53'59.1", E 041°14'19.0", 1880 m, 22.VII.2011, ♀, 1850 m, 23.VII.2007, ♀, 4.VIII.2008, ♂, 14.VIII.1998, 2 ♂♂, 17.VIII.1998, ♂; **Kars:** Sarıkamış, Karakurt, 1500 m, 22.VI.2010, 3 ♀♀.

Distribution in Turkey: Amasya, Edirne (Hoberlandt 1955); Afyonkarahisar, Ankara, Çanakkale, Çankırı, Kayseri, Nevşehir (Önder 1976); Amasya, Edirne, Malatya, Niğde (Lodos et al. 2003; Önder et al. 2006); Diyarbakır (Matocq et al. 2014).

Phoenicocoris obscurellus (Fallén 1829)

Material examined: **Erzurum:** Aziziye, Rizekent, N 40°0'9'49.1", E 041°00'03.5", 2097 m, 20.VII.2011, ♀; Şenkaya, Sindiran, N 40°37'14.2", E 042°22'09.3", 1497 m, 19.VII.2011, ♂; Yakutiye, Atatürk University field, 1850 m, 14.VII.1996, ♂, 15.VII.1998, 6 ♂♂, 16.VII.1996, ♀, ♂, 18.VII.1997, ♂, 31.VII.1998, 3 ♂♂, 1.VIII.1998, 3 ♂♂, 2.VIII.1998, ♀, 4 ♂♂, 3.VIII.1998, 3 ♂♂, 5.VIII.1998, 2 ♂♂, 23.VIII.1998, ♀, 7 ♂♂, 31.VIII.1996, 2 ♂♂.

Distribution in Turkey: Erzurum (Yıldırım 2002).

Pilophorus cinnamopterus (Kirschbaum 1856)

Material examined: **Artvin:** Yusufeli, Demirkent, 5.VI.1991, ♀, Morkaya, 6.VI.1991, ♀; **Erzurum:** İspir, Madenköprübaşı, 24.VII.1991, ♀; Oltu, 11.VII.1991, 2 ♂♂; Tortum, Pehlivanlı, 13.VIII.1991, ♀; Yakutiye, Atatürk University field, 1850 m, 10.VII.1996, ♂, 18.VIII.1997, ♀, 23.VIII.1996, ♀; **Kars:** Kağızman, 19.VII.1991, ♀.

Distribution in Turkey: Artvin, Balıkesir, Bolu, Çanakkale, Kahramanmaraş (Önder 1976; Önder et al. 2006); İstanbul (Önder et al. 1981); İzmir (Tezcan et al. 2010).

Pilophorus clavatus (Linnaeus, 1767)

Material examined: **Artvin:** Yusufeli, Morkaya, 6.VII.1991, ♀; **Erzurum:** Yakutiye, Atatürk University field, 1850 m, 26.VII.1997, ♀.

Distribution in Turkey: Tokat (Çam 1988); Adana, Ankara, Bartın, Çankırı, İçel, Kahramanmaraş, Kayseri, Konya, Nevşehir, Yozgat (Lodos et al. 2003); Nevşehir (Kiyak et al. 2004); Afyonkarahisar, Adana, Amasya, Ankara, Artvin, Balıkesir, Bartın, Bilecik, Bursa, Burdur, Çankırı, Gaziantep, İzmir, Kahramanmaraş, Kırklareli, Konya, Kütahya, Manisa, Nevşehir, Rize, Tekirdağ, Yozgat (Önder et al. 2006).

Pilophorus pusillus Reuter 1878

Material examined: **Antalya:** 22.VIII.1984, ♀, ♂; **Artvin:** Yusufeli, Morkaya, 6.VI.1991, ♀; **Erzurum:** Pazaryolu, 23.VII.1991, ♀; Tortum, Pehlivanlı, 13.VIII.1991, ♀, 2 ♂♂; Yakutiye, Atatürk University field, 1850 m, 21.VIII.1998, ♀; **Iğdır:** Akyumak, 18.VI.1991, 2 ♀♀, Melekli, 18.VII.1991, ♀; **Kars:** Kağızman, 19.VII.1991, 2 ♀♀, ♂.

Distribution in Turkey: İğdir (Hoberlandt 1955); Afyonkarahisar, Artvin, Aydın, Balıkesir, Bilecik, Burdur, Çanakkale, Denizli, Edirne, Isparta, Kırklareli, Kocaeli, Muğla, Sakarya, Samsun, Sinop, Tekirdağ, Trabzon, Uşak (Önder 1976); Bursa (Önder et al. 1981); Antalya (Yayla 1983; Özkan 1984); Tokat (Çam 1988); Adana, Ankara, Bolu, Çorum, Eskişehir, Hatay, İçel, Kahramanmaraş, Karabük, Kastamonu, Kayseri, Kırşehir, Konya, Nevşehir, Yozgat, Zonguldak (Lodos et al.

2003); West, Northwest, North and South Anatolia Regions (Önder et al. 2006); İzmir (Tezcan et al. 2010).

***Plagiognathus (Plagiognathus) bipunctatus* Reuter 1883**

Material examined: **Artvin:** Ardanuç, Ferhatlı, 573 m, 15.VI.2010, 2 ♀♀, Yusufeli, Sebzeciler, 500 m, 16.VI.2010, 2 ♀♀; **Bayburt:** 1650 m, 17.VI.2010, 2 ♂♂, 1550 m, 12.VIII.2009, 3 ♀♀, Kopdağı Pass, 2344 m, 12.VIII.2009, 3 ♀♀, Aydintepe, Arpalı, 1550 m, 17.VI.2010, 2 ♀♀, Çalidere, 1700 m, 17.VI.2010, 3 ♀♀, ♂, Demirözü, 1650 m, 17.VI.2009, 6 ♀♀; **Erzincan:** 1185 m, 10.VI.2010, ♀, ♂, Akyazı, 1202 m, 11.VI.2010, 4 ♀♀, ♂, Mercan, 11.VII.1990, 7 ♀♀, 3 ♂♂, Refahiye, 1500 m, 20.VI.2009, ♀, Tercan, 1408 m, 10.VI.2010, 3 ♀♀; **Erzurum:** Aşkale, Adaçay, 11.VII.1990, 3 ♀♀, ♂, Athikonak, N 39°55'36.5", E 040°56'16.9", 1757 m, 5.VII.2012, ♂, 1760 m, 9.VIII.2011, 4 ♀♀, 4 ♂♂, Büyükgeçit, N 39°57'32.3", E 040°46'17.1", 1730 m, 18.VII.2012, 2 ♀♀, Çayköy, N 39°56'46.6", E 040°48'27.4", 1720 m, 21.VI.2011, ♂, N 39°58'35.9", E 040°55'49.8", 1876 m, 23.VII.2011, ♀, Dereköy, N 39°47'33.3", E 040°43'57.8", 1929 m, 5.VII.2012, ♀, ♂, Gökçebük, N 39°56'55.8", E 040°48'22.6", 1699 m, 23.VII.2011, ♀, ♂, Gölören, N 40°01'28.9", E 040°47'55.1", 1897 m, 18.VII.2012, ♀, Gökçebük, 1699 m, 23.VII.2011, ♀, Hatuncuk, N 39°48'15.6", E 040°42'09.3", 1869 m, 5.VII.2012, ♀, ♂, Kandilli, N 39°54'38.5", E 040°51'19.9", 1710 m, 21.VI.2011, 3 ♀♀, 2 ♂♂, 1737 m, 1.VIII.2010, ♀, Kavurmaçukur, N 39°45'39.9", E 040°51'58.4", 2233 m, 5.VII.2012, 7 ♂♂, Koşapınar, N 40°00'55.4", E 040°47'39.3", 1886 m, 18.VII.2012, ♀, Küçükgeçit, N 39°56'43.2", E 040°47'12.1", 1709 m, 23.VII.2011, 4 ♀♀, 2 ♂♂, Tazegül, 26.VI.1990, ♂; Aziziye, Ağzıaçık Pass, 2300 m, 30.VII.2011, 2 ♀♀, ♂, Alaca, N 39°55'50.7", E 040°58'00.9", 1751 m, 5.VII.2012, 2 ♀♀, Başçakmak, N 40°00'05.6", E 040°54'39.3", 1860 m, 20.VII.2011, 3 ♀♀, ♂, Dallikavak Pass, N 40°10'33.55", E 040°58'16.1", 2349 m, 20.VII.2011, ♀, Eskipolat, N 40°4'10", E 040°56'8.2", 1748 m, 20.VII.2011, ♂, 1857 m, 30.VII.2010, 3 ♀♀, ♂, Ortabayçe, N 39°53'35.3", E 040°51'19.0", 1763 m, 5.VII.2012, 2 ♂♂, Özbek, N 39°48'57.1", E 041°04'39.6", 1930 m, 5.VIII.2011, ♀, Paşayurdu, N 39°59'42.3", E 041°00'18.3", 1836 m, 18.VII.2012, ♀, Rizekent, N 40°09'56.2", E 040°59'53.1", 2088 m, 24.VI.2012, ♂, 2070 m, 30.VII.2010, 9 ♀♀, 2 ♂♂, Tebrizcik, N 39°56'24.6", E 040°59'46.7", 1750 m, 5.VII.2012, ♂, Toprakkale, 2157 m, 30.VII.2010, 8 ♀♀, Yoncalık, 1900 m, 16.VIII.2014, 7 ♀♀, 2 ♂♂; Çat, N 39°35'42.4", E 040°57'59.6", 1913 m, 23.VII.2011, 4 ♀♀, Aşağı Çat, N 39°38'29.5", E 041°00'11.7", 2065 m, 9.VIII.2011, 2 ♂♂, Güzelyurt, N 39°46'4.7", E 041°2'42.6", 2099 m, 5.VIII.2011, ♀, Kom, N 39°43'41.9", E 040°58'10.1", 2206 m, 5.VII.2012, ♂, Sarıkaya, N 39°38'15.1", E 040°52'37.7", 1798 m, 11.VIII.2012, ♀, Yaylasuyu, 2322 m, 29.VI.2009, ♂, N 39°43'45.0", E 040°56'34.0", 2203 m, 5.VII.2012, 2 ♀♀, ♂, Yukarı Çat, N 39°39'04.0", E 041°01'17.1", 2127 m, 18.VII.2012, 2 ♀♀, N 39°39'46.3", E 041°00'56.2", 2162 m, 23.VII.2011, 2 ♀♀; Hinis, N 39°52'33", E 041°16'46", 1955 m, 26.VI.2011, 2 ♂♂, 1742 m, 2.VII.2010, ♀, ♂, Yolüstü, N 39°19'37.2", E 041°44'37.7", 1664 m, 17.VI.2012, ♂; Horasan, Çayırdüzü, N 40°08'21.9", E 042°03'47.4", 1799 m, 22.VI.2012, 3 ♀♀, ♂, Dönertas, N 40°13'33.4",

E 042°06'58.5", 2009 m, 13.VIII.2012, ♂; İspir, 1300 m, 7.VIII.2009, ♀, Akseki, N 40°17'15.7", E 041°00'23.6", 2190 m, 20.VII.2011, 2 ♀♀, 3 ♂♂, Elmalı, N 40°25'31.9", E 040°50'13.6", 1328 m, 20.VII.2011, 2 ♀♀, 3 ♂♂, Güneyköy, N 40°19'59.6", E 040°56'20.5", 2024 m, 24.VII.2011, ♀, İncesu, 2100 m, 20.VIII.2009, 2 ♀♀, İspir Pass, 1950 m, 20.VIII.2009, 3 ♀♀, Kirazlı, N 40°26'47.9", E 040°53'09.2", 1207 m, 24.VII.2011, ♀, Madenköprübaşı, N 40°25'58.4", E 040°50'11.3", 1307 m, 20.VII.2011, ♀, N 40°26'34.3", E 040°50'42.4", 1245 m, 4.VIII.2012, ♂, 1200 m, 20.VIII.2009, ♀, Yukarı Özbağ, 1175 m, 30.VII.2010; Karaçoban, Duman, N 39°18'22", E 041°55'5", 1560 m, 26.VI.2011, ♂; Karayazı, Aşağı İncesu, N 39°40'45.5", E 041°57'05.9", 2098 m, 17.VII.2012, ♀, Çatalören, N 39°46'32.1", E 042°01'43.1", 2313 m, 16.VII.2011, ♀, ♂, Kırgındere, 2215 m, 2.VII.2010, 2 ♀♀, Yukarı Söylemez, 1896 m, 2.VII.2010, 2 ♀♀, 2 ♂♂, N 39°36'46.8", E 041°50'24.3", 1905 m, 17.VII.2012, 2 ♀♀; Köprüköy, 1720 m, 2.VII.2010, ♂, Çatalören, N 39°46'36.1", E 042°01'49.2", 2310 m, 17.VII.2012, ♂, Eğirmez, N 39°58'09.9", E 041°49'20.7", 1670 m, 17.VI.2012, ♂, N 39°58'09.9", E 041°49'20.7", 1670 m, 13.VIII.2011, 6 ♀♀, 6 ♂♂, Eyüpler, N 39°47'0.5", E 041°46'55.4", 1745 m, 26.VI.2011, ♂, Güzelhisar, 1930 m, 2.VII.2010, 2 ♀♀, İlicasu, 2380 m, 17.VII.2010, 2 ♀♀, 2 ♂♂, N 40°5'39.6", E 041°51'57.1", 2111 m, 30.VII.2011, 3 ♀♀, ♂, Örentaş, 2038 m, 29.VII.2010, 3 ♀♀, Pekcik, N 39°53'12.9", E 041°53'08.3", 1722 m, 26.VI.2011, 4 ♂♂, Yücelik, N 39°46'02.7", E 042°02'07.8", 2288 m, 17.VII.2012, ♀; Narman, 2.VII.1987, ♀, 1900 m, 24.VII.2009, 3 ♀♀, 1428 m, 4.VIII.2009, 2 ♂♂, 1557 m, 4.VIII.2009, 3 ♀♀, Çimenli, N 40°09'31.3", E 041°53'50.5", 2326 m, 30.VI.2012, ♀, ♂, N 40°8'47", E 041°53'22.5", 2311 m, 19.VII.2011, 2 ♀♀, ♂, Demirdağ, N 40°19'23.5", E 041°42'45.5", 2028 m, 16.VII.2012, 2 ♂♂, Göllü, N 40°13'52.1", E 041°52'06.0", 1842 m, 30.VI.2012, ♀, ♂, N 40°14'7.9", E 041°51'57.2", 1083 m, 30.VII.2011, 2 ♀♀, İncedere, 1987 m, 21.VII.2010, 2 ♀♀, ♂, Kışlaköy, N 40°19'20.7", E 042°02'08.3", 1892 m, 13.VIII.2012, 2 ♀♀, ♂, Kilimli, N 40°19'50.9", E 041°58'50.3", 1726 m, 16.VII.2012, 4 ♀♀, Mahmutçavuş, N 40°20'18.8", E 041°55'36.3", 1560 m, 16.VII.2012, ♀, ♂, Samikale, N 40°21'57.5", E 041°55'20.6", 1541 m, 30.VII.2011, ♀, Şehitler, N 40°19'54.9", E 041°46'39.2", 1884 m, 16.VII.2012, ♀, Toygarlı, N 40°12'18.9", E 041°53'16.6", 2071 m, 30.VI.2012, 2 ♂♂, 2123 m, 10.VII.2011, ♀, N 40°12'42.9", E 041°53'17.8", 2123 m, 19.VII.2011, 3 ♀♀, ♂, Yanıktaş, N 40°16'52.7", E 041°52'18.1", 1612 m, 30.VI.2012, ♀, ♂; Oltu, 1180 m, 24.VII.2009, ♀, 1750 m, 4.VIII.2009, 4 ♀♀, 3 ♂♂, Ayvalı, 700 m, 16.VI.2010, ♀, Başbağlar, N 40°27'56.1", E 041°41'37.4", 1868 m, 6.VII.2012, 2 ♀♀, N 40°28'1.1", E 041°40'27.6", 1925 m, 13.VII.2011, 2 ♀♀, Çamlıbel, N 40°28'31.3", E 041°46'26.9", 1661 m, 30.VI.2012, ♂, N 40°29'36.1", E 041°45'45.2", 1775 m, 13.VII.2011, ♀, ♂, N 40°29'06", E 041°45'47", 1635 m, 14.VII.2014 m, 3 ♀♀, ♂, N 40°28'34.3", E 041°46'24.1", 1682 m, 31.VII.2011, ♀, ♂, Demirtaş, N 40°24'58.6", E 041°44'14.4", 1888 m, 6.VII.2012, ♀, İğdeli, N 40°32'44.8", E 041°50'27.6", 1660 m, 30.VI.2012, 4 ♀♀, Duralar, N 40°28'31.5", E 041°58'42.4", 1421 m, 30.VII.2011, ♀, ♂, İnanmış, N 40°27'50.1", E 041°42'19.9", 1823 m, 6.VII.2012, ♀, 2 ♂♂, Orcuk, N 40°29'44.9", E 041°49'01.0", 1556 m, 6.VII.2012, ♂, Özdere, N 40°25'19.5", E 041°43'25.1", 1822 m,

6.VII.2012, ♂, N 40°27'57.7", E 041°44'11.9", 1927 m, 16.VII.2012, ♂, Sarısaç, N 40°31'56.1", E 041°54'36.9", 1388 m, 5.VIII.2012, ♀, Toprakkale, N 40°27'55.5", E 041°59'21.1", 1416 m, 23.VI.2011, 3 ♀♀, 3 ♂♂, Tutmaç, N 40°26'46.6", E 041°44'06.5", 1756 m, 6.VII.2012, 4 ♀♀, ♂, N 40°27'23.2", E 041°44'41.4", 1720 m, 16.VII.2012, ♀, Yolboyu, 1229 m, 4.VIII.2009, ♀; Olur, Boğazgören, N 40°49'19.2", E 042°12'42.6", 1168 m, 19.VII.2012, ♀, Kaban, N 40°46'41.1", E 042°12'00.6", 1215 m, 19.VII.2012, ♂, Kaledibi, 1057 m, 4.VIII.2009, ♀, Yeşilbağlar, 1000 m, 16.VI.2010, 6 ♀♀; Palandöken, 2300 m, 21.VII.2010, 5 ♀♀, ♂, 2400 m, 21.VII.2010, 6 ♀♀, 2 ♂♂, N 39°51'56.4", E 041°16'19.7", 2155 m, 22.VII.2012, 4 ♀♀, ♂, 2300 m, 1.VIII.2010, 3 ♀♀, 2000 m, 8.VIII.2009, 4 ♀♀, Kümbet, N 39°48'59.7", E 041°4'32.8", 1832 m, 5.VIII.2011, 3 ♀♀, N 39°48'57.8", E 041°0.4'33", 1836 m, 9.VIII.2011, ♀, ♂, Taşlıgüney, 11.VII.1990, 7 ♀♀, 3 ♂♂, Yağmurcuk, N 39°47'13.2", E 041°06'06.1", 2010 m, 9.VIII.2011, 4 ♀♀, 3 ♂♂; Pasinler, 15.VI.1986, ♀, 29.VI.1987, ♀, ♂, 11.VII.1987, ♀, ♂, 1600 m, 25.VII.2009, ♀, Büyüktuy, 1800 m, 2.VII.2010, 2 ♀♀, ♂, N 39°58'46.7", E 041°27'58", 1819 m, 30.VII.2011, ♀, Çögender, 1740 m, 22.VI.2012, 3 ♀♀, ♂, N 39°59'15.6", E 041°31'07.6", 1768 m, 30.VII.2011, ♀, Kurbançayı, N 40°01'59.3", E 041°37'38.1", 1760 m, 12.VIII.2012, ♀, Ovaköy, N 39°58'57.7", E 041°29'40.8", 1803 m, 30.VII.2011, ♀, ♂, N 39°58'56", E 041°29'38", 1782 m, 31.VII.2014, 3 ♀♀, Sansar Stream, N 40°04'21.5", E 041°43'28.5", 1877 m, 17.VII.2011, ♀, Serçeboğazı, N 40°00'39.4", E 041°38'23.9", 1725 m, 12.VIII.2012, ♂, Üğümü, N 40°00'39.9", E 041°43'57.0", 1724 m, 17.VII.2011, ♀, 3 ♂♂, Yayla, N 40°05'42.7", E 041°44'03.5", 1990 m, 17.VII.2011, 3 ♀♀, Yiğitpinarı, N 40°02'15.6", E 041°35'51.0", 1815 m, 22.VII.2012, ♀; Pazaryolu, N 40°25'12.8", E 040°46'13.5", 1450 m, 24.VII.2011, ♂, N 40°25'12.1", E 040°47'07.7", 1495 m, 24.VII.2011, ♀, ♂, Göztepe, N 40°23'10.1", E 040°43'44.6", 1672 m, 15.VII.2012, ♂, Gülçimen, N 40°24'36.4", E 040°47'28.5", 1663 m, 20.VII.2011, ♂; Şenkaya, 2.VII.1987, 2 ♀♀, ♂, Deliktaş, N 40°41'57.1", E 042°15'52.7", 1219 m, 13.VII.2011, ♀, Esenyurt, N 40°25'40.2", E 042°18'15.2", 1622 m, 14.VII.2012, 4 ♀♀, İçmesu, N 40°25'27.0", E 042°18'32.1", 1656 m, 14.VII.2012, ♀, İlkizpınar, N 40°36'57.4", E 042°21'30.9", 1589 m, 31.VII.2011, 6 ♀♀, 2 ♂♂, Sındiran, N 40°37'20", E 042°21'34", 1409 m, 14.VII.2014, 2 ♀♀, ♂, Taht, N 40°38'29", E 042°20'0.3", 1232 m, 14.VII.2014, ♂, N 40°37'59.4", E 042°20'39.7", 1283 m, 31.VII.2011, ♂; Tekman, Çiçekdağı, N 39°34'40.8", E 041°44'26.8", 1846 m, 26.VI.2011, 5 ♂♂, Geyikli, N 39°47'48.5", E 042°4'31.4", 2185 m, 16.VII.2011, ♂, Güzeldere, N 39°49'17.1", E 042°00'19.8", 1948 m, 16.VII.2011, ♂, Körsu, 1940 m, 2.VII.2010, ♀, ♂, N 39°34'8.2", E 041°44'30.2", 1901 m, 16.VII.2011, 8 ♀♀, 2 ♂♂; Tortum, 1653 m, 15.VI.2010, ♀, 4 ♂♂, N 40°16'37.4", E 041°33'30", 1661 m, 23.VI.2011, ♂, 2.VII.1987, 2 ♀♀, 22.VII.1987, 6 ♀♀, 7 ♂♂, N 40°16'38", E 041°33'29.9", 1660 m, 31.VII.2011, 2 ♀♀, Akbababa, N 40°13'32.6", E 041°29'4.1", 2073 m, 31.VII.2011, 3 ♀♀, ♂, Aksukapı, N 40°23'58.8", E 041°32'46.4", 1648 m, 13.VII.2011, 3 ♀♀, 5 ♂♂, Aktaş, N 40°15'40.5", E 041°32'39.2", 1765 m, 31.VII.2011, 2 ♀♀, ♂, Aşağı Sivri, N 40°19'33.8", E 041°31'46.9", 1546 m, 16.VII.2012, ♀, Derekapı, 950 m, 15.VI.2010, 5 ♀♀, 5 ♂♂, 1247 m, 9.VII.2011, ♀, N 40°24'47.2", E 041°29'19.4", 1247 m,

13.VII.2011, 5 ♀♀, 2 ♂♂, Kireçli Pass, 2400 m, 31.VII.2010, 6 ♀♀, Taşbaşı, N 40°14'01.4", E 041°30'11.9", 1968 m, 6.VII.2012, ♀, 4 ♂♂, N 40°15'26.6", E 041°32'13.8", 1816 m, 13.VII.2011, 5 ♀♀, 2 ♂♂, Yukarı, Sivri, N 40°19'34", E 041°35'51.6", 1734 m, 23.VI.2011, ♂, N 40°20'27.5", E 041°39'01.1", 1962 m, 16.VII.2012, 3 ♀♀; Uzundere, Gölbaşı, N 40°36'04.0", E 041°33'43.1", 1260 m, 7.VIII.2011, ♀, Kirazlı, N 40°34'26.9", E 041°38'52.5", 1231 m, 21.VII.2012, ♀, Yayla, N 40°27'48.6", E 041°39'32.3", 2005 m, 6.VII.2012, 2 ♀♀, ♂, N 40°27'48.6", E 041°39'32.3", 2005 m, 13.VII.2011, 4 ♀♀, ♂; Yakutiye, Atatürk University field, 1850 m, 15.V.2010, ♀, 25.VI.2009, 2 ♀♀, 2.VII.1987, 3 ♀♀, 3.VII.1987, ♀, 4 ♂♂, 9.VII.1986, ♀, ♂, 13.VII.2009, 2 ♀♀, 21.VII.1987, 3 ♀♀, 22.VII.2009, 4 ♀♀, 23.VII.1987, ♀, ♂, 29.VII.2009, ♀, 3.VIII.1998, ♀, 7.VIII.1996, ♂, 19.VIII.1998, ♂, Dumlubaba, Güngörmez, 2500 m, 1.VII.2010, 3 ♂♂, 2200 m, 19.VIII.2009, 3 ♀♀, Güzelyayla, 2000 m, 19.VIII.2009, 6 ♀♀, Karagöbek, N 40°10'15.1", E 041°26'14.9", 2033 m, 13.VII.2011, 4 ♀♀, 2 ♂♂, N 40°09'28.9", E 041°25'58.5", 1961 m, 16.VII.2012, ♀, N 40°09'57.8", E 041°26'8.1", 1993 m, 31.VII.2011, 3 ♀♀, ♂, 1998 m, 3.VIII.2009, 2 ♀♀; **Iğdır:** Melekli, 850 m, 23.VI.2010, ♀, 2 ♂♂, Aralık, 830 m, 23.VI.2010, ♂; **Kars:** Kağızman, 19.VII.1991, ♀, Sarıkamış, 1900 m, 13.VIII.2009, 3 ♀♀, Akkurt, 1650 m, 13.VIII.2009, 3 ♀♀, Karakurt, 1500 m, 13.VIII.2009, 11 ♀♀, 3 ♂♂; **Ordu:** Perşembe, 31.VII.1991, ♂.

Distribution in Turkey: Adana, Ankara (Hoberlandt 1955); Afyonkarahisar, Ağrı, Amasya, Artvin, Balıkesir, Burdur, Çanakkale, Edirne, Elazığ, Kırklareli, Kütahya, Manisa, Muş, Ordu, Sivas, Tekirdağ (Önder 1976); Malatya, Van (Bingöl 1978); Denizli, Uşak (Giray 1980); Bolu, Bursa, İstanbul, Kocaeli, Sakarya (Önder et al. 1981); Diyarbakır (Önder et al. 1995); Yozgat (Tamer et al. 1998); Antalya, Çankırı, Çorum, Eskişehir, Gaziantep, Hatay, İçel, Kahramanmaraş, Karabük, Karaman, Kastamonu, Kayseri, Kırıkkale, Kırşehir, Kilis, Konya, Nevşehir, Niğde, Osmaniye, Sinop (Lodos et al. 2003); almost all regions (Önder et al. 2006); İzmir (Tezcan et al. 2010); Mardin, Siirt (Matocq and Özgen 2010); Diyarbakır (Matocq et al. 2014).

Plagiognathus chrysanthemi (Wolff 1804)

Material examined: **Erzurum:** Aşkale, 1600 m, 8.VII.2009, ♂, Dereköy, N 39°47'33.3", E 040°43'57.8", 1929 m, 5.VII.2012, ♂, Hatuncuk, N 39°48'15.6", E 040°42'09.3", 1869 m, 5.VII.2012, 2 ♂♂, Kavurmaçukur, N 39°45'39.9", E 040°51'58.4", 2233 m, 5.VII.2012, 2 ♂♂, Yaylayolu, N 39°46'53.2", E 040°49'51.4", 2108 m, 5.VII.2012, ♂; Aziziye, Ağzıaçık Pass, 2300 m, 30.VII.2010, 2 ♀♀, 10 ♂, Dallıkavak, N 40°10'33.55", E 040°58'16.1", 2349 m, 20.VII.2011, 2 ♂♂, N 40°10'33.3", E 040°58'14.3", 2243 m, 24.VII.2011, 2 ♂♂, 30.VII.2010, ♂, Eskipolat, N 40°4'10", E 040°56'8.2", 1748 m, 20.VII.2011, 2 ♂♂, Gelinkaya, N 40°01'09.6", E 040°54'48.1", 1796 m, 24.VI.2012, 2 ♂♂, Kümbet, N 39°48'57.8", E 041°0.4'33", 1836 m, 9.VIII.2011, ♂, Paşayurdu, N 39°59'42.3", E 041°00'18.3", 1836, 18.VII.2012, ♂, Rizekent, 2070 m, 30.VII.2010, ♂, Toprakkale, 2157 m, 30.VII.2010, ♀, ♂, Yoncalık, 1900 m, 16.VIII.2014, ♂; Çat, N 39°35'42.4", E 040°57'59.6", 1913 m, 23.VII.2011, ♀, ♂, Yukarı Çat, N 39°39'04.0", E 041°01'17.1", 2127 m, 18.VII.2012,

♂, Hınıs, 1742 m, 2.VII.2010, 2 ♂♂; Horasan, Kırkgözeler, 1760 m, 15.VI.2010, ♂; İspir, Akseki, N 40°17'15.7", E 041°00'23.6", 2190 m, 20.VII.2011, 2 ♂♂, Güneyköy, N 40°19'59.6", E 040°56'20.5", 2024 m, 24.VII.2011, 2 ♀♀, ♂, Madenköprübaşı, N 40°25'58.4", E 040°50'11.3", 1307 m, 20.VII.2011, ♂; Karayazı, Çatalören, N 39°46'32.1", E 042°01'43.1", 2313 m, 16.VII.2011, ♂, Kırgındere, 2215 m, 2.VII.2010, 3 ♂♂, Yukarı Söylemez, 1896 m, 2.VII.2010, 2 ♂♂; Köprüköy, Eğirmez, N 39°58'09.9", E 041°49'20.7", 1670 m, 13.VIII.2011, ♂, Eyüpler, N 39°47'0.5", E 041°46'55.4", 1745 m, 26.VI.2011, ♂, Geyikli, N 39°48'02.1", E 042°04'05.4", 2144 m, 17.VII.2012, 2 ♂♂, Güzelhisar, 1930 m, 2.VII.2010, 2 ♂♂, İlicasu, 2380 m, 17.VII.2010, ♀, N 40°5'39.6", E 041°51'57.1", 2111 m, 30.VII.2011, ♂, Pekcik, N 39°53'12.9", E 041°53'08.3", 1722 m, 26.VI.2011, ♂, Yeşilova, N 39°46'02.7", E 042°02'07.8", 2288 m, 17.VII.2012, 2 ♂♂; Narman, N 40°21'09.9", E 041°55'02.0", 1543 m, 30.VI.2012, ♂, 1900 m, 24.VII.2009, ♀, 2 ♂♂, Çimenli, N 40°8'47", E 041°53'22.5", 2311 m, 19.VII.2011, 2 ♂♂, 2308 m, 4.VIII.2009, 2 ♂♂, Demirdağ, N 40°19'23.5", E 041°42'45.5", 2028 m, 16.VII.2012, 2 ♂♂, Göllü, N 40°13'52.1", E 041°52'06.0", 1842 m, 30.VI.2012, 3 ♂♂, N 40°14'7.9", E 041°51'57.2", 1083 m, 30.VII.2011, ♂, Samikale, N 40°21'57.5", E 041°55'20.6", 1541 m, 30.VII.2011, ♀, Şehitler, N 40°19'54.9", E 041°46'39.2", 1884 m, 16.VII.2012, ♂, Toygarlı, N 40°12'42.9", E 041°53'17.8", 2123 m, 19.VII.2011, 3 ♂♂; Oltu, N 40°32'43.9", E 41°59'44.0", 1455 m, 8.VI.2012, 2 ♂♂, Başbağlar, N 40°27'56.1", E 041°41'37.4", 1868 m, 6.VII.2012, 2 ♂♂, Çamlıbel, N 40°29'36.1", E 041°45'45.2", 1775 m, 13.VII.2011, ♂, N 40°29'06", E 041°45'47", 1635 m, 14.VII.2014 m, 3 ♂♂, Demirtaş, N 40°24'58.6", E 041°44'14.4", 1888, 6.VII.2012, ♂, İğdeli, N 40°32'44.8", E 041°50'27.6", 1660 m, 30.VI.2012, 3 ♂♂, İnanmış, N 40°27'44.4", E 041°42'42", 1812 m, 31.VII.2011, ♀, Orcuk, N 40°29'44.9", E 041°49'01.0", 1556 m, 6.VII.2012, 2 ♂♂, Özdere, N 40°25'19.5", E 041°43'25.1", 1822 m, 6.VII.2012, 3 ♂♂, N 40°27'57.7", E 041°44'11.9", 1927 m, 16.VII.2012, ♂, Toprakkale, N 40°27'55.5", E 041°59'21.1", 1416 m, 23.VI.2011, ♀, Tutmaç, N 40°26'46.6", E 041°44'06.5", 1756 m, 6.VII.2012, 3 ♂♂, N 40°27'23.2", E 041°44'41.4", 1720 m, 16.VII.2012, ♂, Yarbaşı, N 40°26'18.9", E 041°43'50.2", 1784 m, 6.VII.2012, ♂; Olur, Boğazören, N 40°49'19.2", E 042°12'42.6", 1168 m, 19.VII.2012, ♂, Kaban, N 40°46'41.1", E 042°12'00.6", 1215 m, 19.VII.2012, ♀, ♂, Yeşilbağlar, 1000 m, 16.VI.2010, ♀, 2 ♂♂; Palandöken, Nenehatun, 1975 m, 29.VII.2010, ♀, Palandöken Mountain, 2300 m, 21.VII.2010, 2 ♂♂, N 39°51'56.4", E 041°16'19.7", 2155 m, 22.VII.2012, 3 ♂♂, Yağmurcuk, N 39°47'13.2", E 041°06'06.1", 2010 m, 9.VIII.2011, ♂; Pasinler, Büyüktuy, 1800 m, 2.VII.2010, ♀, 3 ♂♂, N 39°58'46.7", E 041°27'58", 1819 m, 30.VII.2011, ♂, Demirdöven, 1727 m, 29.VII.2010, ♂, Ügümü, N 40°00'39.9", E 041°43'57.0", 1724 m, 17.VII.2011, ♂; Pazaryolu, N 40°25'02.8", E 040°46'41.3", 1452, 15.VII.2012, ♀, ♂, N 40°25'13.1", E 040°46'13.9", 1445 m, 20.VII.2011, ♂, N 40°25'12.8", E 040°46'13.5", 1450 m, 24.VII.2011, ♂, Çiftepinar, N 40°22'05.3", E 041°43'07.4", 1742 m, 15.VII.2012, ♂, Göztepe, 1672 m, 15.VII.2012, 2 ♂♂, Gülçimen, N 40°24'36.4", E 040°47'28.5", 1663 m, 20.VII.2011, ♂; Şenkaya, Gözebaşı, N 40°24'39.9", E 042°18'11.5", 1794 m, 14.VII.2012, 2 ♂♂, İkizpınar, N

40°36'57.4", E 042°21'30.9", 1589 m, 31.VII.2011, 2 ♂♂, Sındırın, N 40°37'20", E 042°21'34", 1409 m, 14.VII.2014, ♀; Tekman, Güzeldere, N 39°49'17.1", E 042°00'19.8", 1948 m, 16.VII.2011, ♂; Tortum, N 40°16'37.4", E 041°33'30", 1661 m, 23.VI.2011, 4 ♂♂, ♀, Aksukapı, 1653 m, 15.VI.2010, 2 ♀♀, ♂, N 40°23'58.8", E 041°32'46.4", 1648 m, 13.VII.2011, 3 ♂♂, 1573 m, 21.VII.2010, ♂, Aktaş, N 40°15'40.5", E 041°32'39.2", 1765 m, 31.VII.2011, ♂, Aşağı Sivri, N 40°19'33.8", E 041°31'46.9", 1546 m, 16.VII.2012, ♂, Derekapı, 950 m, 15.VI.2010, 2 ♂♂, N 40°24'47.2", E 041°29'19.4", 1247 m, 13.VII.2011, ♂, Kireçli Pass, 2400 m, 31.VII.2010, ♂, Pehlivanlı, N 40°29'37.1", E 041°30'17.6", 1158 m, 21.VII.2012, ♂, ♀, Suyatağı, N 40°27'20.2", E 041°30'14.0", 1236 m, 21.VII.2012, ♂, Taşbaşı, N 40°15'26.6", E 041°32'13.8", 1816 m, 13.VII.2011, ♀, ♂, Yukarı Sivri, N 40°20'27.5", E 041°39'01.1", 1962 m, 16.VII.2012, ♂; Uzundere, Çamlıyamaç, N 40°36'30.1", E 041°32'51.2", 1282 m, 14.VIII.2012, ♂, Gölbaşı, N 40°35'34.2", E 041°34'40.9", 1260 m, 7.VIII.2011, 2 ♂♂, Kirazlı, N 40°34'26.9", E 041°38'52.5", 1231 m, 21.VII.2012, ♂, Yayla, N 40°27'48.6", E 041°39'32.3", 2005 m, 6.VII.2012, 2 ♂♂, N 40°27'48.6", E 041°39'32.3", 2005 m, 13.VII.2011, ♂; Yakutiye, Atatürk University field, 1800 m, 26.VI.1996, ♂, 2.VII.2008, ♂, 9.VII.1986 m, ♂, 14.VII.1986, ♀, 27.VII.2010, ♀, 1850 m, 3.VIII.1998, ♀, Dumlubaba, Güngörmez, 1950 m, 1.VII.2010, 8 ♂, 2500 m, 1.VII.2010, ♂, Güzelyayla, 2000 m, 19.VIII.2009, 2 ♀♀, 4 ♂♂, Karagöbek, N 40°10'15.1", E 041°26'14.9", 2033 m, 13.VII.2011, 2 ♂♂, N 40°09'28.9", E 041°25'58.5", 1961, 16.VII.2012, ♂, N 40°09'57.8", E 041°26'8.1", 1993 m, 31.VII.2011, ♀, ♂, 1998 m, 3.VIII.2009, ♀, Yeşilyayla, 1950 m, 1.VII.2010, ♀, 3 ♂♂.

Distribution in Turkey: Aydin, Bilecik, Bursa, Edirne, İstanbul, Kütahya, Manisa, Mersin, Muğla, Sakarya, Sivas, Şanlıurfa (Önder 1976); Bolu, Bursa, Kocaeli (Önder et al. 1981); Diyarbakır, Gaziantep (Önder et al. 1995); İçel (Lodos et al. 2003); Adana, Hatay, Kahramanmaraş, Kayseri (Önder et al. 2006); İzmir (Tezcan et al. 2010); Diyarbakır (Matocq et al. 2014).

Plagiognathus fulvipennis (Kirschbaum 1856)

Material examined: Erzurum: Karayazı, Yukarı Söylemez, 1896 m, 2.VII.2010, ♀; Köprüköy, Güzelhisar, 1930 m, 2.VII.2010, 2 ♀♀; Narman, Mahmutçavuş, N 40°20'18.8", E 041°55'36.3", 1560 m, 16.VII.2012, ♂; Oltu, Çamlıbel, N 40°29'06", E 041°45'47", 1635 m, 14.VII.2014 m, ♀; Pazaryolu, 1010 m, 7.VIII.2009, ♀; Tortum, Aksukapı, N 40°23'58.8", E 041°32'46.4", 1648 m, 13.VII.2011, 2 ♂♂; Yakutiye, Karagöbek, N 40°09'57.8", E 041°26'8.1", 1993 m, 31.VII.2011, ♂.

Distribution in Turkey: Ankara, Edirne (Hoberlandt 1955); Denizli, Erzurum, Isparta, Kars, Kırklareli, Mersin, Muğla (Önder 1976); Bolu, Bursa, İstanbul, Kocaeli, Sakarya (Önder et al. 1981); Diyarbakır (Önder et al. 1995); İçel (Lodos et al. 2003); Adana, Antalya, Gaziantep, Hatay, Kahramanmaraş, Konya, Nevşehir, Osmaniye (Önder et al. 2006); İzmir (Tezcan et al. 2010); Diyarbakır (Matocq et al. 2014).

Psallus lepidus Fieber 1858

Material examined: Erzurum: Yakutiye, Atatürk University field, 1850 m, 15.VII.1982, 4 ♀♀, 3 ♂♂.

Distribution in Turkey: Hakkâri, Kırklareli, Tekirdağ, (Önder 1976; Önder et al. 2006); Diyarbakır (Önder et al. 1995); Bartın, Zonguldak (Lodos et al. 2003).

Psallus oleae Wagner, 1963

Material examined: Antalya: Manavgat, 8.VI.1978, ♂.

Distribution in Turkey: Antalya (Yayla 1983); Yozgat (Lodos et al. 2003); Hatay, Yozgat (Önder et al. 2006).

Psallus pinicola Reuter 1875

Material examined: Erzurum: Yakutiye, Atatürk University field, 18.VII.1997, ♂, 15.VIII.1997, 2 ♂♂, 18.VIII.1997, ♀, 19.VIII.1998, ♂.

Distribution in Turkey: Artvin (Önder 1976); Ankara, Çankırı, Eskişehir (Lodos et al. 2003; Önder et al. 2006).

Psallus variabilis Fall.

Material examined: Artvin: Yusufeli, Sebzeciler, 500 m, 16.VI.2010, ♀.

Distribution in Turkey: Bursa (Önder et al. 1981); Antalya (Lodos et al. 2003; Önder et al. 2006).

Sthenarus roseri (Herrich-Schaeffer 1838)

Material examined: Erzurum: Narman, Çimenli, N 40°09'31.3", E 041°53'50.5", 2126 m, 30.VI.2012, ♀; Oltu, 23.VI.1988, ♀.

Distribution in Turkey: Balıkesir, Burdur, Çanakkale, Edirne, Isparta, Kütahya, Mersin, Muğla, Sinop (Önder 1976); Bursa, Kocaeli (Önder et al. 1981); Diyarbakır (Önder et al. 1995); İçel (Lodos et al. 2003); Adana, Ankara, Bartın, Bolu, Eskişehir, Kahramanmaraş, Karabük, Karaman, Kastamonu, Kayseri, Kırıkkale, Kırşehir, Konya, Nevşehir (Önder et al. 2006); İzmir (Tezcan et al. 2010).

REFERENCES

- Altınayar G. 1981. Studies on insect fauna in Cereal fields of Central anatolian region. Plant Protection Bulletin, 21(2):53-88, ISSN 0406-3597.
- Atakan E. 2000. Within-plant distribution of predators *Chrysoperla carnea* (Stephens) (Neuroptera: Chrysopidae), *Deraeocoris pallens* Reut. (Heteroptera: Miridae) and *Orius niger* Wolff (Heteroptera: Anthocoridae) on cotton. Turkish Journal of Entomology, 24(4): 267-277.

- Ayyıldız Y. and Athihan R. 2006. Determination of Aphid Species and Their Natural Enemies in Vegetables at Around Balıkesir Province. Yüzüncü Yıl University Journal of Agricultural Sciences, 16(1): 1-5.
- Basnagala S., Wijesekara G.A.W. and Wijayagunasekara H.N.P. 2002. Review of the Subfamily Bryocorinae (Heteroptera: Miridae) of Sri Lanka. Tropical Agricultural Research Vol. 14: 154-164.
- Beyaz G. 2000. Manisa yöresi kültür kekiği (*Origanum* spp.) üretim alanlarındaki Heteroptera takımına bağlı böcek faunasının belirlenmesi üzerine araştırmalar. Aegean Üniversitesi, Fen Bilimleri Enstitüsü, Yüksek Lisans Tezi, 52.
- Bingöl C. M. 1978. Güneydoğu Anadolu Bölgesi'nde kültür yem bitkilerinde zarar yapan Miridae ve Curculionidae familyalarına ait böcek türleri, tanımları ve zararları üzerinde araştırmalar. Bölge Zirai Mücadele Araştırma Enstitüsü, Diyarbakır, 4: 7 pp.
- Cassis G., Wall M. and Schuh R. T. 2006. Insect biodiversity and industrialising the taxonomic process: the Plant Bug Case Study (Insect:Heteroptera:Miridae). Reconstructing the Tree of Life, The Systematics Association Special Volume Series 72:193-212.
- Cassis G. and Schuh R.T. 2012. Systemetic, biodiversity, biogeography, and host associations of the Miridae (Insecta: Hemiptera: Heteroptera: Cimicomorpha). – The Annual review of Entomology 57: 377-404.
- Çam H. 1988. Tokat ve çevresinde kiraz, vişne ve idris ağaçlarında bulunan Heteroptera türlerinin tanımları ve beslenme rejimleri üzerinde araştırmalar. Aegean Üniversitesi Fen Bilimleri Enstitüsü, Yüksek Lisans Tezi, İzmir, 52 pp.
- Çetin H. and Alaoğlu Ö. 2005. Investigations on the olive pests of secondary importance, their population changes and damages in Mut District (Turkey). Selçuk Journal of Agriculture and Food Sciences, 19 (36): 59-65.
- Çevik T. 1996. Faunistic studies on the harmful and beneficial insects on walnuts in Central Anatolia. Plant Protection Bulletin, 36 (1-2): 55–72, ISSN 0406-3597.
- Giray H. 1980. A list of the injurious insects on anise (*Pirnpinella anisum* L.) in the Aegean region (Turkey). Turkish Journal of Plant Protection., 4(1): 49-57.
- Güçlü Ş., Hayat R. and Özbeğ H. 1995a. An investigation on phytophagous insect species on walnut (*Juglans regia* Linnaeus) in Erzurum and its neighbouring provinces. Turkish Journal of Entomology, 19(2): 137-145.
- Güçlü Ş., Hayat R. and Özbeğ H. 1995b. An investigation on phytophagous and predator insect species on olive trees (*Olea europaea* L.) in Artvin provinces. Turkish Journal of Entomology, 19(3): 231-240.
- Hoberlandt L. 1955. Results of the zoological scientific expedition of the national museum in Praha to Turkey. Acta Entomologica Musei Nationalis Pragae, 3: 261 pp.
- Karaat Ş. 1986. Doğu ve Güneydoğu Anadolu Bölgelerinde tütün (*Nicotiana tabacum* L.)’de zararlı olan böcek türleri, tanımları, yayılış alanları ve zararları üzerinde

- araştırmalar. Diyarbakır Bölge Zirai Mücadele Araştırma Enstitüsü Müdürlüğü, Araştırma Eserleri Serisi, No:4, 74 pp.
- Kiyak S., 1990. Binboğa dağları (Kahramanmaraş- Kayseri) kara Heteropter'lerinin erginlerinin eko-faunistik ve sistematigi üzerine araştırmalar. Doktora Tezi, Gazi Üniversitesi, Fen Bilimleri Enstitüsü, Ankara, 172 pp.
- Kiyak S. Özsaraç Ö. and Salur A. 2004. Additional notes on the Heteroptera fauna of Nevşehir province (Turkey). G.U. Journal of Science, 17(1): 21-29.
- Lehr P. A. 1988. Keys to the insects of the far east of the USSR, Volume II. (Hemiptera). Academy of Sciences of the USSR far east Branch Institute of Biology and Soil Sciences, 211 pp.
- Lodos N., Önder F., Pehlivan E. and Atalay R. 1978. Ege ve Marmara Bölgesinin zararlı bölge faunasının tespiti üzerine çalışmalar. T.C. Gıda-Tarım ve Hayvancılık Bakanlığı Zirai Mücadele ve Zirai Karantina Genel Müdürlüğü, Ankara, 135-136.
- Lodos N., Önder F., Pehlivan E., Atalay R., Erkin E., Karsavuran Y. and Tezcan S. 1989. Karadeniz bölgesinin ziraatta zararlı ve faydalı böcek faunasının tespiti üzerinde araştırmalar [Curculionidae, Scarabaeidae (Coleoptera), Plataspidae, Cydnidae, Acaethosomatidae, Scutelleridae, Pentatomidae, Lygaeidae, Miridae (Hemiptera)]. Aegean Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Bornova, İzmir, 75 pp.
- Lodos N., Önder F., Pehlivan E., Atalay R., Erkin E., Karsavuran Y., Tezcan S. and Aksoy S. 2003. Faunistic studies on Miridae (Hemiptera) of western Black Sea, Central Anatolia and Mediterranean Regions of Turkey. Department of Plant Protection Faculty of Agriculture University of Aegean 35100 Bornova, İzmir, Turkey, 85 pp.
- Matocq A. and Özgen, İ., 2010. A preliminary list of Heteroptera collected in Mardin and Siirt provinces from South-Eastern Anatolia of Turkey (Hemiptera). Mun. Ent. Zool., 5: 1011-1019.
- Matocq, A., Pluot-Sigwalt, D., Özgen, İ., 2014. Terrestrial Hemiptera (Heteroptera) collected in South-East Anatolia (Diyarbakır, Mardin and Elazığ Provinces) (Turkey): Second List. Mun. Ent. Zool., 9(2):884-930.
- Ölmez, M., 2008. Kahramanmaraş İli mısır alanlarında zararlı ve yararlı faunanın belirlenmesi ve önemli zararlıların popülasyon gelişimi takibi. Kahramanmaraş Sütçü İmam Üniversitesi, Fen Bilimleri Enstitüsü, Yüksek Lisans Tezi, Kahramanmaraş, 39 pp.
- Öncüler, C., 1991. Türkiye bitki zararlısı böceklerinin parazit ve predatör kataloğu. Ege University Faculty of Agriculture Publications, No:505, 15-33.
- Önder F. 1976. Türkiye Miridae (Hemiptera) faunası üzerinde sistematik çalışmalar. Ege Üniversitesi Ziraat Fakültesi, Entomoloji ve Zirai Zooloji Kürsüsü, Doçentlik Tezi, İzmir, 506 pp.
- Önder F., Ünal A. and Ünal E. 1981. Hemiptera fauna collected by light traps in some districts of northwestern part of Anatolia. Turkish Journal of Plant Protection, 5(3): 151-169.

- Önder F. and Lodos N. 1987. General evaluation of the predatory species of Heteroptera in the Turkish fauna. *Turkish Journal of Entomology*, 11(2): 117-125.
- Önder F., Karsavuran Y. and Tezcan S., 1990. Yabancı ot savaşında potansiyel önemine sahip Türkiye Hemiptera faunası üzerinde araştırmalar. *Çevre Biyolojisi Sempozyumu*, Ankara, 15.
- Önder F., Karsavuran Y. and Örümlü E. E. 1995. Güneydoğu Anadolu Projesi (GAP) uygulama alanında saptanan Cimicoidea ve Tingoidae (Heteroptera) türlerinin durumu ve gelecek için düşündürdükleri. GAP Bölgesi Bitki Koruma Sorunları ve Çözüm Önerileri Sempozyumu Bildirileri, 27-29 Nisan, Şanlıurfa, 149-158.
- Önder F., Karsavuran Y. and Tezcan S. 1998. Investigations on the vertical distribution in the habitats of plant bugs (Hemiptera: Miridae) of Turkey. *Journal of Agriculture Faculty of Ege University*, 35(1-3): 33-40.
- Önder F., Karsavuran Y., Tezcan S. and Fent M. 2006. *Türkiye Hemiptera Kataloğu*. Meta Basım Matbaacılık Hizmetleri, İzmir, 164.
- Özbek H. and Alaoglu Ö. 1987. Phytophagus Heteroptera on Potato in Erzurum. *Plant Protection Bulletin*, 27(3-4): 227-238, ISSN 0406-3597.
- Özkan A. 1984. Antalya ve çevresi yumuşak çekirdekli meyve ağaçlarının Coleoptera ve Hemiptera takımlarına ait faydalı böcek türleri, tanınmalari, konukcuları ve önemlilerinin etkinlikleri üzerine araştırmalar. *Biyolojik Mücadele Araştırma Enstitüsü*, Antalya, 65.
- Tamer, A., Has, A., Aydemir, M., Çalışkaner, S., 1998. Faunistic survey studies on harmful and beneficial insects on food legumes (lentil, chickpea, bean) in Central Anatolia Region. *Plant Protection Bulletin*, 38(1-2):65-90, ISSN 0406-3597.
- Tezcan S. and Önder F. 1999. Heteropterous insects associated with cherry trees in Kemalpaşa district of İzmir, Turkey. *Journal of Agriculture Faculty of Ege University*, 36 (1-3): 119-124.
- Tezcan S. and Önder F. 2003. Faunistical studies in ecological cherry orchards in Izmir and Manisa Provinces of Turkey: An evaluation on the species of Heteroptera. *Anadolu*, 13(1): 124-131.
- Tezcan S., Tezcan F. and Gülperçin N. 2010. *İzmir'den 4000 Böcek Türü*. Egetan Bas. Yay. Tan. Ltd. Şti., İzmir, 250.
- Yaşarakıncı N. and Hıncal P. 1997. The Research on determining the pests and beneficial species and their population densities on the tomato, cucumber, pepper and lettuce glasshouses in İzmir. *Plant Protection Bulletin*, 37(1-2): 79-89, ISSN 0406-3597.
- Yaşarakıncı N. and Hıncal P. 2000. The studies on the pests, their natural enemies and population developments on protected eggplant in İzmir province. *Plant Protection Bulletin*, 40(1-2): 29-48, ISSN 0406-3597.
- Yayla A. 1983. Antalya ve çevresi zeytin ağaçlarında rastlanan faydalı Hemipter'lerin tanınmalari, konukcuları ve etkinlikleri üzerine araştırmalar. *Biyolojik Mücadele Araştırma Enstitüsü*, Antalya, 58.

- Yıldırım E. and Özbek H. 1992. Erzurum Şeker Fabrikasına bağlı şekerpancarı üretim alanlarındaki zararlı ve yararlı böcek türleri. Türkiye II. Entomoloji Kongresi, 28-31 Ocak 1992, Adana, 621-635.
- Yıldırım E., Özbek H. and Önder F. 1999. Heteropterous species caught in light traps in the Campus of Atatürk University in Erzurum (Türkiye). Turkish Journal of Entomology, 23(3): 225-228.
- Yıldırım E. 2002. *Phoenicocoris obscurellus* (Fallen, 1829) (Heteroptera: Miridae) a new record and a new scotch pine (*Pinus sylvestris* L.) pest for Turkey from Erzurum. Turkish Journal of Entomology, 26(1): 45-49.
- Wagner E. 1955. Die Plagiognathus gruppe (Hemiptera: Heteroptera: Miridae). Acta Entomologica Musei Nationalis Pragae, 30(461): 291-304.