

Compact and noncompact structures of the nonlinearly dispersive $GNLS(m,n,k,l)$ equation

Bulent Kilic

Firat University, Department of Mathematics, 23119 Elazig, Turkey.

Received: 24 July 2014, Revised: 7 August 2014, Accepted: 2 December 2014

Published online: 10 February 2015

Abstract: In this paper, we establish exact-special solutions of the generalized nonlinear dispersion $GNLS(m,n,k,l)$ equation. We use the ansatz method for acquiring the compactons, solitary patterns, solitons and other types of solutions.

Keywords: Generalized nonlinear dispersion, Ansatz method, solitary patterns.

1 Introduction

The mathematical modeling of scientific events usually is expressed by nonlinear evolution equations. So, it is crucial to reach general solutions, which give some behaviors about the character and the structure of the equations for researchers, of these corresponding nonlinear equations. Many effective methods have been improved to provide much information for physicians and engineers to present the actual physical characters of solutions [1]. We recall that most of these methods use the wave variable transformation to reduce the nonlinear PDE to ODE in order to acquire the solution. Several are the generalized Miura [2], Darboux [3], Cole-Hopf [4], Hirota's dependent variable [5], the Backlund [6] transformations and sine-cosine [7], homogeneous balance (HB) [8], similarity reduction [9], automated tanh-function [10], Exp-function [11], (G'/G) -expansion [12] methods and there are many papers about the basis of compactons [13–32].

De Angelis studied [14] the cubic quintic nonlinear Schrödinger equation;

$$ip_x + p_{tt} + \gamma p|p|^2 + \alpha p|p|^4 = 0, \quad (1)$$

Yan [15] studied $NLS(m,n)$ equation;

$$ip_t + \left(p|p|^{m-1}\right)_{xx} + \mu p|p|^{n-1} = 0, \quad (2)$$

and obtained envelope compactons and solitary pattern solutions.

Agrawal and Zeng studied [16],[17] w.r.t. the GNLS equation;

$$ip_t + Ap_{xx} + Bp|p|^2 + iCp_{xxx} + iD\left(p|p|^2\right)_x = 0, \quad (3)$$

* Corresponding author e-mail: bulentkilic@firat.edu.tr

Hirota introduced [18] another GNLS equation in the following form

$$ip_t + Ap_{xx} + Bp|p|^2 + iCp_{xxx} + iD|p|^2 p_x = 0. \quad (4)$$

Yan introduced and studied [19] the GNLS equation with nonlinear dispersion (called GNLS(m, n, k, l) equation) and the equation given by

$$ip_t + A(p|p|^{m-1})_{xx} + Bp|p|^{n-1} + iC(p|p|^{k-1})_{xxx} + iD(p|p|^{l-1})_x = 0, \quad (5)$$

where A, B, C and D are arbitrary constants.

For this paper our aim is to apply the ansatz method [20–23] to Eq. (5) in light of the above-mentioned studies and then we give some explanations about the solutions and conclusions in the last section.

2. Ansatz method

For initially, the solution of the equation is considered

$$\left(\frac{dv}{dy}\right)^2 = A_0 - A_1 v^2 \quad (6)$$

where $A_0 \neq 0$ and $A_1 \neq 0$ are constants. When $A_1 > 0$, Eq. (6) has the solutions:

$$\begin{aligned} v_1 &= \pm \sqrt{\frac{A_0}{A_1}} \sin[\sqrt{A_1}(y+a)], \\ v_2 &= \pm \sqrt{\frac{A_0}{A_1}} \cos[\sqrt{A_1}(y+a)], \end{aligned} \quad (7)$$

where a is constant. If $A_1 < 0$ then the Eq. (6) has the solutions:

$$\begin{aligned} v_3 &= \pm \sqrt{\frac{A_0}{A_1}} \sinh[\sqrt{A_1}(y+a)], \\ v_4 &= \pm i \sqrt{\frac{A_0}{A_1}} \cosh[\sqrt{A_1}(y+a)], \end{aligned} \quad (8)$$

where $i = \sqrt{-1}$. For secondly, if the solutions of the other equation is considered,

$$\left(\frac{dv}{dy}\right)^2 = v^2 (A_2 + A_3 v^2) \quad (9)$$

where $A_2 \neq 0$ and $A_3 \neq 0$ are constants. When $A_2 < 0$, Eq. (9) has the solutions

$$\begin{aligned} v_5 &= \pm \sqrt{\frac{A_2}{A_3}} \sec[\sqrt{-A_2}y], \\ v_6 &= \pm \sqrt{\frac{A_2}{A_3}} \csc[\sqrt{-A_2}y], \end{aligned} \quad (10)$$

If $A_2 > 0$ in Eq. (9), then the solutions are

$$\begin{aligned} v_7 &= \pm \sqrt{\frac{A_2}{A_3}} \operatorname{sech}[\sqrt{-A_2}y], \\ v_8 &= \pm i \sqrt{\frac{A_2}{A_3}} \operatorname{csch}[\sqrt{-A_2}y], \end{aligned} \quad (11)$$

2.1 Nonlinear Dispersive GNLS(m, n, k, l) Equation

If we take the solution with transform $u(x, t) = \phi(\xi) \exp(i\sigma t)$ with

$\xi = c(x - \lambda t)$, (c, λ and σ are constants) we have the following ODE:

$$-\sigma\phi + Ac^2(\phi^m)'' + B\phi^n + i \left[-\lambda c\phi' + Cc^3(\phi^k)''' + Dc(\phi^l)' \right] = 0 \tag{12}$$

By separating real and imaginary parts of the above equation and then by integration of imaginary part we have

$$-\sigma\phi + Ac^2(\phi^m)'' + B\phi^n = 0 \tag{13}$$

$$-\lambda\phi + Cc^2(\phi^k)'' + D\phi^l = 0 \tag{14}$$

To reach compactons, solitary patterns, solitons and periodic solutions of nonlinearly dispersive GNLS equation firstly assume the following constraints:

$$\lambda = \sigma, \quad A = C, \quad B = D, \quad m = k \text{ and } n = l \tag{15}$$

then by the following transformation

$$\frac{d\phi^m}{d\xi} = z, \quad \frac{d^2\phi^m}{d\xi^2} = z \frac{dz}{d\phi^m} \tag{16}$$

substituting (16) into (13) leads to the following equation,

$$\frac{Ac^2}{2} \left(n\phi^{\frac{n-3}{2}} \frac{d\phi}{d\xi} \right)^2 = \frac{n\sigma}{n+1} - \frac{nB}{m+n} \phi^{n-1}. \tag{17}$$

Letting $\phi = w^{\frac{2}{n-1}}$, we have

$$\phi = w^{\frac{2}{n-1}} \Rightarrow d\phi = \frac{2}{n-1} \phi^{\frac{n-2}{n-1}-1} dw \tag{18}$$

which changes Eq. (13) to

$$\frac{2Ak^2n}{(m-1)^2} \left(w^{\frac{n-m}{m-1}} \frac{dw}{d\xi} \right)^2 = \frac{\sigma}{n+1} - \frac{n}{m+n} w^2. \tag{19}$$

Case I. If we take $n = 2m - 1$ in Eq. (19), then we get:

$$p_1(x, t) = \exp[i\sigma t] \left[\pm \frac{1}{A_4} \left(A_5 - A_4^2 k^2 (x - \lambda t)^2 \pm 2A_4^2 k^2 (x - \lambda t)^2 A_6 - A_4^2 A_6^2 \right) \right]^{\frac{1}{m-1}} \tag{20}$$

where

$$A_4 = \frac{B(m-1)^2}{2Ak^2(3m-1)(2m-1)} \text{ and } A_5 = \frac{(m-1)^2}{4Ak^2m(2m-1)}$$

Case II. If $m = n$ we know that Eq. (19) becomes

$$\left(\frac{dw}{d\xi} \right)^2 = \frac{\sigma(m-1)^2}{2Ak^2m(m+1)} - \frac{B(m-1)^2}{4Ak^2m^2} w^2. \tag{21}$$

in which we assume that $m \neq -1$ and $A, B, k, \lambda \neq 0$.

If $AB > 0$, it becomes from Eqs. (6) and (21) that

$$p_2(x, t) = \exp [i\tau t] \left(\frac{2\sigma m}{B(m+1)} \sin^2 \left[\frac{m-1}{2m|k|} \sqrt{\frac{B}{A}} (c(x-\lambda t) + a) \right] \right)^{\frac{1}{m-1}}, \quad (22)$$

$$p_3(x, t) = \exp [i\tau t] \left(\frac{2\sigma m}{B(m+1)} \cos^2 \left[\frac{m-1}{2m|k|} \sqrt{\frac{B}{A}} (c(x-\lambda t) + a) \right] \right)^{\frac{1}{m-1}}, \quad (23)$$

If $AB < 0$, it becomes from Eqs. (7) and (21) that

$$p_4(x, t) = \exp [i\tau t] \left(-\frac{2\sigma m}{B(m+1)} \sinh^2 \left[\frac{m-1}{2m|c|} \sqrt{-\frac{B}{A}} (c(x-\lambda t) + a) \right] \right)^{\frac{1}{m-1}}, \quad (24)$$

$$p_5(x, t) = \exp [i\tau t] \left(\frac{2\sigma m}{B(m+1)} \cosh^2 \left[\frac{m-1}{2m|c|} \sqrt{-\frac{B}{A}} (c(x-\lambda t) + a) \right] \right)^{\frac{1}{m-1}}. \quad (25)$$

Theorem 1. The nonlinear dispersion GNLS equation has solutions under the constraint (15) described as follows.

1. When $AB > 0$ and $m > 1$

$$p(x, t) = \begin{cases} \exp [i\tau t] \left(\frac{2\sigma m}{B(m+1)} \cos^2 \left[\frac{m-1}{2m|c|} \sqrt{\frac{B}{A}} (c(x-\lambda t) + a) \right] \right)^{\frac{1}{m-1}}, & \left| \frac{m-1}{2m|c|} \sqrt{\frac{B}{A}} (c(x-\lambda t) + a) \right| \leq \frac{\pi}{2} \\ 0, & \text{Otherwise} \end{cases} \quad (26)$$

is the solitary wave solution(s.w.s) with compact support of GNLS equation.

2. When $AB > 0$ and $m > 1$

$$p(x, t) = \begin{cases} \exp [i\tau t] \left(\frac{2\sigma m}{B(m+1)} \sin^2 \left[\frac{m-1}{2m|c|} \sqrt{\frac{B}{A}} c(x-\lambda t) \right] \right)^{\frac{1}{m-1}}, & 0 \leq \frac{m-1}{2m|c|} \sqrt{\frac{B}{A}} c(x-\lambda t) \leq \pi \\ 0, & \text{Otherwise} \end{cases} \quad (27)$$

is a compacton solution.

3. When $AB < 0$ and $m < 1$

$$p(x, t) = \exp [i\tau t] \left(\frac{B(m+1)}{2\sigma m} \csc^2 \left[\frac{m-1}{2m|c|} \sqrt{\frac{B}{A}} c(x-\lambda t) \right] \right)^{\frac{1}{1-m}} \quad (28)$$

is a s.w.s for GNLS Eq. for $0 < (x-\lambda t) < \frac{2m|c|}{m-1} \sqrt{\frac{A}{B}} \pi$.

4. When $AB < 0$ and $m < 1$

$$p(x, t) = \exp [i\tau t] \left(\frac{B(m+1)}{2\sigma m} \sec^2 \left[\frac{m-1}{2m|c|} \sqrt{\frac{B}{A}} c(x-\lambda t) \right] \right)^{\frac{1}{1-m}} \quad (29)$$

is a s.w.s for the GNLS equation for $|x - \lambda t| < \frac{2m|c|}{m-1} \sqrt{\frac{A}{B}} \pi$.

Theorem 2. The nonlinear dispersion GNLS equation has the following solutions with $m=n$ under the constraint (19).

1. When $B < 0$, $\sigma m(m+1) > 0$

$$p(x,t) = \exp[i\tau t] \left(-\frac{2\sigma m}{B(m+1)} \sinh^2 \left[\frac{m-1}{2m|c|} \sqrt{-\frac{B}{A}} c(x-\lambda t) \right] \right)^{\frac{1}{m-1}}, \tag{30}$$

is a solitary pattern solution (s.p.s).

1. When $B < 0$, $\sigma m(m+1) < 0$

$$p(x,t) = \exp[i\tau t] \left(\frac{2\sigma m}{B(m+1)} \cosh^2 \left[\frac{m-1}{2m|c|} \sqrt{-\frac{B}{A}} c(x-\lambda t) \right] \right)^{\frac{1}{m-1}}, \tag{31}$$

is a s.p.s, when $m < 1$, Eq. (31) is a bounded solution.

1. $m=n < 1$ solutions (30) and (31) become s.w.s

$$p(x,t) = \exp[i\tau t] \left(-\frac{2\sigma m}{B(m+1)} \operatorname{csch}^2 \left[\frac{m-1}{2m|c|} \sqrt{-\frac{B}{A}} c(x-\lambda t) \right] \right)^{\frac{1}{1-m}}, \tag{32}$$

which is the singular s.w.s. The solution improves a singularity at a finite point, i.e., for any fixed $t = t_0$, there is blow up of the solution [24], [25].

$$p(x,t) = \exp[i\tau t] \left(\frac{2\sigma m}{B(m+1)} \operatorname{sech}^2 \left[\frac{m-1}{2m|c|} \sqrt{-\frac{B}{A}} c(x-\lambda t) \right] \right)^{\frac{1}{1-m}}, \tag{33}$$

which is the bell-shaped s.w.s.

Case III. If $n = 1$ then the Eq. (19) becomes

$$\left(\frac{dw}{d\xi} \right)^2 = w^2 \left(\frac{\sigma(m-1)^2}{4Ac^2m} + \frac{-B(m-1)^2}{2Ac^2(m+1)} w^2 \right). \tag{34}$$

If $\sigma Am < 0$ and $m > 1$ it becomes from Eqs. (9) and (34) that

$$p_6(x,t) = \exp[i\tau t] \left(\frac{\sigma(m+1)}{2Bm} \sec^2 \left[\frac{m-1}{2|c|} \sqrt{-\frac{\sigma}{Am}} (c(x-\lambda t)) \right] \right)^{\frac{1}{m-1}}, \tag{35}$$

$$p_7(x,t) = \exp[i\tau t] \left(\frac{\sigma(m+1)}{2Bm} \operatorname{csc}^2 \left[\frac{m-1}{2|c|} \sqrt{-\frac{\sigma}{Am}} (c(x-\lambda t)) \right] \right)^{\frac{1}{m-1}}. \tag{36}$$

Case IV. If $\sigma Am > 0$ and $m > 1$ it becomes from Eqs. (10) and (34) that

$$p_8(x,t) = \exp[i\tau t] \left(\frac{\sigma(m+1)}{2Bm} \operatorname{sech}^2 \left[\frac{m-1}{2|c|} \sqrt{-\frac{\sigma}{Am}} (c(x-\lambda t)) \right] \right)^{\frac{1}{m-1}}, \tag{37}$$

which is the bell-shaped s.w.s.

$$p_9(x,t) = \exp[i\tau t] \left(-\frac{\sigma(m+1)}{2Bm} \operatorname{csch}^2 \left[\frac{m-1}{2|c|} \sqrt{-\frac{\sigma}{Am}} (c(x-\lambda t)) \right] \right)^{\frac{1}{m-1}}, \quad (38)$$

which is the singular s.w.s. The solution improves a singularity at a finite point, i.e., for any fixed $t = t_0$, there is blow up of the solution [24], [25].

Case V. If $\sigma Am < 0$ and $m > 1$

we have the following compacton solutions with the aid of Eqs. (37) and (38)

$$p_{10}(x,t) = \exp[i\tau t] \left(\frac{2Bm}{\sigma(m+1)} \cos^2 \left[\frac{m-1}{2|c|} \sqrt{-\frac{\sigma}{Am}} c(x-\lambda t) \right] \right)^{\frac{1}{1-m}}, \quad (39)$$

$$\left| (m-1)(x-\lambda t) \sqrt{-\frac{\sigma}{Am}} \right| \leq \pi \text{ and } p = 0 \text{ otherwise.}$$

$$p_{11}(x,t) = \exp[i\tau t] \left(\frac{2Bm}{\sigma(m+1)} \sin^2 \left[\frac{m-1}{2|c|} \sqrt{-\frac{\sigma}{Am}} c(x-\lambda t) \right] \right)^{\frac{1}{1-m}}, \quad (40)$$

$$\text{for } 0 \leq \left| (m-1)(x-\lambda t) \sqrt{-\frac{\sigma}{Am}} \right| \leq 2\pi \text{ and } p = 0 \text{ otherwise.}$$

Case VI. If $\sigma Am > 0$ and $m < 1$

Using $\cosh(x) = \cos(ix)$ and $\sinh(x) = -\sin(ix)$ we have the following s.p.s

$$p_{12}(x,t) = \exp[i\tau t] \left(-\frac{2Bm}{\sigma(m+1)} \sinh^2 \left[\frac{m-1}{2|c|} \sqrt{-\frac{\sigma}{Am}} c(x-\lambda t) \right] \right)^{\frac{1}{1-m}}, \quad (41)$$

$$p_{13}(x,t) = \exp[i\tau t] \left(\frac{2Bm}{\sigma(m+1)} \cosh^2 \left[\frac{m-1}{2|c|} \sqrt{-\frac{\sigma}{Am}} c(x-\lambda t) \right] \right)^{\frac{1}{1-m}} \quad (42)$$

4. Conclusions

We used the ansatz method with the aid of reducing the order of the equation for finding some new exact solutions for GNLS(m, n, k, l) equation. We have acquired different types of exact solutions to the equation. The solutions obtained are expressed in terms of s.w, s.p, compacton, bell shaped and singular.

Conflict Interest

I declare that there is no conflict of interests regarding the publication of this article.

References

- [1] Hereman W., Banerjee P.P., Korpel A., Assanto, G., van Immerzeel A., Meerpoel, A., Exact solitary wave solutions of nonlinear evolution and wave equations using a direct algebraic method. Journal Physics A: Mathematical and General 19(1986)607-628.

- [2] Wang Deng-S., Complete integrability and the Miura transformation of a coupled KdV equation, *Applied Mathematics Letters* 23(2010)665-669.
- [3] Geng X., H. G., Darboux transformation and explicit solutions for the Satsuma–Hirota coupled equation, *Applied Mathematics and Computation* 216(2010)2628-2634.
- [4] Cesar A., Gómez S., Alvaro H. S., The Cole–Hopf transformation and improved tanh–coth method applied to new integrable system (KdV6), *Applied Mathematics and Computation* 204(2008)957-962.
- [5] Lei Y., Fajiang Z., Yinghai W., The homogeneous balance method, Lax pair, Hirota transformation and a general fifth-order KdV equation, *Chaos, Solitons and Fractals* 13(2002)337-340.
- [6] Wang M. L., Wang Y.M., A new Bäcklund transformation and multi-soliton solutions to the KdV equation with general variable coefficients. *Physics Letters A* 287(2001)211-216.
- [7] Taşcan F., Bekir A., Analytic solutions of the $(2 + 1)$ -dimensional nonlinear evolution equations using the sine–cosine method, *Applied Mathematics and Computation* 215(2009)3134-3139.
- [8] Wang M.L., Zhou Y.B., Li Z.B., Application of a homogeneous balance method to exact solutions of nonlinear equations in mathematical physics, *Physics Letters A* 216(1996)67-75.
- [9] Fan E.G., Auto-Bäcklund transformation and similarity reductions for general variable coefficient KdV equations, *Physics Letters A* 294(2002)26-30.
- [10] Parkes E.J., Duffy B.R., An automated tanh-function method for finding solitary wave solutions to non-linear evolution equations. *Computers Physics Communications* 98(1996)288-300.
- [11] Wu, X. H. and He, J. H. Solitary solutions, periodic solutions and compacton-like solutions using the Exp-function method. *Computer Mathematics with Application* 54 (2007) 966–986.
- [12] Zhang S., Ba JM., Sun YN., Dong L., A Generalized (G'/G) -expansion method for the nonlinear Schrödinger equation with variable coefficients, *Z. Naturforsch* 64(2009)691-696.
- [13] U. Goktas, E. Hereman Symbolic computation of conserved densities for systems of nonlinear evolution equations *J. Symb. Comput.* , 24(5)(1997)591–622.
- [14] De Angelis C., Self-trapped propagation in the nonlinear cubic-quintic equation: a variational schrodinger approach, *iEEE Journal of Quantum Electronics* 30(1994)818-821.
- [15] Yan Z, [Envelope compactons and solitary patterns](#), *Physics Letters A* 355(2006)212-215.
- [16] P. G. Agrawal, *Nonlinear Fiber Optics*, (Academic Press, New York, 1989).
- [17] Zheng W, An envelope solitary-wave solution for a generalized nonlinear Schrödinger equation, *Journal of Physics A* 27(1994)931-934.
- [18] Hirota R, [Exact envelope soliton solutions of the wave equation](#), *Journal of Mathematical Physics* 14(1973)805-810.
- [19] Yan Z, Envelope compact and solitary pattern structures for the GNLS(m,n,p,q) equations, *Physics Letters A* 357(2006)196–203.
- [20] Lai S, Wu Y, Zhou Y., Some physical structures for the $(2+1)$ dimensional Boussinesq water equation with positive and negative exponents, *Computers & Mathematics with Applications* 56(2008)339–345.
- [21] Lai S, Wu Y., The compact and noncompact structures for two types of generalized Camassa-Holm-KP equations, *Mathematical and Computer Modelling* 47(2008)1089–1098.
- [22] Lai S, Wu Y., Wiwatanapataphee B., On exact travelling wave solutions for two types of nonlinear $K(n,n)$ equations and a generalized KP equation, *Journal of Computational and Applied Mathematics* 212(2008)291–299.
- [23] Lai SJ., Different physical structures of solutions for a generalized Boussinesq wave equation, *Computational & Applied Mathematics* 231(2009)311–318.
- [24] Clarkson PA, Mansfield E. Symmetry reductions and exact solutions of a class of nonlinear heat equations, *Physica D* 70(1993)250-288.
- [25] Kudryashov N.A, E. D. Zargaryan, Solitary waves in active dissipative media, *Journal of Physics A* 29(1996)8067-8077.
- [26] He JH. Asymptotic methods for solitary solutions and compactons, *Abstract and Applied Analysis* (2012)916793.
- [27] Inc M, On new exact special solutions of GNLS(m,n,p,q) equations, *Modern Physics Letter B* 24(2010)1769-1783.
- [28] Inc M, New compact and noncompact solutions of the $K(k,n)$ equations, [Chaos, Solitons & Fractals](#) 29(2006)895–903.
- [29] Inc M, New exact solitary pattern solutions of the nonlinearly dispersive $R(m,n)$ equations, [Chaos, Solitons & Fractals](#) 29(2006)499–505.

- [30] Abdou M.A., New exact travelling wave solutions for the generalized nonlinear Schroedinger equation with a source, *Chaos, Solitons and Fractals* 38(2008)949–955.
- [31] Sun C, Gao H, Exact solitary and periodic wave solutions for a generalized nonlinear Schrödinger equation, *Chaos, Solitons and Fractals* 39(2009)2399–2410.
- [32] Yomba E, The sub-ODE method for finding exact travelling wave solutions of generalized nonlinear Camassa–Holm, and generalized nonlinear Schrödinger equations, *Physics Letters A* 372(2008)215–222.