



**The Use of Amelogenin Gene in
Sex Determination from Human Skeletal
Fragments and Teeth Specimens**
İnsan İskeleti Parçalarından ve Diş Örneklerinden Amelogenin
Genini Kullanarak Cinsiyet Belirleme

Abdullahi Daudu Zagga¹, Hamidu Ahmed², Aziz Abdo Tadros¹,
Somaia Muhammad Ismail³

¹Department of Anatomy and ²Department of Paediatrics, Usmanu Danfodiyo University, Sokoto, Nigeria.

³Department of Medical Molecular Genetics, National Research Centre, Cairo, Egypt.

ABSTRACT

Alternative approaches to sex determination of DNA samples involve investigation of regions of the amelogenin gene. This is the gene that encodes tooth enamel and is present on both the X and Y chromosomes. A review composed via Medline Internet search of literature and contributions from our experiences as well as experiences from colleagues. The rareness of failures in sex determination provides confidence in current techniques, but amelogenin gene method (singly) of sex determination is not without failures. Amelogenin PCR method/system of sex determination should not, at the moment, completely replace traditional methods of sex identification. Hence, sex identification with amelogenin gene, of subjects for forensic purposes should be conducted as much as possible through a multiple morphological-molecular combined methods to avoid fallibility of *amelogenin* gene.

Key words: Amelogenin, bone, tooth, sexing.

ÖZET

DNA örneklerindeki cinsiyet tayini için alternatif yaklaşımlardan biri de Amelogenin gen bölgeleridir. X ve Y kromozomlarının üzerinde bulunan bu gen diş minesini de kodlar. Medline araştırma sitesinde makale ve deneyimler taranarak kendi deneyimlerimizin dışında farklı araştırmalarda gözden geçirilmektedir. Mevcut tekniklerle cinsiyet tayini başarısızlıkları seyreklik gösterirken, Amelogenin



gen yöntemi tek başına başarılı değildir. Amelogenin PCR yöntemi/sisteminde cinsiyet tayini şu anda geleneksel yöntemin yerini almamalıdır. Bu nedenle özellikle adli amaçlı konularda Amelogenin geniyle cinsiyet tayininin yanılabilirliğini önlemek için bir çok morfolojik-moleküler yöntemle birleştirilerek yapılmalıdır.

Anahtar Kelimeler: Amelogenin geni, kemik, diş, cinsiyet

Amelogenin

Amelogenin is a low-molecular-weight protein found in developing tooth enamel, and it belongs to a family of extracellular matrix (ECM) proteins. Developing enamel contains about 30% protein, and 90% of this is comprised of *amelogenins*¹.

In the early stages of tooth development, internal enamel epithelial cells differentiate into ameloblasts which synthesize and secrete specific proteins as enamel matrix. Protein concentrations as high as 25-30% can be found in newly-secreted enamel. The matrix proteins then decrease during the maturation of the enamel and are replaced by apatite crystals. These proteins are thought to play important roles in the process of enamel mineralization¹.

There are two classes of proteins in the early enamel, *amelogenin* and enamelin. *Amelogenin*, which was termed by Eastoe² is a discrete and major constituent unique to the developing enamel. This protein contains high concentrations of proline, glutamine, leucine and histidine. *Amelogenin* is now well characterized from amino acid sequencing data and the gene structure¹. Another protein, enamelin, termed by Termine *et al.*³ is a minor protein component present in a form associated with the mineral phase in the developing enamel.

Identification of the Amelogenin Gene

The *amelogenin* genes were identified on sex chromosomes in both mouse and man first by Lau *et al.*⁴. They constructed a recombinant cDNA for mouse *amelogenin* and used it as a specific hybridization probe. They found that there is only one *amelogenin* gene in the mouse genome, which is located on the X-chromosome. The analysis established the locus of the *amelogenin* gene to the distal portion of the mouse X-chromosome. On the other hand, two copies of the *amelogenin* genes were detected in the human male genome by Southern blot analysis of human-rodent cell hybrids. One copy of the *amelogenin* genes was located on the

distal short arm of the X-chromosome in the p22.1-p22.3 region. The second copy of the *amelogenin* gene was localized near the centromere of the Y-chromosome. However, they did not determine DNA sequences for human and mouse *amelogenin* genes. At about the same time, human *amelogenin* gene, assumed to be derived from X-chromosome, was sequenced by Shimokawa *et al.*⁵.

Thereafter, it has been confirmed that there are two *amelogenin* genes, one on the X-chromosome and the other on the Y-chromosome in some animals and a single gene resides exclusively on the X-chromosome in other animals¹.

In mammals, *amelogenin* is said to possess seven exons, although exon 4 is lacking in most species studied, and two extra exons (8 and 9) are found in human and rodents⁶. The five first exons (exon1—exon5) are small (42—56 bp) and exon 6 (426 bp) and 7 (160 bp) are large. Exon 1 is uncoding, most exon 2 sequence codes the signal peptide, and the only first three nucleotides of exon 7 are coding for the protein⁶. The C- and N-terminal regions are hydrophilic, while the central region (most of exon 6) is hydrophobic. *Amelogenin* is subjected to extensive alternative splicing⁷. The N-terminal region contains amino acids that are suspected to be involved in various functions: a phosphorylation site⁸, alpha helices⁹, a binding site to *N*-acetylglucosamine and keratins¹⁰. The hydrophilic C- and N-terminal regions interact with the surrounding environment (cell processes and hydroxyapatite crystals) and the hydrophobic region (exon 6) forms the core of the nanospheres, which are responsible for the enamel microstructure¹¹. The C-terminal region plays an important role in contributing to stabilize these nanospheres¹².

Nucleotide Sequences of Human Amelogenin Genes

Human *amelogenin* gene was first isolated and sequenced by Shimokawa *et al.*⁵. A human genomic DNA library in the phage-vector was screened by plaque hybridization using a bovine *amelogenin* cDNA probe. One clone containing a 13-kb insert, probably from X-chromosome, was obtained. Restriction enzyme fragments were subcloned and sequenced by the dideoxy chain termination method. Some segments of the isolated human genomic DNA corresponded to the bovine *amelogenin* sequence. At that time, the presence of at least four exons in the human gene structure was suggested. First coding exon was not identified, although nucleotides of three putative exons (corresponding to exons 3, 5 and 6 by Salido *et al.*¹³) were sequenced and the deduced amino acid sequence was reported¹.

Nakahori *et al.*¹⁴ reported two genomic sequences for human *amelogenin* from the X- and Y-chromosomes. They isolated a Y-specific clone encoding *amelogenin* which was used as a probe. Southern hybridization analysis of male and female genomic DNAs detected a male-specific band on the Y-chromosome and a band common to male and female on the X-chromosome. By digestion analysis with restriction enzymes, the difference between the X- and the Y-chromosome sequences was confirmed. They sequenced the nucleotides in the homologous regions from the X- and Y-chromosomes and aligned them with the best possible match. There was 88.9% homology between the X and Y nucleotide sequences. First exon was not identified and three presumed exons (exons 3, 5 and 6 by Salido *et al.*¹³) both on the X and Y (93% homology) were recognized and amino acid sequences were deduced¹.

Fincham *et al.*¹⁵ reported the differences in human *amelogenin* components according to the sex of the individual. They extracted enamel protein samples from fetal and postnatal human deciduous dentitions and the proteins were analyzed by electrophoresis. *Amelogenin* bands specific for male-derived specimens were detected and a sex-linked dimorphism of *amelogenin* was indicated. Later, in 1992, Salido *et al.*¹³ reported that human *amelogenin* genes are expressed from both the X- and Y-chromosomes. A human genomic library was screened with a probe of mouse *amelogenin* cDNA, and positive clones from the X- and Y chromosomes were isolated. Both genes spanned more than 9 kb and 7 exons were identified in the sequences of both the X- and Y derived clones of *amelogenin* genes. Complete sequences of human *amelogenin* protein including signal peptide of 16 residues on both X- and Y-chromosomes were identified¹.

The mature human *amelogenin* from the X-chromosome skipping exon 4 is a protein of 19.8 kilo Daltons with 175 amino acid residues and which from Y is a 20 kDa protein with 176 amino acids. The methionine at residue 29 of the Y -chromosome *amelogenin* is absent from the X *amelogenin*, due to a 3-base deletion in the human X *amelogenin* gene. The overall sequence similarity between the X- and Y derived cDNAs is 91%. The protein coding regions (exons 2-6) are highly conserved, with a similarity index between 93% and 100%, whereas the 5' and 3' untranslated regions (exons 1 and 7) are much less conserved¹.

A major transcript was shown to be derived from the X-chromosome by the technique of Real Time (RT)-PCR. However, faint Y-derived transcripts which hybridized only with the Y-specific probe were detected. The result indicated that the *amelogenin* sequence on the Y-

chromosome is also transcribed, although at a level much lower (about 10% of the total) than that of the X-homolog¹.

Mapping of Human Amelogenin Gene Loci on the X- and Y-Chromosomes

Lau *et al.*⁴ mapped human *amelogenin* genes on the short arm of the X- chromosome in the p22.1-p22.3 region and in the pericentric region of the Y-chromosome. The *amelogenin* sequence on the human Y-chromosome was tentatively mapped onto the proximal long arm in the Yq11 region, on the basis of retention of the Y locus in a fibroblast cell line GM-1709 which was derived from a phenotypic female with an intact X-chromosome and an isochromosome for the long arm of the Y-chromosome⁴. However, a controversy arose over the localization of human *amelogenin* gene on the Y-chromosome. Nakahori *et al.*¹⁶ mapped the *amelogenin* sequence on the short arm at Y-chromosome using newly cloned Y-specific fragment probes. Bailey *et al.*¹⁷ also reported that the location of the Y *amelogenin* sequence was found in the short arm, Yp.

According to Salido *et al.*¹³ aberrant Y-chromosome in GM-1709 was redefined as dicentric with a breakpoint in the p11.2 region, thus raising the possibility that the Y locus may be on the short arm of the Y-chromosome. They mapped the Y locus by fluorescent *in situ* hybridization using a Y-genomic clone as a probe. Metaphase Y-chromosomes with hybridization signals were examined and the *amelogenin* locus was found clearly on the short arm of the Y-chromosome in the p11.2 region, and this is generally agreed upon^{13, 17}. Precise mapping of human *amelogenin* gene (AMG) in the p22 region on the X-chromosome¹⁸ and the homolog (AMGL) on the Y-chromosome¹⁹ were reported.

Phylogeny and Evolution of Amelogenin Gene

As described above, it is well established that there are two *amelogenin* genes, one on the X- and the other on the Y-chromosome in some mammals, with a single gene residing exclusively on the X-chromosome in other vertebrates. Male and female genomic DNA samples from a variety of animals were investigated using the "Noah's ark blot" by Nakahori *et al.*¹⁴ and also using the PCR method by Bailey *et al.*¹⁷.

It seems to be reasonable that the *amelogenin* genes reside on both the X- and Y-chromosomes in anthropoids (gorilla, chimpanzee and orangutan) as well as in man. Some old-world monkeys (Japanese monkey, rhesus monkey and crab-eating macaque) have *amelogenin* genes on the X- and Y-chromosomes, while other old-world monkeys (baboon, patas monkey, green monkey and talapoin) have no Y-homolog. A new-world monkey, tamarin, has only one *amelogenin* gene on the X-chromosome, while the capuchin monkey, another new-world variety, has both the X and Y genes. Only one *amelogenin* gene locus resides on the X-chromosome in the mouse and rat, but bovine genes are on the X and Y-chromosomes. The data of chromosomal distribution of *amelogenin* gene(s) cannot simply be attributed to single gene duplication or translocation via phylogenetic lineage. Multiple duplication or deletion is required to conform to the phylogenetic distribution of the species¹.

It is of particular interest that the *amelogenin* gene homologs are located on the autosomes but not on sex chromosomes in the primeval forms of mammals, monotreme (platypus) and marsupial (wallaby), according to the study of Watson *et al.*²⁰. These authors determined that *amelogenin* gene homologs were located on chromosomes 5q and 1q in wallaby, and on chromosomes 1 and 2 in platypus (duckbill) by Southern blot analysis. The adult duckbill has no true teeth, although the hatchling has an egg tooth and very young duckbills transiently possess teeth with typically mammalian characteristics. And this primitive egg-laying mammal has *amelogenin* gene-like loci on autosomes. However, the authors were not able to determine whether these homologous sequences are active *amelogenin* genes or pseudogenes. They surmised that the *amelogenin* gene region must have been translocated to X- and Y-chromosomes before the divergence of primates and other mammals¹.

Amelogenesis Imperfecta

Amelogenesis imperfecta is an inherited disease affecting the formation of tooth enamel in primary and permanent dentition. In general, there are two different phenotypes in clinical manifestations, the hypoplastic and the hypomineralization. The former is characterized by enamel that does not attain normal thickness during development, whereas in the latter the enamel is softer than normal. Genetically, amelogenesis imperfecta contains heterogeneous and autosomal dominant, autosomal recessive and X-linked forms which have been included in the different classification schemes with subdivision of these categories according to the perceived phenotype²¹. Affected (hemizygous) males and (heterozygous) females in the same

family have different clinical manifestations, with the teeth of heterozygous females exhibiting vertical markings of the enamel. Therefore, it was suggested that this type of amelogenesis imperfecta is caused by a structural alteration in the *amelogenin* gene¹.

The review is aimed at bringing out latest works and challenges experienced in using *amelogenin* gene primers in sex determination from human skeletal fragments and teeth specimens. This review is composed via Medline Internet search of literature and contributions from our experiences as well as experiences from colleagues.

The Application of Amelogenin Gene in Sex Determination from Human Skeletal Fragments and Teeth Specimens

Sex determination is not only a useful tool but of extreme necessity in forensic investigations specifically in cases of sexual assault, aged blood stains and human skeletal remains whereby rapid, sensitive, accurate and reliable methods of investigation are of the utmost importance²². The genetic difference between males and females is defined by the presence or absence of the Y-chromosome. The majority of the DNA of the sex chromosomes is specific to either the X or Y form. Closer inspection of *amelogenin* gene on each of the sex chromosomes identified regions of true homology, but also identified a number of sequence differences between the two forms¹⁴. Because of these homologous regions, PCR systems could now be designed to amplify both X- and Y-chromosomes in a single reaction by placing primers in homologous positions on the *amelogenin* gene that flank areas of difference between the X and Y forms. This dual amplification from a single set of primers provides an internal PCR control, meaning reaction failure will affect both X- and Y-chromosomes, removing the possibility of false-positive results that could be encountered by separate amplification strategies. Early systems designed to amplify regions of the *amelogenin* gene focused around an 189bp deletion present in the first intron of the X homolog of the gene¹⁴ and required large amounts of template DNA to be entered into the amplification reaction²³. The starting amount of template DNA was quickly reduced by optimization of the reaction leading to reported success rates from quantities equivalent to a single diploid cell²⁴.

The biggest breakthrough in DNA sex typing came with the publication of an alternative target sequence within *amelogenin* gene²⁵. This system was again designed to co-amplify X- and Y-chromosomes in a single reaction, but was based around a shorter, 6bp X-chromosome

deletion, allowing for amplification of shorter target sequences. Instead of long-length products (977bp and 788bp), amplified fragments were a mere 106bp and 112bp for the X- and Y-chromosomes respectively, and these targets could also be amplified from minute amounts of template DNA. This system was compatible with both low copy number and degraded DNA, and as such quickly became the system of choice within both forensic and archaeological disciplines. The short product lengths of these redesigned amplicons also had the advantage of falling within the size range of forensically important STR loci, and as such were perfectly suited to multiplexing within DNA profiling kits. Today, forensic DNA profiling is primarily carried out using commercially produced kits, designed to amplify up to 16 STR loci simultaneously. All kit primers for amplification of the *amelogenin* loci are designed around the 6bp deletion of intron 1 of this gene²⁶.

An additional bonus of this particular system comes with the detection method of choice. Because the *amelogenin* DNA fragments can be co-amplified along with autosomal STR loci within commercial DNA profiling kits, the fluorescent labeling and visualization method used for this process provides a degree of quantitative information along with fragment-length measurement. The quantitative data can then be used to determine the ratio of X- to Y-chromosomes present in the analyzed sample. Male/female DNA mixtures are often encountered during forensic investigation, especially in sexual assault investigations. The ability to calculate the relative ratio of male to female DNA present in a given mixture can be extremely useful for downstream analysis. This quantification will also detect chromosome duplication events of the sex chromosomes, for example, to detect the presence of an extra X-chromosome in XXY males. This system is not useful for correct sex determination in sex-reversal syndromes such as XY females and XX males where a mutational event or translocation results in the loss or gain of the *amelogenin* gene²⁷.

Since the paper by Hochmeister *et al.*²⁸ reporting the use of DNA extracted from human femoral bone of a corpse submerged 18 months under water and a 11-year-old mummified-corpse, using restriction fragment length polymorphisms (RFLP) and variable tandem repeats (VNTR) loci amplified by PCR, as well HLA DQ A1, other studies in the literature report using restriction fragment length polymorphisms (RFLP), short tandem repeat (STR) and *amelogenin* typing kits, PCR amplified sequences of the HLA DRB1 gene and PM Amplitype kits with bones using different methods of extraction²⁹.

From Genova in Italy, Berks in UK and Moscow in Russia, Manucci *et al.*³⁰ reported that gender identification of forensic samples was determined by amplifying a segment of the X-Y homologous gene *amelogenin*. Using a single pair of primers spanning part of the first intron, 106 and 112 by PCR products were generated from the X and Y homologous respectively, which were then resolved by agarose gel electrophoresis.

Faerman *et al.*³¹ reported from Hadassah, Jerusalem, on the establishment of a sensitive and reliable method of sex determination in skeletal remains based on amplification of single-copy *amelogenin*-encoding gene (*AMG*). The Y allele carries a small deletion in the first intron, facilitating the design of distinct X- and Y-specific polymerase chain reactions. Using this method, the sex of 18 individuals, including young children, out of 22 examined from periods ranging from 200 to around 8000 years ago. In no case was there a discrepancy between the DNA and the morphometric studies³¹.

A study by Alvaraz Garcia *et al.*³² used the homologous gene *amelogenin* to determine sex in dental pulp subjected to various environmental factors.

The applicability of the amplification of 9 STR loci and *amelogenin* was shown by typing bones and teeth, from few months to up to 3,000 years of age, using the Ampf/STR Profiler Plus. This method proved to be efficient in the analysis of ancient degraded DNA, in addition to the positive aspect of the utilization of minimal amounts of the sample³³.

From the Pennsylvania State University, Pennsylvania, and the University of Munich, Germany, Stone *et al.*³⁴ reported on the accurate genetic sex determination of 19 out of 20 ancient human skeletons dating to A. D. 1300. For sex determination using DNA, primers for PCR were designed to amplify a small fragment (112 bp) in exon 6 of the *amelogenin* gene³⁴.

In 1997, Cattaneo *et al.*³⁵ analyzed the genetic sex of 32 human skeletons using PCR amplification for HLA DPB1 (327 pb), *amelogenin* (106/112 pb), and mtDNA loci. The best amplification was that of the mtDNA, followed by HLA DPB1 that amplified at the same frequency of the *amelogenin* loci; and amplification was possible in 11 of a total of 32 skeletons³⁵.

Evison *et al.*³⁶ analyzed human bone samples from 1986 to 1994 in forensic cases or exhumed material that had been buried from 1904 to 1984, as well as from human teeth and blood stains that were 3 months to 91 years old. The amplification and analysis of *amelogenin*, HLA-

DPB1 gene and mtDNA sequences were performed. There was no correlation between the ages of specimen and the degree of DNA preservation.

Faerman *et al.*³⁷ from the Hebrew University, Hadassah, Jerusalem, Israel, determined the sex of infanticide victims from the late Roman era through ancient DNA analysis. They applied a highly sensitive method based on PCR amplification of the X and Y *amelogenin* alleles. The reaction yielded distinguishable X- and Y-chromosome products by the simultaneous use of three primers (M4, M5 and M6). Amplification was successful for 19 out of the 43 ancient specimen tested. Fourteen specimens were found to be males and five females giving a significantly higher frequency of boys than girls. There were no inconsistencies or conflicting data for any of the specimen.

To determine the effect of environmental factors on the preservation of DNA, 38 archaeological teeth from 18 individuals of approximately similar age (900 to 2000 years BC), but greatly differing in terms of site milieu, were analyzed for aDNA content. The STR loci HUMvWA31A, HUMTH01, HUMFES/FPS and *amelogenin* were analyzed. Differences in quantity and quality of the results are attributed to the respective prevailing environmental factors or to respective storage conditions³⁸.

From the United States of America, Seo *et al.*³⁹ reported that they analyzed the maternity of two newborns who were murdered and abandoned >5 and 10 years, by amplification and direct sequencing of mitochondrial DNA control regions. Sexes of the remains were determined to be female and male by amplifying a segment of the X-Y homologous gene, *amelogenin*.

Hummel *et al.*⁴⁰ conducted a study on 33 human skeletons to evaluate the morphological sex determination based on traits of the ossa coxae and the cranium. The evaluation criterion was genetic analysis of the *amelogenin* gene, which represents part of X-and Y-chromosome. They found that 88% of the morphological determination matched the genetic sex⁴⁰.

Mays and Faerman⁴¹ reported from English Heritage Centre for Archaeology, Fort Cumberland, Eastney, Portsmouth, UK and Laboratory of Biological Anthropology and Ancient DNA, Hadassah School of Dental Medicine, Hebrew University, Jerusalem Israel, on sex identification in some putative infanticide victims from Roman Britain using ancient DNA. Sex identification was performed using a method based on PCR amplification of the X and Y *amelogenin* alleles. The reaction yielded distinguishable PCR products of the X and Y *amelogenin* alleles by the

simultaneous use of three primers. The primers M4, M5 and M6 amplify PCR products of 329-bp (X allele) and 235-bp (Y allele). Altogether nine specimens were identified as male, four as female. Again, the sex of 11 infants could be identified: nine males and two females. Importantly, only the Y *amelogenin* allele could be amplified in all nine male samples, thus reflecting a relative poor state of DNA preservation in the bone specimen⁴¹.

Matheson and Loy⁴² reported from the University of Queensland, Brisbane, Australia, on the use of *amelogenin* gene in the genetic sex identification of 9400-year-old ten human skull samples from Çayönü Tepesi, Turkey. The X chromosome amplification with the M4 and M5 primers produced a 331bp fragment while the Y chromosome amplified with M4 and M6 to produce a 237bp product. However, the *amelogenin* X-specific detection system only amplified three of the samples. Generally, the *amelogenin* amplifications did not amplify efficiently and reliably, this could be due to degree of fragmentation of the DNA in these same samples⁴².

From the University of Göttingen, Germany, Diane *et al.*⁴³ introduced a polymerase chain reaction-based multiplex approach to improve the certainty of molecular sex identification on archaeological skeletal material. They coamplified *amelogenin*, two X-chromosomal short tandem repeats (STRs) (DXS6789 and DXS9898) and two Y-chromosome specific STRs (DYS391 and DYS392). The amplification results of this multiplex approach back each other up, and enable reliable sex identification.

From the Masaryk University, Czech Republic, Vaňharová and Drozdová⁴⁴ reported on the sex determination of 53 skeletal remains of 4000 year old children and juveniles from Hoštice 1 za Hanou by ancient DNA analysis. They amplified the SRY locus and the *amelogenin*. Concordance between the genetic and archeological determinations occurred in 13 cases out of 21 (~62%)⁴⁴.

From Johannesburg, South Africa, Gibbon *et al.*⁴⁵ designed two novel methods of molecular sex identification from 30 skeletal tissues, using *amelogenin* gene. The classification of sex using the first system was determined by analysing the sequence for 10 sex-specific single nucleotide polymorphisms (SNPs) to determine whether both the X and Y (male) or only the X (female) was represented. System two spans an indel (6 bp) in addition to 10 sex-specific SNPs⁴⁵. Using both systems of sex identification 14 (46.66%) specimens produced results, and of these 12 (85.71%) were males and 2 (14.28%) were females. Based on this data, 4 (28.57%) of the specimens in the sample possess the same result using both methods.

Method one produced 2 (14.28%) specimens and 8 (57.14%) specimens were achieved using system two⁴⁵.

From the Aristotle University of Thessaloniki, Greece, Kovatsi *et al.*⁴⁶ reported on the successful genetic sex identification from 10 human teeth using *amelogenin* gene locus, through DNA repair. The amplification of a single 106 bp segment of the *amelogenin* gene corresponding to the X chromosome indicates the female origin of those samples. The double PCR bands (106 bp and 112 bp), corresponding to the X and Y chromosomes, indicate the male origin of the samples⁴⁶. The *amelogenin* gene was amplified by PCR in 8 out of 10 DNA extracts (amplification success rate 80%) following DNA repair, while prior to this repair treatment no DNA extract yielded PCR products for this gene. Four out of the ten teeth examined were found to belong to females, another four were found to belong to males and two could not be typed⁴⁶.

From Malay Peninsular, Shahrul Hisham and his colleagues⁴⁷ reported from the University of Kebangsaan Malaysia, Bangi, Selangor and Kuala Lumpur, Malaysia, on the molecular gender determination of 2 ancient human bones, using *amelogenin* gene. They used a cocktail of three primers, consisting of the common forward primer and both the human X-specific reverse primer (*AMELX*) and human Y-specific (*AMELY*) reverse primer. In male samples, the *amelogenin* gene was located on both the X and Y chromosomes and, thus, two bands of PCR products were produced, i.e., 320 bp (*AMELX*) and 235 bp (*AMELY*). On the other hand, as female samples have two X chromosomes, PCR generated only a single band of 235 bp, but in higher concentration than the X chromosome band from males⁴⁷.

A study from Czech Republic by Eliášová *et al.*⁴⁸ reported that comparison of genetic methods for sex determination, amplification of *amelogenin* XY and short tandem repeats (STR) analysis, showed 100% accordance in all 11 cases of Knezeves bone samples.

In a recent forensic study for genetic sex determination of burnt powdered skeletal fragments from Sokoto, Northwestern Nigeria, by Zagga *et al.*⁴⁹ a success rate of 33.3% was recorded for correct genetic sex identification using *amelogenin* gene.

In a research study on the polymerase chain reaction (PCR)-based sex determination using unembalmed human cadaveric skeletal fragments from Sokoto, Northwestern Nigeria, Zagga *et al.*⁵⁰ reported that with *amelogenin* gene, genetic sex identification was achieved in four only out of twelve samples.

In a comparative analysis on the efficiency of *amelogenin* gene and *alphoid* repeats in sex determination of embalmed cadaveric skeletal fragments from Sokoto, Nigeria, Zagga *et al.*⁵¹ recorded a success rate of 43% was using *amelogenin* gene.

The Reliability of Amelogenin Gene in Sex Determination of Human Skeletal Fragments and Teeth Specimens

The phenotypic nature of the information provided by investigation of the *amelogenin* locus could potentially present massive problems if an error occurs that is not identified⁵². The failure of a forensic DNA profiling technique to correctly resolve an issue as simple as sex could seriously misdirect the police investigation, as well as cast doubts on the techniques' reliability for criminal prosecution²⁶.

The reliability of *amelogenin*-based sex testing on hard tissues was first questioned in 1996, by Stone and his colleagues, in a study of ancient human skeletal remains of twenty individuals from Germany, where one case (5%) which was classified as a male based on morphology, but PCR products hybridized only with the X oligonucleotide. This may be the result of insufficient DNA, or mutations in the priming or oligonucleotide binding sites on the Y chromosome copy of the *amelogenin* gene³⁴.

In a study on sex determination of skeletal remains of 4000 year old children from the Czech Republic, Vanharova and Drozdova⁴⁴ reported seven cases in which the results were divergent between archaeological and genetic analyses. About 67% failure rate of genetic sex identification using *amelogenin* gene was reported by Zagga *et al.*⁴⁹ in a recent forensic study for genetic sex determination of burnt powdered skeletal fragments from Sokoto, Northwestern Nigeria. A similar failure rate (67%) of *amelogenin* gene sex identification was reported by Zagga *et al.*⁵⁰ in a different research study on the polymerase chain reaction (PCR)-based sex determination using unembalmed human cadaveric skeletal fragments from Sokoto, Northwestern Nigeria. Zagga *et al.*⁵¹ reported in a comparative analysis on the efficiency of *amelogenin* gene and *alphoid* repeats in sex determination of embalmed cadaveric skeletal fragments from Sokoto, Nigeria, that 57% of the samples was wrongly identified as males instead of females with *amelogenin* gene.

Recently, in a single blind study for genetic sex identification using *amelogenin* gene multiplex primers on dry human teeth specimens grouped into embalmed, unembalmed and

deciduous from Sokoto, Northwestern Nigeria, Zagga *et al.*⁵³ reported that no genetic sex identification was achieved in any of the dry human teeth samples.

Conclusion

Amelogenin PCR method/system of sex determination should not, at the moment, completely replace traditional methods of sex identification. Hence, sex identification with *amelogenin* gene, of subjects for forensic purposes should be achieved as much as possible through a multiple morphological-molecular combined approach.

References

1. Sasaki S, Shimokawa H. The amelogenin gene, *Int J Dev Biol.* 1995;39:127-33.
2. Eastoe JE. The chemical composition of bone and teeth. *Adv Fluorine Res Dent Caries Prevent.* 1965;3: 5-17.
3. Termine JD, Belcourt AB, Christner PJ, Conn KM, Nylen MU. Properties of dissociatively extracted fetal tooth matrix proteins 1. Principal molecular species in developing bovine enamel. *J Biol Chem.* 1980;255:9760-68.
4. Lau EC, Mohandas TK, Shapiro LJ, Slavkin HC, Snead ML. Human and mouse amelogenin gene loci are on the sex chromosomes. *Genomics.* 1989;4:162-8.
5. Shimokawa H, Tamura H, Ibaraki K., Sasaki S. Human amelogenin gene. In *Tooth Enamel V* (Ed, RW Fearnhead):301-5. Yokohama, Florence Pub, 1989.
6. Sire J-Y, Delgado S, Fromentin D, Grindot M. Amelogenin: lessons from evolution. *Arch of Oral Biol.* 2005;50:205-12.
7. Yuan ZA, Chen E, Gibson CW. Model system for evaluation of alternative splicing: exon skipping. *DNA Cell Biol.* 2001;20:807-13.
8. Fincham AG, Bessem CC, Lau EC, Pavlova Z, Shuller C, Slavkin HC et al. Human developing enamel proteins exhibit a sex-linked dimorphism. *Calcif Tissue Int* 1991;48:288-90.
9. Toyosawa S, O'HUigin C, Figueroa F, Tichy H, Klein J. Identification and characterization of amelogenin genes in monotremes, reptiles, and amphibians. *Proc Natl Acad Sci USA.* 1998;95:13056-61.
10. Ravindranath RMH, Basilrose RM, Ravindranath NH, Vaitheesvaran B. Amelogenin interacts with cytokeratin-5 in ameloblasts during enamel growth. *J Biol Chem.* 2003;278:20293-302.
11. Snead M. Amelogenin protein exhibits a modular design: Implications for form and function. *Connect Tissue Res.* 2003;44:47-51.

12. Moradian-Oldak J, Simmer JP, Sarte PE, Zeichner-David M, Fincham AG. Specific cleavage of a recombinant murine amelogenin at the carboxy-terminal region by a proteinase fraction isolated from developing bovine tooth enamel. *Arch Oral Biol.* 1994;39:647–56.
13. Salido EC, Yen PH, Koprivnikar K, Yu LC, Shapiro LJ. The human enamel protein gene amelogenin is expressed from both the X and the Y chromosomes. *Am J Hum Genet.* 1992;50:303–16.
14. Nakahori Y, Takenaka O, Nakagome Y. A human X-Y homologous region encodes amelogenin. *Genomics.* 1991;9:264-9.
15. Fincham AG, Hu Y, Lau EC, Slavkin HC, Snead ML. Amelogenin post-secretory processing during biomineralization in the postnatal mouse molar tooth. *Arch Oral Biol.* 1991;36:305–17.
16. Nakahori Y, Tamura S, Nagafuchi S, Fujieda K, Minowada S, Fukutani K et al. Molecular cloning and mapping of 10 new probes on the human Y chromosome. *Genomics.* 1991; 9: 765-9.
17. Bailey DM, Affara NA, Ferguson-Smith MA. The X-Y homologous gene amelogenin maps to the short arms of both the X and Y chromosomes and is highly conserved in primates. *Genomics.* 1992;14:203-5.
18. Schaefer L, Ferrero GB, Grillo A, Bassi MT, Roth EJ, Wapenaar MC et al. A high resolution deletion map of human chromosome Xp22. *Nature Genet.* 1993;4:272-9.
19. Foote S, Vollrath D, Hilton A, Page DC. The human Y chromosome: overlapping DNA clones spanning the euchromatic region. *Science.* 1992;258:60-6.
20. Watson JM, Spencer JA, Graves JA, Snead ML, Lau EC. Autosomal localization of the amelogenin gene in monotremes and marsupials: implications for mammalian sex chromosome evolution. *Genomics.* 1992;14:785-9.
21. Witkop CJ Jr. Amelogenesis imperfecta, dentinogenesis imperfect and dentin dysplasia revisited: problems in classification. *J Oral Pathol.* 1989;17:547-53.
22. Butler JM. Chromosomal duplications along the Y-chromosome and their potential impact on Y-STR interpretation. *J Forensic Sci.* 2005;50:853-9.
23. Akane A, Shiono H, Matsubara K, Nakahori Y, Seki S, Nagafuchi S et al. Sex identification of forensic specimen by polymerase chain reaction (PCR): two alternative methods. *Forensic Sci Int.* 1991;49:81-8.
24. Akane A, Seki S, Shiono H, Nakamura H, Hasegawa M, Kagawa M et al. Sex determination of forensic samples by dual PCR amplification of an X-Y homologous gene. *Forensic Sci Int.* 1992;52:143-8.
25. Sullivan KM, Mannucci A, Kimpton CP, Gill P. A rapid and quantitative DNA sex test: fluorescence-based PCR analysis of X–Y homologous gene amelogenin. *Biotechniques.* 1993;15:636–8.
26. Graham E A M. Sex determination. *Forensic Sci Med Pathol.* 2006;2:283-6.
27. Mohammed F, Tayel SM. Sex identification of normal persons and sex reverse cases from bloodstains using FISH and PCR. *J Clin Forensic Med.* 2005;12:122–7.

28. Hochmeister MN, Budowle B, Borer UV, Rudin O, Bohnert M, Dirnhofer, R. Confirmation of the human skeletal remains using multiplex PCR amplification and typing kits. *J Forensic Sci.* 1994;40:701-5.
29. Alonso A, Andenovic S, Martin P, Sutlovic D, Erceg I, Huffine E. DNA typing from skeletal remains evaluation of multiplex and megaplex STR systems on DNA isolated from bone and teeth samples. *Croat Med J.* 2001;42:260-6.
30. Manucci A, Sullivan KM, Ivanov PL, Gill P. Forensic application of a rapid and quantitative DNA sex test by amplification of the X – Y homologous gene amelogenin. *Int J Legal Med.* 1994;106:190-3.
31. Faerman M, Filon D, Kahila G, Greenblatt CL, Smith P, Oppenheim A. Sex identification of archaeological human remains based on amplification of the X and Y amelogenin alleles. *Gene.* 1995;167:327-32.
32. Alvarez Garcia A, Munoz I, Pestoni C, Lareu MV, Rodriguez-Calvo MS, Carracedo AA. Effect of environmental factors on PCR-DNA analysis of dental pulp. *Int J Legal Med.* 1996;109:125-9.
33. Handt O, Krings M, Ward RH, Paabo S. The retrieval of ancient human DNA sequences. *Am J Hum Genet.* 1996;59:368-76.
34. Stone AC, Milner GR, Paabo S, Stoneking M. Sex Determination of ancient human skeletons using DNA. *Am J of Phys Anthropol.* 1996;99:231- 8.
35. Cattaneo C, Craig OE, James NT, Sokol RJ. Comparison of three DNA extraction methods on bone and blood stains up to 43 years old and amplification of three different gene sequences. *J Forensic Sci* 1997;42:1126-35.
36. Evison MP, Smillie DM, Chamberlain AT. Extraction of single-copy nuclear DNA from forensic specimen with a variety of post-mortem histories. *J Forensic Sci.* 1997;42:1032-8.
37. Faerman M, Bar-Gal KG, Filon D, Greenblatt CL, Stager L, Oppenheim A et al. Determining the sex of infanticide victims from the late roman era through ancient DNA analysis. *J Archaeol Sci.* 1998;25:861-5.
38. Burger J, Hummel S, Herrmann B, Henke W. DNA preservation: a microsatellite-DNA study on ancient skeletal remains. *Electrophoresis.* 1999;20:1722-8.
39. Seo Y, Uchiyama T, Shimizu K, Takahama K. Identification of remains by sequencing of mitochondrial DNA control region. *Am JForensic Med Pathol.* 2000;21:138-43.
40. Hummel S, Bramanti B, Finke T, Hermann B. Morphological sex determination based on traits of the ossa coxae and the cranium. *Anthropologischer Anzeiger; Bericht uber die biologisch-anthropologische Literatur.* 2000;58:9-13.
41. Mays S, Faerman M. Sex Identification in some putative infanticide victims from roman britain using ancient DNA. *J Archaeol Sci.* 2001;28:555-9.

42. Matheson DE, Loy TH. Genetic sex identification of 9400-old human skull samples from Cayönü Tepesi, Turkey. *J Archaeol Sci.* 2001;28:569-75.
43. Diane S, Susanne H, Bernd H. Multiplex X/Y-PCR improves sex identification in a DNA analysis. *Am J Phys Anthropol.* 2003;121:337-41.
44. Vanharova M, Drozdova E. Sex determination of skeletal remains of 4000 year old children and juveniles from Hostice 1 za Hanou (Czech Republic) by ancient DNA analysis. *Anthropol Rev.* 2008;71:63-70.
45. Gibbon V, Paximadis M, Štrkalj G, Ruff P, Penny C. Novel methods of molecular sex identification from skeletal tissue using the amelogenin gene. *Forensic Sci Int.* 2009;3:74-9.
46. Kovatsi L, Nikou D, Triantaphyllou S, Njau SN, Voutsaki S, Koidou S. DNA repair enables sex identification in genetic material from human teeth. *Hippokratia* 2009;13:165-8.
47. Shahrul Hisham ZA, Sahidan S, Rohaya MAW, Siti Ateefah MY, Intan Zarina ZA, Nor Hidayah JAN, et al. Molecular gender determination of ancient human from Malay Peninsular. *Am J Appl Sci.* 2009;6:1770-75.
48. Eliášová I, Mazura I, Smejtek L. DNA analysis of ancient skeletal remains. *Fol Biol (Praha).* 2010;56:47-50.
49. Zagga AD, Ismail SM, Ahmed H, OON, Tadros AA. Forensic study for genetic sex determination of burnt powdered skeletal fragments from Sokoto, Northwestern Nigeria. *IOSR Journal of Dental and Medical Sciences.* 2013;7:47-54.
50. Zagga AD, Tadros AA, Ismail SM, Ahmed H OON. Polymerase chain reaction (PCR)-based sex determination using unembalmed human cadaveric skeletal fragments from Sokoto, Northwestern Nigeria, *IOSR Journal of Pharmacy and Biological Sciences.* 2013;7: 01-08.
51. Zagga AD, Tadros AA, Ismail SM, Ahmed H OON. Comparative analysis on the efficiency of *Amelogenin* gene and aliphoid repeats in sex determination of embalmed cadaveric skeletal fragments from Sokoto, Northwestern Nigeria. *Int J Biomed Health Sci.* 2013;9:127-38.
52. Sinclair AH, Berta P, Palmer MS, Hawkins JR, Griffiths BL, Smith MJ et al. A gene from the human sex-determining region encodes a protein with homology to a conserved DNA-binding motif. *Nature.* 1990;346:240-4.
53. Zagga AD, Ahmed H OON, Tadros AA, Ismail SM. *Amelogenin* gene failure in sex determination of dry human teeth specimens from Sokoto, Northwestern Nigeria. *IOSR Journal of Pharmacology.* 2013;3:32-40.

Correspondence Address / Yazışma Adresi

Abdullahi D. Zagga,
Department of Anatomy,
College of Health Sciences,
Usmanu Danfodiyo University,
Sokoto, Sokoto State, Nigeria.
E-mail: adauduzagga@gmail.com