




Eurasian Journal of Soil Science

Journal homepage : <http://ejss.fesss.org>



Soil properties and growth of yellow bell pepper (*Capsicum annum*) as influenced by compost and arbuscular mycorrhizal fungi

Tolulope Yetunde Akande ^{a,b,*}, Kehinde Olajide Erinle ^{b,c}, Tope Daniel Bitire ^{b,d}

^a College of Resources and Environment, Northeast Agricultural University, Harbin Heilongjiang Province, China

^b Agronomy Department, University of Ibadan Nigeria

^c School of Agriculture, Food and Wine, University of Adelaide, Adelaide SA 5005, Australia

^d Department of Microbiology, Northwest University, Mafikeng Campus, South Africa

Abstract

Compost is an inexpensive agricultural waste which improves soil health and quality. The experiment was carried out to assess the influence of compost and mycorrhizal inoculation (*Glomus mosseae*) on soil properties and growth of yellow bell pepper in pots under screen house conditions, in a completely randomized design with three replicates. The treatments included mycorrhizal inoculation only (C0M1), compost at 20 t ha⁻¹ only (C1M0), compost at 30 t ha⁻¹ only (C2M0), compost and mycorrhizal inoculation at 20 t ha⁻¹ (C1M1), compost and mycorrhizal inoculation at 30 t ha⁻¹ (C2M1) and control (no amendment / uninoculated). Compost and mycorrhizal inoculation (C1M1 and C2M1) significantly improved soil N, P and K compared to control. Inoculation with mycorrhizal only (C0M1) increased uptake of N, P, K, Ca and Mg compared to uninoculated. Co-utilization of compost and mycorrhizal inoculation significantly increased root and shoot dry biomass compared to uninoculated. The highest fruit yield was obtained at C2M1 followed by C1M1 in comparison to compost application only. Treatment C2M1 recorded the highest prevalence of percent root colonization. This suggests that compost and *Glomus mosseae* could be considered to have a sustainable potential for better growth and yield performance in the production of yellow bell pepper in an Alfisol.

Keywords: Arbuscular mycorrhizal fungi, Compost, Nutrient uptake, Soil fertility, Soil nutrient

© 2023 Federation of Eurasian Soil Science Societies. All rights reserved

Article Info

Received : 07.12.2021

Accepted : 06.12.2022

Available online : 15.12.2022

Author(s)

T.Y.Akande

K.O.Erinle

T.D.Bitire



* Corresponding author

Introduction

Many agricultural lands require an intervention strategy to improve the nutrients contents of the soil. The organically managed soil systems in the tropics are increasingly utilized to combat constraint to food security (Chukwuka and Omotayo, 2009). Major factors that constrain agricultural soil fertility are caused by excessive erosion, nutrient run-off, loss of organic matter and low soil biodiversity (Santos et al., 2011). Compost, an effective organic fertilizer produced from organic waste biomasses (Patchaye et al., 2018). The use of organic biomass as soil conditioner is now rapidly gaining importance in tropical agriculture as it promotes soil physical, chemical and biological properties (Zong et al., 2010; Gautam and Pathak, 2014). Furthermore, evidence suggest that application of compost to soil increase soil organic matter, promotes soil fertility and activities of soil microbial communities (González et al., 2010; Carlson et al., 2015; Cozzolino et al., 2016). The application of compost resulted to increased soil available macro and micro nutrients that stimulates microorganism development in soils (Biala, 2000); moreover, compost significantly increases crop growth and activities of soil microorganisms (Gosling et al., 2006; Weber et al., 2007). Compost is a potential habitat for arbuscular mycorrhizal fungi (AMF) functionality (Warnock et al., 2010). Arbuscular mycorrhizal fungus is important in sustainable agriculture since mycorrhizal association hastens nutrient acquisition for crop growth and development (Hu et al., 2009; Smith et al., 2015). According to Hodge and Fitter (2010), the co-

amendment of compost and mycorrhizal inoculation increased crop growth better than either sole amendment due to arbuscular mycorrhizal fungi ability to exploits nutrients released by mineralization of organic matter. One of the benefits derived by host of arbuscular mycorrhizal fungi is that it enhances the absorption of soil mineral nutrients, which includes the major and micro nutrients (Linderman, 1992; Perner et al., 2007). The availability of these nutrients is usually affected by soil property, climate and crop type (Cely et al., 2016). Phosphorus has been described in literature as poorly accessible plant nutrient because of its immobility in soils (Smith and Read, 2008; Neumann and George, 2010). Evidence by research has confirmed AM fungi provides multifunctional benefits to soil by improving aggregate stability, creating more resistance to water stress and diseases which mostly depends on origin and species compatibility (Jung et al., 2012; Verbruggen et al., 2013). Horticultural crop such as yellow bell pepper is one of the major components of traditional food in Nigeria. It is known to be rich in vitamins and minerals (Malik et al., 2011) and play significant roles in disease prevention including diabetes, fever, heart and cancerous diseases (Serpeloni et al., 2015; Shahidi and Ambigaipalan, 2015). They are commercially produced due to its high demand and value which is mostly nutritive and medicinal purpose (Tanwar et al., 2021). However, yield and fruit quality are affected by a number of environmental factors such as type of soil, chemical fertilizer, pests and diseases (López et al., 2014). There is need to meet rising demands by adopting sustainable practices to assist in improving the soil, increasing crop growth and enhance mineral nutrient uptake. Several authors reported positive outcome of utilizing arbuscular mycorrhizal fungi on horticultural crops most especially *Glomus species* (Tanwar et al., 2011; Krüger et al., 2012). *Glomus mosseae* is known to provide specific benefits with horticultural crops by improving fruit yield and quality (Franco et al., 2013). It can facilitate uptake of mineral nutrients, promote growth and develop symbiotic association in rhizospheric region of pepper plants (Nadeem et al., 2014; Rouphael et al., 2015; Smith et al., 2015). Therefore, the objective of this study is to evaluate the effect of compost and *Glomus mosseae* inoculation on improving soil fertility, growth and yield of yellow bell pepper.

Material and Methods

Experimental site and design

The experiment was performed at the screen house of the Department of Agronomy University of Ibadan Nigeria (7°24'N; 3°48'E) between June and August, 2020. Soil samples were collected at 0-20 cm depth at Teaching and Research farm and transported in plastic bags to the screen house. A completely randomized design with three replicates involving yellow bell pepper (*Capsicum annum* L) at two seedlings per pot, and two treatment factors: compost and mycorrhizal inoculation. Three levels of compost applied in this study: 0, 20 and 30 t ha⁻¹ (equivalent to 0, 50 and 75 g compost per pot); two rates of mycorrhizal inoculation at 0 and 20 gram per pot and each treatment with 5 kg soil per pot.

Soil and plant analysis

Soil pH in water measured using pH meter. The particle size analysis was carried out by Gee and Or (2002) indicating loamy sand texture and classified as an Alfisol according to US soil taxonomy, total nitrogen by Kjeldahl method (Bremner, 1996) and organic carbon was determined by Walkley-Black method (Nelson and Sommers, 1996). Available phosphorus was by Bray extraction method (Frank et al., 1998). The Exchangeable bases K⁺, Ca²⁺ and Mg²⁺ was extracted using 1N ammonium acetate, where K⁺ was determined by flame photometer and Ca²⁺ and Mg²⁺ were determined by atomic absorption spectrophotometer (Okalebo et al., 2002). Plant samples were oven-dried (70°C) and milled. Then, digested milled samples were used to determine N content by Kjeldahl method (Bremner, 1996). The samples were dry ashed in a furnace (500°C for 6 h) and extracted using nitric-perchloric-sulfuric acid mixture which was used to detect phosphorus (P), potassium (K), calcium (Ca) and magnesium (Mg) according to Tel and Hagarty (1984). Afterwards, P was finally determined by vanadomolybdate colorimetry method, K by flame photometer while a method of titration (EDTA) was used to determine Ca and Mg (AOAC, 2005).

Plant, compost and mycorrhizal treatment

The pepper seedlings used in this study was yellow bell pepper (*Capsicum annum* L.) collected from Institute of Agriculture and Research Training Ibadan Nigeria. Animal waste compost was collected from Organic farm of the Federal University of Agriculture Abeokuta Nigeria. Arbuscular mycorrhizal fungi (*Glomus mosseae*) inoculum was originally and commercially obtained from Biocult Pty Limited, South Africa and collected from soil microbiology laboratory of Department of Agronomy, University of Ibadan Nigeria. It was multiplied in pot cultures using maize as host plant. The inoculum was applied to the third central part of the soil a day before transplanting yellow bell pepper seedlings (Ubah et al., 2012). Percent root colonization was done by observing stained roots under a dissected microscope (Giovannetti and Mosse, 1980).

Table 1. Soil physical and chemical characteristics and nutrient composition of compost

Parameters	Units	Soil	Units	Compost
pH(H ₂ O)		7.40	pH	8.03
TN	g kg ⁻¹	0.42	TN	%
AP	mg kg ⁻¹	7.23	AP	%
OC	g kg ⁻¹	6.20	OC	%
K ⁺	cmol kg ⁻¹	2.00	K	%
Ca ²⁺	cmol kg ⁻¹	7.80	Ca	%
Mg ²⁺	cmol kg ⁻¹	2.80	Mg	%
Particle Size Distribution				
Sand	%	73.80		
Silt	%	26.10		
Clay	%	2.00		
Textural Class		Loamy Sand		

TN = Total nitrogen, AP = Available phosphorus, OC = Organic carbon

Statistical analysis

The results of this experiment were obtained by one-way ANOVA using SPSS v 19.0. Differences were separated by Duncan Multiple Range Test (DMRT) at significance level of 0.05. All data presented are means of three replicates.

Results

Effect of compost and AMF on nutrient concentration

The effect of compost with or without AMF inoculation on the improvement of soil nutrients is shown in Table 2. The AMF inoculation only (C0M1) led to 29.1 % increase in total nitrogen (TN) concentration compare to control. Compost and AMF inoculation (C1M1) at 20 t ha⁻¹ significantly increase (3.49 %) compared to C1M0 (compost application only) and control. Similarly, compost and AMF inoculation at 30 t ha⁻¹ (C2M1) was higher by 3.92 % compare to compost application only (C2M0) and significantly higher than control (C0M0). Soil Ca content increased with or without AMF inoculation. Treatment with AMF inoculation only (C0M1) significantly increase compare to control. Both levels of compost and AMF inoculation (C1M1 and C2M1) were higher by 4.28 % and 3.57 % when compared to C1M0 and C2M0 respectively. Mg concentration showed AMF inoculation only (C0M1) increase by 6.95 % compare to control. C1M1 treatment at 20 t ha⁻¹ was higher by 4.83 % when compared to compost application only (C1M0). Compost and AMF inoculation (C2M1) significantly increase compare to C2M0 at 30 t ha⁻¹ and control. Application of compost with or without AMF inoculation influenced changes in K concentration. C0M1 increased by 7.63 % compare to control. Treatment C1M1 at 20 t ha⁻¹ led to significant increase compare to compost application only (C1M0). The application of compost in combination with AMF inoculation (C2M1) showed no significant difference compare to C2M0. AMF inoculation only (C0M1) showed no significant effect compare to control in soil organic carbon (Figure 1). Compost and AMF inoculation at 20 t ha⁻¹ was significantly higher by 5.23 % compared to the application of compost only (C1M0). Similarly, C2M1 significantly increase compare to compost application only and control. Available P showed significant differences (Figure 2). AMF inoculation only (C0M1) was higher by 28.67 % compared to control. Compost and AMF inoculation at 20 t ha⁻¹ significantly increase compare to C1M0 and control. Similarly, C2M1 treatment at 30 t ha⁻¹ significantly increase compare C2M0 and control.

Table 2. Nutrient concentration in soil as affected by compost and AMF inoculation

	TN (g.kg ⁻¹)	Ca (cmol.kg ⁻¹)	Mg (cmol.kg ⁻¹)	K (cmol.kg ⁻¹)
C0M0	0.95±0.06d	1.13±0.01d	1.07±0.01d	1.09±0.01e
C0M1	1.34±0.04c	1.25±0.02c	1.15±0.01d	1.18±0.02d
C1M0	1.38±0.02bc	1.34±0.02c	1.38±0.02c	1.24±0.02c
C1M1	1.43±0.01abc	1.40±0.03b	1.45±0.03c	1.31±0.01b
C2M0	1.47±0.02ab	1.62±0.03a	1.62±0.04b	1.38±0.01a
C2M1	1.53±0.01a	1.68±0.02a	1.94±0.03a	1.40±0.02a

Mean ± SE of three replicates are shown; different letters on numerical values indicate significant differences ($P < 0.05$) along the columns. C0M0=control, C0M1= mycorrhizal inoculation at 20g only; C1M0= compost at 20tha⁻¹ only; C2M0=compost at 30tha⁻¹only; C1M1=compost and mycorrhizal inoculation at 20tha⁻¹; C2M1=compost and mycorrhizal inoculation at 30tha⁻¹.




Figure 1. Organic carbon as affected by compost and AMF inoculation. Bars represent the mean \pm SE of three replicates ($P < 0.05$). COM0=control, COM1= mycorrhizal inoculation at 20g only; C1M0= compost at 20 tha⁻¹ only; C2M0=compost at 30 tha⁻¹only; C1M1=compost and mycorrhizal inoculation at 20 tha⁻¹; C2M1=compost and mycorrhizal inoculation at 30 tha⁻¹.




Figure 2. Available P as affected by compost and AMF inoculation. Bars represent the mean \pm SE of three replicates ($P < 0.05$). COM0=control, COM1= mycorrhizal inoculation at 20g only; C1M0= compost at 20 tha⁻¹ only; C2M0=compost at 30tha⁻¹only; C1M1=compost and mycorrhizal inoculation at 20 tha⁻¹; C2M1=compost and mycorrhizal inoculation at 30 tha⁻¹.

Effect of compost and AMF on nutrient uptake

Table 3 presents trends of compost application with or without AMF inoculation in uptake of nutrients. AMF inoculation only (COM1) significantly increased N uptake compared to control. Compost and AMF inoculation (C1M1) at 20 t ha⁻¹ significantly increase compare to compost application only (C1M0) and control. Treatment C2M1 and C2M0 (30 t ha⁻¹) showed no significant difference but both significantly higher than control (COM0) for N uptake. Treatment application significantly differs in P uptake as shown in Table 3. AMF inoculation (COM1) significantly increase compared to control. The application of compost and AMF inoculation at 20 t ha⁻¹ significantly increase compared to C1M0. Similarly, C2M1 was significantly higher compared to compost application only at 30 t ha⁻¹. Ca uptake showed AMF inoculation only (COM1) significantly increase compare to control. Treatment C1M1 was higher by 11.1 % compare to compost application only (C1M0). However, C2M1 at 30 t ha⁻¹ significantly increase compare to compost application only (C2M0) and control. Mg uptake indicated compost application with or without AMF inoculation at both levels was significantly higher than control except AMF inoculation only (COM1). The treatment C1M1 and C2M1 were significantly higher by 14.8 % and 7.46 % compared to C1M0 and C2M0 respectively. K uptake showed AMF inoculation only (COM1) was 13.3 % higher compared to control. C1M1 was significantly higher compare to compost application only at 20 t ha⁻¹ (C1M0). Compost in combination with AMF inoculation at 30 t ha⁻¹ was higher by 5.8 % compare to C2M0.

Table 3. Nutrient uptake as affected by compost and AMF inoculation

	N (gkg ⁻¹)	P (mg kg ⁻¹)	Ca (mg kg ⁻¹)	Mg (mg kg ⁻¹)	K (mg kg ⁻¹)
COM0	0.10 \pm 0.01c	1.11 \pm 0.01f	0.11 \pm 0.01e	0.12 \pm 0.01d	0.26 \pm 0.02d
COM1	0.13 \pm 0.01bc	1.22 \pm 0.01e	0.20 \pm 0.02d	0.18 \pm 0.01d	0.30 \pm 0.01d
C1M0	0.17 \pm 0.01b	1.36 \pm 0.02d	0.24 \pm 0.01cd	0.46 \pm 0.03c	0.38 \pm 0.02c
C1M1	0.25 \pm 0.02a	1.56 \pm 0.02c	0.27 \pm 0.01c	0.54 \pm 0.04b	0.46 \pm 0.02b
C2M0	0.26 \pm 0.01a	1.66 \pm 0.02b	0.35 \pm 0.02b	0.62 \pm 0.02a	0.48 \pm 0.02ab
C2M1	0.27 \pm 0.01a	1.72 \pm 0.12a	0.47 \pm 0.03a	0.67 \pm 0.02a	0.51 \pm 0.01a

Mean \pm SE of three replicates are shown; different letters on numerical values indicate significant differences ($P < 0.05$) along the columns. COM0=control, COM1= mycorrhizal inoculation at 20g only; C1M0= compost at 20tha⁻¹ only; C2M0=compost at 30tha⁻¹only; C1M1=compost and mycorrhizal inoculation at 20tha⁻¹; C2M1=compost and mycorrhizal inoculation at 30tha⁻¹.

Effect of compost and AMF on vegetative growth and percent root colonization

As shown in Table 4, the plant height basically increased after soil amendment. AMF inoculation (COM1) significantly increase compare to control. Application of compost at 20 and 30 t ha⁻¹ increase by 7.65 % and 7.83 % compared to compost and AMF inoculation respectively. Treatment with AMF inoculation showed no significant difference compare to control in stem diameter. The application of compost in combination with AMF inoculation (C1M1) was higher by 18.03 % compare to compost application only. Similarly, compost and AMF inoculation at 30 t ha⁻¹ increase by 13 % compared to C2M0. Root dry biomass for AMF inoculation (COM1) was significantly higher compared to control. Compost and AMF inoculation at 20 and 30 t ha⁻¹ increased by 5.69 % and 6.08 % compare to compost application only. Shoot dry biomass revealed significant

differences of treatment application. Inoculation with AMF only (C0M1) was significantly higher compared to control. Compost in combination with AMF inoculation at 20 and 30 t ha⁻¹ significantly increase compare to C1M0 and C2M0 respectively. Significant increases in fruit yield were recorded with compost and AMF inoculation in comparison to compost only. Compost and AMF inoculation (C2M1) at 30 t ha⁻¹ significantly increased (8.17%) compared to compost application only (C2M0). Similarly, compost and AMF inoculation (C1M1) at 20 t ha⁻¹ was significantly higher compared to compost application only (C1M0). Inoculation with AMF only (C0M1) significantly increased compared to control. Percentage root colonization is presented in Figure 3. AMF inoculation (C0M1) significantly increase compared to control. Treatment C1M1 at 20 t ha⁻¹ significantly increase compared to compost application only and control. Similarly, compost and AMF inoculation (C2M1) significantly increase compared to compost application only (C2M0).

Table 4. Vegetative growth of pepper plant as affected by compost and AMF inoculation

	Plant Height (cm)	Stem Diameter (cm)	Root Dry Biomass (g/pot)	Shoot Dry Biomass (g/pot)	Fruit Yield (g/pot)
C0M0	12.83±0.57d	0.46±0.02b	0.67±0.03e	1.07±0.02f	0.71±0.31e
C0M1	16.03±0.42c	0.47±0.02b	1.04±0.02d	1.31±0.03e	2.37±0.05d
C1M0	17.90±0.12bc	0.50±0.03b	1.16±0.01c	2.14±0.02d	4.01±0.06c
C1M1	16.53±0.31bc	0.61±0.03ab	1.23±0.02c	2.23±0.01c	5.71±0.02b
C2M0	20.43±0.56a	0.60±0.01ab	1.39±0.02b	2.31±0.02b	5.84±0.19b
C2M1	18.83±0.24ab	0.69±0.01a	1.48±0.04a	2.40±0.01a	6.36±0.04a

Mean ± SE of three replicates are shown; different letters on numerical values indicate significant differences ($P < 0.05$) along the columns. C0M0=control, C0M1= mycorrhizal inoculation at 20g only; C1M0= compost at 20tha⁻¹ only; C2M0=compost at 30tha⁻¹only; C1M1=compost and mycorrhizal inoculation at 20tha⁻¹; C2M1=compost and mycorrhizal inoculation at 30tha⁻¹.




Figure 3. Percent root colonisation as affected by compost and AMF inoculation. Bars represent the mean ± SE of three replicates ($P < 0.05$). C0M0=control, C0M1= mycorrhizal inoculation at 20g only; C1M0= compost at 20 tha⁻¹ only; C2M0=compost at 30 tha⁻¹only; C1M1=compost and mycorrhizal inoculation at 20 tha⁻¹; C2M1=compost and mycorrhizal inoculation at 30 tha⁻¹.

Discussion

We found evidence that in this study compost and AMF inoculation increase soil total N concentration. This result corresponds with that of Xu et al. (2015) who reported that AMF can access nitrogen from organic sources and also mineralization of compost in the soil help increase nitrogen content as declared by Aylaj et al. (2019). Astiko et al. (2013) argued that organic amendments with AMF inoculation increase soil N, P, K and organic carbon. It is important to note that AMF inoculation and organic amendment such as compost, manure, plant residue positively impact soil fertility status and biological activities (Warnock et al., 2007). The available P concentration in our study increased with compost and AMF inoculation compared to uninoculated. Ortas et al. (2012) reported that inoculation with mycorrhizal fungi increased phosphorus concentration compared to uninoculated plants which might be due to production of phosphatase enzyme. Also, Fischer and Glaser (2012) stated that organic additives contributed to increased concentration of phosphorus as a result of various nutrient releases such as nitrogen, phosphorus and potassium from compost application. In our study, the application of compost and AMF inoculation increased K concentration in the soil. This trend did not differ from Astiko et al. (2013) who reported organic amendment with AMF inoculation increased the concentration of K in the soil. In our study, compost and AMF inoculation significantly increase concentration of soil carbon. Quilliam et al. (2010) and Olsson et al. (2014) discovered that inoculation with symbiotic fungi aided nutrient exchange which promoted increase in soil carbon content. More so, incorporation of organic amendment will help improve the functions of AM fungi (Warnock et al., 2010) and

increase soil N, P and K content as discussed by Demir and Gülser (2015). One of the most positive effects of mycorrhizal colonization is the increase in nutrient uptake. In this study, treatments showed significant increases in the nutrient uptake. The application of compost and AMF inoculation showed a positive response in N uptake. Earlier studies suggested that introduction of AM fungi enhanced the uptake of nitrogen (Hodge and Fitter, 2010; Whiteside et al., 2012). Mycorrhizal inoculated plants in combination with compost significantly influenced uptake of Ca, Mg and K. This conforms to Akande et al. (2018) who reported that AM fungi in combination with organic amendment increased Ca, Mg and K uptake. P uptake was promoted with compost and AMF inoculation in this study which was possibly due to phosphatase activity from external hyphal of AMF in rhizosphere (Widiastuti et al., 2003). It is ascertained that increased P uptake caused nutrients balance in plant and therefore promote uptake of other available soil nutrients (Boonlue et al., 2012). Furthermore, several mechanisms are responsible for increased P uptake, including soil exploration, absorption and release of phosphorus through fungal hyphae (Nazeri et al., 2014). Ortas (2012) proved that inoculation with arbuscular mycorrhizal fungi proved effective in soils with low phosphorus concentration than with higher content in a field research. In addition, compost in combination with AMF inoculation increased K uptake than uninoculated with compost. Perner et al. (2007) in earlier works concluded that arbuscular mycorrhizal fungi enhanced phosphorus and potassium uptake in pelargonium plant. The result of this study showed compost and AMF significantly contributed to increase in yellow bell pepper growth and yield compared to uninoculated. This affirms with Tanwar and Aggarwal (2013) who stated that inoculation with AMF especially *Glomus mosseae* promotes growth of red bell pepper due to its compatibility and ability to absorb nutrients from the soil. Also, Aminifard et al. (2013) in a research reported that *Capsicum species* exhibited rapid growth after incorporation of compost which might be related to the decomposition of compost materials and nitrogen mineralization. Furthermore, other authors reported that organic amendment and AMF relatively increased vegetative growth of tomato (Maaitah et al., 2014), red bell pepper (Tanwar et al., 2013b), green bell pepper (Tanwar et al., 2013a), Lentil (Hanif et al., 2013) and improved varieties of *Capsicum species* (Olawuyi et al., 2014). More so, improvement in plant growth might be attributed to supply of sufficient nutrients to soil and mutual connection of AMF and function in root zones as reported by Akyol et al. (2019). Another trending reason as stated by Tanwar et al. (2011) and Kim et al. (2010) is the ability of AMF to acquire and translocate nutrients via mycelia throughout the roots. In our study compost and AMF inoculation increased shoot and root dry biomass compared to uninoculated. This connotes the study of Olawuyi et al. (2014) who reported that organic amendment and AMF inoculation positively influenced shoot and root biomass of pepper plants in a field experiment. Çekic et al. (2012) stated that inoculation with *Glomus mosseae* significantly influenced shoot and root biomass yield of *Capsicum annum* due to enhanced nutrient acquisition. An increase in biomass might be achieved through improved plant growth which suggests improved fresh shoot and root development (Tanwar et al., 2012). In addition, compost and AMF improved vegetative growth and yield of tomato and pepper plant as a result of symbiotic association between plant root and advanced uptake of mineral nutrients as investigated by Fawole et al. (2016) and Oni (2015). As documented in this study, AMF inoculated plants significantly increased root colonization compared to uninoculated. This relates to findings of other studies that AMF root colonization increased as a result of acquisition of phosphorus by the mycelia network of host root plant in the soil (Tanwar et al., 2014; Kim et al., 2017). Also, conforms to a report that increase in percent root colonization is as a result of increase in soil nitrogen content (Hodge and Fitter, 2010; Smith and Smith, 2011). Contrarily, a non-effective colonization has been attributed to increased level of phosphorus in the soil as stated by Prasad et al. (2012) and Naghashzadeh et al. (2013). The extent of development and colonization of fungi to roots is dependent of host plant, origin and strain of mycorrhiza as discussed by Mau et al. (2014) and Verbruggen et al. (2013). Yield is considered an important factor in production of horticultural crops. In this study, compost and AMF positively contributed to yield of yellow bell pepper of inoculated than uninoculated. This is relevant to findings of Copetta et al. (2011) who reported that green compost and AMF inoculation assisted in improving yield of tomato plants due to increased level of colonization and nutrients accumulation. AyanfeOluwa (2019) and Hossain et al. (2017) affirmed that utilization of compost and mycorrhizal inoculation enhanced growth and yield of horticultural crops by mutual association of AMF and steady release of nutrients from applied compost. Investigations from similar studies for Onion (Ortega-García et al., 2015), Celosia (Komolafe et al., 2021) and Okra (Kayode et al., 2018) have proven increase in crop growth and yield may be assigned to symbiotic associations of AMF and compost mineralization.

Conclusion

Short-term application of compost with and without mycorrhizal inoculation at transplanting enhanced plant growth, uptake of essential nutrient and improved soil productive capacity. The effect of combined application

of compost and AMF inoculation positively influence macronutrients in the soil, plant growth, nutrient uptake and root colonization. Since compost is an organic amendment, combination with AMF could be a management strategy to improve soil productive quality. This could be recommended to farmers since it is environmental friendly and requires no specialized skill for its application. Therefore a long-term experiment should be conducted since longer time allows more decomposition of compost.

Acknowledgements

The authors acknowledge the Organic Agriculture Farm of the Federal University of Agriculture Abeokuta and Institute of Agricultural Research and Training Ibadan, Nigeria.

References

- Akande, T.Y., Fagbola, O., Erinle, K.O., Bitire, T.D., Urhie, E.J., 2018. Effect of organic manure and mycorrhizal on the growth and yield of *Capsicum annum* (Hot Pepper). *New York Science Journal* 11(5): 1-9.
- Akyol, T.Y., Niwa, R., Hirakawa, H., Maruyama, H., Sato, T., Suzuki, T., Fukunaga, A., Sato, T., Yoshida, S., Tawaraya, K., 2019. Impact of introduction of arbuscular mycorrhizal fungi on the microbial community in agricultural fields. *Microbes and Environments* 34(1): 23–32.
- Aminifard, M.H., Aroiee, H., Azizi, M., Nemati, H., Jaafar, H.Z.E., 2013. Effect of compost on antioxidant components and fruit quality of sweet pepper (*Capsicum annum L.*). *Journal of Central European Agriculture* 14(2): 47-56.
- AOAC, 2005. Official Methods of Analysis of the AOAC International. Association of Official Analytical Chemists International. 18th Edition. Horwitz, W., Latimer, G.W., (Eds.), AOAC (Association of Official Agricultural Chemists) International, USA.
- Astiko, W., Ika, R.S., Syamsuddin, D., Anton, M., 2013. The role of indigenous mycorrhiza in combination with cattle manure in improving maize yield (*Zea Mays L*) on sandy loam of Northern Lombok, Eastern of Indonesia. *Journal of Tropical Soils* 18: 53-58.
- AyanfeOluwa, O.E., 2019. Availability of nutrients from an accelerated compost for maize (*Zea mays*) production in two soil types. *Communications in Soil Science and Plant Analysis* 50(12): 1476-1486.
- Aylaj, M., Lhadi, E., Adani, F., 2018. Municipal waste and poultry manure compost affect biomass production, nitrate reductase activity and heavy metals in tomato plants. *Compost Science and Utilization* 27(1): 11-23.
- Biala, J., 2000. The use of composed organic waste in viticulture – A review of the international literature and experience, Sustainable Industries Branch, Canberra act 2601, Environment Australia, Canberra.
- Boonlue, S., Surapat, W., Pukahuta, C., Suwanarit, P., Suwanarit, A., Morinaga, T., 2012. Diversity and efficiency of arbuscular mycorrhizal fungi in soils from organic chili (*Capsicum frutescens*) farms. *Mycoscience* 53(1): 10-16.
- Bremner, J.M., 1996. Nitrogen-total. In: *Methods of Soil Analysis: Part 3 Chemical Methods*, 5.3. Sparks, D.L. Page, A.L., Helmke, P.A., Loeppert, R.H., Soltanpour, P.N., Tabatabai, M.A., Johnston, C.T., Sumner, M.E. (Eds.). SSSA Book Series No. 5. ASA-SSSA Madison WI, USA, pp. 1085– 1121.
- Carlson, J., Saxena, J., Basta, N., Hundal, L., Busalacchi, D., Dick, R.P., 2015. Application of organic amendments to restore degraded soil: effects on soil microbial properties. *Environmental Monitoring and Assessment* 187: 109.
- Cely, M.V.T., de Oliveira, A.G., de Freitas, V.F., de Luca, M.B., Barazetti, A.R., dos Santos, I.M.O., Gionco, B., Garcia, G.V., Prete, C.E.C., Andrade, G., 2016. Inoculant of arbuscular mycorrhizal fungi (*Rhizophagus clarus*) increase yield of soybean and cotton under field conditions. *Frontiers in Microbiology* 7: 720.
- Chukwuka, K.S., Omotayo, O.E., 2009. Soil fertility restoration potentials of tithonia green manure and water hyacinth compost on a nutrient depleted soil in South Western Nigeria using *Zea mays L.* as test crop. *Research Journal of Soil Biology* 1: 20-30.
- Copetta, A., Bardi, L., Bertolone, E., Berta, G., 2011. Fruit production and quality of tomato plants (*Solanum lycopersicum L.*) are affected by green compost and arbuscular mycorrhizal fungi. *Plant Biosystems - An International Journal Dealing with all Aspects of Plant Biology* 145(1): 106–115.
- Cozzolino, V., Di Meo, V., Monda, H., Spaccini, R., Piccolo, A., 2016. The molecular characteristics of compost affect plant growth, arbuscular mycorrhizal fungi, and soil microbial community composition. *Biology and Fertility of Soils* 52: 15–29.
- Çekic, F.Ö., Ünyayar, S., Ortaş, I., 2012. Effects of arbuscular mycorrhizal inoculation on biochemical parameters in *Capsicum annum* grown under long term salt stress. *Turkish Journal of Botany* 36(1): 63–72.
- Demir, Z., Gülser, C., 2015. Effects of rice husk compost application on soil quality parameters in greenhouse conditions. *Eurasian Journal of Soil Science* 4(3): 185-190.
- Fawole, O.B., Alori, E.T., Ojo, O.A., 2016. Evaluation of two composts for the improvement of crop yield using tomato (*Lycopersicon esculentum*) as test crop. *Journal of Agricultural Sciences, Belgrade* 61(1): 37-44.
- Fischer, D., Glaser, B., 2012. Synergisms between compost and biochar for sustainable soil amelioration In: *Management of Organic Waste*. Kumar, S., Bharti, S. (Eds.). pp.167-198. IntechOpen.
- Franco, A.D., Carrillo, M.A., Chairez, F.O., Cabrera, O.G., 2013. Plant nutrition and fruit quality of pepper associated with arbuscular mycorrhizal in greenhouse. *Revista Mexicana de Ciencias Agrícolas* 4(2): 315–321.
- Frank, K., Beegle, D., Denning, J., 1998. Phosphorus. In: *Recommended Chemical Soil Test Procedures for the North Central Region*. Brown, J.R. (Ed.), North Central Regional Research Publication No. 221 (revised) Missouri Agriculture

- Experimental Station Columbia, MO, USA. pp. 21–26. Available at [Access date: 07.12.2021]: <https://extension.missouri.edu/media/wysiwyg/Extensiondata/Pub/pdf/specialb/sb1001.pdf>
- Gautam, S.S., Pathak, N., 2014. Effect of organic fertilizers on soybean yield in Bundelkhand. *TECHNOFAME-A Journal of Multidisciplinary Advance Research* 3(2): 84-87.
- Gee, G.W., Or, D., 2002. Particle-size analysis. In: *Methods of Soil Analysis: Part 4 Physical Methods*, 5.4. Dane, J.H., Topp G.C. (Eds.), SSSA Book Series. Soil Science Society of America, Madison, Wisconsin, USA. pp. 255–293.
- Giovannetti, M., Mosse, B., 1980. An evaluation of techniques for measuring vesicular arbuscular mycorrhizal infection in roots. *New Phytologist* 84(3): 489–500.
- González, M., Gomez, E., Comese, R., Quesada, M., Conti, M., 2010. Influence of organic amendments on soil quality potential indicators in an urban horticultural system. *Bioresource Technology* 101(22): 8897-8901.
- Gosling, P., Hodge, A., Goodlass, G., Bending, G.D., 2006. Arbuscular mycorrhizal fungi and organic farming. *Agriculture, Ecosystems & Environment* 113(1-4): 17-35.
- Hanif, H., Akhtar, N., Cheema, T.A., Achakzai, A.K.K., 2013. Effect of dung, leaf litter and urea on growth of VA mycorrhizae in *Lens culinaris* medik. cv. massur 95 grown under field conditions. *Biologia (Pakistan)* 59(2): 221-226.
- Hodge, A., Fitter A.H., 2010. Substantial nitrogen acquisition by arbuscular mycorrhizal fungi from organic material has implications for N cycling. *PNAS - Proceedings of the National Academy of Sciences* 107(31): 13754–13759.
- Hossain, M.Z., Fragstein, V., Niemsdorff, P., Heß, J., 2017. Effect of different organic wastes on soil properties and plant growth and yield: a review. *Scientia Agriculturae Bohemica* 48(4):224–237.
- Hu, J., Lin, X., Wang, J., Dai, J., Cui, X., Chen, R., Zhang, J., 2009. Arbuscular mycorrhizal fungus enhances crop yield and P-uptake of maize (*Zea mays* L.): A field case study on a sandy loam soil as affected by long-term P-deficiency fertilization. *Soil Biology and Biochemistry* 41(12): 2460–2465.
- Jung, S.C., Martinez-Medina, A., Lopez-Raez, J.A., Pozo, M.J., 2012. Mycorrhiza-induced resistance and priming of plant defenses. *Journal of Chemical Ecology* 38: 651-664.
- Kayode, C.O., Adeoye, G.O., Ezekiel-Adewoyin, D.T., AyanfeOluwa, O.E., Ogunleti, D.O., Adekunle, A.F., 2018. Influence of cocoa pod husk-based compost on nutrient uptake of okra (*Abelmoschus esculentus* (L.) MOENCH) and soil properties on an Alfisol. *Communications in Soil Science and Plant Analysis* 48(17): 2113-2122.
- Kim, S.J., Eo, J.K., Lee, E.H., Park, H., Eom, A.H., 2017. Effects of arbuscular mycorrhizal fungi and soil conditions on crop plant growth. *Mycobiology* 45(1): 20-24.
- Kim, K, Yim, W., Trivedi, P., Madhaiyan, M., Boruah, H.P.D., Islam, R., Lee, G., Sa, T., 2010. Synergistic effects of inoculating arbuscular mycorrhizal fungi and *Methylobacterium oryzae* strains on growth and nutrient uptake of red pepper (*Capsicum annum* L.). *Plant and Soil* 327: 429-440.
- Komolafe, A.F., Kayode, C.O., Ezekiel-Adewoyin, D.T., AyanfeOluwa, O.E., Ogunleti, D.O., Makinde, A.I., 2021. Soil properties and performance of celosia (*Celosia Argentea*) as affected by compost made with *Trichoderma asperellum*. *Eurasian Journal of Soil Science* 10(3): 199-206.
- Krüger, M., Krüger, C., Walker, C., Stockinger, H., Schüßler, A., 2012. Phylogenetic reference data of systematics and phylotaxonomy of arbuscular mycorrhizal fungi from phylum to species level. *New Phytologist* 193(4): 970-984.
- Linderman, R.G., 1992. Vesicular-arbuscular mycorrhizae and soil microbial interactions. In: *Mycorrhizae in Sustainable Agriculture*. Bethlenfalvay, G.J., Linderman, R.G. (Eds.), ASA Special Publication 54, Madison, WI, USA. pp. 45–70.
- López, A., Fenoll, J., Hellín, P., Flores, P., 2014. Cultivation approach for comparing the nutritional quality of two pepper cultivars grown under different agricultural regimes. *LWT - Food Science and Technology* 58(1): 299-305.
- Maaitah, S.A., Atif, Y.M., Ghazi, N.A., 2014. Response of potted-grown tomato (*Lycopersicon esculentum* Mill var. *Lorely F1*) to AMF inoculation with organic and inorganic fertilizer. *American-Eurasian Journal of Agriculture and Environment* 14(3): 225-262.
- Malik, A.A., Chattoo, M.A., Sheemar, G., Rashid, R., 2011. Growth, yield and fruit quality of sweet pepper hybrid SH-SP-5 (*Capsicum annum* L.) as affected by integration of inorganic fertilizers and organic manures (FYM). *Journal of Agricultural Technology* 7(4): 1037-1048.
- Mau, A.E., Utami, S.R., 2014. Effects of biochar amendment and arbuscular mycorrhizal fungi inoculation on availability of soil phosphorus and growth of maize. *Journal of Degraded and Mining Lands Management* 1(2):69-74.
- Nadeem, S.M., Ahmad, M., Zahir, Z.A., Javaid, A., Ashraf, M.M., 2014. The role of mycorrhizae and plant growth promoting rhizobacteria (PGPR) in improving crop productivity under stressful environments. *Biotechnology Advances* 32(2): 429-448.
- Naghashzadeh, M., Modaberi, A., Sharafizad, I., Sharafizad, M., 2013. Phosphorus concentration and root colonization as affected by arbuscular mycorrhizal fungi in maize (*Zea mays* L.) under drought stress conditions. *Advances in Environmental Biology* 7(14): 4623-4629
- Nazeri, N.K., Lambers, H., Tibbett, M., Ryan, M.H., 2014. Moderating mycorrhizas: arbuscular mycorrhizas modify rhizosphere chemistry and maintain plant phosphorus status within narrow boundaries. *Plant, Cell and Environment* 37(4): 911–921.
- Nelson, D.W., Sommers, L.E., 1996. Total carbon organic carbon and organic matter. In: *Methods of Soil Analysis: Part 3 Chemical Methods*, 5.3. Sparks, D.L. Page, A.L., Helmke, P.A., Loeppert, R.H., Soltanpour, P.N., Tabatabai, M.A., Johnston, C.T., Sumner, M.E. (Eds.). SSSA Book Series No. 5. ASA-SSSA Madison WI, USA, pp. 961–1010.

- Neumann, E., George, E., 2010. Nutrient uptake: the arbuscular mycorrhiza fungal symbiosis as a plant nutrient acquisition strategy. In: *Arbuscular Mycorrhizas: Physiology and Function*. Koltai, H., Kapulnik, Y. (Eds.). Springer, Dordrecht. pp.137-167.
- Okalebo, J.R., Gathua, K.W., Woomer, P.L., 2002. *Laboratory Methods of Soil and Plant Analysis. A Working Manual*. 2nd Ed. TSBF-CIAT, SACRED Africa, KARI, Soil Science Society of East Africa, Nairobi, Kenya. 128p.
- Olawuyi, O.J., Jonathan, S.G., Babatunde, F.E., Babalola, B.J., Yaya, O.O.S., Agbolade, J.O., Aina, D.A., Egun, C.J., 2014. Accession × Treatment Interaction, Variability and Correlation Studies of Pepper (*Capsicum spp.*) under the Influence of Arbuscular Mycorrhiza Fungus (*Glomus clarum*) and Cow Dung. *American Journal of Plant Sciences* 4(5): 683-690.
- Olsson, O., Olsson, P.A., Hammer, E.C., 2014. Phosphorus and carbon availability regulate structural composition and complexity of AM fungal mycelium. *Mycorrhiza* 24: 443-451.
- Oni, S.A., 2015. Growth and fruit yield of Pepper (*Capsicum frutescens L.*) as influenced by Arbuscular mycorrhizal (AM) inoculation and fertilizers under greenhouse and field conditions. *Applied Tropical Agriculture* 15: 126-133.
- Ortas, I., 2012. The effect of mycorrhizal fungal inoculation on plant yield, nutrient uptake and inoculation effectiveness under long-term field conditions. *Field Crops Research* 125: 35-48.
- Ortega-García, J.G., Montes-Belmont, R., Rodríguez-Monroy, M., Ramírez-Trujillo, J.A., Suárez-Rodríguez, R., Sepúlveda-Jiménez, G., 2015. Effect of *Trichoderma asperellum* applications and mineral fertilization on growth promotion and the content of phenolic compounds and flavonoids in onions. *Scientia Horticulturae* 195: 8-16.
- Patchaye, M., Sundarkrishnan, B., Tamilselvan, S., Sakthivel, N., 2018. Microbial management of organic waste in agroecosystem. In: *Microorganisms for green revolution. Microorganisms for sustainability*. Panpatte, D., Jhala, Y., Shelat, H., Vyas, R. (Eds.). vol 7, Springer, Singapore. pp.45-73.
- Perner, H., Schwarz, D., Bruns, C. Mäder, P.C., George, E., 2007. Effect of arbuscular mycorrhizal colonization and two levels of compost supply on nutrient uptake and flowering of pelargonium plants. *Mycorrhiza* 17: 469-474.
- Prasad, K., Aggarwal, A., Yadav, K., Tanwar, A., 2012. Impact of different levels of superphosphate using arbuscular mycorrhizal fungi and *Pseudomonas fluorescens* on *Chrysanthemum indicum L.* *Journal of Soil Science and Plant Nutrition* 12(3): 451-462.
- Quilliam, R.S., Hodge, A., Jones, D.L., 2010. Sporulation of arbuscular mycorrhizal fungi in organic-rich patches following host excision. *Applied Soil Ecology* 46(2): 247-250.
- Rouphael, Y., Franken, P., Schneider, C., Schwarz, D., Giovannetti, M., Agnolucci, M., De Pascale, S., Bonini, P., Colla, G., 2015. Arbuscular mycorrhizal fungi act as biostimulants in horticultural crops. *Scientia Horticulture* 196: 91-108.
- Santos, L.Z.H., Myrna, S.O.M., Wenndy, L.W., Andrea, V.R., Manuel, G.P.J.M., 2011 Effects of compost made with sludge and organic residues on bean (*Phaseolus vulgaris L.*) crop and arbuscular mycorrhizal fungi density. *African Journal of Agricultural Research* 6: 1580-1585.
- Serpeloni, J.M., Leal Specian, A.F., Ribeiro, D.L., Tuttis, K., Vilegas, W., Martínez- López, W., Dokkedal, A.L., Saldanha, L.L., Cólus, I.M.S., Varanda, E.A., 2015. Antimutagenicity and induction of antioxidant defense by flavonoid rich extract of *Myrcia bella* Cambess, in normal and tumor gastric cells. *Journal of Ethnopharmacology* 176: 345-355.
- Shahidi, F., Ambigaipalan, P., 2015. Phenolics and polyphenolics in foods, beverages and spices: antioxidant activity and health effects - A review. *Journal of Functional Foods* 18(B): 820-897.
- Smith, S.E., Read, D.J., 2008. *Mycorrhizal symbiosis*, 3rd Edition. Academic Press. New York, 800p.
- Smith, S.E., Smith, F.A., 2011. Roles of arbuscular mycorrhizas in plant nutrition and growth: new paradigms from cellular to ecosystem scales. *Annual Review of Plant Biology* 62: 227-250.
- Smith, S.E., Anderson, I.C., Smith, A.F., 2015. Mycorrhizal associations and phosphorus acquisition: from cells to ecosystems. In: *Annual Plant Reviews Volume 48: Phosphorus Metabolism in Plants*. Plaxton, W., Lambers, H. (Eds.). John Wiley & Sons, Ltd. pp. 409-440.
- Tanwar, A., Aggarwal, A., Yadav, A., Kadian, N., 2011. Enhanced growth and yield of *Capsicum annum L.* with two endomycorrhizal fungi and other bioinoculants. *Journal of Indian Botany Society* 90: 351-359.
- Tanwar, A., Aggarwal, A., Karishma, A., Neetu, A., 2012. Effectiveness of endomycorrhizal fungi and *Pseudomonas fluorescens* under different phosphorus levels on *Capsicum annum L.* *Kasetsart Journal (Natural Science)* 46: 769-782.
- Tanwar, A., Aggarwal, A., Kadiyan, N., Gupta, A., 2013a. Arbuscular mycorrhizal inoculation and super phosphate application influence plant growth and yield of *Capsicum annum*. *Journal of Soil Science and Plant Nutrition* 13(1): 55-66.
- Tanwar, A., Aggarwal, A., Yadav, A., Parkash, V., 2013b. Screening and selection of efficient host and sugarcane bagasse as substrate for mass multiplication of *Funneliformis mosseae*. *Biological Agriculture & Horticulture* 29(2): 107-117.
- Tanwar, A., Aggarwal, A., 2014. Multifaceted potential of bioinoculants on red bell pepper (F1 hybrid, Indam Mamatha) production. *Journal of Plant Interactions* 9(1): 82-91.
- Tanwar, A., Singh, A., Aggarwal, A., Jangra, E., Pichardo, S.T., 2021. Evaluation of municipal sewage sludge for Arbuscular mycorrhizal fungi inoculum production. *Eurasian Journal of Soil Science* 10(4): 343-353.
- Tel, D.A., Hagarty, M., 1984. *Soil and Plant Analysis-Study Guide for Agricultural Laboratory, Directors and Technologist Working in Tropical Regions*. International Institute of Tropical Agriculture, Ibadan, Nigeria & University of Guelph, Canada Publications. 277p.

- Ubah, J., Fagbola, O., Aladele, S.E., 2012. Growth of two cowpea [*Vigna unguiculata* (L.) Walp.] varieties as influenced by arbuscular mycorrhizal fungi and *Tithonia* (*Tithonia diversifolia* Hemsl.) application under screen house conditions. *Crop Research* 44(3): 338-343.
- Verbruggen, E., van der Heijden, M.G.A., Rillig, M.C., Kiers, T., 2013. Mycorrhizal fungal establishment agricultural soils: factors determining, inoculation success. *New Phytologist* 197(4): 1104-1109.
- Warnock, D.D., Lehmann, J., Kuyper, T.W., Rillig, M.C., 2007. Mycorrhizal responses to biochar in soil concepts and mechanisms. *Plant and Soil* 300: 9-20.
- Warnock, D.D., Daniel, L., Mummey, D.D., Mcbride, B., Julie-Major, J., Lehmann, J., Rillig M.C., 2010. Influences of non-herbaceous biochar on arbuscular mycorrhizal fungal abundances in roots and soils: Results from growth-chamber and field experiments. *Applied Soil Ecology* 46: 450-456.
- Weber, J., Karczewska, A., Drozd, J., Licznar, M., Licznar, S., Jamroz, E., Kocowicz, A., 2007. Agricultural and ecological aspects of a sandy soil as affected by the application of municipal solid waste composts. *Soil Biology and Biochemistry* 39(6): 1294-1302.
- Whiteside, M.D., Digman, M.A., Gratton, E., Treseder, K.K., 2012. Organic nitrogen uptake by arbuscular mycorrhizal fungi in a boreal forest. *Soil Biology and Biochemistry* 55: 7-13.
- Widiastuti, H., Sukarno, N., Darusman, L.K., Goenadi, D.H., Smith, S., dan Guhardja, E., 2003. Phosphatase activity and organic acid production in rhizosphere and hyphosphere of mycorrhizal oil palm seedling. *Journal of Menara Perkebunan* 71: 70-81. [in Indonesian].
- Xu, P., Liang, L.Z., Dong, X.Y., Shen, R.F., 2015. Effect of arbuscular mycorrhizal fungi on aggregate stability of a clay soil inoculating with two different host plants. *Acta Agriculturae Scandinavica, Section B-Soil & Plant Science* 65(1): 23-29.
- Zong, W., Gu, T., Wang, W., Zang, B., Lin, X., Huang, Q., Shen, W., 2010. The effects of mineral fertilizer and organic manure on soil microbial community and diversity. *Plant and Soil* 326: 511-522.