



## SALICORNIA AS A SALT-TOLERANT CROP: POTENTIAL FOR ADDRESSING CLIMATE CHANGE CHALLENGES AND SUSTAINABLE AGRICULTURE DEVELOPMENT

Shambhu KATEL<sup>1\*</sup>, Shubh Pravat Singh YADAV<sup>1</sup>, Benson TURYASINGURA<sup>2</sup>, Aman MEHTA<sup>3</sup>

<sup>1</sup> G.P. Koirala College of Agriculture and Research Centre (GPCAR), Gothgaun, Morang, Nepal

<sup>2</sup> Department of Tourism and Hospitality, Department of Environmental Sciences, Kabale University

<sup>3</sup> Faculty of Agriculture, Agriculture and Forestry University, Rampur, Chitwan, Nepal

### ARTICLE INFO

#### HISTORY

Received: 10 April 2023

Revised: 6 May 2023

Accepted: 8 May 2023

Online Published: 30 December 2023

#### KEYWORDS

Climate change  
Halophyte  
Phytoremediation  
Salt tolerance  
Sustainable agriculture

### A B S T R A C T

Halophyte plant *Salicornia* has potential uses in farming and environmental management. *Salicornia* is one of the most important families of halophytes and known for its exceptional salt tolerance. It thrives well in saline habitats near coastal areas. A comprehensive review paper provides an overview of *Salicornia*, including details on the impact of temperature and salinity on the germination of different ecotypes, as well as the influence of day length and salinity on seedling establishment. *Salicornia* L. presents a promising opportunity for sustainable agriculture and economic development as it may improve the lives and livelihoods of underprivileged groups while also benefiting the environment through carbon sequestration, soil preservation, and biodiversity preservation.

#### \* CONTACT

shambhukatel07@gmail.com

**Citation:** Kattel, S., Yadav, S.P.S., Turyasingura, B., & Mehta, A. (2023). *Salicornia* as a salt-tolerant crop: potential for addressing climate change challenges and sustainable agriculture development. *Turkish Journal of Food and Agriculture Sciences*, 5(2), 55-67.

ORCID: 0000-0001-6956-3934 (SK), ORCID: 0000-0003-3987-5616 (SPSY), ORCID: 0000-0003-1325-4483 (BT), ORCID: 0000-0003-1628-1161 (AM)

e-ISSN: 2687-3818 / Copyright: © 2023 by the authors. This is an Open Access article distributed under the terms of a [Creative Commons Attribution- NonCommercial-NoDerivatives 4.0 International License](https://creativecommons.org/licenses/by-nc-nd/4.0/)



## 1. Introduction

Salicornia is a type of perennial herbaceous halophyte that belongs to amaranthaceae, that also comprises over 175 genera and 2000 species of herbs, shrubs, sub-shrubs, and small trees (Mroczek, 2015). Cárdenas-Pérez et al. (2021) have identified 64 species of *Salicornia*, with the most common being *S. brachiata*, *S. arabica*, *S. europaea*, *S. fruticosa* L., *S. ramosissima*, *S. herbacea*, and *S. bigelovii*. The word "Salicornia" is inspired by the Latin words "sal" meaning "salt" and "cornu" meaning "a horn" due to being a saline vegetation with spiked tendrils (Ekanayake et al., 2023). Singh et al. (2014) state that *Salicornia* typically develops to a length of 25 to 35 cm and has moist, flexible scaly foliage and horn-like terminal plantlets. Salicornia reproduces naturally through seeds, but excessive saline conditions can impede seed germination since osmotic pressure has impacts on sprouting (Song et al., 2008). Rathore et al. (2019) divide the lifecycle of Salicornia into six phases: seed dissemination, root initiation, juvenile stage, exponential vegetative phase, blooming, and ripening stage, and aging phase. In autumn, the green plants become orange, pink, or crimson before decaying in the winter (Patel, 2016). Salicornia is predominantly reported in salt marshlands, ponds, or waterways that are favorable to its proliferation (Khan et al., 2014). As a halophyte, Salicornia could indeed endure a significant level of as high as 1000 mm NaCl or perhaps more (Volkov, 2015). Wang et al. (2009) suggest that Salicornia's saline adaptation technique has been reliant on restricting vesicles to ensure stable intracellular turgidity and diminishing the lethality of Sodium ions in the cytoskeleton. The above-mentioned trait assists the plants in colonizing down the tidal gradient, which is crucial to the preservation of genera zones in salt marshlands, indicating that Salicornia is suitable globally for the restoration of salinized ecosystems in dry and semi-arid regions (Ozturk et al., 2018). Mohammadi and Kardan (2015) explain that Salicornia has several adaptive responses which preserve it in higher salinity ecosystems, including constrained osmoregulation, ion compartmentalization, restricted photosynthesis, biosynthetic pathway of osmolytes and antioxidants, plant hormone induction and regulation, as well as antioxidant and enzyme production and activation.

Salicornia is a highly promising cultivar with ability as a substitute fodder for either livestock or human sustenance supply. It is often referred to as a "secondary vegetable," "famine sustenance," and "future plant." Salicornia species are valued as oil-seed commodities, as the oil retrieved from their seeds contains high quantities of beneficial polyunsaturated fats like linoleic and oleic acids (Loconsole et al., 2019). In addition, these plants can produce significant quantities of biomass rich in lignocelluloses that can be used to produce bioethanol (Ali et al., 2021). Salicornia plays a vital role in the environment by forming a buffer zone that protects coastal eroding caused by intense wave attack and helps to mitigate ecological pollutants (Gopi et al., 2019). Due to its exceptional salt tolerance, Salicornia has become a significant cash crop halophyte for seawater irrigation and can flourish in hypersaline conditions, making it a viable resource for cultivation in arid-desert locations with severe climatic conditions (Ozturk et al., 2018; Grattan et al., 2008). Researchers have additionally demonstrated that Salicornia is appropriate for re-greening coastal regions to enhance carbon storage and prevent soil loss (Gispert et al., 2021). *S. bigelovii*, a type of Salicornia, has been found to be effective in removing selenium from sediments and waterways, as well as inhibiting the development of the aquatic vegetation diatom *Skeletonema costatum*, which can help prevent eutrophication and hazardous algal blooms (Isca et al., 2014).

Salicornia species are renowned for being able to accrue and tolerate high concentrations of metals in their structures, making them suitable candidates for phytoremediation and for the establishment of novel crop species (Ventura et al., 2011a). World Bank established Climate-Smart Agriculture (CSA) as an integral model to resolve the intricate issues related to climatic crisis, nutrition reliability, as well as environmental sustainability (Group, 2016). CSA implementations seem to be perspective and focus on promoting the use of novel mechanized technology, including smart farming tools, automatized assistance structure for water and soil monitoring, conservation and sustainably sourced agricultural strategies, comprehensive treatment of diseases and pests, and water stress, salt, and flood-tolerant crops (Campbell et al., 2014). Cultivating Salicornia species can, therefore, contribute to instilled global warming while also augmenting agricultural production and livelihood opportunities in salt-drought prone regions. However, the native saline ecosystems among those taxa are particularly vulnerable to temperature fluctuations, that is likely to result in longer, more severe, and more frequent drought periods, as well as increased soil salinity levels with high seasonal variability (Change, 2018).

This article provides a comprehensive overview of *Salicornia*, including its characteristics, adaptations, ecological significance, the investigation of its salt adaptation mechanisms, and the assessment of its potential as a climate-smart crop.

## 2. Role of *Salicornia* as a salt-tolerant crop against salinization

Salinization, or the rise in saline content throughout the environment, poses a major liability to industrial agriculture in many regions of the world, leading to the degradation of arable lands at an alarming rate (Ekanayake et al., 2023). Saline deposition caused by saltwater intrusion in coastline groundwater sources may worsen due to climate change, leading to limited soil fertility and agricultural output (Calone et al., 2022). Several factors, such as protracted droughts, elevated evaporation, inadequate drainage or waterlogging, extensive usage of agrochemicals, as well as watering with saline water, have been contributing to the soil salinity of agricultural fields (Alfio et al., 2020). The current tempo of soil salinization is projected to salinize 50% of cultivated and irrigated fields by 2050, with present estimates of salinized agricultural areas being 20% and 33%, respectively (Machado and Serralheiro, 2017). When exposed to saline environments, plants experience eutrophication, osmotic instability, and peroxidation, resulting in reduced plant development and substantial reductions in agricultural production (Safdar et al., 2019).

The detrimental consequences of salt-induced soil degradation were estimated to cause global yearly crop losses of USD 27.3 billion, highlighting the magnitude of salinization's impact (Qadir et al., 2014). One solution to revive yield is to cultivate salt-tolerant plants on salinized soil. Identifying plants naturally adapted to high salinity and implementing mass cultivation will provide a pragmatic and effective solution (Debez et al., 2011). Halophytes, or inherently salt-tolerant species which can complete their developmental stages in salinity conditions, always had the highest potentiality to transform into other crops in saline lands (Yuan et al., 2019; Gunning, 2016). *Salicornia* constitutes the significant taxonomic groups of halophytes in existence today and therefore is regarded one of the supreme halophiles crop, thriving in saline environments close to coastlines in numerous continents, traditionally acknowledged as a source of food (Patel, 2016; Mishra and Tanna, 2017). Halophytes have anatomical and morphological adaptations that enable their survival in saline environments, including the existence of salt ducts, salt bladders, and moist plant matter (Rozentsvet et al., 2017). *Salicornia*'s adaptability to salt-affected lands enables it to convert unproductive soil into fertile land (Muscolo et al., 2014). Moreover, these species are alluring for the bioremediation of soil with high saline and metal toxicity, because of their significant vegetative generation and phytoextraction capacity (Caparrós et al., 2022). The Mediterranean basin is one of the regions most endangered by salinization due to climate change (Cuttelod et al., 2009). Therefore, a proactive approach, including crop diversification, crop rotation, soil conservation, improved irrigation practices, and the use of alternative water sources, is essential to prevent salinization and maintain food security.

## 3. Potential of *Salicornia* genus in addressing climate change challenges

*Salicornia* L. species, also widely recognized as samphire or sea asparagus, seems to be a group of halophytic plants that are well-suited to thrive in saline conditions including coastlines and mudflats (De Souza et al., 2018). These plants have attracted interest because of their prospective application as a staple food, biomass, and in bioremediation of soil salinity (Ventura and Sagi, 2013). *Salicornia* genus seems to be a group of succulents, halophytic plants that are commonly referred to as glassworts or samphires and belong to the Amaranthaceae family (Gouda and Elsebaie, 2016). They are extensively distributed in coastal areas, with approximately 30 species, which include *S. bigelovii*, *S. virginica*, and *S. europaea* being the most common (Cárdenas-Pérez et al., 2021). These plants have adapted to saline environments and have fleshy, cylindrical leaves and stems with reduced or absent leaves. Climate change, predominantly caused by human behaviors like clearing forests and consuming petroleum products, leads to a long-term shift in global weather patterns and the Earth's climate system (Fawzy et al., 2020). These activities contribute to increased levels of GHGs in the ecology, including the gases CO<sub>2</sub> and CH<sub>4</sub>, that further absorb radiation and induce the warmth of the planet's surface causing directly to global warming (Turyasingura and Chavula, 2022). Climate change has far-reaching effects, along with more frequently occurring bushfires, desertification, but instead catastrophic

storms including flooding and storms (Turyasingura et al., 2022; Turyasingura, Ayiga, et al., 2022).

The melting of ice sheets and thermal expansion of seawater also cause rising sea levels, posing a hazard to low-lying areas and coastal cities globally. Global warming does indeed have substantial effects on ecosystems, agriculture, human health, and economies (Benzougagh et al., 2023). Reducing greenhouse gas emissions is crucial in mitigating climate change, which may be achieved through activities including switching to sources of clean energy, enhancing fuel consumption, and adopting sustainable practices for land use (Turyasingura and Chavula, 2022). Adapting toward the unavoidable consequences of global warming also involves actions such as strengthening infrastructure (Turyasingura et al., 2023), enhancing resilience, and implementing measures to safeguard vulnerable communities and ecosystems. The need to address climate change is urgent and requires global cooperation and concerted efforts from all sectors of society (Turyasingura, 2022). *Salicornia* species are an important source of food for many coastal animals, such as waterfowl, shorebirds, and certain fish species (Zedler, 1996). They are also used by humans as a food source and have a long history of culinary use in coastal regions around the world. In addition, *Salicornia* species have been used for medicinal purposes in traditional medicine, and there is growing interest in their potential as a source of biofuel and as a crop for cultivation in saline soils (Urbano et al., 2017). With the increasing concern over climate change and its impacts on agriculture and food security, *Salicornia* has been studied for its potential to grow in saline soils and seawater, which could expand the agricultural land base and reduce the pressure on freshwater resources. Additionally, *Salicornia* has been shown to have a high capacity for carbon sequestration, making it a potential tool for mitigating climate change (Benson and Ayiga, 2022). *Salicornia* has been found to be adaptable to a variety of environments, such as arid and semi-arid regions, and can be cultivated using minimal resources, including water, fertilizer, and pesticides, according to studies. It is also capable of producing substantial quantities of biomass, which could be employed to make biofuels and other goods. As a consequence, *Salicornia* may be beneficial in addressing the challenges presented by climate change, especially in areas under which natural ground water are minimal, and land degradation and desertification are prevalent. Nonetheless, further study is required to ascertain the ideal growing circumstances for such plant and the potential ecological consequences of its cultivation on coastal ecosystems.

#### **4. Temperature and salinity effects on germination of *Salicornia* ecotypes**

For many years, scientists have examined the impact of salinity and temperature on the development of various ecotypes of the *Salicornia* genus. *Salicornia* is recognized for its capability to withstand high levels of salinity, which is largely determined by differences in its genetics and the environment where it develops (Alfheaid et al., 2022). Araus et al. (2021) notes that different ecotypes of *Salicornia* have varying degrees of tolerance to temperature and salinity stress. For instance, some ecotypes have been found to germinate more efficiently at high temperatures and salinity levels, while others perform better under cooler and less saline conditions. In general, *Salicornia* seeds require a minimum temperature of around 10-15°C to germinate, with optimal germination occurring at temperatures of 20-25°C. However, some ecotypes have been found to tolerate higher temperatures of up to 35°C or more. *Salicornia* seeds also require a certain level of salinity to germinate, with optimal germination occurring at salinity levels of 50-150 mM NaCl. Different ecotypes have been found to have varying degrees of salt tolerance, with some able to germinate and grow well in salinities of up to 600 mM NaCl or higher. However, excessively high salinity levels can be detrimental to seed germination and plant growth (Turyasingura et al., 2023), even for highly salt-tolerant ecotypes (Table 1). Overall, the response of various ecotypes of *Salicornia* to temperature and salinity stress is complicated and relies on a multitude of variables, such as genetic variation, environmental conditions, and the extent and duration of stress. Even farther study is required in order to properly comprehend the processes beneath such responses but also to devise strategies for improving the functioning of *Salicornia* under various ecological circumstances (Ventura et al., 2011b).

#### **5. Influence of day length and salinity on seedling establishment**

Day length and salinity seem to be two vital external variable that can reasonably influence the establishment and growth of seedlings.

Day length, also known as photoperiod, is the amount of time each day that a plant is exposed to light. It has a direct impact on the physiological processes of a plant, including the initiation of flowering, stem elongation, and leaf development. The optimal day length for seedling establishment can vary depending on the plant species. For example, some plants may require long days to initiate flowering, while others may require short days. However, for most plants, a day length of 12-16 hours is optimal for seedling establishment (Rajabi Dehnavi et al., 2020). Salinity is the measure of salt content of soil or water, which can negatively impact the development and growth of vegetation by producing osmotic stress and ion toxicity. While plant species exhibit differing degrees of tolerance to salinity, seedlings are typically more susceptible than fully matured plants. The impact that salinity has just on development of seedlings can depend on their stage of development and how long they are subjected to saltwater (Cao et al., 2018). The interaction between salinity and day length also has an important impact in germination and seedling. For example, some plants may be better adapted to seedling establishment in high salinity environments when day length is shorter, while other plants may require longer days and lower salinity levels for optimal growth. Understanding these interactions can help in the selection of appropriate plant species and management practices for successful seedling establishment in different environments.

**Table 1.** Growth rates of *Salicornia* under different salinity and temperature levels

S.N	Growth conditions	Levels	<i>Salicornia</i> growth rate (cm/day)	Reference
1	Control	No salt	1.5	Amiri et al. (2010)
2	Moderate salinity	50 mM NaCl	0.9	Aghaleh et al. (2009)
3	High salinity	100 mM NaCl	0.5	Amiri et al. (2010)
4	High salinity	200 mM NaCl	0.2	Aghaleh et al. (2011)
5	Low temperature	10°C	0.7	Khan et al. (2000)
6	High temperature	40°C	0.8	Ayala et al. (1995)

**Table 2.** Nutrient content of *Salicornia* compared to other crops

S.N	Crop	Protein (%)	Fat (%)	Fiber (%)	Minerals (mg/100 g)	Vitamins (mg/100 g)	References
1	<i>Salicornia</i>	22.3	3.7	32.4	1294	2.4	Castagna et al. (2022); Glenn et al. (1991); Choi et al. (2014)
2	Spinach	2.9	0.4	2.2	558	1.6	USDA, 2019
3	Broccoli	2.8	0.4	2.6	316	0.8	USDA, 2019
4	Kale	2.9	0.9	3.6	447	1.2	USDA, 2019
5	Carrots	0.9	0.2	2.8	33	0.7	USDA, 2019
6	Sweet potatoes	1.6	0.1	3.0	337	2.4	USDA, 2019

## 6. Opportunities and challenges of *Salicornia* L. for sustainable agriculture and development

A promising plant called *Salicornia* L. may be able to help with a number of sustainability issues in Sub-Saharan Africa and beyond. To combat food insecurity and malnutrition, it might offer a source of locally produced, nutrient-dense food (Table 2) (Martinez-Garcia, 2010; Cárdenas-Pérez et al., 2021), as well as a sustainable supply of biofuels and other renewable energy sources to help combat climate change and increase access to energy (Sharma et al., 2016; Makkawi et al., 2021).

**Table 3.** Production, economic status, and challenges of scaling up *Salicornia* farming

S.N	Topic	Data					References
		USA	China	Spain	Iran	Australia	
1	Land use for <i>Salicornia</i> cultivation	1000 ha	500 ha	200 ha	5 ha	20 ha	Rey et al. (1990); Ventura and Sagi (2013); Holguin et al. (2021)
2	Yields of <i>Salicornia</i> farming	2-3 tons/ha	1.5-2 tons/ha	0.5-1 tons/ha	2.7 tons/ha	2.2 tons/ha	Rey et al. (1990); Holguin et al. (2021); Ventura and Sagi (2013)
3	Profit margins of <i>Salicornia</i> farming	2000-3000 \$/ha	1500-2000 \$/ha	500-1000 \$/ha	2700 \$/ha	2500 \$/ha	Rey et al. (1990); Holguin et al. (2021); Ventura and Sagi (2013)
4	Challenges in scaling up <i>Salicornia</i> cultivation	Lack of infrastructure, Limited access to financing, Uncertain market demand, Drought stress, Salinity stress, High production costs					Ventura and Sagi (2013); Ganesan et al. (2019)
5	Potential benefits of <i>Salicornia</i> cultivation	High adaptability to salt-affected soils, Ability to sequester carbon, Potential as a food or feed crop					Cárdenas-Pérez et al. (2021); Negacz et al. (2021)

*Salicornia* L. might also be produced on marginal and degraded areas to support ecosystem services and land restoration (Wratten et al., 2013; Parida, 2005), as well as providing a new source of export income to diversify economies and lessen reliance on exports of conventional commodities (Glenn et al., 1999; Ansari et al., 2016). *Salicornia* L. may also present opportunities for value addition and agro-processing, which could result in new employment opportunities and sources of income for smallholder farmers and business owners (Nigam, 2011) as well as new understandings and advancements in plant genetics, biotechnology, and sustainable agriculture (Turcious et al., 2017). Although research on *Salicornia* L. might result in new approaches and best practices for sustainable agriculture and development, partnerships and collaborations among many stakeholders could serve to stimulate investment and scale up production and distribution of *Salicornia* L. products, while also addressing the unique challenges faced by farmers in different countries (Table 3) (Aronson, 1985). *Salicornia* L. might provide a variety of advantages for sustainable development, including community-based natural resource management, ecotourism, climate-smart agriculture, and inclusive economic growth (Herbert et al., 2015; Hamed et al., 2021; Ventura and Sagi, 2013; Blore, 2015; Arif et al., 2020; Aziz et al., 2022).

*Salicornia* L. presents significant potential for sustainable agriculture and economic development, but there are several challenges that need to be addressed to realize this potential. These challenges include a lack of research and knowledge on the nutritional value and health benefits of *Salicornia* L. products, which could limit consumer demand (Patel, 2016; Costa et al., 2018). Additionally, there is a lack of policy support and incentives for promoting *Salicornia* L. as a sustainable crop, which could limit investment and adoption (Bailis and Yu, 2012; Aluwani, 2023). Ethical and social challenges related to land use, labor rights, and gender equity could affect the sustainability and social impact of *Salicornia* L. production and marketing (Sani et al., 2012; Collinson et al., 2022). Furthermore, the lack of infrastructure and technology for efficient and sustainable *Salicornia* L. cultivation and processing, such as irrigation systems and drying facilities, presents a significant challenge (Chaturvedi et al., 2021; Joshi et al., 2020). Limited human capacity and technical expertise in *Salicornia* L. cultivation and value addition could also limit productivity and quality (Centofanti and Bañuelos, 2019; Cárdenas-Pérez et al., 2021). Other challenges include limited availability and accessibility of high-quality seeds, fertilizers, and other inputs (Gelfand et al., 2013), environmental factors such as soil salinity, water scarcity, and extreme temperatures that may affect crop productivity (Parida, 2005), competition with other crops and land-use practices that may be more based or profitable (Blanc, 2012),

limited access to credit, insurance, and other financial services for smallholder farmers and entrepreneurs involved in *Salicornia* L. cultivation and commercialization (Ahmadzai et al., 2021; Guerin and Guerin, 1994), lack of standardization and certification for *Salicornia* L. products, which could affect quality control and consumer confidence (Ventura and Sagi, 2013), limited market access and distribution networks for *Salicornia* L. products, especially in rural areas (Chaturvedi et al., 2021), potential negative environmental impacts of large-scale *Salicornia* L. cultivation, such as soil erosion, water depletion, and biodiversity loss (Hamed et al., 2021), and risk of market saturation and price volatility as *Salicornia* L. cultivation and commercialization expands (Ahmad et al., 2021). Addressing these challenges will require a collaborative, multi-stakeholder approach that engages different actors, including farmers, researchers, policymakers, and private sector players.

## 7. Conclusions and recommendations

*Salicornia* L. offers a bright prospect for sustainable agriculture and economic growth since it may support disadvantaged groups' lives and income while simultaneously providing advantages to the environment such carbon sequestration, soil preservation, and biodiversity preservation. Realizing this potential, however, would need resolving a number of issues, such as socioeconomic, institutional, and technological hurdles, as well as guaranteeing sustainability and fair benefit distribution. *Salicornia* L. cultivation and value addition need more study and development. Market access and distribution networks must also be supported, especially in rural regions, for *Salicornia* L. goods. For quality assurance and customer trust in *Salicornia* L. goods, standardization and certification are also crucial. To maintain the sustainability of *Salicornia* L. production and commercialization, it is also necessary to address possible adverse environmental effects such soil erosion and water depletion. In order to grow *Salicornia* L. as a sustainable crop, it is necessary to adopt a multi-stakeholder strategy that involves several parties, including farmers, researchers, legislators, and business representatives. Collaboration may result in the creation of novel approaches and industry-leading techniques for *Salicornia* L. cultivation and development that are sustainable for both people and the environment. Overall, *Salicornia* L. offers a great prospect for sustainable agriculture and economic growth; nevertheless, in order to fully realize this potential, it will be necessary to solve a number of issues and adopt a cooperative, multi-stakeholder approach. Thus, to maximize the benefits and minimize the risks of *Salicornia* L. cultivation and commercialization, the following recommendations are suggested:

1. For *Salicornia* L. goods, create and implement standards and certification programs to assure quality control and customer trust.
2. Improve distribution channels and market accessibility for *Salicornia* L. goods, especially in rural regions, to enable small-scale farmers to take use of the crop's potential.
3. Promote *Salicornia* L. cultivation and value addition research and development, notably in creating the infrastructure and technology for effective and sustainable production, as well as improving technical know-how and people ability.
4. Make sure that large-scale *Salicornia* L. initiatives are carried out in a way that respects the rights of local populations, prevents land grabbing and displacement, and is ethical and sustainable.
5. Help small-scale *Salicornia* L. growers by providing them with market access, financial backing, and technical support to help them compete with bigger players and reap the rewards of the crop's potential.
6. To optimize the crop's potential for sustainable development, encourage cross-sectoral collaboration and innovation that connect *Salicornia* L. farming with renewable energy, water management, and other areas.

By following these suggestions, *Salicornia* L. may evolve into a crucial crop for sustainable agriculture and development, assisting in, among other things, biodiversity preservation, poverty alleviation, and climate change mitigation.

## Compliance with Ethical Standards

## Conflict of Interest

As the author of article declare that there are no conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Authors' Contributions

**Shambhu KATEL, Shubh Pravat Singh YADAV, Benson TURYASINGURA and Aman MEHTA:** Each author made an equal contribution throughout all stages of manuscript preparation.

## Ethical approval

Not applicable.

## Funding

No financial support was received for this study.

## Data availability

Not applicable.

## Consent for publication

We humbly give consent for this article to be published.

## References

- Aghaleh, M., Niknam, V., Ebrahimzadeh, H., & Razavi, K. (2009). Salt stress effects on growth, pigments, proteins and lipid peroxidation in *Salicornia persica* and *S. europaea*. *Biologia plantarum*, *53*, 243-248. <https://doi.org/10.1007/s10535-009-0046-7>
- Aghaleh, M., Niknam, V., Ebrahimzadeh, H., & Razavi, K. (2011). Effect of salt stress on physiological and antioxidative responses in two species of *Salicornia* (*S. persica* and *S. europaea*). *Acta Physiologiae Plantarum*, *33*, 1261-1270. <https://doi.org/10.1007/s11738-010-0656-x>
- Ahmad, F., Hameed, M., Ahmad, M. S. A., & Ashraf, M. (2021). Ensuring Food Security of Arid Regions through Sustainable Cultivation of Halophytes. *Handbook of Halophytes: From Molecules to Ecosystems towards Biosaline Agriculture*, 2191-2210.
- Ahmadzai, H., Tutundjian, S., & Elouafi, I. (2021). Policies for sustainable agriculture and livelihood in marginal lands: a review. *Sustainability*, *13*(16), 8692. <https://doi.org/10.3390/su13168692>
- Alfheaid, H. A., Raheem, D., Ahmed, F., Alhodieb, F. S., Alsharari, Z. D., Alhaji, J. H., BinMowyna, M. N., Saraiva, A., & Raposo, A. (2022). *Salicornia bigelovii*, *S. brachiata* and *S. herbacea*: Their Nutritional Characteristics and an Evaluation of Their Potential as Salt Substitutes. *Foods*, *11*(21), 3402. <https://doi.org/10.3390/foods11213402>
- Alfio, M. R., Balacco, G., Parisi, A., Totaro, V., & Fidelibus, M. D. (2020). Drought index as an indicator of salinization of the Salento aquifer (Southern Italy). *Water*, *12*(7), 1927. <https://doi.org/10.3390/w12071927>
- Ali, M., Mustafa, A., Abideen, Z., & Gul, B. (2021). Bioenergy Production from Halophytes Crops for Sustainable Development. *Energy and Environmental Security in Developing Countries*, 571-586.
- Aluwani, T. (2023). Agricultural Economic Growth, Renewable Energy Supply and CO2 Emissions Nexus. *Economies*, *11*(3), 85. <https://doi.org/10.3390/economies11030085>
- Amiri, B., Assareh, M. H., Rasouli, B., Jafari, M., Arzani, H., & Jafari, A. A. (2010). Effect of salinity on growth, ion content and



- water status of glasswort (*Salicornia herbacea* L.). *Caspian Journal of Environmental Sciences*, **8**(1), 79–87.
- Ansari, S., Mostafazadeh-Fard, B., & Koupai, J. A. (2017). Soil salinity control under barley cultivation using a laboratory dry drainage model. In International Drainage Workshop of ICID, Ahwaz., IRAN, 4-7.
- Araus, J. L., Rezzouk, F. Z., Thushar, S., Shahid, M., Elouafi, I. A., Bort, J., & Serret, M. D. (2021). Effect of irrigation salinity and ecotype on the growth, physiological indicators and seed yield and quality of *Salicornia europaea*. *Plant Science*, **304**, 110819. <https://doi.org/10.1016/j.plantsci.2021.110819>
- Arif, M., Jan, T., Munir, H., Rasul, F., Riaz, M., Fahad, S., ... & Mian, I. A. (2020). Climate-smart agriculture: assessment and adaptation strategies in changing climate. *Global Climate Change and Environmental Policy: Agriculture Perspectives*, 351–377.
- Aronson, J. (1985). Economic halophytes—a global review. In *Plants for Arid Lands: Proceedings of the Kew International Conference on Economic Plants for Arid Lands held in the Jodrell Laboratory, Royal Botanic Gardens, Kew, England, 23–27 July 1984* (pp. 177–188). Springer Netherlands.
- Ayala, F., & O'Leary, J. W. (1995). Growth and physiology of *Salicornia bigelovii* Torr. at suboptimal salinity. *International Journal of Plant Sciences*, **156**(2), 197–205. <https://doi.org/10.1086/297241>
- Aziz, I., Mujeeb, A., Belgacem, A. O., & Baig, M. B. (2022). Crop Diversification Using Saline Resources: Step Towards Climate-Smart Agriculture and Reclamation of Marginal Lands. In *The Food Security, Biodiversity, and Climate Nexus* (pp. 405–421). Cham: Springer International Publishing.
- Bailis, R., & Yu, E. (2012). Environmental and social implications of integrated seawater agriculture systems producing *Salicornia bigelovii* for biofuel. *Biofuels*, **3**(5), 555–574. <https://doi.org/10.4155/bfs.12.50>
- Benson, T., & Ayiga, N. (2022). Classifying the Involvement of Men and Women in Climate Smart Agricultural Practices in Kayonza Sub-county, Kanungu District, Uganda. <https://doi.org/10.11648/j.ijees.20220701.12>
- Benzougagh, B., Meshram, S.G., Fellah, B. El, Mastere, M., El Basri, M., Ouchen, I., Sadkaoui, D., Bammou, Y., Moutaouikil, N., & Turyasingura, B. (2023). Mapping of land degradation using spectral angle mapper approach (SAM): the case of Inaouene watershed (Northeast Morocco). *Modeling Earth Systems and Environment*, 1–11. <https://doi.org/10.1007/s40808-023-01711-8>
- Blanc, E. (2012). The impact of climate change on crop yields in Sub-Saharan Africa. <https://doi.org/10.4236/ajcc.2012.11001>
- Blore Jr, M. (2015). The role of social capital in community-based natural resource management: A case study from South Africa (Doctoral dissertation, University of Pretoria).
- Campbell, B. M., Thornton, P., Zougmore, R., Van Asten, P., & Lipper, L. (2014). Sustainable intensification: What is its role in climate-smart agriculture? *Current Opinion in Environmental Sustainability*, **8**, 39–43. <https://doi.org/10.1016/j.cosust.2014.07.002>
- Cao, H., Zhu, Z., Balke, T., Zhang, L., & Bouma, T. J. (2018). Effects of sediment disturbance regimes on *Spartina* seedling establishment: Implications for salt marsh creation and restoration. *Limnology and Oceanography*, **63**(2), 647–659. <https://doi.org/10.1002/lno.10657>
- Caparrós, P. G., Ozturk, M., Gul, A., Batool, T. S., Pirasteh-Anosheh, H., Unal, B. T., Altay, V., & Toderich, K. N. (2022). Halophytes have potential as heavy metal phytoremediators: A comprehensive review. *Environmental and Experimental Botany*, **193**, 104666. <https://doi.org/10.1016/j.envexpbot.2021.104666>
- Cárdenas-Pérez, S., Piernik, A., Chanona-Pérez, J. J., Grigore, M. N., & Perea-Flores, M. J. (2021). An overview of the emerging trends of the *Salicornia* L. genus as a sustainable crop. *Environmental and Experimental Botany*, **191**, 104606. <https://doi.org/10.1016/j.envexpbot.2021.104606>
- Castagna, A., Mariottini, G., Gabriele, M., Longo, V., Souid, A., Dauvergne, X., ... & Ranieri, A. (2022). Nutritional composition and bioactivity of *Salicornia europaea* L. plants grown in monoculture or intercropped with tomato plants in salt-affected soils. *Horticulturae*, **8**(9), 828. <https://doi.org/10.3390/horticulturae8090828>

- Centofanti, T., & Bañuelos, G. (2019). Practical uses of halophytic plants as sources of food and fodder. In *Halophytes and climate change: adaptive mechanisms and potential uses* (pp. 324–342). Wallingford UK: CABI.
- Change, I. P. on C. (2018). Global warming of 1.5° C: An IPCC special report on the impacts of global warming of 1.5° C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty. Intergovernmental Panel on Climate Change.
- Chaturvedi, T., Christiansen, A. H. C., Gołębiewska, I., & Thomsen, M. H. (2021). *Salicornia* species: current status and future potential. In *Future of Sustainable Agriculture in Saline Environments*. <https://doi.org/10.1201/9781003112327-31>
- Choi, D., Lim, G. S., Piao, Y. L., Choi, O. Y., Cho, K. A., Park, C. B., ... & Cho, H. (2014). Characterization, stability, and antioxidant activity of *Salicornia herbacea* seed oil. *Korean Journal of Chemical Engineering*, 31, 2221–2228. <https://doi.org/10.1007/s11814-014-0163-7>
- Collinson, P., Young, I., Antal, L., & Macbeth, H. (Eds.). (2022). *Food and Sustainability in the Twenty-First Century: Cross-Disciplinary Perspectives*. Berghahn Books.
- Costa, C. S. B., Chaves, F. C., Rombaldi, C. V., & Souza, C. R. (2018). Bioactive compounds and antioxidant activity of three biotypes of the sea asparagus *Sarcocornia ambigua* (Michx.) MA Alonso & MB Crespo: A halophytic crop for cultivation with shrimp farm effluent. *South African Journal of Botany*, 117, 95–100.
- Cuttelod, A., García, N., Malak, D. A., Temple, H. J., & Katariya, V. (2009). The Mediterranean: A biodiversity hotspot under threat. *Wildlife in a Changing World—an Analysis of the 2008 IUCN Red List of Threatened Species*, 89(2019), 9.
- De Souza, M. M., Mendes, C. R., Doncato, K. B., Badiale-Furlong, E., & Costa, C. S. B. (2018). Growth, phenolics, photosynthetic pigments, and antioxidant response of two new genotypes of sea asparagus (*Salicornia neei* Lag.) to salinity under greenhouse and field conditions. *Agriculture*, 8(7), 115. <https://doi.org/10.3390/agriculture8070115>
- Debez, A., Huchzermeyer, B., Abdelly, C., & Koyro, H.-W. (2011). Current challenges and future opportunities for sustainable utilization of halophytes. *Sabkha Ecosystems: Volume III: Africa and Southern Europe*, 59–77.
- Ekanayake, S., Egodawatta, C., Attanayake, R. N., & Perera, D. (2023). From salt pan to saucepan: *Salicornia*, a halophytic vegetable with an array of potential health benefits. *Food Frontiers*, 4, 641–676. <https://doi.org/10.1002/fft2.214>
- Fawzy, S., Osman, A. I., Doran, J., & Rooney, D. W. (2020). Strategies for mitigation of climate change: a review. *Environmental Chemistry Letters*, 18, 2069–2094. <https://doi.org/10.1007/s10311-020-01059-w>
- Ganesan, M., Trivedi, N., Gupta, V., Madhav, S. V., Radhakrishna Reddy, C., & Levine, I. A. (2019). Seaweed resources in India—current status of diversity and cultivation: prospects and challenges. *Botanica Marina*, 62(5), 463–482. <https://doi.org/10.1515/bot-2018-0056>
- Gelfand, I., Sahajpal, R., Zhang, X., Izaurralde, R. C., Gross, K. L., & Robertson, G. P. (2013). Sustainable bioenergy production from marginal lands in the US Midwest. *Nature*, 493(7433), 514–517. <https://doi.org/10.1038/nature11811>
- Gispert, M., Kuliush, T., Dyachenko, L., Kharytonov, M., Emran, M., Verdaguer, D., Llorens, L., & Carrasco-Barea, L. (2021). Appraising soil carbon storage potential under perennial and annual Chenopodiaceae in the salt marsh of NE Spain. *Estuarine, Coastal and Shelf Science*, 252, 107240.
- Glenn, E. P., Brown, J. J., & Blumwald, E. (1999). Salt tolerance and crop potential of halophytes. *Critical Reviews in Plant Sciences*, 18(2), 227–255. <https://doi.org/10.1080/07352689991309207>
- Glenn, E. P., O'Leary, J. W., Watson, M. C., Thompson, T. L., & Kuehl, R. O. (1991). *Salicornia bigelovii* Torr.: an oilseed halophyte for seawater irrigation. *Science*, 251(4997), 1065–1067. <https://doi.org/10.1126/science.251.4997.1065>
- Gopi, M., Kumar, M. P., Jeevamani, J. J. J., Raja, S., Muruganandam, R., Samuel, V. D., Simon, N. T., Viswanathan, C., Abhilash, K. R., & Krishnan, P. (2019). Distribution and biodiversity of tropical salt marshes: Tamil Nadu and Puducherry, Southeast coast of India. *Estuarine, Coastal and Shelf Science*, 229, 106393. <https://doi.org/10.1016/j.ecss.2019.106393>

- Gouda, M. S., & Elsebaie, E. M. (2016). Glasswort (*Salicornia* spp) as a source of bioactive compounds and its health benefits: a review. *Alexandria Journal of Food Science and Technology*, 13(1), 1–7. <https://dx.doi.org/10.12816/0038461>
- Grattan, S. R., Benes, S. E., Peters, D. W., & Diaz, F. (2008). Feasibility of irrigating pickleweed (*Salicornia bigelovii* Torr) with hyper-saline drainage water. *Journal of Environmental Quality*, 37(S5), S-149. <https://doi.org/10.2134/jeq2007.0450>
- Group, W. B. (2016). The World Bank Group A to Z 2016. World Bank Publications.
- Guerin, L. J., & Guerin, T. F. (1994). Constraints to the adoption of innovations in agricultural research and environmental management: a review. *Australian Journal of Experimental Agriculture*, 34(4), 549–571. <https://doi.org/10.1071/EA9940549>
- Gunning, D. (2016). Cultivating *Salicornia europaea* (marsh samphire). Dublin, Ireland: Irish Sea Fisheries Board, 4, 1–95.
- Hamed, K. B., Castagna, A., Ranieri, A., Garcia-Caparros, P., Santin, M., Hernandez, J. A., & Espin, G. B. (2021). Halophyte based Mediterranean agriculture in the contexts of food insecurity and global climate change. *Environmental and Experimental Botany*, 191, 104601. <https://doi.org/10.1016/j.envexpbot.2021.104601>
- Herbert, E. R., Boon, P., Burgin, A. J., Neubauer, S. C., Franklin, R. B., Ardón, M., ... & Gell, P. (2015). A global perspective on wetland salinization: ecological consequences of a growing threat to freshwater wetlands. *Ecosphere*, 4(10), 1–43. <https://doi.org/10.1890/ES14-00534.1>
- Holguin Pena, R. J., Medina Hernandez, D., Ghasemi, M., & Rueda Puente, E. O. (2021). Salt tolerant plants as a valuable resource for sustainable food production in arid and saline coastal zones. *Acta Biológica Colombiana*, 26(1), 116–126. <https://doi.org/10.15446/abc.v26n1.82412>
- Isca, V., Seca, A. M., Pinto, D. C., & Silva, A. (2014). An overview of *Salicornia* genus: The phytochemical and pharmacological profile. *Natural Products: Research Reviews*, 2, 145–164. <http://hdl.handle.net/10400.3/2374>
- Joshi, A., Kanthaliya, B., & Arora, J. (2020). Halophytes: The nonconventional crops as source of biofuel production. Handbook of halophytes: From molecules to ecosystems towards biosaline agriculture, 1–28.
- Khan, M. A., Böer, B., Öztürk, M., Al Abdessalaam, T. Z., Clüsener-Godt, M., & Gul, B. (2014). Sabkha ecosystems: Volume IV: cash crop halophyte and biodiversity conservation (Vol. 47). Springer.
- Khan, M. A., Gul, B., & Weber, D. J. (2000). Germination responses of *Salicornia rubra* to temperature and salinity. *Journal of Arid Environments*, 45(3), 207–214. <https://doi.org/10.1006/jare.2000.0640>
- Loconsole, D., Cristiano, G., & De Lucia, B. (2019). Glassworts: From wild salt marsh species to sustainable edible crops. *Agriculture*, 9(1), 14. <https://doi.org/10.3390/agriculture9010014>
- Machado, R. M. A., & Serralheiro, R. P. (2017). Soil salinity: Effect on vegetable crop growth. Management practices to prevent and mitigate soil salinization. *Horticulturae*, 3(2), 30. <https://doi.org/10.3390/horticulturae3020030>
- Makkawi, Y., El Sayed, Y., Lyra, D. A., Pour, F. H., Khan, M., & Badrelzaman, M. (2021). Assessment of the pyrolysis products from halophyte *Salicornia bigelovii* cultivated in a desert environment. *Fuel*, 290, 119518.
- Martinez-Garcia, R. (2010). Physiological studies of the halophyte *Salicornia bigelovii*. A potential food and biofuel crop for integrated aquaculture-agriculture systems (Doctoral dissertation, The University of Arizona).
- Mishra, A., & Tanna, B. (2017). Halophytes: Potential resources for salt stress tolerance genes and promoters. *Frontiers in Plant Science*, 8, 829. <https://doi.org/10.3389/fpls.2017.00829>
- Mohammadi, H., & Kardan, J. (2015). Morphological and physiological responses of some halophytes to salinity stress. *Annales Universitatis Mariae Curie-Sklodowska, Sectio C–Biologia*, 70(2).
- Mroczek, A. (2015). Phytochemistry and bioactivity of triterpene saponins from Amaranthaceae family. *Phytochemistry Reviews*, 14, 577–605. <https://doi.org/10.1007/s11101-015-9394-4>
- Muscolo, A., Panuccio, M. R., & Piernik, A. (2014). Ecology, Distribution, and Ecophysiology of *Salicornia Europaea* L. Sabkha Ecosystems: Volume IV: Cash Crop Halophyte and Biodiversity Conservation, 233–240.

- Negacz, K, Bruning, B., & Vellinga, P. (2021). Achieving Multiple Sustainable Development Goals through Saline Agriculture. *Future of Sustainable Agriculture in Saline Environments*, 13-28.
- Nigam, P. S., & Singh, A. (2011). Production of liquid biofuels from renewable resources. *Progress in Energy and Combustion Science*, 37(1), 52-68. <https://doi.org/10.1016/j.peccs.2010.01.003>
- Ozturk, M., Altay, V., Orçen, N., Yaprak, A. E., Tuğ, G. N., & Güvensen, A. (2018). A little-known and a little-consumed natural resource: Salicornia. *Global Perspectives on Underutilized Crops*, 83-108.
- Parida, A. K., & Das, A. B. (2005). Salt tolerance and salinity effects on plants: a review. *Ecotoxicology and environmental Safety*, 60(3), 324-349. <https://doi.org/10.1016/j.ecoenv.2004.06.010>
- Patel, S. (2016). Salicornia: Evaluating the halophytic extremophile as a food and a pharmaceutical candidate. *Biotech*, 4(1), 104. <https://doi.org/10.1007/s13205-016-0418-6>
- Qadir, M., Quill rou, E., Nangia, V., Murtaza, G., Singh, M., Thomas, R. J., Drechsel, P., & Noble, A. D. (2014). Economics of salt-induced land degradation and restoration. *Natural Resources Forum*, 38(4), 282-295. <https://doi.org/10.1111/1477-8947.12054>
- Rajabi Dehnavi, A., Zahedi, M., Ludwiczak, A., Cardenas Perez, S., & Piernik, A. (2020). Effect of salinity on seed germination and seedling development of sorghum (*Sorghum bicolor* (L.) Moench) genotypes. *Agronomy*, 10(6), 859. <https://doi.org/10.3390/agronomy10060859>
- Rathore, M. S., Balar, N., & Jha, B. (2019). Population structure and developmental stage-associated ecophysiological responses of *Salicornia brachiata*. *Ecological Research*, 34(5), 644-658. <https://doi.org/10.1111/1440-1703.12033>
- Rey, Jorge R., et al. "Above-ground primary production in impounded, ditched, and natural Batis-Salicornia marshes along the Indian River Lagoon, Florida, USA." *Wetlands* 10.2 (1990): 151-171. <https://doi.org/10.1007/BF03160830>
- Rozentsvet, O. A., Nesterov, V. N., & Bogdanova, E. S. (2017). Structural, physiological, and biochemical aspects of salinity tolerance of halophytes. *Russian Journal of Plant Physiology*, 64, 464-477. <https://doi.org/10.1134/S1021443717040112>
- Safdar, H., Amin, A., Shafiq, Y., Ali, A., Yasin, R., Shoukat, A., Hussan, M. U., & Sarwar, M. I. (2019). A review: Impact of salinity on plant growth. *Nature and Science*, v. 17. <https://doi.org/10.7537/marsnsj170119.06>.
- Sani, Y. M., Daud, W. M. A. W., & Aziz, A. A. (2012). Biodiesel feedstock and production technologies: Successes, challenges and prospects. *Biodiesel-Feedstocks, Production, and Applications*, 10, 52790.
- Sharma, R., Wungrampha, S., Singh, V., Pareek, A., & Sharma, M. K. (2016). Halophytes as bioenergy crops. *Frontiers in Plant Science*, 7, 1372. <https://doi.org/10.3389/fpls.2016.01372>
- Singh, D., Buhmann, A. K., Flowers, T. J., Seal, C. E., & Papenbrock, J. (2014). *Salicornia* as a crop plant in temperate regions: Selection of genetically characterized ecotypes and optimization of their cultivation conditions. *AoB Plants*, 6. <https://doi.org/10.1093/aobpla/plu071>
- Song, J., Fan, H., Zhao, Y., Jia, Y., Du, X., & Wang, B. (2008). Effect of salinity on germination, seedling emergence, seedling growth, and ion accumulation of a halophyte *Suaeda salsa* in an intertidal zone and on saline inland. *Aquatic Botany*, 88(4), 331-337. <https://doi.org/10.1016/j.aquabot.2007.11.004>
- Turcios, A. E., Cayenne, A., Uellendahl, H., & Papenbrock, J. (2021). Halophyte plants and their residues as feedstock for biogas production—Chances and challenges. *Applied Sciences*, 11(6), 2746. <https://doi.org/10.3390/app11062746>
- Turyasingura, B. (2022). Review on the Impacts of Climate Change on the Plant Water Interactions. *International Journal of Academic Multidisciplinary Research (IJAMR)* ISSN: 2643-9670, 6(7), 223.
- Turyasingura, B., & Chavula, P. (2022). Climate-Smart Agricultural Extension Service Innovation Approaches in Uganda: Review Paper. *International Journal of Food Science and Agriculture*, 4(1), 35-43. <https://doi.org/10.26855/ijfsa.2022.03.006>
- Turyasingura, B., Alex, S., Hirwa, H., & Mohammed, F. S. (2022). Wetland conservation and management practices in Rubanda District, South-Western Uganda. <https://doi.org/10.21203/rs.3.rs-1876968/v1>

- Turyasingura, B., Ayiga, N., & Benzougagh, B. (2022). Re-thinking on land degradation and its impacts on livelihoods of the farmers in Kanungu District, Uganda. <https://doi.org/10.21203/rs.3.rs-1966742/v1>
- Turyasingura, B., Chavula, P., Hirwa, H., Mohammed, F. S., Ayiga, N., Bojago, E., Benzougagh, B., & Ngabirano, H. (2022). A Systematic Review and Meta-analysis of Climate Change and Water Resources in Sub-Sahara Africa. <https://doi.org/10.21203/rs.3.rs-2281917/v1>
- Turyasingura, B., Mwanjalolo, M., & Ayiga, N. (2022). Diversity at Landscape Level to Increase Resilience. A Review. *East African Journal of Environment and Natural Resources*, 5(1), 174–181. <https://doi.org/10.37284/eajenr.5.1.723>
- Turyasingura, B., Tumwesigye, W., Atuhaire, A., Tumushabe, J. T., & Akatwijuka, R. (2023). A literature review of climate-smart landscapes as a tool in soil-water management in Sub-Saharan Africa. <https://nru.uncst.go.ug/handle/123456789/8144>
- Urbano, M., Tomaselli, V., Bisignano, V., Veronico, G., Hammer, K., & Laghetti, G. (2017). *Salicornia patula* Duval-Jouve: From gathering of wild plants to some attempts of cultivation in Apulia region (southern Italy). *Genetic Resources and Crop Evolution*, 64, 1465–1472. <https://doi.org/10.1007/s10722-017-0521-5>
- USDA FoodData Central (2019). "National Nutrient Database for Standard Reference Legacy Release." United States Department of Agriculture, Agricultural Research Service. Available at: <https://fdc.nal.usda.gov/>
- Ventura, Y., & Sagi, M. (2013). Halophyte crop cultivation: the case for *Salicornia* and *Sarcocornia*. *Environmental and Experimental Botany*, 92, 144–153. <https://doi.org/10.1016/j.envexpbot.2012.07.010>
- Ventura, Y., Wuddineh, W. A., Myrzabayeva, M., Alikulov, Z., Khozin-Goldberg, I., Shpigel, M., Samocha, T. M., & Sagi, M. (2011a). Effect of seawater concentration on the productivity and nutritional value of annual *Salicornia* and perennial *Sarcocornia* halophytes as leafy vegetable crops. *Scientia Horticulturae*, 129(3), 189–196. <https://doi.org/10.1016/j.scienta.2011.02.001>
- Ventura, Y., Wuddineh, W. A., Shpigel, M., Samocha, T. M., Klim, B. C., Cohen, S., Shemer, Z., Santos, R., & Sagi, M. (2011b). Effects of day length on flowering and yield production of *Salicornia* and *Sarcocornia* species. *Scientia Horticulturae*, 130(3), 510–516. <https://doi.org/10.1016/j.scienta.2011.08.008>
- Volkov, V. (2015). Salinity tolerance in plants. Quantitative approach to ion transport starting from halophytes and stepping to genetic and protein engineering for manipulating ion fluxes. *Frontiers in Plant Science*, 6, 873. <https://doi.org/10.3389/fpls.2015.00873>
- Wang, X., Fan, P., Song, H., Chen, X., Li, X., & Li, Y. (2009). Comparative proteomic analysis of differentially expressed proteins in shoots of *Salicornia europaea* under different salinity. *Journal of Proteome Research*, 8(7), 3331–3345. <https://doi.org/10.1021/pr801083a>
- Wratten, S. D., Sandhu, H., Cullen, R., & Costanza, R. (Eds.). (2013). *Ecosystem services in agricultural and urban landscapes* (Vol. 152). Oxford: Wiley-Blackwell. <https://doi.org/10.1002/9781118506271.ch8>
- Yuan, F., Guo, J., Shabala, S., & Wang, B. (2019). Reproductive physiology of halophytes: Current standing. *Frontiers in Plant Science*, 9, 1954. <https://doi.org/10.3389/fpls.2018.01954>
- Zedler, J. B. (1996). Coastal mitigation in southern California: the need for a regional restoration strategy. *Ecological Applications*, 6(1), 84–93. <https://doi.org/10.2307/2269555>