

- Lake R, Hudson A, Cressey P, Gilbert S, 2007: Risk profile Campylobacter Jejuni/coli in red meat. New Zealand Food Safety Authority, http://www.foodsafety.govt.nz/elibrary/industry/Risk_Profile_Campylobacter_Jejuni_Coli-Science_Research.pdf, Accessed: 12.11.2014.
- Lawley R, 2013: Campylobacter. Food Safety Watch, <http://www.foodsafetywatch.org/factsheets/campylobacter/> Accessed: 06.10.2014.
- Leonard EE, Tompkins LS, Falkow S, Nachamkin I, 2004: Comparison of Campylobacter jejuni isolates implicated in Guillain-Barre´ Syndrome and strains that cause enteritis by a DNA microarray. *Infect Immun*, 72 (2), 1199-1203.
- Liu L, Hussain SK, Miller RS, Oyarzabal OA, 2009: Efficacy of Mini VIDAS for the detection of Campylobacter spp. from retail broiler meat enriched in bolton broth with or without the supplementation of blood. *J Food Protect*, 72 (11), 2428-2432.
- Madden RH, Espie WE, Moran L, McBride J, Scates P, 2001: Occurrence of Escherichia coli O157:H7, Listeria monocytogenes, Salmonella and Campylobacter spp. on beef carcasses in Northern Ireland. *Meat Sci*, 58 (4), 343-6.
- OIE, 2008: Manuel of diagnostic tests and vaccines for terrestrial animals. 5th ed. 1185-1191.
- Oktay İ, Heperkan D, 2006: Listeria ve salmonella tayininde ISO yönteminin VIDAS otomatik sistemi ile karşılaştırılması. *Mikrobiyol Derg*, 4, 12.
- Omar A, Oyarzabal OA, Battie C, 2012: Biochemistry, Genetics and Molecular Biology Trends in Immunolabelled and Related Techniques. In, , "Immunological Methods for the Detection of Campylobacter spp. Current Applications and Potential Use in Biosensors" Ed., Eltayb Abuelzein., InTech.
- Paulsen P, Kanzler P, Hilbert F, Mayrhofer S, Baumgartner S, Smulders FGM, 2005: Comparison of three methods for detecting Campylobacter spp. in chilled or frozen meat. *Int J Food Microbiol*, 103 (2), 229-233.
- Pearson AD, Greenwood M, Healing TD, Rollins D, Shahamet M, Donaldson J, Colwell RR, 1993: Colonization of broiler chickens by waterborne Campylobacter jejuni. *Appl Environ Microbiol*, 59 (4), 987-996.
- Reiter GM, Lopez C, Jordano R, Medina LM, 2010: Comparative study of alternative methods for food safety control in poultry slaughterhouses. *Food Anal Meth*, 3 (3), 253-260.
- Reiter MG, Bueno CM, Lopez C, Jordano R, 2005: Occurrence of Campylobacter and Listeria monocytogenes in a poultry processing plant. *J Food Protec*, 68 (9), 1903-1906.
- Zhao C, Ge B, Villena JD, Sudler R, Yeh E, Zhao S, White DG, Wagner D, Meng J, 2001: Prevalence of Campylobacter spp., Escherichia coli, and Salmonella serovars in retail chicken, turkey, pork, and beef from the Greater Washington, D.C., Area. *Appl Environ Microbiol*, 67 (12), 5431-5436.

***Yazışma Adresi:** Semra GÜRBÜZ

Mardin Artuklu Üniversitesi, Gastronomi ve Mutfak Sanatları Bölümü, Mardin, Türkiye.

e-mail:semragurbuz@gmail.com