

**Gaziosmanpaşa Bilimsel Araştırma Dergisi**

Dergiye Geliş Tarihi: 19.03.2015

Yayına Kabul Tarihi: 02.12.2015

Baş Editör: Bilge Hilal ÇADIRCI

Alan Editörü: İzzet KADIOĞLU

İstanbul Park ve Bahçelerindeki Tetranychidae Türleri

Ayşe YEŞİLAYER^{a,1} (ayse.yesilayer@gop.edu.tr)
Sultan COBANOĞLU^b (coban.sultan@gmail.com)

^aGOP Üniversitesi, Ziraat Fakültesi Bitki Koruma Bölümü, 60216 Tokat
^bAnkara Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, 06116 Ankara

Özet – Bu çalışma, İstanbul ili park ve bahçelerinde, yaprağını döken ağaçlar, ibrelierler ve çalılarda bulunan fitofag tetranychidleri belirlemek amacıyla yapılmıştır. 2006-2008 yıllarında haftalık olarak yapılan sörveyler sonucunda, 10 adet tetranychid tür tespit edilmiştir. Bu türlerden bir tanesi, *Platyteanychus libocedri* McGregor (Acar: Tetranychidae) Türkiye faunası için yeni kayıt niteligidendir.

Anahtar Kelimeler –
Platyteanychus libocedri,
Eotetranychus bryobia,
süs bitkileri, İstanbul, Türkiye

Gaziosmanpaşa Journal of Scientific Research 11 (2015) 90-98

Tetranychidae (Acari: Prostigmata) species from parks and ornamental plants in Istanbul, Turkey

Abstract – This research was conducted to determine phytophagous tetranychid mites occurring on deciduous trees, conifers and shrubs in the parks and ornamental plants of İstanbul province. Weekly surveys were carried out between 2006 and 2008. As a result of the surveys, 10 Tetranychidae species were identified. One of these species, *Platyteanychus libocedri* McGregor (Tetranychidae) was the first record for the Tetranychidae fauna of Turkey.

Received: 19.03.2015

Keywords –
Platyteanychus libocedri,
Eotetranychus bryobia,
ornamental plants,
Istanbul, Turkey

Accepted: 02.12.2015

1. Introduction

There are many reports available on mites that are parasitic to plants and cause economic damage on the ornamental plants throughout the world (Jeppson *et al.*, 1975; Kropczynska *et al.*, 1992; Ripka, 1997, 1998; Ripka *et al.*, 2005; Mesa *et al.*, 2009). In Turkey, different phytophagous mite species were reported from ornamental plants (Yüksel, 1999; Uysal *et al.*, 2001; Cobanoğlu *et al.*, 2003; Elma and Alaoglu, 2008; Yeşilayer and Cobanoğlu, 2009, 2011, 2013).

Tetranychidae (Acari: Prostigmata), spider mites, is the most important families of the Acari because many species can be serious pests for ornamental and agricultural crops

(Jeppson *et al.*, 1975). There are 70 genera of tetranychid mites in the world, and these contain 1275 species (Hoy, 2011; Migeon and Dorkeld, 2006-2013). The serious economic damage caused by spider mites is because of their large host plant range, high fecundity and rapid developmental rates (Smiley and Baker, 1995).

However, prior to this study, there was no detail data available on the tetranychid mites of the Istanbul province, located on the Marmara coast. There are many studies about predatory mites in different locations in Turkey and Istanbul (Uysal *et al.*, 2001; Cobanoğlu *et al.*, 2003; Yeşilayer and Cobanoğlu, 2011, 2013). The objective of this study was to determine and make a quantitative assessment of phytophagous tetranychid mites occurring on deciduous trees, conifers and shrubs in the parks and ornamental plants of Istanbul province between 2006 and 2008.

2. Materials and Methods

Tetranychids were surveyed in parks and urban areas of Istanbul at weekly intervals between April to October in 2006 and 2008. Samples were taken in different localities (Kadıköy, Küçükçekmece, Tuzla, Beşiktaş, Bakırköy, Florya, Fatih, Sultanahmet, Büyükçekmece) mainly from unsprayed areas during the growing seasons in Istanbul. They were transferred to the laboratory in an icebox. In total, 1200 samples were taken from branches and leaves of ornamental plants. Mites were removed from the leaves under a stereomicroscope and extracted using Berlese funnels. The specimens were preserved in 70% alcohol. After they were cleaned in lactophenol solution, they were mounted in Hoyer's medium for identification. The slides were kept for 2-4 weeks at 35°C in an incubator (Düzungüneş, 1980). The slides of the mounted specimens were deposited in both authors' collection at the University of Ankara, Department of Plant Protection, Ankara and University of GOP (University of Gazi Osman Paşa, Tokat) Department of Plant Protection, Tokat. For identification, following references were used: Pritchard and Baker (1955), Jeppson *et al.* (1975) and Seeman and Beard (2011).

3. Results and Discussion

Ten species belonging to the family Tetranychidae were identified; *Tetranychus urticae* Koch, *Panonychus ulmi* Koch, *Oligonychus ununguis* Jacobi, *Oligonychus coniferarum* McGregor, *Eotetranychus populi* Koch, *Eotetranychus uncatus* Garman, *Eotetranychus carpini* Oudemans, *Bryobia rubrioculus* (Scheuten), *Bryobia praetiosa* Koch, and *Platyteanychus libocedri* McGregor. The latter one is a new record for Turkey. *Tetranychus urticae*, *O. ununguis* and *Platyteanychus* sp. most common harmful mite species in Istanbul parks (Yeşilayer and Cobanoğlu 2011).

Magnolia soulangeana L. (Magnoliaceae) was the most preferred and populated host plant (40.7%) while *Picea pungens* (Engelm) Pinaceae was rarely populated by mites (0.39%).

3.1.Tetranychidae

3.1.1. *Tetranychus urticae* Koch, 1836

Material examined: Yıldız Parkı: *Quercus robur* L. 01.11.2006 (3♀♀); *Acer negundo* L. (15♀♀); İstanbul Sabahattin Zaim University: *Pinus pinea* L. 13.09.2007 (1♂); *Cupressus arizonica* Green 25.09.2007 (1♀); Halkalı: *Pitoporum tobira* Thunb 01.10.2007 (3♀♀V); *Prunus avium* L. 10.11.2008 (1 nymph); Aliağaoğlu Nursery: *Chamaerops excelsa* Hort 17.07.2007 (1♀); Ada Park: *Lagostermia indica* L. 23.07.2007 (1♀), *Cupressus*

macrocarpha Hartw. 23.10.2007 (1♀); Haliç Hospital Garden: *Erobryria japonica* L. (Thunb.) 21.10.2007 (1♀); Zirai Karantina Province: *Magnolia soulangeana* L. 17.06.2008 (17♀, 2♂, 3 nymphs, 1 larva), *Prunus domestica* L. 10.11.2008 (1♀, 2♂); Özgürk parkı: *Ulmus* sp. 04.07.2008 (5♀, 3♂, 4 nymphs).

Comments: *T. urticae* is a very common and well known harmful mite species globally. It has been reported from Afghanistan, Algeria, America, Australia, Belgium, Bulgaria, Canada, Denmark, Finland, France, Germany, Holland, Hungary, Iran, Iraq, Italy, Korea, Lebanon, Lithuania, Mexico, Morocco, New Zealand, Norway, Pakistan, Portugal, Sri Lanka, Syria, Spain, Yugoslavia, Yemen, Turkey (Zhang and Henderson, 2002; Migeon and Dorkeld, 2006-2013). This species was reported from *Buddleia* sp., *Capsicum annum*, *Chrysanthemum* sp., *Citrus* spp., *Cucumis sativus*, *Dianthus caryophyllus*, *Fuchsia* sp., *Gardenia* sp., *Glycine max*, *Hydrangea* sp., *Juglans regia*, *Prunus* sp., *Salix* sp., *Solanum* sp., *Ulmus* spp. (Zhang and Henderson, 2002). It was previously reported from cultural, ornamental plants and weeds. This is a very common and cosmopolitan species in Turkey (Düzungüneş 1954; Aydemir and Toros, 1990; Uysal et al., 2001).

3.1.2. *Bryobia rubrioculus* (Scheuten), 1857

Material examined: Bahçeşehir: *Pinus nigra* L. 11.12.2007 (1♀); Dostlar Park: *Nerium oleander* L. (1♀).

Comments: *B. rubrioculus* was reported from Austria, Belgium, Cyprus, England, Finland, France, Germany, Ireland, Italy, Netherlands, Poland, Portugal, Spain, Sweden, Switzerland, and Turkey (Anonymous, 2008). This species was previously recorded on *Malus*, *Prunus*, *Juglans*, *Medicago sativa*, *Trifolium* sp., *Hedera helix* and *Rubus* sp. (İyriboz, 1940; Düzungüneş, 1954; Yiğit and Uygun, 1982; Erol and Yaşar, 1996). *B. rubrioculus* was also recorded on *Cotoneaster horizontalis* and *Malus floribunda* (Uysal et al., 2001),

3.1.3. *Bryobia pretiosa* Koch, 1836

Material examined: Dostlar Parkı: *Picea pungens* L., 11.08.2006 (1♀).

Comments: This species was previously recorded on deciduous trees in USA, Europe, Greece and Japan (Anonymous, 2008). *B. pretiosa* was also recorded on *Muscari longipes* in Ankara, Turkey (Bayram and Cobanoğlu, 2006). It was recorded on *Fragaria ananassa* Duch. in Izmir (Madanlar and Yoldaş 1996); *Malus communis* L. in Van (Erol and Yaşar, 1996), *Ficus carica* in Izmir (İyriboz, 1940)

3.1.4. *Panonychus ulmi* Koch, 1929

Material examined: Sultanahmet: *Pittosporum tobira* (Thunb) 01.10.2007 (1♀).

Comments: *P. ulmi* was reported from deciduous trees in the USA, Argentina, Bermuda, Canada, China, Georgia, Hungary, India, Iran, Japan, Morocco, Russia, South Africa, Spain, Sweden, Switzerland, Uruguay, Venezuela and Yugoslavia (Jeppson et al., 1975; Zhang and Henderson, 2002).

It was recorded on *M. communis*, *Prunus persica* L., *Prunus domestica* L., *Pyrus communis* L., *Cydonia oblonga* Miller, *Prunus avium* L., *Prunus cerasus* L., *Corylus avellana* L., *Ficus carica* L., in Amasya, İçel, Bursa, Yalova, Thrace region, Van, Adana and Niğde (Düzungüneş, 1963; Göksu and Atak 1968; Yiğit and Uygun, 1982; Ulusoy et al., 1999;

İncekulak and Ecevit, 2002; Akşit *et al.*, 2003; Çakmak and Akşit, 2003; Gencer *et al.*, 2002; Kasap *et al.*, 2004)

3.1.5. *Oligonychus ununguis* (Jacobi, 1920)

Material examined: Ozgürlik Parkı: *P. pungens* 06.06.2006, 31.08.2006 (1♀, 1 larva); *Euonymus fortunei* L. (Celastraceae) 03.07.2006 (1♀); *Cedrus atlantica* Endl. 31.08.2006 (1♀); *E. fortunei* 29.09.2006 (1♀); İstanbul Sabahattin Zaim University: *Berberis* sp. 13.09.2006 (1♀); Başakşehir Hospital: *Picea orientalis* L. 01.10.2007 (2♀); İkitelli Housing: *Aesculus hippocastanum* L. 27.09.2007 (1♂); *Ulmus* sp. 04.07.2008 (1♀); Halkalı: *Prunus domestica* L. 10.11.2008 (1♀).

Comments: *O. unungius* was recorded on *P. pungens*, *Pinus*, *Juniperus*, *Thuja*, *Alnus*, *Taxus*, *Cedrus*, *Abies* spp., *A. hippocastanum*, *Buxus sempervirens* L. and, *Quercus* sp. (Jeppson *et al.*, 1975). This species was reported from *Picea* sp. in the East Blacksea region (Yüksel and Ulusoy, 1999).

3.1.6. *Oligonychus coniferarum* (McGregor), 1950

Material examined: Atatürk Park: *P. pinea* 19.10.2006 (1♀); İkitelli: *A. hippocastaneum* 27.09.2007 (1♀).

Comments: *O. coniferarum* was found on *Cupressus sempervirens* L., *Cupressus lusitanica* Mill., *Juniperus depeana*, *Juniperus* sp. (Pritchard and Baker, 1955; Tuttle and Baker, 1968; Smiley and Baker, 1995).

This species was recorded on *P. sylvestris* in Erzurum (Ecevit, 1977); *Cornus alba* Sibirica, *Juniperus horizontalis* Moench, *Juniperus sabina* L. in Ankara (Uysal *et al.*, 2001). It was also recorded on *P. nigra* (needle), *P. sylvestris*, *P. pungens* in Turkey (Bayram and Çobanoğlu, 2007) and *Quercus* sp. in Konya (Elma and Alaoglu, 2008).

3.1.7. *Eotetranychus populi* (Koch), 1838

Material examined: Dostlar Park: *Ligustrum vulgare* Aurea 11.07.2006 (1♀); Yıldız Park: *Laurus nobilis* L. 01.11.2006 (1♀); Yeniçiftlik: *Salix babylonica* L. 13.10.2007 (1♀).

Comments: *E. populi* was recorded on poplar and willow trees in Russia, Serbia and England (Jeppson *et al.*, 1975). This species was found on *Salix* sp. and *Platanus orientalis* L. in Konya, Turkey (Elma and Aloğlu, 2008).

3.1.8. *Eotetranychus carpini* Oudemans, 1905

Material examined: Belgrad Forest: *Carpinus betulus* L. 23.10.2007 (2♀, 1♂); İstanbul Sabahattin Zaim University: *P. domestica* 10.11.2008 (1♀).

Comments: This species was recorded on *Alnus* sp., *Carpinus* sp., *Fagus* sp., *Quercus* sp., and *Vitis vinifera* L. (Hatzinikolis, 1970; Jeppson *et al.*, 1975). *E. carpini* was found on *Ficus* sp. in Gaziantep (Düzgüneş, 1965), Chestnut in West Anatolia (Önuçar and Ulu, 1988), and on *Berberis thunbergii* DC in Ankara (Uysal *et al.*, 2001).

3.1.9. *Eotetranychus uncatus* (Garman), 1952

Material examined: Sabahattin Zaim University: *P. domestica*, 10.11.2008 (1♀).

Comments: *E. uncatus* was found *Acer campestre* Binazzi and Blackman, *Acer negundo* L., *Acer platanoides* L., *Alnus* sp., *Betula alba* L., *Malus domestica* L. (Pritchard and

Baker, 1952; Reeves, 1963, Dobosz *et al.*, 1995). In Turkey, this species was recorded on Chestnut and apple trees (Düzungüneş 1963; Önuçar and Ulu, 1988; Yanar and Ecevit, 2005).

3.1.10. *Platytranychus libocedri* (McGregor, 1936)

Synonyms:

Eotetranychus libocedri (McGregor, 1936)

Mononychus libocedri (McGregor, 1936)

Tetranychus libocedri McGregor, 1936

Platytranychus libocedri has short dorsocentral hysterosomals and dorsolateral hysterosomals are longer than dorsocentrals (Figure 1). On females, Tibia I nine; tarsus I five and tibia II five bear tactile setae.

The aedeagus has a long and narrowing rounded and emarginated at the tip (Figure 2)

P. libocedri is similar *P. thujae* (McGregor). *P. thujae*'s dorsal setae longer than *P. libocedri*.




Figure 1. *Platytranychus libocedri* McGregor female




Figure 2. *Platyteranychus libocedri* McGregor aedeagus

Material examined University of Istanbul Sabahattin Zaim: *E. fortunei* 29.09.2006 (2♀♀); Özgür Park: *Thuja occidentalis* Aurea 06.06.2007 (1♀), *C. arizonica* 18.10.2007 (1♀, 1 larva, 1 nymph); Park of Ada: *C. macrocarpha* 23.10.2007 (3♀♀, 1 nymph); Kurtköy Nursery: *Cupressocyparis leylandii* L., 12.12.2007 (1♀); Istanbul Sabahattin Zaim University: *C. arizonica* (litter) 10.11.2008 (2♂♂, 2 nymph); Özgür Park: *E. fortunei* 29.09.2006 (1♀).

This is a new record for Turkey.

Comments: Specimens were recorded on *Thuja occidentalis* L. and it was also found in Utah on junipers (Pritchard and Baker, 1955). This species was recorded on *C. arizonica* *Cupressus* sp., *C. sempervirens* *Juniperus* sp., *Libocedrus decurrens* Torrey, *Thuja* sp., *Pinus* sp., *Abies religiosa* (Kunth.) Schltdl. and Cham., *Tamarix* sp. (Pritchard and Baker, 1955; Tuttle and Baker, 1964).

CONCLUSION

From the surveys of the parks and ornamental plants of Istanbul, 10 Tetranychidae species were identified. *P. libocedri* was first reported for the mite fauna of Turkey. It is necessary to research on the biology, morphology, distribution, symptomatology and control measures of this species. In this study *T. urticae* and *O. unungius* are the most common phytophagous mite species on the parks and ornamental plants in Istanbul. It is also very important to study natural enemies, and their importance in the integrated pest control of this phytophagous mite species..

References

- Anonymous 2008. <http://www.faunaeur.org>, 2008.02.08.2014.
- Akşit T., F. Özsemerci, and İ. Çakmak 2003. Aydin ilinde incir ağaçlarında saptanan zararlı türler. Türk. entomol. derg., 27: 181-189.

- Aydemir, M. and S. Toros 1990. Erzincan ili koşullarında fasulyelerde zararlı *Tetranychus urticae*'nin doğal düşmanları. Türkiye II. Biyolojik Mücadele Kongresi, 26-29 Eylül 1990, Ankara, s. 533-540.
- Bayram, Ş. and S. Çobanoğlu 2006. Astigmata and Prostigmata (Acari) of bulbaceous ornamental plants in Ankara-Turkey. Acta Phytopathologica et entomologica Hungarica, 41: 367-381.
- Bayram, Ş. and S. Çobanoğlu 2007. Mite fauna (Acari: Prostigmata, Mesostigmata, Astigmata) of coniferous plants in Turkey. Turkish Entomology, 31: 279-290.
- Çakmak, İ. and T. Akşit 2003. Aydin ili incir ağaçlarında zararlı akar türleri, doğal düşmanları ve önemlilerinin populasyon değişimleri üzerinde çalışmalar. Türkiye Entomoloji Dergisi, 27: 27-38.
- Çobanoğlu, S., C. Uysal, and E. Ökten 2003. The complex of beneficial mite fauna of ornamental trees and shrubs in Ankara Turkey. Entomologist's Monthly Magazine, 139: 7-12.
- Dobosz, R., A. Skorupska, and C. Blaszak 1995. The appearance of spider mites (Tetranychidae) in parks of Poznan. Boczek, J. and Ignatowicz, S., Materiały z Sympozjum na temat: "Osiagniecia Akarologii w Polsce", Siedlce, 26-27 wrzesnia 1995, Komitet Ochrony Roslin Polska Akademia Nauk: p. 39-42.
- Düzungüneş, Z. 1954. Orta Anadolu meyve ağaçlarına zarar veren Tetranychidae familyası türleri üzerinde sistematik ve biyolojik çalışmalar ve mücadele denemeleri. Ziraat Vekâleti Neşriyat ve Haberleşme Müdürlüğü 706, 104.
- Düzungüneş, Z. 1963. Türkiyede yeni bulunan akarlar. Türkiye Bitki Koruma Bülteni, 3: 237-246.
- Düzungüneş, Z. 1965. Türkiye'de bitkilerde zarar veren Tenuipalpidae Sayed familyası türleri üzerine incelemeler. Ank. Üni. Ziraat Fakültesi Yıllığı 3: 120-148.
- Düzungüneş, Z. 1980. Küçük arthropodların toplanması, saklanması ve mikroskopik reparatlarının hazırlanması. Zir. Müc. ve Zir. Kar. Gen. Müd. Ankara 77 .
- Ecevit, O. 1977. Oligonichus coniferarum (McGregor) (Acarina: Tetranychidae) üzerinde morfolojik çalışmalar. Atatürk Univ. Zir. Fak. Dergisi, 81: 29-33.
- Elma, N.F. and Ö. Alaoğlu 2008. Konya ilinde peyzaj alanlarındaki ağaç ve çalılarda bulunan zararlı akar türleri ve doğal düşmanları. Turk. Entomol. Derg., 2008, 32 : 115-129.
- Erol, T. and B. Yaşar 1996. Van ili elma bahçelerinde bulunan zararlı türler ile doğal düşmanları. Türkiye Entomol. Derg., 20:281-293.
- Gencer, N.S., K.S. Coşkuncu and N.A. Kumral 2002. Bursa ilinde Bursa Siyahı incirlerinde bulunan zararlı akar türleri ve doğal düşmanları üzerinde araştırmalar. Türk. entomol. derg., 26: 229-239.
- Göksu, M.E. and E.D. Atak 1968. Marmara ve Trakya Bölgesinde avrupa kırmızı örümceği (*Panonychus ulmi* Clk.) üzerinde araştırmalar. Bitki Koruma Bülteni, 9: 19-36.
- Hatzinikolis, E.N. 1970. Sept espèces d'acariens phytophages signalées pour la première fois en Grèce pendant l'année 1969. Geponika: 184-187.
- Hoy, A.M. 2011. Agricultural Acarology. Introduction to Integrated Mite Management. University of Florida Gainesville, U.S.A.. CRC. Press. 410 .
- İncekulak, R. and O. Evevit 2002. Amasya elma bahçelerinde bulunan zararlı ve yararlı akar türleri ile poulation yoğunlıklarının saptanması üzerinde bir araştırma. Türkiye V. Biyolojik Mücadele Kongresi Bildirileri (4-7 Eylül), Erzurum, 297-314.
- İyriboz, N. 1940. İncir Hastalıkları. Zirai Mücadele İstasyonu, İzmir, 10-13 .
- Jeppson, L.R., H.H. Keifer, and E.W. Baker 1975. Mites Injurious to Economic Plants. University of California Press, Berkeley, 646 .
- Kasap, İ., S. Çobanoğlu, Y. Aktuğ, and E. Denizhan 2004. Van Gölü çevresi elma bahçelerinde saptanan zararlı ve yararlı akar türleri. Türkiye I. Bitki Koruma Kongresi Bildirileri (8-10 Eylül), Samsun, 104-104.
- Kropczynska, M.D., Van de Vrie and A. Tomycyk 1992. Woody Ornamentals. Spider Mites Their Biology. Natural Enemies and Control. Helle, W. and Sabelis, M.W. (eds). Univ. of Amsterdam, Lab. Of Exp. Ent., The Netherlands. 684-687.
- Madanlar, N. and Z. Yoldaş 1996. Menemen (İzmir)'de açık alanlarda çilek bitkisinin toprak üstü böcek ve akar faunası ile bunların populasyon gelişimi üzerinde araştırmalar. Türkiye III. Entomoloji Kongresi Bildirileri (24-28 Eylül), Ankara, 52-59.

- Mesa, N.C., R. Ochoa, C.W. Welbourn, Evans A. Gregory, and G.J. D. Moraes 2009. A catalog of the Tenuipalpidae (Acari) of the world with a key to genera, Zootaxa, 1-4 .
- Migeon, A. and D. Dorkeld 2006-2013. Spider Mites Web: a comprehensive database for the Tetranychidae. <http://www.montpellier.inra.fr/CBGP/spmw>
- Önuçar, A. and O Ulu. 1988. Kestane ağaçlarında bulunan akar türleri hakkında kısa bilgiler. Türk. entomol. derg., 12: 33-38.
- Pritchard, A.E. and E.W. Baker 1952. A guide to the spider mites of deciduous fruit trees. Hilgardia, 21: 253-287.
- Pritchard, A.E. and E.W. Baker 1955. A revision of the spider mite family Tetranychidae. Memoirs Series, San Francisco, Pacific Coast Entomological Society, 2: 472.
- Reeves, R.M. 1963. Tetranychidae infesting woody plants in New York State, and a life history study of the elm mite *Eotetranychus matthyssei* n. sp.. Cornell University Agricultural Station Mem.: 99.
- Ripka, G. 1997. Aphid and mite fauna of ornamental trees and shrubs. Pro. Dissertations. Budapest.209 .
- Ripka, G. 1998. New data to the knowledge on the phytoseiid fauna in Hungary (Acari: Mesostigmata). Acta Phytopathologica et Entomologica Hungarica, 33: 395–405.
- Ripka, G., A. Fain, A. Kaz'mierski, S. Kreiter, and W. Magowski 2005. New data to the knowledge of the mite fauna of Hungary (Acari: Mesostigmata, Prostigmata and Astigmata). 40: 159-176.
- Seeman, O.D. and J.J. Beard 2011. Identification of exotic pest and Australian native and naturalised species of *Tetranychus* (Acari: Tetranychidae): Zootaxa 2961: 1–72.
- Smiley, R.L. and Baker, E.W. 1995. A report on some tetranychid mites (Acari: Prostigmata) from Yemen. International Journal of Acarology, 21: 135-164.
- Tuttle, D.M. and E.W. Baker 1964. The spider mites of Arizona (Acarina : Tetranychidae). Agricultural Experiment Station, University of Arizona, Technical Bulletin, 158: 1-41.
- Tuttle, D.M. and E.W. Baker 1968. Spider mites of southwestern United States and a revision of the family Tetranychidae. Tuscon, Usa, The University of Arizona Press: 143.
- Ulusoy, R., G. Vatansever and N. Uygun 1999. Ulukışla (Niğde) ve Pozantı (Adana) yöresi kiraz ağaçlarında zararlı olan türler, doğal düşmanları ve önemlileri üzerindeki gözlemler. Türk. Entomol. Derg., 23: 111-120.
- Uysal, C., S. Çobanoglu and M.E. Ökten 2001. Determination of Tetranychoidea (Acarina: Prostigmata) species harmful in the park areas of Ankara. Turkiye Entomoloji Dergisi, 25: 147-160.
- Yanar, D. and O. Ecevit 2005. Tokat ilinde elma (*Malus communis* L.) bahçelerinde görülen bitki zararlısı ve predator akar türleri [Plant injurious and predatory mite species in apple (*Malus communis* L.) orchards in Tokat province]. Ondokuz Mayıs Üniversitesi, Ziraat Fakultesi Dergisi, 20: 18-23 (Turkish, with English summary).
- Yüksel, B. 1999. Doğu Karadeniz Bölgesinde Ladin Örülü Akarı (Jacobi)'nın Zararı ve Mücadelesi, Investigations on damage of Spruce Spider Mite in the Eastern Black Sea Region and Its Control, Orman Mühendisliği Dergisi, Ankara, 36: 28-31.
- Yesilayer, A., S. Çobanoğlu 2009. Major Mites Listed in Turkey's External Quarantine. International Journal of Acarology, International Journal of Acarology. 36: 483–486.3.
- Yesilayer, A. and S. Çobanoğlu 2011. Predatory mite species Phytoseiidae (Acari) distribution on ornamental plants and parks of Istanbul, Turkey. Turkiye Entomoloji Bülteni, 1: 3-15.
- Yesilayer, A. and S. Çobanoğlu 2013. İstanbul (Türkiye) park ve süs bitkilerinde tespit edilen Raphignathoid akarları (Acari and: Prostigmata: Raphignathoidea). Turkiye Entomoloji Dergisi. 37: 93-103.
- Yiğit, A. and N. Uygun 1982. Adana, İçel, Kahramanmaraş illeri elma bahçelerinde zararlı ve yararlı faunasının saptanması üzerinde çalışmalar. Bitki Koruma Bülteni, 22(4):163-178.
- Yüksel, B. 1999. Doğu Karadeniz Bölgesinde Ladin Örülü Akarı (Jacobi)'nın zararı ve mücadelesi, Investigations on damage of Spruce Spider Mite in the Eastern Black Sea Region and Its Control, Orman Mühendisliği Dergisi, Ankara, 36:28-31.

Zhang, Z-Q. and R. Henderson 2002. Key to Tetranychidae of New Zealand. Landcare Research Private Bag 92170 Auckland New Zealand, 62 p.