

APPLICATIONS OF REDUCED AND COREDUCED MODULES I

David Ssevviiri

Received: 10 October 2022; Revised: 6 February 2023; Accepted: 13 March 2023

Communicated by Abdullah Harmanci

ABSTRACT. This is the first in a series of papers highlighting the applications of reduced and coreduced modules. Let R be a commutative unital ring and I be an ideal of R . We show that I -reduced R -modules and I -coreduced R -modules provide a setting in which the Matlis-Greenless-May (MGM) Equivalence and the Greenless-May (GM) Duality hold. These two notions have been hitherto only known to exist in the derived category setting. We realise the I -torsion and the I -adic completion functors as representable functors and under suitable conditions compute natural transformations between them and other functors.

Mathematics Subject Classification (2020): 13D07, 13C12, 13B35, 13C13

Keywords: Reduced and coreduced module, torsion and adic completion

1. Introduction

Let R be a commutative unital ring and I be an ideal of R . The additive endo functors $\Gamma_I : R\text{-Mod} \rightarrow R\text{-Mod}$, $M \mapsto \Gamma_I(M) := \varinjlim \text{Hom}_R(R/I^k, M)$ and $\Lambda_I : R\text{-Mod} \rightarrow R\text{-Mod}$, $M \mapsto \Lambda_I(M) := \varprojlim (M/I^k M)$; called the I -torsion functor and the I -adic completion functor respectively have been widely studied. Grothendieck was the first to study the torsion functor in the algebraic geometry setting of sheaves, [10]. This functor is central to the well known notions of local cohomology, local duality, [6]; Cousin complexes, [11]; and their generalizations, [8,21,24]. Sharp in [19] introduced their algebraic avatars. Λ_I the dual to the functor Γ_I has been used widely to study local homology, see [7,9,18] among others. The functors Γ_I and Λ_I and their derived functors were key in [1,4,5,9,16,18,22,25] where notions such as Greenless-May (GM) Duality and Matlis-Greenless-May (MGM) Equivalence were studied in different settings.

The author was supported by the International Science Programme through the Eastern Africa Algebra Research Group and also by the EPSRC GCRF project EP/T001968/1, Capacity building in Africa via technology-driven research in algebraic and arithmetic geometry (part of the Abram Gannibal Project).

The functors Γ_I and Λ_I are not adjoint to each other. However, under suitable conditions, their corresponding derived functors

$$\mathbf{R}\Gamma_I \text{ and } \mathbf{L}\Lambda_I : D(R) \rightarrow D(R)$$

where $D(R)$ is the derived category of R -modules are adjoint. This is what is known as the Greenlees-May (GM) Duality. Our first main result, is Theorem 1.1 which shows that in the setting of I -reduced modules and I -coreduced modules, the functors Γ_I and Λ_I form an adjoint pair.

Let I be an ideal of a ring R and $(R\text{-Mod})_{I\text{-red}}$ (resp. $(R\text{-Mod})_{I\text{-cor}}$) denote a full subcategory of $R\text{-Mod}$ consisting of all I -reduced (resp. I -coreduced) R -modules.

Theorem 1.1. [GM Duality in $R\text{-Mod}$] *For any ideal I of a ring R ,*

(1) *The functor*

$$\Gamma_I : (R\text{-Mod})_{I\text{-red}} \rightarrow (R\text{-Mod})_{I\text{-cor}}$$

is idempotent and for any $M \in (R\text{-Mod})_{I\text{-red}}$, $\Gamma_I(M) \cong \text{Hom}_R(R/I, M)$.

(2) *The functor*

$$\Lambda_I : (R\text{-Mod})_{I\text{-cor}} \rightarrow (R\text{-Mod})_{I\text{-red}}$$

is idempotent and for any $M \in (R\text{-Mod})_{I\text{-cor}}$, $\Lambda_I(M) \cong R/I \otimes M$.

(3) *For any $N \in (R\text{-Mod})_{I\text{-red}}$ and $M \in (R\text{-Mod})_{I\text{-cor}}$,*

$$\text{Hom}_R(\Lambda_I(M), N) \cong \text{Hom}_R(M, \Gamma_I(N)).$$

Let $D(R)$ denote the derived category of the R -module category $R\text{-Mod}$. A complex $M \in D(R)$ is called a *derived I -torsion complex* if the canonical morphism $\mathbf{R}\Gamma_I(M) \rightarrow M$ is an isomorphism. The complex M is called a *derived I -adically complete complex* if the canonical morphism $M \rightarrow \mathbf{L}\Lambda_I(M)$ is an isomorphism. Denote by $D(R)_{I\text{-tor}}$ and $D(R)_{I\text{-com}}$ the full subcategories of $D(R)$ consisting of derived I -torsion complexes and derived I -adically complete complexes, respectively. These are triangulated subcategories. One version of the MGM Equivalence is Theorem 1.2 which appears as Theorem 7.11 in [16]. Recently, its version in the noncommutative setting was given, see [23].

Theorem 1.2. *Let R be a ring, and let I be a weakly proregular ideal in it.*

(1) *For any $M \in \mathbf{D}(R)$, $\mathbf{R}\Gamma_I(M) \in \mathbf{D}(R)_{I\text{-tor}}$ and $\mathbf{L}\Lambda_I(M) \in \mathbf{D}(R)_{I\text{-com}}$.*

(2) *The functor*

$$\mathbf{R}\Gamma_I : \mathbf{D}(R)_{I\text{-com}} \rightarrow \mathbf{D}(R)_{I\text{-tor}}$$

is an equivalence, with quasi-inverse $\mathbf{L}\Lambda_I$.

Our second main result is Theorem 1.3 which realises the MGM Equivalence in the setting of I -reduced and I -coreduced R -modules. An R -module M is I -torsion (resp. I -complete) if $\Gamma_I(M) \cong M$ (resp. $\Lambda_I(M) \cong M$). The full subcategory of $R\text{-Mod}$ consisting of I -torsion (resp. I -complete) R -modules is denoted by $(R\text{-Mod})_{I\text{-tor}}$ (resp. $(R\text{-Mod})_{I\text{-com}}$).

Theorem 1.3. [The MGM Equivalence in $R\text{-Mod}$] *Let I be any ideal of a ring R ,*

- (1) *For any $M \in (R\text{-Mod})_{I\text{-red}}$,*

$$\Gamma_I(M) \in (R\text{-Mod})_{I\text{-com}} \cap (R\text{-Mod})_{I\text{-cor}} =: \mathfrak{E}.$$

- (2) *For any $M \in (R\text{-Mod})_{I\text{-cor}}$,*

$$\Lambda_I(M) \in (R\text{-Mod})_{I\text{-tor}} \cap (R\text{-Mod})_{I\text{-red}} =: \mathfrak{D}.$$

- (3) *The functor $\Gamma_I : (R\text{-Mod})_{I\text{-red}} \rightarrow (R\text{-Mod})_{I\text{-cor}}$ restricted to \mathfrak{D} is an equivalence between \mathfrak{D} and \mathfrak{E} with quasi-inverse Λ_I .*

Reduced modules were introduced by Lee and Zhou in [14] and later studied in [17]. In both papers, the method of study was mainly element-wise. It was however observed in [20] that “reduced modules” is a categorical property. Therefore, it is amenable to study by use of category theory. The use of this method to study reduced modules first appeared in [12] and [13]. In this paper, we continue with this approach; it allows us to benefit from the powerful machinery of category theory. The key idea that makes all that we do possible is the fact that the functor Γ_I when restricted to I -reduced R -modules is representable, i.e., for all I -reduced R -modules M , $\Gamma_I(M) \cong \text{Hom}_R(R/I, M)$. Dually, for any I -coreduced R -module M , $\Lambda_I(M)$ is naturally isomorphic to $R/I \otimes M$.

The paper is organised as follows. It consists of five sections. In Section 2, we give the basic properties of I -reduced and I -coreduced R -modules (given an ideal I of R) necessary for the subsequent sections. In Sections 3 and 4, we prove the GM Duality and the MGM Equivalence respectively. Weakly proregular ideals have been hitherto known to be the most general condition under which the GM Duality and the MGM Equivalence hold (although in the derived category setting). Since in this paper we have another condition of I -reduced and I -coreduced R -modules serving the same purpose, part of Section 4 aims at comparing the two conditions. In general, none of the conditions implies the other. The last section; Section 5, explores more ways (beyond those in [12] for I -reduced R -modules) of realising the functors Γ_I and Λ_I as representable functors. Furthermore, we demonstrate

that doing so is important by utilising the Yoneda Lemma to compute natural transformations.

2. Basic properties

Definition 2.1. Let I be an ideal of a ring R , M be an R -module and f_a be the endomorphism of M given by $m \mapsto am$ for $a \in R$. M is

- (1) I -reduced if for every $a \in I$, $\text{Ker } f_a \cong \text{Ker } f_a^2$;
- (2) I -coreduced if for every $a \in I$, $\text{Coker } f_a \cong \text{Coker } f_a^2$.

An R -module M is *reduced* (resp. *coreduced*) if for every ideal I of R , M is I -reduced (resp. I -coreduced). A ring R is reduced (resp. coreduced) if and only if it is reduced (resp. coreduced) as an R -module. Since a ring is reduced if and only if it has no nonzero nilpotent ideals and it is coreduced if and only if all its ideals are idempotent, it follows that idempotent ideals dualise having no nonzero nilpotent ideals.

For any ideal I of a ring R , the class of I -reduced R -modules is quite large. It contains reduced R -modules which also contain the well studied class of prime R -modules. Dually, the I -coreduced R -modules contain the class of coreduced R -modules which in turn contain the second (also called coprime) R -modules.

Proposition 2.2. For any R -module M and an ideal I of R , the following statements are equivalent:

- (1) M is I -reduced,
- (2) $(0 :_M I) = (0 :_M I^2)$,
- (3) $\text{Hom}_R(R/I, M) \cong \text{Hom}_R(R/I^2, M)$,
- (4) $\Gamma_I(M) \cong \text{Hom}_R(R/I, M)$,
- (5) $I\Gamma_I(M) = 0$.

Proof. (1) \Rightarrow (2) Since in general $(0 :_M I) \subseteq (0 :_M I^2)$, let $m \in (0 :_M I^2)$. Then $I^2m = 0$ and $a^2m = 0$ for all $a \in I$. So, $m \in (0 :_M a^2)$ for all $a \in I$. $\text{Ker } f_a^2 = (0 :_M a^2)$ and by hypothesis, $\text{Ker } f_a = \text{Ker } f_a^2$. So, we get $m \in (0 :_M a)$ for all $a \in I$ and hence $m \in (0 :_M I)$.

(2) \Rightarrow (3) This is immediate since $\text{Hom}_R(R/I, M) \cong (0 :_M I)$.

(3) \Rightarrow (4) By definition, $\Gamma_I(M) := \varinjlim \text{Hom}_R(R/I^k, M)$. It follows by statement (3) that $\text{Hom}_R(R/I^k, M) \cong \text{Hom}_R(R/I, M)$ for all $k \in \mathbb{Z}^+$. So, $\Gamma_I(M) \cong \text{Hom}_R(R/I, M)$.

(4) \Rightarrow (5) If $\Gamma_I(M) \cong \text{Hom}_R(R/I, M)$, then $I\Gamma_I(M) \cong I\text{Hom}_R(R/I, M) \cong 0$.

(5) \Rightarrow (1) Since for any $a \in I$, $\text{Ker } f_a \subseteq \text{Ker } f_a^2$, we prove the reverse inclusion.

Let $m \in \text{Ker } f_a^2 = (0 :_M a^2)$ for all $a \in I$. $a^2m = 0$ and $(a)^2m = 0$ for all $a \in I$, where (a) is the ideal of R generated by a . This implies that $m \in \Gamma_{(a)}(M)$ for all $a \in I$ and $m \in \bigcap_{a \in I} \Gamma_{(a)}(M) = \Gamma_I(M)$. Therefore, $am \in Im \subseteq I\Gamma_I(M) = 0$ for all $a \in I$ and hence $m \in (0 :_M a) = \text{Ker } f_a$ for all $a \in I$. \square

Proposition 2.3. *For any R -module M and an ideal I of R , the following statements are equivalent:*

- (1) M is I -coreduced,
- (2) $IM = I^2M$,
- (3) $R/I \otimes M \cong R/I^2 \otimes M$,
- (4) $\Lambda_I(M) \cong R/I \otimes M$,
- (5) $I\Lambda_I(M) = 0$.

Proof. (1) \Rightarrow (2) Suppose that $M/a^2M \cong M/aM$ for all $a \in I$. So, $a^2M = aM$ for all $a \in I$. To see this, note that the natural epimorphisms $M \rightarrow M/a^2M$ and $M \rightarrow M/aM$ have kernels a^2M and aM respectively. The isomorphism $M/a^2M \cong M/aM$ implies that the submodules a^2M and aM of M should coincide for all $a \in I$. Therefore, $I^2M = IM$.

(2) \Rightarrow (3) $R/I \otimes M \cong M/IM = M/I^2M \cong R/I^2 \otimes M$.

(3) \Rightarrow (4) $\Lambda_I(M) := \varprojlim (R/I^k \otimes M) \cong \varprojlim (R/I \otimes M) = R/I \otimes M$.

(4) \Rightarrow (5) Given statement (4), we have $I\Lambda_I(M) \cong I(R/I \otimes M) \cong (0 \otimes M) \cong 0$.

(5) \Rightarrow (1) $0 \cong I\Lambda_I(M) = I \varprojlim (M/I^k M)$ implies that $IM = I^k M$ for all $k \in \mathbb{Z}^+$. Since $a^2M \subseteq aM$ for all $a \in I$, let $m \in aM$ for all $a \in I$. Then $m \in IM = I^k M$ for all $k \in \mathbb{Z}^+$. So, $m \in a^2M$ for all $a \in I$. This establishes the equality $a^2M = aM$ for all $a \in I$. Thus $M/aM = M/a^2M$ for all $a \in I$. \square

Remark 2.4. Proposition 2.2 holds when in the place of the ideal I of R one has an element $a \in R$ (or the principal ideal aR) in which case M is called a -reduced, see [12, Proposition 2.2].

Proposition 2.5. *Every coreduced ring is reduced.*

Proof. If a ring R is coreduced, then every ideal I of R is idempotent since $I^2R = IR$. If $I^2 = 0$, then $I = 0$ and R is reduced. \square

The converse of Proposition 2.5 is not true in general. The ring of integers is reduced but it is not coreduced. A coreduced module also need not be reduced. The \mathbb{Z} -module \mathbb{Q}/\mathbb{Z} is coreduced but it is not reduced.

Proposition 2.6. *Let R be a ring.*

- (1) *For any R -module N and a coreduced (resp. I -coreduced) R -module M , the R -module $\text{Hom}_R(M, N)$ is reduced (resp. I -reduced).*
- (2) *Let N be an injective cogenerator of $R\text{-Mod}$. If $\text{Hom}_R(M, N)$ is a reduced (resp. I -reduced) R -module for some $M \in R\text{-Mod}$, then M is a coreduced (resp. I -coreduced) R -module.*
- (3) *Let N be an injective cogenerator R -module. The R -module M is coreduced (resp. I -coreduced) if and only if $\text{Hom}_R(M, N)$ is a reduced (resp. I -reduced) R -module.*

Proof. We prove the I -reduced and I -coreduced cases. The reduced and coreduced versions follow immediately.

(1) Let I be an ideal of R and M an I -coreduced R -module. By the Hom-Tensor duality and Proposition 2.3, we have $\text{Hom}_R(R/I, \text{Hom}_R(M, N)) \cong \text{Hom}_R(R/I \otimes M, N) \cong \text{Hom}_R(R/I^2 \otimes M, N) \cong \text{Hom}_R(R/I^2, \text{Hom}_R(M, N))$. By Proposition 2.2, $\text{Hom}_R(M, N)$ is I -reduced.

(2) Let I be an ideal of R and $\text{Hom}_R(M, N)$ be an I -reduced R -module. By the Hom-Tensor duality and Proposition 2.2,

$$\begin{aligned} \text{Hom}_R(R/I \otimes M, N) &\cong \text{Hom}_R(R/I, \text{Hom}_R(M, N)) \\ &\cong \text{Hom}_R(R/I^2, \text{Hom}_R(M, N)) \cong \text{Hom}_R(R/I^2 \otimes M, N). \end{aligned}$$

Since N is an injective cogenerator, the functor $\text{Hom}_R(-, N)$ reflects isomorphisms. So, $R/I \otimes M \cong R/I^2 \otimes M$ and by Proposition 2.3, M is I -coreduced.

(3) This is immediate from (1) and (2) above. \square

Proposition 2.7. *Let M be an S - R -bimodule and N a left R -module. If J is an ideal of S and ${}_S M$ is a J -coreduced (resp. coreduced) S -module, then so is the S -module $M \otimes N$.*

Proof. If J is an ideal of S , then $S/J \otimes_S (M \otimes_R N) \cong (S/J \otimes_S M) \otimes_R N \cong (S/J^2 \otimes_S M) \otimes_R N \cong S/J^2 \otimes_S (M \otimes_R N)$. \square

Corollary 2.8. *Let L be an R -module. If $\text{Hom}_R(L, R)$ is a coreduced R -module, then so is the R -module $\text{Hom}_R(L, M)$ for any $M \in R\text{-Mod}$.*

Proof. Follows from Proposition 2.7 and $\text{Hom}_R(L, M) \cong \text{Hom}_R(L, R) \otimes M$. \square

Let I be an ideal of a ring R . We denote by $(R\text{-Mod})_{I\text{-cor}}$ (resp. $(R\text{-Mod})_{I\text{-red}}$) the subcategory of $R\text{-Mod}$ consisting of I -coreduced (resp. I -reduced) R -modules.

Proposition 2.9. *For any ideal I of a ring R , we have:*

(1) $R/I \otimes -$ and $\text{Hom}_R(R/I, -)$ are idempotent functors from $R\text{-Mod}$ to

$$(R\text{-Mod})_{I\text{-cor}} \cap (R\text{-Mod})_{I\text{-red}}.$$

(2) For any R -module M ,

$$R/I \otimes \text{Hom}_R(R/I, M) \cong \text{Hom}_R(R/I, M). \quad (1)$$

$$\text{Hom}_R(R/I, R/I \otimes M) \cong R/I \otimes M. \quad (2)$$

(3) For any R -module M , the R -modules $\text{Hom}_R(R/I, M)$ and $R/I \otimes M$ are both I -torsion and I -complete.

(4) The set $\mathfrak{A} := \{\text{Hom}_R(R/I, -), R/I \otimes -\}$ forms a noncommutative semi-group where the operation is composition of functors.

Proof. (1) Idempotency holds because

$$R/I \otimes (R/I \otimes M) \cong (R/I \otimes R/I) \otimes M \cong R/I \otimes M$$

and

$$\text{Hom}_R(R/I, \text{Hom}_R(R/I, M)) \cong \text{Hom}_R(R/I \otimes R/I, M) \cong \text{Hom}_R(R/I, M).$$

For any R -module M , $R/I \otimes M \cong M/IM$ and $\text{Hom}_R(R/I, M) \cong (0 :_M I)$. Also, $I(M/IM) = 0$ and $I(0 :_M I) = 0$. So, the R -modules M/IM and $(0 :_M I)$ are I -coreduced. It is also easy to see that

$$(0 :_{(0 :_M I)} I) = (0 :_{(0 :_M I)} I^2) = (0 :_M I)$$

and

$$(\bar{0} :_{M/IM} I) = (\bar{0} :_{M/IM} I^2) = M/IM.$$

This shows that the R -modules M/IM and $(0 :_M I)$ are I -reduced.

(2)

$$R/I \otimes \text{Hom}_R(R/I, M) \cong \frac{\text{Hom}_R(R/I, M)}{I \text{Hom}_R(R/I, M)} \cong \frac{\text{Hom}_R(R/I, M)}{0} \cong \text{Hom}_R(R/I, M).$$

$$\text{Hom}_R(R/I, R/I \otimes M) \cong \text{Hom}_R(R/I, M/IM) \cong (\bar{0} :_{M/IM} I) = M/IM \cong R/I \otimes M.$$

(3) The following maps always hold¹

$$\text{Hom}_R(R/I, \text{Hom}_R(R/I, M)) \hookrightarrow \Gamma_I(\text{Hom}_R(R/I, M)) \hookrightarrow \text{Hom}_R(R/I, M) \quad (3)$$

¹ \hookrightarrow denotes a monomorphism and \twoheadrightarrow denotes an epimorphism.

$$\mathrm{Hom}_R(R/I, R/I \otimes M) \hookrightarrow \Gamma_I(R/I \otimes M) \hookrightarrow R/I \otimes M \quad (4)$$

$$\mathrm{Hom}_R(R/I, M) \twoheadrightarrow \Lambda_I(\mathrm{Hom}_R(R/I, M)) \twoheadrightarrow R/I \otimes \mathrm{Hom}_R(R/I, M) \quad (5)$$

$$R/I \otimes M \twoheadrightarrow \Lambda_I(R/I \otimes M) \twoheadrightarrow R/I \otimes (R/I \otimes M) \quad (6)$$

Idempotency of the functors $\mathrm{Hom}_R(R/I, -)$ and $R/I \otimes -$ shows that the maps in (3) and (6) are all isomorphisms in which case, $\mathrm{Hom}_R(R/I, M)$ and $R/I \otimes M$ become I -torsion and I -complete respectively. Invariance of $R/I \otimes M$ and $\mathrm{Hom}_R(R/I, M)$ under the functor $\mathrm{Hom}_R(R/I, -)$ and $R/I \otimes -$ respectively shows that the morphisms in (4) and (5) are all isomorphisms. This shows that $R/I \otimes M$ and $\mathrm{Hom}_R(R/I, M)$ are I -torsion and I -complete respectively.

(4) From (1) and (2) above, we get Figure 1 which shows that the set \mathfrak{A} is a noncommutative semigroup.

	$\mathrm{Hom}_R(R/I, -)$	$R/I \otimes -$
$\mathrm{Hom}_R(R/I, -)$	$\mathrm{Hom}_R(R/I, -)$	$R/I \otimes -$
$R/I \otimes -$	$\mathrm{Hom}_R(R/I, -)$	$R/I \otimes -$

FIGURE 1. Multiplication table

□

Corollary 2.10. *Every R -module M has a submodule and a quotient module which are both I -reduced and I -coreduced as R -modules.*

Proof. Immediate from Proposition 2.9(1). They are $(0 :_M I)$ and M/IM respectively. □

Corollary 2.11. *If M is I -reduced (resp. I -coreduced), then*

$$\mathrm{Hom}_R(R/I, \Gamma_I(M)) \cong \mathrm{Hom}_R(R/I, M) \cong \Gamma_I(M)$$

and

$$(\text{resp. } R/I \otimes \Lambda_I(M) \cong R/I \otimes M \cong \Lambda_I(M)).$$

Proof. By Proposition 2.2, $\Gamma_I(M) \cong \text{Hom}_R(R/I, M)$. So,

$$\text{Hom}_R(R/I, \Gamma_I(M)) \cong \text{Hom}_R(R/I, \text{Hom}_R(R/I, M)) \cong \text{Hom}_R(R/I, M) \cong \Gamma_I(M).$$

Also by Proposition 2.3, $\Lambda_I(M) \cong R/I \otimes M$ and therefore $R/I \otimes \Lambda_I(M) \cong R/I \otimes R/I \otimes M \cong R/I \otimes M \cong \Lambda_I(M)$. \square

Proposition 2.12. *Let I be an ideal of a ring R . An inverse limit of I -reduced (resp. reduced) R -modules is an I -reduced (resp. reduced) R -module and a direct limit of I -coreduced (resp. coreduced) R -modules is an I -coreduced (resp. coreduced) R -module.*

Proof. We prove the cases for I -reduced and I -coreduced. The other cases become immediate. It is well known that the functor $\text{Hom}_R(R/I, -)$ (resp. $R/I \otimes -$) preserves inverse limits (resp. direct limits). A property possessed by right adjoint (resp. left adjoint) functors, see for instance [15, V. 5]. So, if each R -module M_k is I -reduced, then $\text{Hom}_R(R/I, \varprojlim M_k) \cong \varprojlim \text{Hom}_R(R/I, M_k) \cong \varprojlim \text{Hom}_R(R/I^2, M_k) \cong \text{Hom}_R(R/I^2, \varprojlim M_k)$ which shows that $\varprojlim M_k$ is an I -reduced R -module. Dually, suppose that each R -module M_k is I -coreduced. $R/I \otimes \varinjlim M_k \cong \varinjlim (R/I \otimes M_k) \cong \varinjlim (R/I^2 \otimes M_k) = R/I^2 \otimes \varinjlim M_k$ so that $\varinjlim M_k$ is an I -coreduced R -module. \square

Corollary 2.13. *Let I be an ideal of R and M be an R -module. M is I -reduced if and only if so is the R -submodule $\Gamma_I(M)$ of M .*

Proof. M is I -reduced if and only if $I\Gamma_I(M) = 0$ (Proposition 2.2). Idempotency of Γ_I implies that $I\Gamma_I(M) = 0$ if and only if $I\Gamma_I(\Gamma_I(M)) = 0$. This is the case if and only if $\Gamma_I(M)$ is I -reduced (Proposition 2.2). \square

By a similar proof, we obtain:

Corollary 2.14. *Let I be an ideal of R , M be an R -module and Λ_I be an idempotent functor, then M is I -coreduced if and only if so is $\Lambda_I(M)$.*

3. Greenless-May Duality

For an arbitrary ideal I of a ring R and R -modules M and N ,

$$\text{Hom}_R(\Lambda_I(M), N) \cong \text{Hom}_R(M, \Gamma_I(N)).$$

So, the functors Γ_I and Λ_I are not in general adjoint to each other. However, in the setting of derived categories, we have Theorem 3.1 which is called the Greenless-May Duality (GM Duality for short). It was first proved by Alonso Tarrío, Jeremias Lopez and Lipman [1] but it also appears in [16, Theorem 7.1.2].

Theorem 3.1. [GM-Duality in $\mathbf{D}(R)$] Let I be a weakly proregular ideal of a ring R and $M, N \in \mathbf{D}(R)$. Then there is a natural isomorphism in $\mathbf{D}(R)$ given by

$$\mathbf{R}\mathrm{Hom}_R(\mathbf{R}\Gamma_I(M), N) \cong \mathbf{R}\mathrm{Hom}_R(M, \mathbf{L}\Lambda_I(N)).$$

This theorem implies that local cohomology is derived left adjoint to local homology. Let I be an ideal of a ring R and let $(R\text{-Mod})_{I\text{-red}}$ (resp. $(R\text{-Mod})_{I\text{-cor}}$) denote a full subcategory of $R\text{-Mod}$ consisting of all I -reduced (resp. I -coreduced) R -modules.

Definition 3.2. Let I be an ideal of a ring R . An R -module M is *I -reduced* if $I\Gamma_I(M) = 0$.

Definition 3.3. Let I be an ideal of a ring R . An R -module M is *I -reduced* if $I\Gamma_I(M) = 0$.

Theorem 3.4. [GM Duality in $R\text{-Mod}$] For any ideal I of a ring R ,

(1) The functor

$$\Gamma_I : (R\text{-Mod})_{I\text{-red}} \rightarrow (R\text{-Mod})_{I\text{-cor}}$$

is idempotent and for any $M \in (R\text{-Mod})_{I\text{-red}}$, $\Gamma_I(M) \cong \mathrm{Hom}_R(R/I, M)$.

(2) The functor

$$\Lambda_I : (R\text{-Mod})_{I\text{-cor}} \rightarrow (R\text{-Mod})_{I\text{-red}}$$

is idempotent and for any $M \in (R\text{-Mod})_{I\text{-cor}}$, $\Lambda_I(M) \cong R/I \otimes M$.

(3) For any $N \in (R\text{-Mod})_{I\text{-red}}$ and $M \in (R\text{-Mod})_{I\text{-cor}}$,

$$\mathrm{Hom}_R(\Lambda_I(M), N) \cong \mathrm{Hom}_R(M, \Gamma_I(N)).$$

Proof. (1) From Proposition 2.2, $\Gamma_I(M)$ is naturally isomorphic to $\mathrm{Hom}_R(R/I, M)$ for any I -reduced R -module M . Idempotency is due to Proposition 2.9(1) and the fact that $\Gamma_I(M) \cong \mathrm{Hom}_R(R/I, M)$ for all I -reduced R -modules M .

(2) Follows from Proposition 2.3(2). Idempotency is due to Proposition 2.9(1) and the fact that $\Lambda_I(M) \cong R/I \otimes M$ for all I -coreduced R -modules M .

(3) Consider the functor $\Gamma_I : (R\text{-Mod})_{I\text{-red}} \rightarrow (R\text{-Mod})_{I\text{-cor}}$. For any module $M \in (R\text{-Mod})_{I\text{-red}}$, $\Gamma_I(M) \cong \mathrm{Hom}_R(R/I, M)$. However, the functor $R/I \otimes -$ is left-adjoint to $\mathrm{Hom}_R(R/I, -)$. By uniqueness of adjoints, the functor $\Lambda_I : (R\text{-Mod})_{I\text{-cor}} \rightarrow (R\text{-Mod})_{I\text{-red}}$ which has the property that for all $M \in (R\text{-Mod})_{I\text{-cor}}$, $\Lambda_I(M) \cong R/I \otimes M$ is the left adjoint of Γ_I . \square

Corollary 3.5. If $M \in (R\text{-Mod})_{I\text{-red}} \cap (R\text{-Mod})_{I\text{-cor}}$, then $\Gamma_I(M) = 0$ if and only if $\Lambda_I(M) = 0$.

Proof. By the adjunction in Theorem 3.4, if $M = N$, we get

$$\mathrm{Hom}_R(\Lambda_I(M), M) \cong \mathrm{Hom}_R(M, \Gamma_I(M)).$$

If $\Lambda_I(M) = 0$, $\mathrm{Hom}_R(M, \Gamma_I(M)) = 0$, and every R -homomorphism $f : M \rightarrow \Gamma_I(M)$ is a zero homomorphism. Applying Γ_I gives $\Gamma_I(f) : \Gamma_I(M) \rightarrow \Gamma_I(\Gamma_I(M))$ which is zero. However, Γ_I is idempotent. So $\Gamma_I(f)$ is an isomorphism and therefore $\Gamma_I(M) = 0$. Conversely, suppose that $\Gamma_I(M) = 0$. Then $\mathrm{Hom}_R(\Lambda_I(M), M) = 0$ and therefore every R -homomorphism $g : \Lambda_I(M) \rightarrow M$ is zero. It follows that $\Lambda_I(g) : \Lambda_I(\Lambda_I(M)) \rightarrow \Lambda_I(M)$ is also a zero R -homomorphism. By idempotency of Λ_I on I -coreduced R -modules $\Lambda_I(g)$ is an isomorphism and hence $\Lambda_I(M) = 0$. \square

Example 3.6. We have the following examples.

- (1) If $I^2 = I$, then $R\text{-Mod} = (R\text{-Mod})_{I\text{-red}} = (R\text{-Mod})_{I\text{-cor}}$.
- (2) For any simple R -module M , $M \in (R\text{-Mod})_{I\text{-red}} \cap (R\text{-Mod})_{I\text{-cor}}$.
- (3) If R is an Artinian ring with an ideal I , then there exists a positive integer k such that every R -module is both I^k -reduced and I^k -coreduced, i.e., $R\text{-Mod} = (R\text{-Mod})_{I^k\text{-red}} \cap (R\text{-Mod})_{I^k\text{-cor}}$.

The first two examples are easy to see. Suppose that R is Artinian. There exists some positive integer k such that $I^k = I^{k+t}$ for all positive integers t . So, $(0 :_M I^k) = (0 :_M I^{k+t})$ and $M/I^k M = M/I^{k+t} M$ for all positive integers t . Accordingly, $\Gamma_{I^k}(M) \cong (0 :_M I^k)$ and $\Lambda_{I^k}(M) \cong M/I^k M$. So, M is both I^k -reduced and I^k -coreduced.

Corollary 3.7. *Let I be an ideal of a ring R ,*

- (1) *For any $M \in (R\text{-Mod})_{I\text{-red}}$, $\Lambda_I(\Gamma_I(M)) = \Gamma_I(M)$.*
- (2) *For any $M \in (R\text{-Mod})_{I\text{-cor}}$, $\Gamma_I(\Lambda_I(M)) = \Lambda_I(M)$.*

Proof. (1) By Theorem 3.4, for any $M \in (R\text{-Mod})_{I\text{-red}}$, $\Gamma_I(M) \in (R\text{-Mod})_{I\text{-cor}}$ and $\Gamma_I(M) \cong \mathrm{Hom}_R(R/I, M)$. So,

$$\Lambda_I(\Gamma_I(M)) \cong R/I \otimes \mathrm{Hom}_R(R/I, M) \cong \mathrm{Hom}_R(R/I, M) \cong \Gamma_I(M).$$

(2) By Theorem 3.4, for any $M \in (R\text{-Mod})_{I\text{-cor}}$, $\Lambda_I(M) \in (R\text{-Mod})_{I\text{-red}}$ and $\Lambda_I(M) \cong R/I \otimes M$. It follows that $\Gamma_I(\Lambda_I(M)) \cong \mathrm{Hom}_R(R/I, R/I \otimes M) \cong R/I \otimes M \cong \Lambda_I(M)$. \square

4. MGM Equivalence

Let R be a ring and $\mathbf{D}(R)$ denote the derived category of the abelian category $R\text{-Mod}$. A complex $M \in \mathbf{D}(R)$ is called *derived I -torsion* [16, Definition 3.11] (resp. *derived I -adically complete* [16, Definition 3.8]) if the morphism

$$\sigma_M^R : \mathbf{R}\Gamma_I(M) \rightarrow M \text{ (resp. } \tau_M^L : M \rightarrow \mathbf{L}\Lambda_I(M))$$

is an isomorphism, where $\mathbf{R}\Gamma_I$ and $\mathbf{L}\Lambda_I$ denote the right derived and left derived functors of Γ_I and Λ_I respectively. The full subcategory of $\mathbf{D}(R)$ consisting of derived I -torsion (resp. derived I -adically complete) complexes is denoted by $\mathbf{D}(R)_{I\text{-tor}}$ (resp. $\mathbf{D}(R)_{I\text{-com}}$).

Theorem 4.1. [MGM Equivalence [16, Theorem 7.11]] Let R be a ring, and let I be a weakly proregular ideal in it.

- (1) For any $M \in \mathbf{D}(R)$, $\mathbf{R}\Gamma_I(M) \in \mathbf{D}(R)_{I\text{-tor}}$ and $\mathbf{L}\Lambda_I(M) \in \mathbf{D}(R)_{I\text{-com}}$.
- (2) The functor

$$\mathbf{R}\Gamma_I : \mathbf{D}(R)_{I\text{-com}} \rightarrow \mathbf{D}(R)_{I\text{-tor}}$$

is an equivalence, with quasi-inverse $\mathbf{L}\Lambda_I$.

Lemma 4.2. *If I is an ideal of a ring R and M an I -reduced (resp. I -coreduced) R -module, then $\Gamma_I(M)$ (resp. $\Lambda_I(M)$) is an I -complete (resp. I -torsion) R -module.*

Proof. If M is I -reduced, $\Gamma_I(M) \cong \text{Hom}_R(R/I, M)$ which is both an I -reduced and I -coreduced R -module by Proposition 2.9. So,

$$\Lambda_I(\Gamma_I(M)) \cong R/I \otimes \text{Hom}_R(R/I, M) \cong \text{Hom}_R(R/I, M) \cong \Gamma_I(M).$$

This proves that $\Gamma_I(M)$ is I -complete. If M is I -coreduced, $\Lambda_I(M) \cong R/I \otimes M$ which is also both an I -reduced and I -coreduced R -module by Proposition 2.9. So, $\Gamma_I(\Lambda_I(M)) \cong \text{Hom}_R(R/I, \Lambda_I(M)) \cong \text{Hom}_R(R/I, R/I \otimes M) \cong R/I \otimes M \cong \Lambda_I(M)$. This proves that $\Lambda_I(M)$ is I -torsion. \square

Theorem 4.3. [The MGM Equivalence in $R\text{-Mod}$] *Let I be any ideal of a ring R ,*

- (1) *For any $M \in (R\text{-Mod})_{I\text{-red}}$,*

$$\Gamma_I(M) \in (R\text{-Mod})_{I\text{-com}} \cap (R\text{-Mod})_{I\text{-cor}} =: \mathfrak{E}.$$

- (2) *For any $M \in (R\text{-Mod})_{I\text{-cor}}$,*

$$\Lambda_I(M) \in (R\text{-Mod})_{I\text{-tor}} \cap (R\text{-Mod})_{I\text{-red}} =: \mathfrak{D}.$$

- (3) *The functor $\Gamma_I : (R\text{-Mod})_{I\text{-red}} \rightarrow (R\text{-Mod})_{I\text{-cor}}$ restricted to \mathfrak{D} is an equivalence between \mathfrak{D} and \mathfrak{E} with quasi-inverse Λ_I .*
- (4) *If I is an idempotent ideal, then the subcategory $(R\text{-Mod})_{I\text{-com}}$ is equivalent to $(R\text{-Mod})_{I\text{-tor}}$.*

Proof. (1) By Theorem 3.4, $M \in (R\text{-Mod})_{I\text{-red}}$ implies that $\Gamma_I(M) \in (R\text{-Mod})_{I\text{-cor}}$. If $M \in (R\text{-Mod})_{I\text{-red}}$, then by Lemma 4.2, $\Gamma_I(M) \in (R\text{-Mod})_{I\text{-com}}$. So, $\Gamma_I(M) \in \mathfrak{E}$.
 (2) By Theorem 3.4, $\Lambda_I(M) \in (R\text{-Mod})_{I\text{-red}}$ for any $M \in (R\text{-Mod})_{I\text{-cor}}$. If $M \in (R\text{-Mod})_{I\text{-cor}}$, then by Lemma 4.2, $\Lambda_I(M) \in (R\text{-Mod})_{I\text{-tor}}$. So, $\Lambda_I(M) \in \mathfrak{D}$.
 (3) If $M \in \mathfrak{D}$, then $\Gamma_I(M)$ is I -complete (Lemma 4.2). So $\Lambda_I(\Gamma_I(M)) \cong \Gamma_I(M) \cong M$ since by hypothesis M is I -torsion. On the other hand, if $M \in \mathfrak{E}$, then $\Lambda_I(M)$ is I -torsion (Lemma 4.2). So $\Gamma_I(\Lambda_I(M)) \cong \Lambda_I(M) \cong M$ since by hypothesis M is I -complete.
 (4) If $I^2 = I$, then $R\text{-Mod} = (R\text{-Mod})_{I\text{-red}} = (R\text{-Mod})_{I\text{-cor}}$. So, $\mathfrak{E} = (R\text{-Mod})_{I\text{-com}}$ and $\mathfrak{D} = (R\text{-Mod})_{I\text{-tor}}$. The rest follows from part (3). \square

Remark 4.4. It is tempting to think that $\mathfrak{D} = \mathfrak{E}$ is a small subcategory (in the sense that it contains few modules). However, this is not the case. For any module M , the R -modules M/IM and $(0 :_M I)$ belong to \mathfrak{D} . In particular, every module has a submodule and a quotient which is contained in \mathfrak{D} .

4.1. Comparison with weak proregularity. The conditions: 1) weak proregularity (which is well studied in the literature) and 2) I -reduced and I -coreduced modules (being studied in this paper) are both necessary for the GM Duality and MGM Equivalence to hold. For the former, the aforementioned results hold in the derived category setting whereas for the later they hold in the module category setting. It is therefore not unreasonable to compare these two conditions. This subsection aims at achieving this.

Let $\mathbf{r} = (r_1, \dots, r_n)$ be a sequence of elements of a ring R . To this sequence, we associate the Koszul complex $K(R; \mathbf{r})$. For each $i \geq 1$, let \mathbf{r}^i be the sequence (r_1^i, \dots, r_n^i) . There is a corresponding Koszul complex $K(R; \mathbf{r}^i)$. Recall that an inverse system of R -modules $\{M_i\}_{i \geq 1}$ is called *pro-zero* if for every i there is some $j \geq i$ such that the R -homomorphism $M_j \rightarrow M_i$ is zero.

Definition 4.5. A finite sequence $\mathbf{r} = (r_1, \dots, r_n)$ in a ring R is *weakly proregular* if for every $p < 0$ the inverse system of R -modules $\{H^p(K(R; \mathbf{r}^i))\}_{i \geq 1}$ is pro-zero.

Definition 4.6. An ideal is *weakly proregular* if it is generated by a weakly proregular sequence.

If I is an idempotent ideal of a ring R , we know that every R -module is both I -reduced and I -coreduced. However, I need not be weakly proregular. On the other hand, every ideal I of a Noetherian ring R is weakly proregular but not all R -modules in this case are either I -reduced or I -coreduced. Instead, if the R -modules

are finitely generated and therefore also Noetherian, then there exists a positive integer k such that all such R -modules are I^k -reduced.

Motivated by [3] where strongly idempotent ideals were defined for an Artin algebra, strongly idempotent ideals for an arbitrary ring were defined, see [23, Definition 4.7]. In our setting, we have:

Definition 4.7. An ideal I of a ring R is *strongly idempotent* if for every $i \geq 1$,

$$\mathrm{Tor}_i^R(R/I, R/I) = 0.$$

Note that an ideal I of a ring R is idempotent precisely when $\mathrm{Tor}_1^R(R/I, R/I) = 0$. It then follows that, a strongly idempotent ideal is idempotent.

Proposition 4.8. *Let I be an idempotent ideal generated by a finite sequence in a ring R . I is strongly idempotent if and only if it is weakly proregular.*

Proof. By [23, Proposition 4.10], an idempotent ideal I is strongly idempotent if and only if the associated torsion class \mathcal{T}_I is weakly stable. However, by [22, Theorem 4.13], \mathcal{T}_I is weakly stable if and only if I is weakly proregular. \square

5. Further representability and applications

Following [18], let $E_R(M)$ denote the injective hull of an R -module M and E be the direct product of the injective hulls $E_R(R/\mathfrak{m})$ of the R -modules R/\mathfrak{m} , where \mathfrak{m} runs through the set of maximal ideals of R . E is an injective cogenerator of $R\text{-Mod}$. The general Matlis duality functor is given by $D(-) := \mathrm{Hom}_R(-, E)$.

Proposition 5.1. *Let R be a Noetherian ring and M be an R -module. M is reduced (resp. coreduced) if and only if the R -module $D(M)$ is coreduced (resp. reduced).*

Proof. Suppose that M is reduced. By [18, Lemma 1.4.6], for any ideal I of R , $R/I \otimes \mathrm{Hom}_R(M, E) \cong \mathrm{Hom}_R(\mathrm{Hom}_R(R/I, M), E) \cong \mathrm{Hom}_R(\mathrm{Hom}_R(R/I^2, M), E) \cong R/I^2 \otimes \mathrm{Hom}_R(M, E)$ so that $D(M) := \mathrm{Hom}_R(M, E)$ is coreduced. For the converse, suppose that the R -module $D(M)$ is coreduced. By [18, Lemma 1.4.6],

$\mathrm{Hom}_R(\mathrm{Hom}_R(R/I, M), E) \cong R/I \otimes \mathrm{Hom}_R(M, E) \cong R/I^2 \otimes \mathrm{Hom}_R(M, E) \cong \mathrm{Hom}_R(\mathrm{Hom}_R(R/I^2, M), E)$ for every ideal I of R . Since E is an injective cogenerator, we have for every ideal I of R , $\mathrm{Hom}_R(R/I, M) \cong \mathrm{Hom}_R(R/I^2, M)$ and this shows that M is reduced. The second part is immediate from Proposition 2.6. \square

Example 5.2. Let (R, \mathfrak{m}) be a Gorenstein local ring of dimension d and M be a finitely generated R -module. For any $0 \leq i \leq d$, the local cohomology R -module $H_{\mathfrak{m}}(M)$ is reduced (resp. coreduced) if and only if the R -module $\mathrm{Ext}_R^{d-i}(M, R)$ is

coreduced (resp. reduced). This is immediate from the local duality theorem and Proposition 5.1.

5.1. Representability of Γ_I .

Proposition 5.3. *Let I be any ideal of a ring R . For any two R -modules M and N , where M is I -coreduced we have*

$$\Gamma_I(\text{Hom}_R(M, N)) \cong \text{Hom}_R(\Lambda_I(M), N) \text{ and } \Gamma_I(D(M)) \cong D(\Lambda_I(M)).$$

Proof. For any I -coreduced R -module M : 1) $\Lambda_I(M) \cong R/I \otimes M$ and; 2) the R -module $\text{Hom}_R(M, N)$ is I -reduced and therefore $\text{Hom}_R(R/I, \text{Hom}_R(M, N)) \cong \Gamma_I(\text{Hom}_R(M, N))$. This together with the Hom-Tensor adjunction, we get

$$\begin{aligned} \text{Hom}_R(\Lambda_I(M), N) &\cong \text{Hom}_R(R/I \otimes M, N) \cong \text{Hom}_R(R/I, \text{Hom}_R(M, N)) \cong \\ &\Gamma_I(\text{Hom}_R(M, N)). \end{aligned}$$

By taking $N = E$, the second isomorphism is obtained. \square

Corollary 5.4. *Let I be any ideal of a ring R , M be an I -coreduced R -module and N any R -module.*

- (1) *The R -module $\text{Hom}_R(\Lambda_I(M), N)$ is I -torsion.*
- (2) *If M is I -complete, then $\text{Hom}_R(M, N)$ (and hence $\mathcal{D}(M)$) is I -torsion.*
- (3) *If $I^2 = I$, then $\Gamma_I(N) \cong \text{Hom}_R(\Lambda_I(R), N)$ and $\Gamma_I(E) = \mathcal{D}(\Lambda_I(R))$.*
- (4) *If $I^2 = I$ and R is I -complete (resp. $\Lambda_I(R) = 0$), then N is I -torsion (resp. I -torsion-free).*

Proof. (1) By Proposition 5.3, $\Gamma_I(\text{Hom}_R(M, N)) \cong \text{Hom}_R(\Lambda_I(M), N)$. Taking $M = \Lambda_I(M)$ and the fact that the functor $\Lambda_I(-)$ is idempotent on I -coreduced modules, we get $\Gamma_I(\text{Hom}_R(\Lambda_I(M), N)) \cong \text{Hom}_R(\Lambda_I(M), N)$.

(2) Immediate from part (1).

(3) If I is idempotent, then R is an I -coreduced R -module. By Proposition 5.3, $\Gamma_I(N) \cong \Gamma_I(\text{Hom}_R(R, N)) \cong \text{Hom}_R(\Lambda_I(R), N)$.

(4) This is immediate from part (3). \square

5.2. Representability of Λ_I .

Proposition 5.5. *Let I be a finitely generated ideal of a ring R , M an I -reduced R -module and N an injective R -module. We have*

$$\Lambda_I(\text{Hom}_R(M, N)) \cong \text{Hom}_R(\Gamma_I(M), N) \text{ and } \Lambda_I(D(M)) \cong D(\Gamma_I(M)).$$

Proof. By the fact that $\Gamma_I(M) \cong \text{Hom}_R(R/I, M)$ and [18, Lemma 1.4.6], we have $\text{Hom}_R(\Gamma_I(M), N) \cong \text{Hom}_R(\text{Hom}_R(R/I, M), N) \cong R/I \otimes \text{Hom}_R(M, N) \cong \Lambda_I(\text{Hom}_R(M, N))$, since $\text{Hom}_R(M, N)$ is I -coreduced under the conditions given in the hypothesis. Let $N := E$; we get $\Lambda_I(\text{Hom}_R(M, E)) \cong \text{Hom}_R(\Gamma_I(M), E)$. So that, $\Lambda_I(D(M)) \cong D(\Gamma_I(M))$. \square

Corollary 5.6. *Let I be a finitely generated ideal of a ring R and N an injective R -module. If M is an I -reduced R -module, then*

- (1) *the R -modules $\text{Hom}_R(\Gamma_I(M), N)$ and $\mathcal{D}(\Gamma_I(M))$ are I -complete.*
- (2) *M I -torsion implies that $\text{Hom}_R(M, N)$ (and hence $\mathcal{D}(M)$) is I -complete.*
- (3) *If R is an I -reduced R -module, $\Lambda_I(N) \cong \text{Hom}_R(\Gamma_I(R), N)$ and $\Lambda_I(E) \cong \mathcal{D}(\Gamma_I(R))$.*
- (4) *If R is I -reduced as an R -module and I -torsion (resp. $\Gamma_I(R) = 0$), then N is I -complete (resp. $\Lambda_I(N) = 0$).*

Proof. (1) By Proposition 5.5, $\Lambda_I(\text{Hom}_R(M, N)) \cong \text{Hom}_R(\Gamma_I(M), N)$. Taking M to be the module $\Gamma_I(M)$ and the fact that the functor $\Gamma_I(-)$ is idempotent, we get $\Lambda_I(\text{Hom}_R(\Gamma_I(M), N)) \cong \text{Hom}_R(\Gamma_I(M), N)$.

(2) Immediate from part (1).

(3) By Proposition 5.5, $\Lambda_I(N) \cong \Lambda_I(\text{Hom}_R(R, N)) \cong \text{Hom}_R(\Gamma_I(R), N)$.

(4) Immediate from part (3). \square

Proposition 5.7. *If I is an idempotent finitely generated ideal of a ring R , then the functor Λ_I is exact on a full subcategory of injective R -modules.*

Proof. By Corollary 5.6, if M is an injective R -module, then

$$\Lambda_I(M) \cong \text{Hom}_R(\Gamma_I(R), M),$$

i.e., on a full subcategory of injective R -modules the functors $\Lambda_I(-)$ and $\text{Hom}_R(\Gamma_I(R), -)$ are isomorphic. So, Λ_I is left exact since it is isomorphic to a left exact functor $\text{Hom}_R(\Gamma_I(R), -)$. However, it is also true that since $I^2 = I$, $\Lambda_I(-) \cong R/I \otimes -$ which is right exact. \square

Corollary 5.8. *For any finitely generated ideal I of a ring R , the diagram in Figure 2 is commutative. In this diagram, \mathcal{A}_I (resp. \mathcal{B}_I) denotes the full subcategory of R -Mod consisting of I -reduced (resp. I -coreduced) R -modules.*

Proof. By Proposition 5.3, $\Gamma_I(D(M)) \cong D(\Lambda_I(M))$. This establishes the first commutative square. The second commutative square is established by Proposition

$$\begin{array}{ccccccc}
 \cdots & \longrightarrow & \mathcal{A}_I & \xrightarrow{D(-)} & \mathcal{B}_I & \xrightarrow{D(-)} & \mathcal{A}_I & \xrightarrow{D(-)} & \mathcal{B}_I & \longrightarrow & \cdots \\
 & & \downarrow \Gamma_I & & \downarrow \Lambda_I & & \downarrow \Gamma_I & & \downarrow \Lambda_I & & \\
 \cdots & \longrightarrow & \mathcal{B}_I & \xrightarrow{D(-)} & \mathcal{A}_I & \xrightarrow{D(-)} & \mathcal{B}_I & \xrightarrow{D(-)} & \mathcal{A}_I & \longrightarrow & \cdots
 \end{array}$$

FIGURE 2. Commutative diagram I

5.5 which asserts that $\Lambda_I(D(M)) \cong D(\Gamma_I(M))$. Repeating this process leads to the required commutative diagram. \square

Corollary 5.9. *For any finitely generated idempotent ideal I of a ring R and for any R -module M , we have*

- (1) $\text{Hom}_R(R/I, \mathcal{D}(M)) \cong \mathcal{D}(R/I \otimes M)$.
- (2) $R/I \otimes \mathcal{D}(M) \cong \mathcal{D}(\text{Hom}_R(R/I, M))$.
- (3) *The following diagram in Figure 3 is commutative.*

$$\begin{array}{ccccccc}
 \cdots & \longrightarrow & R\text{-Mod} & \xrightarrow{D(-)} & R\text{-Mod} & \xrightarrow{D(-)} & R\text{-Mod} & \xrightarrow{D(-)} & R\text{-Mod} & \longrightarrow & \cdots \\
 & & \downarrow \text{Hom}_R(R/I, -) & & \downarrow R/I \otimes - & & \downarrow \text{Hom}_R(R/I, -) & & \downarrow R/I \otimes - & & \\
 \cdots & \longrightarrow & R\text{-Mod} & \xrightarrow{D(-)} & R\text{-Mod} & \xrightarrow{D(-)} & R\text{-Mod} & \xrightarrow{D(-)} & R\text{-Mod} & \longrightarrow & \cdots
 \end{array}$$

FIGURE 3. Commutative diagram II

5.3. Computation of natural transformations. In this subsection, we demonstrate that representability of Γ_I and Λ_I which is facilitated by I -reduced and I -coreduced modules makes it easy (by use of Yoneda Lemma) to compute natural transformations from the functors Γ_I and Λ_I to other functors under some suitable conditions.

Proposition 5.10. *For any ideal I of a ring R , and functors*

$$\Gamma_I : (R\text{-Mod})_{I\text{-red}} \rightarrow (R\text{-Mod})_{I\text{-cor}} \text{ and } I \otimes - : (R\text{-Mod})_{I\text{-red}} \rightarrow \mathbf{Set},$$

we have $\text{Nat}(\Gamma_I(-), \Gamma_I(-)) \cong R/I$ and $\text{Nat}(\Gamma_I(-), I \otimes -) \cong 0$.

Proof. Recall that $R/I \in (R\text{-Mod})_{I\text{-red}}$. Since the functor Γ_I is representable on $(R\text{-Mod})_{I\text{-red}}$, Yoneda Lemma asserts that for any functor $F : (R\text{-Mod})_{I\text{-red}} \rightarrow \mathbf{Set}$,

$$\text{Nat}(\text{Hom}_R(R/I, -), F(-)) \cong F(R/I).$$

It follows that $\text{Nat}(\Gamma_I(-), \Gamma_I(-)) \cong \Gamma_I(R/I) \cong (0 :_{R/I} I) \cong R/I$. For the second part, $\text{Nat}(\text{Hom}_R(R/I, -), I \otimes -) \cong I \otimes R/I \cong 0$. \square

Let R be a finite dimensional algebra over a field k and $R\text{-mod}$ a category of all finitely generated R -modules. Let $\mathcal{D} := \text{Hom}_k(-, k)$. The Nakayama functor \mathcal{V} is the composition of two contravariant functors $\text{Hom}_R(-, R)$ and \mathcal{D} , i.e.,

$$\mathcal{V} := \mathcal{D}\text{Hom}_R(-, R) : R\text{-mod} \rightarrow R\text{-mod}.$$

If $R\text{-proj}$ (resp. $R\text{-inj}$) denotes the full subcategory of $R\text{-mod}$ consisting of all projective R -modules (resp. injective R -modules), then the restriction of \mathcal{V} on $R\text{-proj}$ defines an equivalence $\mathcal{V} : R\text{-proj} \rightarrow R\text{-inj}$ between $R\text{-proj}$ and $R\text{-inj}$ with quasi-inverse $\mathcal{V}^{-1} := \text{Hom}_R(\mathcal{D}(R), -)$, see [2, Definition 2.8 & Proposition 2.10].

Proposition 5.11. *Let R be a Noetherian von-Neumann regular finite dimensional algebra over a field k . If R is self-injective, then for any $M \in R\text{-inj}$ and functors $\Lambda_I : R\text{-inj} \rightarrow \mathbf{Set}$ and $\mathcal{V}^{-1} : R\text{-inj} \rightarrow R\text{-proj}$, $\text{Nat}(\Lambda_I(M), \mathcal{V}^{-1}(M)) \cong k$.*

Proof. By hypothesis, $R \in R\text{-inj}$. By [6], Γ_I preserves injective modules defined over Noetherian rings. So, $\Gamma_I(R) \in R\text{-inj}$. By Corollary 5.6(3), $\Lambda_I(M) \cong \text{Hom}_R(\Gamma_I(R), M)$. Since R is Noetherian, there exists a positive integer t such that $\Gamma_I(R) \cong \text{Hom}_R(R/I^t, R)$. We can see that the functor $\Lambda_I : R\text{-inj} \rightarrow \mathbf{Set}$ is representable. So, given the functor $\mathcal{V}^{-1} : R\text{-inj} \rightarrow R\text{-proj}$, we invoke Yoneda Lemma to have $\text{Nat}(\Lambda_I(M), \mathcal{V}^{-1}(M)) \cong \text{Nat}(\text{Hom}_R(\Gamma_I(R), M), \mathcal{V}^{-1}(M)) \cong \mathcal{V}^{-1}(\Gamma_I(R))$. However, $\mathcal{V}^{-1}(\Gamma_I(R)) \cong \text{Hom}_R(\mathcal{D}(R), \Gamma_I(R))$. Substituting for $\mathcal{D}(R)$ and $\Gamma_I(R)$; and applying Yoneda embedding, we get the following functors

$$\mathcal{V}^{-1}(\Gamma_I(R)) \cong \text{Hom}_R(\text{Hom}_k(R, k),$$

$$\text{Hom}_R(R/I^t, R)) \cong \text{Hom}_k(k, R) \cong k. \quad \square$$

Lemma 5.12. *Let I be an ideal of a ring R . If R is a reduced (resp. I -reduced) R -module, then the full subcategory $R\text{-proj}$ of $R\text{-Mod}$ which consists of all projective R -modules has all modules reduced (resp. I -reduced).*

Proof. If R is I -reduced, then so is any free R -module; since such a module is isomorphic to R^n for some positive integer n and I -reduced modules are closed under taking direct sums. Since every projective R -module is a direct summand of

a free R -module, it follows that, all projective R -modules in this case are I -reduced. For the reduced case, the proof is similar. \square

Proposition 5.13. *Let I be an ideal of a ring R such that R is an I -reduced R -module and R/I is a projective R -module. Then for any $M \in R\text{-proj}$ we have $\Gamma_I : R\text{-proj} \rightarrow \mathbf{Set}$ and $\mathcal{V} : R\text{-proj} \rightarrow R\text{-inj}$; $\text{Nat}(\Gamma_I(M), \mathcal{V}(M)) \cong \mathcal{D}(0 :_R I)$.*

Proof. If $M \in R\text{-proj}$, then by Lemma 5.12, it is I -reduced and therefore $\Gamma_I(M) \cong \text{Hom}_R(R/I, M)$. The hypothesis satisfies conditions of the Yoneda Lemma. So, we have $\text{Nat}(\Gamma_I(M), \mathcal{V}(M)) \cong \text{Nat}(\text{Hom}_R(R/I, M), \mathcal{V}(M)) \cong \mathcal{V}(R/I)$. However, $\mathcal{V}(R/I) \cong \mathcal{D}\text{Hom}_R(R/I, R) \cong \mathcal{D}(0 :_R I)$. \square

Acknowledgement. The author would like to thank the referee for the valuable suggestions and comments. Part of this work was written while the author was visiting University of Glasgow and University of Oxford. He is grateful to Kobi Kremnizer, Balazs Szendroi and Michael Wemyss for the stimulating discussions and for the hospitality during his stay in the United Kingdom. He is also grateful to Annet Kyomuhangi and Amanuel Senbato Mamo for the comments.

References

- [1] L. Alonso Tarrío, A. Jeremías López and J. Lipman, *Local homology and cohomology on schemes*, Ann. Sci. École Norm. Sup., 30(1) (1997), 1-39.
- [2] I. Assem, D. Simson and A. Skowronski, *Elements of the Representation Theory of Associative Algebras, Volume 1: Techniques of Representation Theory*, Cambridge University Press, 2006.
- [3] M. Auslander, M. I. Platzeck and G. Todorov, *Homological theory of idempotent ideals*, Trans. Amer. Math. Soc., 332(2) (1992), 667-692.
- [4] T. Barthel, D. Heard and G. Valenzuela, *Local duality in algebra and topology*, Adv. Math., 335 (2018), 563-663.
- [5] T. Barthel, D. Heard and G. Valenzuela, *Derived completion for comodules*, Manuscripta Math., 161 (2020), 409-438.
- [6] M. P. Brodmann and R. Y. Sharp, *Local Cohomology: An Algebraic Introduction with Geometric Applications*, Cambridge Studies in Advanced Mathematics, 136, Cambridge University Press, Cambridge, Second Edition, 2013.
- [7] K. Divaani-Aazar, H. Faridian and M. Tousi, *Local homology, finiteness of Tor modules and cofiniteness*, J. Algebra Appl., 16(12) (2017), 1750240 (10 pp).

- [8] T. H. Freitas, V. H. Jorge-Pérez, C. B. Miranda-Neto and P. Schenzel, *Generalized local duality, canonical modules, and prescribed bound on projective dimension*, J. Pure Appl. Algebra, 227(2) (2023), 107188, (17 pp).
- [9] J. P. C. Greenlees and J. P. May, *Derived functors of I -adic completion and local homology*, J. Algebra, 149(2) (1992), 438-453.
- [10] R. Hartshorne, *Local Cohomology*, a seminar given by A. Grothendieck, Harvard University, Fall, 1961, Lecture Notes in Mathematics, 41, Springer-Verlag, Berlin-New York 1967.
- [11] G. Kempf, *The Grothendieck-Cousin complex of an induced representation*, Adv. in Math., 29(3) (1978), 310-396.
- [12] A. Kyomuhangi and D. Ssevviiri, *The locally nilradical for modules over commutative rings*, Beitr. Algebra Geom., 61(4) (2020), 759-769.
- [13] A. Kyomuhangi and D. Ssevviiri, *Generalised reduced modules*, Rend. Circ. Mat. Palermo (2), 72(1) (2023), 421-431.
- [14] T. K. Lee and Y. Zhou, *Reduced modules, rings, modules, algebras and abelian groups*. Lecture Notes in Pure and Applied Math, 236, 365-377, Marcel Dekker, New York, 2004.
- [15] S. MacLane, *Categories for the Working Mathematician*, Berlin, Heidelberg, New York, Springer-Verlag, 1971.
- [16] M. Porta, L. Shaul and A. Yekutieli, *On the homology of completion and torsion*, Algebr. Represent. Theory, 17(1) (2014), 31-67.
- [17] M. B. Rege and A. M. Buhphang, *On reduced modules and rings*, Int. Electron. J. Algebra, 3 (2008), 58-74.
- [18] P. Schenzel and A. M. Simon, *Completion, Čech and Local Homology and Cohomology*, Springer Monographs in Mathematics, 2018.
- [19] R. Y. Sharp, *Local cohomology theory in commutative algebra*, Quart. J. Math. Oxford Ser. (2), 21 (1970), 425-434.
- [20] D. Ssevviiri, *Nilpotent elements control the structure of a module*, arXiv:1812.04320 [math.RA], (2018).
- [21] N. Suzuki, *On the generalized local cohomology and its duality*, J. Math. Kyoto Univ., 18 (1978), 71-85.
- [22] R. Vyas and A. Yekutieli, *Weak proregularity, weak stability, and the noncommutative MGM equivalence*, J. Algebra, 513 (2018), 265-325.
- [23] R. Vyas, *Weakly stable torsion classes*, Algebr. Represent. Theory, 22(5) (2019), 1183-1207.

- [24] S. Yassemi, *Generalized section functors*, J. Pure Appl. Algebra, 95(1) (1994), 103-119.
- [25] A. Yekutieli, *On flatness and completion for infinitely generated modules over Noetherian rings*, Comm. Algebra, 39(11) (2011), 4221-4245.

David Ssevviiri

Department of Mathematics

College of Natural Sciences

Makerere University

P.O. Box 7062, Kampala, Uganda

e-mail: david.ssevviiri@mak.ac.ug or ssevviiridaudi@gmail.com