



# Determination of phenolic acids in pomegranate juices by HPLC-DAD

Mürüvvet Karakaplan<sup>1\*</sup>, Mustafa Özcan<sup>1</sup>

<sup>1</sup> Department of Chemistry, Faculty of Science, Istanbul Technical University, Maslak, 34469 Istanbul, Turkey

(Dergiye gönderilme tarihi: 29 Haziran 2017, Kabul Tarihi: 19 Temmuz 2017)

## Abstract

In this study, five phenolic acids, gallic, chlorogenic, caffeic, p-coumaric and ferulic acid were determined in freshly squeezed pomegranate juice, commercial pomegranate juice and pomegranate juice concentrate beverages. For this purpose, an analytical method using high-performance liquid chromatography (HPLC), with a combination of photodiode array detection (DAD), was developed for the characterization and quantification of phenolic acid in pomegranate juices. Then the analyses of pomegranate juices were performed. The limit of detection (LOD, S/N=3) of individual compounds was gallic acid 0.1 mg/L, chlorogenic acid 0.01 mg/L, caffeic acid 0.03mg/L, p-coumaric acid 0.02 mg/L and ferulic acid 0.04 mg/L. The limit of quantitation (LOQ, S/N=10) of individual compounds was gallic acid 0.4 mg/L, chlorogenic acid 0.05 mg/L, caffeic acid 0.1 mg/L, p-coumaric acid 0.1 mg/L and ferulic acid 0.1 mg/L. The relative standard deviation (% RSD) in commercial pomegranate juice was found to be gallic acid 0.08%, chlorogenic acid 0.32%, caffeic acid 0.21%, p-coumaric acid 0.08% and ferulic acid 0.38%. The relative standard deviation (% RSD) in freshly squeezed pomegranate juice was found to be gallic acid 0.01%, chlorogenic acid 0.46%, caffeic acid 0.17%, p-coumaric acid 0.04% and ferulic acid 0.8%. The relative standard deviation (% RSD) in pomegranate concentrate juice was found to be gallic acid 0.78%, chlorogenic acid 0.25%, caffeic acid 0.49%, p-coumaric acid 0.36% and ferulic acid 0.17%.

**Keywords:** HPLC; gallic acid; chlorogenic acid; caffeic acid; p-coumaric acid; ferulic acid.

## 1. Introduction

Phenolic acids are widely distributed in the plant cause tailing and broadening of their peaks. In this kingdom and are present in, e.g. tea, red wine, fruits, paper, we report the results of a simple and rapid beverages and various medicinal plants [1].

The Pomegranate juice was reported to be effective in the prevention of attherosclerosis, low-density lipoprotein oxidation, prostate cancer, platelet aggregation and various cardiovascular diseases [2]. Pomegranate fruit juice has potential chemopreventive agents for prostate cancer [3]. Sumner et al. showed that daily consumption of pomegranate juice for 3 months may decrease myocardial is chemia and improve myocardial perfusionin patients who have is chemic coronary heart disease [4]. Anoosha et al. conclude that consumption of pomegranate juice may be proven to be beneficial in attenuation of atherosclerosis development [5].

Pomegranate juice has gained lots of attention in the last years due to its nutritional value and antioxidant properties. Pomegranate fruit has three-fold higher antioxidant activity than that of red wine or green tea [6] and two-, six- and eight-fold higher levels than grape/cranberry, grape fruit, and orange juice,

respectively [7]. Pomegranate (*Punica granatum*) is an important source of bioactive compounds. These compounds are primarily phenolic or polyphenolic, flavonoids (flavonois, flavoneflavanones, catechins and isoflavones) and related compounds (phenolic acids, chalcones and isoflavones) [8].

The term “phenolic acids” refers to phenols that have one carboxylic acid functionality. But, when describing plant metabolites, it refers to a distinct group of organic acids. In the view of the importance of these substances for health, accurate analysis methods for their determination in foodstuffs are required. In the literature on the determination of phenolic acids in foodstuffs, including UV spectrophotometry, gas chromatography, thin-layer chromatography, high-performance liquid chromatography (HPLC), liquid chromatography mass spectrometry and capillary electrophoresis systems are used [9].

(*Punica granatum*) has gainedlots ofattention for its health benefits in the past couple years.


Pomegranate (*Punica granatum* Linn.) is a very rich source of anthocyanins (cyanidin 3,5-di and 3-O-glucoside, delphinidin 3,5-di and 3-O-glucoside, pelargonidin 3,5-di and 3-O-glucoside), punicalagin isomers, different flavanols (catechins ascatechins

<sup>1</sup> Corresponding Author: Department of Chemistry, Faculty of Science, Istanbul Technical University, Maslak, 34469 Istanbul, Turkey, [muruvvetkarakaplan@hotmail.com](mailto:muruvvetkarakaplan@hotmail.com)



and gallic acids with isocratic elution. The mobile phase composition was optimized to obtain a good resolution between the different peaks detected for juice samples by HPLC-DAD.

In this study, we tried to determine the optimum isocratic HPLC conditions for determination of the five phenolic acids. As a consequence, the sample solution was chromatographed on an column ODS2 with a mobile phase water:acetic acid (99.5:0.5)-methanol at a flow-rate of 0.8 ml/min. Using these proposed HPLC conditions, a well resolved chromatogram between the different peaks detected for the standard by HPLC was obtained (Figure 1).



**Figure 1.** Chromatograms of HPLC analyses of standards using DAD at 280 nm (A) and 320 nm (B). Peaks: (1) gallic acid; (2) chlorogenic acid; (3) caffeic acid; (4) p-coumaric acid; (5) ferulic acid.

The within-day repeatability (n=3) and between day precision (n=3) of retention times were within 0.01 and 0.86 % relative standard deviation (RSD), drink respectively. RSD values are given in Table 2. The limit of detection (LOD, S/N=3) of individual compounds at 280 nm and 320 nm are given in Table 3.

**Table 2.** RSD values

	%RSD (HPLC-DAD)		
	commercial pomegranate juice	freshly squeezed pomegranate juice	pomegranate concentrate juice
Caffeic acid	0.08	0.01	0.78
p-coumaric acid	0.32	0.46	0.25
Ferulic acid	0.21	0.17	0.49
Gallic acid	0.08	0.04	0.36
Chlorogenic acid	0.38	0.8	0.17

**Table 3.** The limit of detection (LOD), the limit of quantitation (LOQ), Coefficient

	LOD (mg/L)	LOQ (mg/L)	Correlation Coefficient
Gallic acid	0.1	0.4	0.9997
Chlorogenic acid	0.01	0.05	0.9996
Caffeic acid	0.03	0.1	0.9999
p-coumaric acid	0.02	0.1	0.9999
Ferulic acid	0.04	0.1	0.9999

a Correlation coefficients of the regression equation  $y = a + bx$ , where x is the phenolic compound concentration (ppm) and y the peak area.


Table 3, where a, b and r were the coefficients of the regression equation  $y = ax + b$  x being the concentration of the phenolic compound (mg/kg), y the peak area, and r the correlation coefficient of the equation. All the phenolic compounds showed good linearity ( $r^2 \geq 0.9996$ ) and obeyed Beer's law in the concentration range investigated [16].

In this work, phenolic compounds of freshly squeezed pomegranate juice, commercial pomegranate juice and pomegranate juice concentrate were determined by using HPLC-DAD.

Phenolic compounds identified in freshly squeezed pomegranate juice, commercial pomegranate juice and pomegranate juice concentrate were gallic, chlorogenic, caffeic, p-coumaric and ferulic acid.




73.3mg/L of caffeic acid, 30.8 mg/L of p-coumaric acid, 27.9 mg/L of ferulic acid, 20.1 of mg/L gallic acid and 15.2 mg/L of chlorogenic acid were quantified in commercial pomegranate juice by using HPLC-DAD (Figure 2).




**Figure 2.** Chromatograms of HPLC analyses of commercial pomegranate juice using DAD at 280 nm (A) and 320 nm (B). Peaks: (1) gallic acid; (2) chlorogenic acid; (3) caffeic acid; (4) p-coumaric acid; (5) ferulic acid.

422.8 mg/L of caffeic acid, 45.5 mg/L of p-coumaric acid, 34.7 mg/L of ferulic acid, 24.3 of mg/L gallic acid and 17.1 mg/L of chlorogenic acid were quantified in freshly squeezed pomegranate juice by using HPLC-DAD (Figure 3).



**Figure 3.** Chromatograms of HPLC analyses of freshly squeezed pomegranate juice using DAD at 280 nm (A) and 320 nm (B). Peaks: (1) gallic acid; (2) chlorogenic acid; (3) caffeic acid; (4) p-coumaric acid; (5) ferulic acid.

61 mg/L of caffeic acid, 21.4 mg/L of p-coumaric acid, 20.6 mg/L of ferulic acid, 15.9 of mg/L gallic acid and 11.5 mg/L of chlorogenic acid were quantified in pomegranate juice concentrate by using HPLC-DAD (Figure 4).



**Figure 4.** Chromatograms of HPLC analyses of pomegranate juice concentrate using DAD at 280 nm (A) and 320 nm (B): Peaks: (1) gallic acid; (2) chlorogenic acid; (3) caffeic acid; (4) *p*-coumaric acid; (5) ferulic acid.

The combination of DAD has been shown to be a powerful technique for the fast finger printing characterization of fruit extracts [14], food-based products [15], food plant and food industry by-products [17].

We analyzed three juices using this method and the experimental result indicated that freshly squeezed pomegranate juice contained an especially high concentration of caffeic acid (422.8 mg/L) in 5 of its phenolic acids. It was also shown that the main acid of the other juices was caffeic acid (73.3 mg/L) in the commercial pomegranate juice, caffeic acid (61 mg/L) in the pomegranate juice concentrate.

The experimental results indicated that freshly squeezed pomegranate juice contained a high concentration of caffeic (422.8 mg/L), which can be hydrolyzed to caffeic acid. It was shown also that the phenolic acids occur mainly in bound forms in commercial pomegranate juice and pomegranate juice and that their bound caffeic acid contents were high (73.3 and 61 mg/L, respectively) compared to those of the other phenolic acids.

The phenolic acids, components of human foods, have shown interesting activities as inhibitors of mutagenic and carcinogenic

processes. However, these studies were performed with pure crystalline substances, while in foods the substances are usually present as bound phenolic acids and little is known about the fate of most plant phenolics after ingestion. In order to utilize the anticarcinogenic properties of the phenolic acids for reduction of risk of human cancer, further studies are necessary. However, we know that chlorogenic acid, a bound caffeic acid, is similar to caffeic acid, an inhibitor of carcinogenesis in animal studies and in vitro [12].

Polyphenolic compounds in fruit been analyzed by using liquid-liquid extraction (LLE) followed by high performance liquid chromatography with ultraviolet detection [18].

Phenolic acid contents of pomegranate juices are given in Table 4.

**Table 4.** Phenolic acid contents of pomegranate juices

Phenolic Acid (mg/L)	Commercial pomegranate juice	Freshly squeezed pomegranate juice	Pomegranate juice concentrate
Caffeic acid	73.3±0.06	422.8±0.06	61±0.05
<i>p</i> -coumaric acid	30.8±0.1	45.5±0.2	21.4±0.05
Ferulic acid	27.9±0.06	34.7±0.06	20.6±0.1
Gallic acid	20.1±0.02	24.3±0.01	15.9±0.06
Chlorogenic acid	15.2±0.06	17.1±0.1	11.5±0.02

## 4. Conclusion

A convenient and rapid DAD-HPLC method for the identification and quantification of the main phenolic compounds in pomegranate in a single chromatographic run was developed, validated and successfully performed.

In this study, for the first time, phenolic contents of freshly squeezed pomegranate juice and pomegranate juice concentrate obtained from the Denizli in Turkey were examined. The results indicated that phenolic contents of freshly squeezed pomegranate juice were higher than that of pomegranate juice concentrate. Phenolic compounds of boiled and pasteurized juices were decreased. The methanol extraction with posterior acid hydrolysis which was practiced has shown to be suitable both for phenolic standards and pomegranate extracts. A methanol-based mobile phase was employed, and the convenience of this organic solvent can substitute acetonitrile as the most commonly used solvent. An exhaustive method validation was achieved, and it showed that the proposed method is linear in the studied concentration range, sensible, precise, and accurate. The phenolic contents obtained for three pomegranate juices from Turkey was in agreement with those reported in literature.

## References

- [1] Amaruka, Y., Okada, M., Tsuji, S., Tonogai, Y. (2000). Determination of phenolic acids in fruit juices by isocratic column liquid chromatography. *Journal of Chromatography A*, 891, 183-188.

- [2] Naveena, B.M., Sen, A.R., Vaithyanathan, S., Patil, G., Kondarah, N. (2010). Antioxidant Potential of Pomegranate Juice Incooked Chicken Patties. *Journal Of Musclefoods*, 21:557-569.
- [3] Naiki-Ito, A., Chewonarin, T., Tang, M., Pitchakarn, P., Kuno, T., Ogawa, K., Asamoto, M., Shiraiand, T., Takahashi, S. (2015). Ellagic Acid, A Component of Pomegranate Fruit Juice, Suppresses Androgen-Dependent Prostate Carcinogenesis. *Induction of Apoptosis The Prostate*, 75:151-160.
- [4] Sumner, M.D., Elliott-eller, M., Weidner, G., Daubenmier, J.J., Chew, H.M., Marlin, R., Raisin, C.J., Ornish, D. (2005). Effects of Pomegranate Juice Consumption on Myocardial Perfusion in Patients With Coronary Heart Disease. *The American Journal of Cardiology*, 96:810-814.
- [5] Anoosha, E., Mojtabaa, E., Fatemeha, S. (2010). Study The Effect of Juice of Two Variety of Pomegranate Ondecreasing Plasma LDL Cholesterol. *Procedia Social and Behavioral Sciences*, 2:620-623.
- [6] Gil, M.I., Tomas-Barberan, F.A., Hess-Pierce, B., Holcroft, D.M., Kader, A.A., (2000). Antioxidant Activity of Pomegranate Juice and Its Relationship with Phenoliccomposition and Processing. *J. Agric. Food Chem.*, 48:4581-4589.
- [7] Rosenblat, M., Aviram, M., Seeram, N.P., Heber, D. (Eds.) (2006). *In Vitro Studies in Pomegranates: Ancient Roots to Modern Medicine*. Taylor and Francis Group, New York, 31-43.
- [8] Aviram, M., Dornfeld, L., Rosenblat, M., Volkova, N., Kaplan, M., Coleman, R., Hayek, T., Presser, D., Fuhrman, B. (2000). Pomegranate Juice Consumption Reduces Oxidative Stress, Atherogenic Modification To Ldl And Platelet Aggregation: Studies In Human And In Atherosclerotic Apolipoprotein Deficient Mice. *Am. J. Clin.Nutr.*,71:1062-1076.
- [9] Chen, H., Zuo, Y., Deng, Y. (2001). Separation and determination of flavanoids and other phenolic compounds in cranberry juice by high-performance liquid chromatography. *Journal of Chromatography A*, 913:387-395.
- [10] Kelebek, H., Selli, S., Canbas, A., Cabaroğlu, T. (2009). HPLC determination of organic acids, sugars, phenolic compositions and antioxidant capacity of orange juice and orange wine made from a Turkish cv. Kozan. *Microchemical Journal*, 91:187-192.
- [11] Kermasha, S., Goetghebeur, M., Dumont, J., Couture, R. (1995). Analyses of phenolic and furfural compounds in concentrated and non-concentrated apple juices. *Food Research International*, 28: 245-252.
- [12] Shahrzad, S., Irmgard, B. (1996) Determination of some pharmacologically active phenolic acids in juices by high-performance liquid chromatography. *Journal of Chromatography A*, 741:223-231.
- [13] Hakkinen, S., Heinonen, M., Karenlampi, S., Mykkanen, H., Ruuskanen, J., Torronen, R. (1999). Screening of selected flavonoids and phenolic acids in 19 berries. *Food Research International*, 32:345-353.
- [14] Fukumoto, L.R., Mazza, G. (2000). Assessing antioxidant and prooxidant activities of phenolic compounds. *J. Agric. Food Chem.*, 48:3597-3604.
- [15] Alasalvar, C., Grigor, J.M., Zhang, D., Quantick, P.C., Shahidi, F. (2002). Comparison of volatiles, phenolics, sugars, antioxidant vitamins, and sensory quality of different colored carrot varieties. *J. Agric. Food Chem.*, 50:2039-2041.
- [16] Escarpe, A., Gonzalez, M.C.(2000). Evaluation of High-Performans Liquid Chromatography for Determination of Phenolic Compounds in Pear Horticultural Curtivars. *Chromatographia*, 51:37-43.
- [17] Simirgiotis, Mario, J., Schmeda-Hirschmann, G. (2010). Determination of phenolic composition and antioxidant activity in fruits, rhizomes and leaves of the white strawberry (*Fragaria chiloensis* spp. *Chiloensis* form *chiloensis*) using HPLC-DAD-ESI-MS and free radical quenching techniques. *Journal of Food Composition and Analysis*, 23:545-553.
- [18] Dennis, F., Giovanni, C., Gianni, S., Filippo, M., Sauro, V., Enrico, M., Roberto, B. (2014). Quantitative Profiling of Volatile and Phenolic Substances in the Wine Vernaccia di Serrapetrona by Development of an HS-SPME-GC-FID/MS Method and HPLC-MS. *Food Anal. Methods*, 7:1651-1660.