



Research Article

## Intuitionistic fuzzy hypersoft topology and its applications to multi-criteria decision-making

Adem YOLCU<sup>1,\*</sup>

<sup>1</sup>Department of Mathematics, Kafkas University, Kars, 36000, Türkiye

### ARTICLE INFO

#### Article history

Received: 17 Mar 2021

Accepted: 10 May 2021

#### Keywords:

Intuitionistic Fuzzy Hypersoft Set; Intuitionistic Fuzzy Hypersoft Topology; IFH Interior(Closure); IFH Basis; MCDM

### ABSTRACT

The aim of this paper is to introduce the concept of intuitionistic fuzzy hypersoft topology. Certain properties of intuitionistic fuzzy hypersoft (IFH) topology like IFH basis, IFH subspace, IFH interior and IFH closure are investigated. Furthermore, the multicriteria decision making (MCDM) algorithms with aggregation operators based on IFH topology are developed. In Algorithm 1 and Algorithm 2, MCDM problem is applied for IFH sets and IFH topology, respectively. Any real-life implementations of the proposed MCDM algorithms are demonstrated by numerical illustrations.

**Cite this article as** Adem Y. Intuitionistic fuzzy hypersoft topology and its applications to multi-criteria decision-making. Sigma J Eng Nat Sci 2023;41(1):106–118.

### INTRODUCTION

Decision-making is an everyday task that can be seen as a method of rating a set of alternatives or choosing the best one(s) from them based on the knowledge given by the decision. Multi criteria decision making (MCDM) refers to a decision-making mechanism in which alternatives are measured on the basis of many attributes, representing the success of alternatives from an individual viewpoint. Its goal is to discern the most precise choice from potential alternatives. A provided option needs to be assessed by the person making the decision by different forms of assessment conditions, such as numbers, intervals, etc. However, it is difficult for one person in a variety of cases, as there are different uncertainties within the results, to choose the

right one due to lack of competence or violation. As a result, a wide variety of hypotheses have been proposed to quantify those threats and to track the operation.

Fuzzy set theory, initiated by Zadeh [32] in 1965, is an important mathematical method for modeling and controlling uncertainty based on an incremental approach. The idea of fuzzy sets plays a key role in the field of soft computing, which deals with complexity, partial truth, imprecision and approximation in order to achieve durability, robustness and low solution costs. In 1986, Atanassov [4] suggested the notion of intuitionistic fuzzy sets, defined by both membership and non-membership functions. Intuitionistic fuzzy sets expand fuzzy sets in a meaningful

\*Corresponding author.

\*E-mail address: yolcu.adem@gmail.com

This paper was recommended for publication in revised form by Regional Editor Vildan ÇETKİN



manner that are more suited for catching the confusion created by indecisiveness and lack of attention to human cognition. Intuitionistic fuzzy sets expand fuzzy sets in a practical manner, and are more suited for catching the confusion created by indecisiveness and lack of attention to human cognition. Xu and Zhao [29] offered a straightforward and intuitionistic viewpoint on the fusion of knowledge for intuitionistic fuzzy decision-making. Many other hypotheses have been established, such as cubic intuitionistic fuzzy sets [13], interval-valued intuitionistic fuzzy sets [5], linguistic interval-valued intuitionistic fuzzy sets [14], etc. used by researchers. Recently, Alcantud et al. [2] introduced a method for aggregating an infinite series of intuitionistic fuzzy sets; they also built scores and accuracy functions for temporary intuitionistic fuzzy sets and used the suggested decision-making functions. As a matter of principle, the ideas listed above are considered by researchers, and the total of their two memberships and non-membership values cannot exceed one.

Atanassov's intuitionistic fuzzy set deals only with incomplete data due to membership and non-membership values, but intuitionistic fuzzy sets cannot deal with conflicting and imprecise knowledge. In 1999, Molodtsov [19] suggested soft set theory, which provides a general mathematical structure for dealing with uncertainty. Maji et al. [18] presented an intuitionistic fuzzy soft set (IFSS) with simple operations and properties. It should be remembered that the parameters will not always be crisp, but may be intuitionistic fuzzy in nature. Various algebraic structures of intuitionistic fuzzy soft sets are studied in [10, 11, 12, 16]. Coker [8] has presented and researched the concept of intuitionistic fuzzy topological spaces. Li et al. [17] initiated intuitionistic fuzzy soft topology constructs on intuitionistic fuzzy soft sets. They discussed the notions of intuitionistic fuzzy soft open (closed) sets, intuitionistic fuzzy soft interior (closure) and intuitionistic fuzzy soft base in intuitive fuzzy soft topological spaces. Many different studies have been done on these hybrid topological structures [3, 6, 7, 21, 22, 23, 27, 28].

Smarandache [26] generalized the concept of soft sets to hypersoft sets by replacing the function of a single parameter with a multi-parameter (sub-attributes) function specified in the cartesian product with  $n$  different attributes. The developed HSS is more versatile than soft sets and is more suited for decision-making environments. He also introduced further extensions of hypersoft sets, such as crisp hypersoft sets, fuzzy hypersoft sets, intuitionistic fuzzy hypersoft sets, neutrosophic hypersoft sets, and plithogenic hypersoft sets. Nowadays, the principle of hypersoft sets and its extensions is making rapid progress [1, 9, 15, 20, 25, 31, 33, 34].

In this paper, using the notion of an intuitionistic fuzzy hypersoft set, we describe intuitionistic fuzzy hypersoft topological spaces. The purpose of this paper is to explore some of the essential properties of intuitionistic fuzzy

hypersoft topological spaces. Later we define the operations of the intuitionistic fuzzy hypersoft closure, intuitionistic fuzzy hypersoft interior, intuitionistic fuzzy hypersoft basis and intuitionistic fuzzy hypersoft sub-space topology. The features of all these defined concepts were examined and the study was enriched with appropriate examples. The main contribution in this paper is the implementation of a decision-making algorithm that uses intuitionistic fuzzy hypersoft and we demonstrate the suitability of this algorithm in real-life scenarios. Intuitionistic fuzzy hypersoft topology was used for the first time in the decision-making method applied to a problem in the field of health. Thus, it is presented how effective intuitionistic fuzzy hypersoft topology is in decision making methods.

**PRELIMINARIES**

**Definition 1** [4] An intuitionistic fuzzy set  $H$  in  $U$  is  $H = \{(u, \theta_H(u), \sigma_H(u)) : u \in U \text{ where } \theta_H : U \rightarrow [0, 1], \sigma_H : U \rightarrow [0, 1] \text{ with the condition } 0 \leq \theta_H(u) + \sigma_H(u) \leq 1, \forall u \in U. \theta_H, \sigma_H \in [0, 1] \}$  denote the degree of membership and non-membership of  $u$  to  $H$ , respectively. The set of all intuitionistic fuzzy sets over  $U$  will be denoted by  $IFP(U)$ .

**Definition 2** [19] Let  $U$  be an initial universe,  $E$  be a set of parameters and  $P(U)$  be a power set of  $U$ . A pair  $(H, E)$  is called a soft set over  $U$ , where  $H$  is a mapping  $H : E \rightarrow P(U)$ . In other words, the soft set is a parameterized family of sub-sets of the set  $U$ .

**Definition 3** [18] Let  $U$  be an initial universe and  $E$  be a set of parameters. A pair  $(H, E)$  is called an intuitionistic fuzzy soft set over  $U$ , where  $H$  is a mapping given by,  $H : E \rightarrow IFP(U)$ .

In general, for every  $e \in E$ ,  $H(e)$  is an intuitionistic fuzzy set of  $U$  and it is called intuitionistic fuzzy value set of parameter  $e$ . Clearly,  $H(e)$  can be written as a intuitionistic fuzzy set such that  $H(e) = \{(u, \theta_H(u), \sigma_H(u)) : u \in U\}$ .

**Definition 4** [26] Let  $U$  be the universal set and  $P(U)$  be the power set of  $U$ . Consider  $e_1, e_2, e_3, \dots, e_n$  for  $n \geq 1$ , be  $n$  well-defined attributes, whose corresponding attribute values are respectively the sets  $E_1, E_2, \dots, E_n$  with  $E_i \cap E_n = \emptyset$ , for  $i \neq j$  and  $i, j \in \{1, 2, \dots, n\}$ , then the pair  $(H : E_1 \times E_2 \times \dots \times E_n \rightarrow P(U))$  is said to be Hypersoft set over  $U$  where  $H : E_1 \times E_2 \times \dots \times E_n \rightarrow P(U)$ .

**Definition 5** [30] Let  $U$  be the universal set and  $IFP(U)$  be a family of all intuitionistic fuzzy set over  $U$ . Consider  $e_1, e_2, e_3, \dots, e_n$  for  $n \geq 1$ , be  $n$  well-defined attributes, whose corresponding attribute values are respectively the sets  $E_1, E_2, \dots, E_n$  with  $E_i \cap E_j = \emptyset$ , for  $i \neq j$  and  $i, j \in \{1, 2, \dots, n\}$ . Let  $A_i$  be the nonempty subset of  $E_i$  for each  $i = 1, 2, \dots, n$ . An intuitionistic fuzzy hypersoft set defined as the pair  $(H, A_1 \times A_2 \times \dots \times A_n)$  where;  $H : A_1 \times A_2 \times \dots \times A_n \rightarrow IFP(U)$  and

$$H(A_1 \times A_2 \times \dots \times A_n) = \left\{ \left\langle \alpha, \left( \frac{u}{\theta_{H(\alpha)}(u), \sigma_{H(\alpha)}(u)} \right) \right\rangle : u \in U, \alpha \in A_1 \times A_2 \times \dots \times A_n \subseteq E_1 \times E_2 \times \dots \times E_n \right\}$$

where  $\theta$  and  $\sigma$  are the membership and non-membership value, respectively such that  $0 \leq \theta_{H(\alpha)}(u), \sigma_{H(\alpha)}(u) \leq 1$  and  $\theta_{H(\alpha)}(u), \sigma_{H(\alpha)}(u) \in [0, 1]$ . For sake of simplicity, we write the symbols  $\Delta$  for  $E_1 \times E_2 \times \dots \times E_n$ ,  $\Omega$  for  $A_1 \times A_2 \times \dots \times A_n$  and  $\alpha$  for an element of the set  $\Gamma$ .

**Definition 6** [30] i) An intuitionistic fuzzy hypersoft set  $(H, \Delta)$  over the universe  $U$  is said to be null intuitionistic fuzzy hypersoft set and denoted by  $0_{(U, IFF, \Delta)}$  if for all  $u \in U$  and  $\xi \in \Delta$ ,  $\theta_{H(\xi)}(u) = 0$  and  $\sigma_{H(\xi)}(u) = 1$ .

ii) An intuitionistic fuzzy hypersoft set  $(H, \Delta)$  over the universe  $U$  is said to be absolute intuitionistic fuzzy hypersoft set and denoted by  $1_{(U, IFF, \Delta)}$  if for all  $u \in U$  and  $\xi \in \Delta$ ,  $\theta_{H(\xi)}(\alpha) = 1$  and  $\sigma_{H(\xi)}(u) = 0$ .

**Definition 7** [30] Let  $U$  be an initial universe set and  $(H, \Omega_1), (H, \Omega_2)$  be two intuitionistic fuzzy hypersoft sets over the universe  $U$ . We say that  $(H, \Omega_1)$  is an intuitionistic fuzzy hypersoft subset of  $(G, \Omega_2)$  and denote  $(H, \Omega_1) \subseteq (G, \Omega_2)$  if

- i)  $\Omega_1 \subseteq \Omega_2$ ,
- ii) For any  $\alpha \in \Omega_1, H(\alpha) \subseteq G(\alpha)$ ,

That is for all  $u \in U$  and  $\alpha \in \Omega_1$ ,  $\theta_{H(\alpha)}(u) \leq \theta_{G(\alpha)}(u)$  and  $\sigma_{H(\alpha)}(u) \geq \sigma_{G(\alpha)}(u)$ .

**Definition 8** [30] The complement of intuitionistic fuzzy hypersoft set  $(H, \Omega)$  over the universe  $U$  is denoted by  $(H, \Omega)^c$  and defined as  $(H, \Omega)^c = (H^c, \Omega)$ , where  $H^c: E_1 \times E_2 \times \dots \times E_n = \Delta \rightarrow IFF(U)$  and  $H^c(\Omega) = (H(\Omega))^c$  for all  $\Omega \subseteq \Delta$ . Thus if

$$(H, \Omega) = \left\langle \alpha, \left( \frac{u}{\theta_{H(\alpha)}(u), \sigma_{H(\alpha)}(u)} \right) \right\rangle : u \in U, \alpha \in \Omega \Bigg\}, \text{ then}$$

$$(H, \Omega)^c = \left\langle \alpha, \left( \frac{u}{\sigma_{H(\alpha)}(u), \theta_{H(\alpha)}(u)} \right) \right\rangle : u \in U, \alpha \in \Omega \Bigg\}.$$

**Definition 9** [30] Let  $U$  be an initial universe set,  $\Omega_1, \Omega_2 \subseteq \Delta$  and  $(H, \Omega_1), (G, \Omega_2)$  be two intuitionistic fuzzy hypersoft sets over the universe  $U$ . The union of  $(H, \Omega_1)$  and  $(G, \Omega_2)$  is denoted by  $(H, \Omega_1) \cup (G, \Omega_2) = (K, \Omega_3)$  where  $\Omega_3 = \Omega_1 \cup \Omega_2$  and

$$\theta_{K(\alpha)}(u) = \begin{cases} H(\alpha) & \text{if } \alpha \in \Omega_1 - \Omega_2 \\ G(\alpha) & \text{if } \alpha \in \Omega_2 - \Omega_1 \\ \max(H(\alpha), G(\alpha)) & \text{if } \alpha \in \Omega_1 \cap \Omega_2 \end{cases}$$

$$\sigma_{K(\alpha)}(u) = \begin{cases} H(\alpha) & \text{if } \alpha \in \Omega_1 - \Omega_2 \\ G(\alpha) & \text{if } \alpha \in \Omega_2 - \Omega_1 \\ \min(H(\alpha), G(\alpha)) & \text{if } \alpha \in \Omega_1 \cap \Omega_2 \end{cases}$$

**Definition 10** [30] Let  $U$  be an initial universe set,  $\Omega_1, \Omega_2 \subseteq \Delta$  and  $(H, \Omega_1), (G, \Omega_2)$  be two intuitionistic fuzzy hypersoft sets over the universe  $U$ . The intersection of  $(H, \Omega_1)$  and  $(G, \Omega_2)$  is denoted by  $(H, \Omega_1) \cap (G, \Omega_2) = (K, \Omega_3)$  where  $\Omega_3 = \Omega_1 \cap \Omega_2$ ,

$$(K, \Omega_3) = \left\langle \xi, \left( \frac{u}{\left( \begin{matrix} \min\{\theta_{H(\xi)}(u), \theta_{G(\xi)}(u)\} \\ \max\{\theta_{H(\xi)}(u), \theta_{G(\xi)}(u)\} \end{matrix} \right)} \right) \right\rangle : u \in U, \xi \in \Omega \Bigg\}$$

**Definition 11** Let  $U$  be an initial universe set,  $\Omega_1, \Omega_2 \subseteq \Delta$  and  $(H, \Omega_1), (G, \Omega_2)$  be two intuitionistic fuzzy hypersoft sets over the universe  $U$ . The difference of  $(H, \Omega_1)$  and  $(G, \Omega_2)$  is denoted by  $(H, \Omega_1) \setminus (G, \Omega_2) = (K, \Omega_3)$  where  $(H, \Omega_1) \setminus (G, \Omega_2)^c = (K, \Omega_3)$ .

**Definition 12** [30] Let  $U$  be an initial universe set,  $\Omega_1, \Omega_2 \subseteq \Delta$  and  $(H, \Omega_1), (G, \Omega_2)$  be two intuitionistic fuzzy hypersoft sets over the universe  $U$ . The “AND” operation on them is denoted by  $(H, \Omega_1) \wedge (G, \Omega_2) = (K, \Omega_3 \times \Omega_2)$  is given as;

$$(K, \Omega_1 \times \Omega_2) = \left\langle (\alpha_1, \alpha_2), \left( \frac{u}{\theta_{K(\alpha_1, \alpha_2)}(u), \sigma_{K(\alpha_1, \alpha_2)}(u)} \right) \right\rangle : u \in U, (\alpha_1, \alpha_2) \in \Omega_1 \times \Omega_2 \Bigg\}$$

where

$$\theta_{K(\alpha_1, \alpha_2)}(u) = \min\{\theta_{H(\alpha_1)}(u), \theta_{G(\alpha_2)}(u)\}$$

$$\sigma_{K(\alpha_1, \alpha_2)}(u) = \max\{\sigma_{H(\alpha_1)}(u), \sigma_{G(\alpha_2)}(u)\}$$

**Definition 13** [30] Let  $U$  be an initial universe set,  $\Omega_1, \Omega_2 \subseteq \Delta$  and  $(H, \Omega_1), (G, \Omega_2)$  be two intuitionistic fuzzy hypersoft sets over the universe  $U$ . The “OR” operation on them is denoted by  $(H, \Omega_1) \vee (G, \Omega_2) = (K, \Omega_1 \times \Omega_2)$  is given as;

$$(K, \Omega_1 \times \Omega_2) = \left\langle (\alpha_1, \alpha_2), \left( \frac{u}{\theta_{K(\alpha_1, \alpha_2)}(u), \sigma_{K(\alpha_1, \alpha_2)}(u)} \right) \right\rangle : u \in U, (\alpha_1, \alpha_2) \in \Omega_1 \times \Omega_2 \Bigg\}$$

where

$$\theta_{K(\alpha_1, \alpha_2)}(u) = \max\{\theta_{H(\alpha_1)}(u), \theta_{G(\alpha_2)}(u)\}$$

$$\sigma_{K(\alpha_1, \alpha_2)}(u) = \min\{\sigma_{H(\alpha_1)}(u), \sigma_{G(\alpha_2)}(u)\}$$

**Theorem 1** [30] Let  $U$  be an initial universe set,  $\Omega_1, \Omega_2 \subseteq \Delta$  and  $(H, \Omega_1), (G, \Omega_2)$  be two intuitionistic fuzzy hypersoft sets over the universe  $U$ . Then De-Morgan Laws are hold.

- i)  $((H, \Omega_1) \cup (G, \Omega_2))^c = (H, \Omega_1)^c \cap (G, \Omega_2)^c$
- ii)  $((H, \Omega_1) \cap (G, \Omega_2))^c = (H, \Omega_1)^c \cup (G, \Omega_2)^c$

**INTUITIONISTIC FUZZY HYPERSOFT TOPOLOGICAL SPACES**

**Definition 14** Let  $IFHS(U, \Delta)$  be the set of all intuitionistic fuzzy hypersoft subsets over the universe  $U$  and  $\tilde{\tau} \subseteq IFHS(U, \Delta)$ . Then  $\tilde{\tau}$  is called a intuitionistic fuzzy hypersoft topology on  $U$  if the following condition hold.

1.  $0_{(U,IFHS,\Delta)}, 1_{(U,IFHS,\Delta)}$  belong to  $\tilde{\tau}$ ,
2.  $(H_1, \Omega_1) \tilde{\cap} (G_2, \Omega_2) \in \tilde{\tau}$  implies  $(H_1, \Omega_1) \tilde{\cap} (G_2, \Omega_2)$
3.  $\{(H_i, \Omega_i) : i \in I\} \subseteq \tilde{\tau}$  implies  $\tilde{\bigcap}_{i \in I} (H_i, \Omega_i) \in \tilde{\tau}$ .

Then  $(U, \tilde{\tau}, \Delta)$  is called an intuitionistic fuzzy hypersoft topological space over  $U$ . The members of  $\tilde{\tau}$  are said to be intuitionistic fuzzy hypersoft open sets in  $U$ .

An intuitionistic fuzzy hypersoft set  $(H, \Omega)$  over  $U$  is said to be an intuitionistic fuzzy hypersoft closed set if its complement  $(H, \Omega)^c$  belongs to  $\tilde{\tau}$ .

**Definition 15** Let  $IFHS(U, \Delta)$  be the set of all intuitionistic fuzzy hypersoft subsets over the universe  $U$ . Then,

1. If  $\tilde{\tau} = \{0_{(U,IFHS,\Delta)}, 1_{(U,IFHS,\Delta)}\}$ , then  $\tilde{\tau}$  is called to be intuitionistic fuzzy hypersoft indiscrete topology and  $(U, \tilde{\tau}, \Delta)$  is called to be intuitionistic fuzzy hypersoft indiscrete topological space over the universe  $U$ .
2. If  $\tilde{\tau} = IFHS(U, \Delta)$ , then  $\tilde{\tau}$  is called to be intuitionistic fuzzy hypersoft discrete topology and  $(U, \tilde{\tau}, \Delta)$  is called to be intuitionistic fuzzy hypersoft discrete topological space over the universe  $U$ .

**Example 1** Let  $U = \{u_1, u_2, u_3\}$  be the universe set and  $E_1, E_2, E_3$  be sets of attributes.  $E_1, E_2, E_3$  are defined as follows;

$$\begin{aligned} E_1 &= \{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}, \\ E_2 &= \{\beta_1, \beta_2, \beta_3\}, \\ E_3 &= \{\gamma_1, \gamma_2, \gamma_3\}. \end{aligned}$$

Suppose that

$$\begin{aligned} A_1 &= \{\alpha_2, \alpha_3\}, A_2 = \{\beta_1, \beta_3\}, A_3 = \{\gamma_2\}, \\ B_1 &= \{\alpha_2\}, B_2 = \{\beta_2, \beta_3\}, B_3 = \{\gamma_1, \gamma_2\} \end{aligned}$$

are subset of  $E_i$  for each  $i = 1, 2, 3$ . Let

$$\tilde{\tau} = \{0_{(U,IFHS,\Delta)}, 1_{(U,IFHS,\Delta)}, (H_1, \Omega_1), (H_2, \Omega_2), (H_3, \Omega_3), (H_4, \Omega_4)\}$$

be a subfamily of  $IFHS(U, \Delta)$  where  $(H_1, \Omega_1), (H_2, \Omega_2), (H_3, \Omega_3), (H_4, \Omega_4)$ , are intuitionistic fuzzy hypersoft sets and defined as follows;

$$(H_1, \Omega_1) = \left\{ \begin{aligned} &\langle (\alpha_2, \beta_1, \gamma_2), \left\{ \frac{u_1}{(0.2, 0.7)}, \frac{u_2}{(0.5, 0.6)}, \frac{u_3}{(0.4, 0.2)} \right\} \rangle, \\ &\langle (\alpha_2, \beta_3, \gamma_2), \left\{ \frac{u_2}{(0.6, 0.3)}, \frac{u_3}{(0.5, 0.4)} \right\} \rangle, \\ &\langle (\alpha_3, \beta_1, \gamma_2), \left\{ \frac{u_1}{(0.6, 0.6)}, \frac{u_2}{(0.8, 0.5)} \right\} \rangle, \\ &\langle (\alpha_3, \beta_3, \gamma_2), \left\{ \frac{u_1}{(0.5, 0.7)}, \frac{u_2}{(0.6, 0.2)}, \frac{u_3}{(0.3, 0.5)} \right\} \rangle \end{aligned} \right\},$$

$$(H_2, \Omega_2) = \left\{ \begin{aligned} &\langle (\alpha_2, \beta_2, \gamma_1), \left\{ \frac{u_1}{(0.5, 0.2)}, \frac{u_2}{(0.7, 0.3)}, \frac{u_3}{(0.3, 0.8)} \right\} \rangle, \\ &\langle (\alpha_2, \beta_2, \gamma_2), \left\{ \frac{u_2}{(0.1, 0.2)}, \frac{u_3}{(0.3, 0.9)} \right\} \rangle, \\ &\langle (\alpha_2, \beta_3, \gamma_1), \left\{ \frac{u_1}{(0.9, 0.8)}, \frac{u_2}{(0.5, 0.6)}, \frac{u_3}{0.8, 0.2} \right\} \rangle, \\ &\langle (\alpha_2, \beta_3, \gamma_2), \left\{ \frac{u_1}{(0.8, 0.1)}, \frac{u_2}{(0.5, 0.2)} \right\} \rangle \end{aligned} \right\},$$

$$(H_3, \Omega_3) = \left\{ \begin{aligned} &\langle (\alpha_2, \beta_1, \gamma_2), \left\{ \frac{u_1}{(0.2, 0.7)}, \frac{u_2}{(0.5, 0.6)}, \frac{u_3}{(0.4, 0.2)} \right\} \rangle, \\ &\langle (\alpha_2, \beta_3, \gamma_2), \left\{ \frac{u_1}{(0.8, 0.1)}, \frac{u_2}{(0.6, 0.2)}, \frac{u_3}{(0.5, 0.4)} \right\} \rangle, \\ &\langle (\alpha_3, \beta_1, \gamma_2), \left\{ \frac{u_1}{(0.6, 0.6)}, \frac{u_2}{(0.8, 0.5)} \right\} \rangle, \\ &\langle (\alpha_3, \beta_3, \gamma_2), \left\{ \frac{u_1}{(0.5, 0.7)}, \frac{u_2}{(0.6, 0.2)}, \frac{u_3}{(0.3, 0.5)} \right\} \rangle, \\ &\langle (\alpha_2, \beta_2, \gamma_1), \left\{ \frac{u_1}{(0.5, 0.2)}, \frac{u_2}{(0.7, 0.3)}, \frac{u_3}{(0.3, 0.8)} \right\} \rangle, \\ &\langle (\alpha_2, \beta_2, \gamma_2), \left\{ \frac{u_2}{(0.1, 0.2)}, \frac{u_3}{(0.3, 0.9)} \right\} \rangle, \\ &\langle (\alpha_2, \beta_3, \gamma_1), \left\{ \frac{u_1}{(0.9, 0.8)}, \frac{u_2}{(0.5, 0.6)}, \frac{u_3}{0.8, 0.2} \right\} \rangle \end{aligned} \right\},$$

$$(H_4, \Omega_4) = \left\{ \langle (\alpha_2, \beta_3, \gamma_2), \left\{ \frac{u_2}{(0.5, 0.3)} \right\} \rangle \right\},$$

Then  $\tilde{\tau}$  is a intuitionistic fuzzy hypersoft topology and hence  $(U, \tilde{\tau}, \Delta)$  is an intuitionistic fuzzy hypersoft topological space over the universe  $U$ .

**Remark 1** It is clear that each intuitionistic fuzzy hypersoft topology is also intuitionistic fuzzy soft topology. We consider that Example-1. If we select the parameters from a single attribute set such as  $E_2$  while creating fuzzy hypersoft topology, then the resulting topology becomes intuitionistic fuzzy soft topology. So intuitionistic fuzzy hypersoft topology is generalized version of intuitionistic fuzzy soft topology. Therefore intuitionistic fuzzy hypersoft topology is also intuitionistic fuzzy soft topology. But the reverse is not true.

**Proposition 1** Let  $(U, \tilde{\tau}_1, \Delta)$  and  $(U, \tilde{\tau}_2, \Delta)$  be two intuitionistic fuzzy hypersoft topologies over  $U$ .

$$\tilde{\tau}_1 \tilde{\cap} \tilde{\tau}_2 = \{(H, \Omega) : (H, \Omega) \in \tilde{\tau}_1 \text{ and } (H, \Omega) \in \tilde{\tau}_2\}$$

Then  $\tilde{\tau}_1 \tilde{\cap} \tilde{\tau}_2$  an intuitionistic fuzzy hypersoft topology on  $U$ .

**Proof.** Obviously  $0_{(U,IFHS,\Delta)}, 1_{(U,IFHS,\Delta)} \in \tilde{\tau}_1 \tilde{\cap} \tilde{\tau}_2$ . Let  $(H_1, \Omega_1), (H_2, \Omega_2) \in \tilde{\tau}_1$  and  $(H_1, \Omega_1), (H_2, \Omega_2) \in \tilde{\tau}_2$ . Note that  $\tilde{\tau}_1$  and  $\tilde{\tau}_2$

are two intuitionistic fuzzy hypersoft topologies on  $U$ . Then  $(H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2) \in \widetilde{\tau}_1$  and  $(H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2) \in \widetilde{\tau}_2$ .  $(H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2) \in \widetilde{\tau}_1 \widetilde{\cap} \widetilde{\tau}_2$ . Let  $\{(H_p, \Omega_p); i \in I\} \subseteq \widetilde{\tau}_1 \widetilde{\cap} \widetilde{\tau}_2$ . Then  $(H_p, \Omega_p) \in \widetilde{\tau}_1$  and  $(H_p, \Omega_p) \in \widetilde{\tau}_2$  for any  $i \in I$ . Since  $\widetilde{\tau}_1$  and  $\widetilde{\tau}_2$  are two intuitionistic fuzzy hypersoft topologies on  $U$ ,  $\cup_{i \in I} \{(H_p, \Omega_p); i \in I\} \in \widetilde{\tau}_1$  and  $\cup_{i \in I} \{(H_p, \Omega_p); i \in I\} \in \widetilde{\tau}_2$ . Thus  $\cup_{i \in I} \{(H_p, \Omega_p); i \in I\} \in \widetilde{\tau}_1 \widetilde{\cap} \widetilde{\tau}_2$ .

**Remark 2** The union of two intuitionistic fuzzy hypersoft topologies over  $U$  may not be a intuitionistic fuzzy hypersoft topology. This claim is proven by the following example.

**Example 2** We consider that attributes in Example 1. Let

$$\begin{aligned} \widetilde{\tau} &= \{0_{(U_{IFH}, \Delta)}, 1_{(U_{IFH}, \Delta)}, (H_1, \Omega_1)\} \\ \widetilde{\tau} &= \{0_{(U_{IFH}, \Delta)}, 1_{(U_{IFH}, \Delta)}, (G_1, \Omega_1), (G_2, \Omega_2)\} \end{aligned}$$

where

$$(G_1, \Omega_1) = \left\langle \begin{aligned} &<(\alpha_4, \beta_1, \gamma_2), \left\langle \frac{u_1}{(0.3, 0.8)}, \frac{u_2}{(0.4, 0.7)} \right\rangle \rangle, \\ &<(\alpha_4, \beta_3, \gamma_2), \left\langle \frac{u_2}{(0.5, 0.8)}, \frac{u_3}{(0.2, 0.8)} \right\rangle \rangle, \\ &<(\alpha_1, \beta_3, \gamma_2), \left\langle \frac{u_1}{(0.3, 0.9)}, \frac{u_3}{(0.5, 0.2)} \right\rangle \rangle \end{aligned} \right\rangle$$

$$(G_2, \Omega_2) = \left\langle \begin{aligned} &<(\alpha_4, \beta_1, \gamma_2), \left\langle \frac{u_1}{(0.5, 0.6)}, \frac{u_2}{(0.6, 0.3)} \right\rangle \rangle, \\ &<(\alpha_1, \beta_3, \gamma_2), \left\langle \frac{u_1}{(0.7, 0.1)}, \frac{u_2}{(0.5, 0.5)} \right\rangle \rangle, \\ &<(\alpha_4, \beta_1, \gamma_2), \left\langle \frac{u_1}{(0.5, 0.4)}, \frac{u_2}{(0.7, 0.2)} \right\rangle \rangle, \\ &<(\alpha_4, \beta_3, \gamma_2), \left\langle \frac{u_1}{(0.6, 0.2)}, \frac{u_3}{(0.7, 0.3)} \right\rangle \rangle \end{aligned} \right\rangle$$

It is clear that  $\widetilde{\tau}_1 \widetilde{\cap} \widetilde{\tau}_2$  is a intuitionistic fuzzy hypersoft topology. But  $(H_1, \Omega_1) \widetilde{\cap} (G_2, \Omega_2) \notin \widetilde{\tau}_1 \widetilde{\cap} \widetilde{\tau}_2$ , then  $\widetilde{\tau}_1 \widetilde{\cap} \widetilde{\tau}_2$  is not a intuitionistic fuzzy hypersoft topology over the universe  $U$ .

**Proposition 2** Let  $(U, \widetilde{\tau}, \Delta)$  be an intuitionistic fuzzy hypersoft topological spaces over  $U$ . Then, for any  $\alpha \in \Omega$ ,

$$\tau = \{H(\alpha) : (H, \Omega) \in \widetilde{\tau}\}$$

is an intuitionistic fuzzy topology on  $U$ .

**Proof.** (1)  $0_{(U_{IFH}, \Delta)}, 1_{(U_{IFH}, \Delta)} \in \widetilde{\tau}$ . In the intuitionistic fuzzy sets, null set  $0_{(u, (\theta, \sigma))} = (u, (0, 1))$  and absolute set  $1_{(u, (\theta, \sigma))} = (u, (0, 1))$ . It is clear that the values of null set and absolute set in intuitionistic fuzzy sets equal to the values of null set and absolute set in intuitionistic fuzzy hypersoft sets. Therefore  $0_{(u, (\theta, \sigma))}, 1_{(u, (\theta, \sigma))} \in \widetilde{\tau}$ .

(2) Let  $G_1, G_2 \in \widetilde{\tau}$ . Then there exist  $(H_1, \Omega_1), (H_2, \Omega_2) \in \widetilde{\tau}$  such that  $G_1 = H_1(\alpha_1)$  and  $G_2 = H_2(\alpha_2)$ . By  $\widetilde{\tau}$  be an intuitionistic fuzzy hypersoft topologies on  $U$ ,  $(H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2) \in \widetilde{\tau}$ . Put  $(H_3, \Omega_3) = (H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2)$ . Then  $(H_3, \Omega_3) \in \widetilde{\tau}$ . Note that  $G_1 \cap G_2 = H_1(\alpha_1) \widetilde{\cap} H_2(\alpha_2) = (H_3, \Omega_3)$  and  $\tau = \{H(\alpha) : (H, \Omega) \in \widetilde{\tau}\}$ . Then  $G_1 \cap G_2 \in \tau$ .

(3) Let  $\{G_i; i \in I\} \subseteq \widetilde{\tau}$ . Then for every  $i \in I$ , there exist  $(H_i, \Omega_i) \in \widetilde{\tau}$  such that  $G_i = H_i(\alpha_i)$ . By  $\widetilde{\tau}$  be an intuitionistic fuzzy hypersoft topology on  $U$ ,  $\cup_{i \in I} \{(H_i, \Omega_i); i \in I\} \in \widetilde{\tau}$ . Put  $(H, \Omega) = \cup_{i \in I} \{(H_i, \Omega_i); i \in I\}$ . Then  $(H, \Omega) \in \widetilde{\tau}$ . Note that  $\cup_{i \in I} G_i = \cup_{i \in I} \{(H_i, \Omega_i); i \in I\} = (H, \Omega)$  and  $\tau = \{H(\alpha) : (H, \Omega) \in \widetilde{\tau}\}$ . Then  $\cup_{i \in I} G_i \in \tau$ .

Therefore  $\tau = \{H(\alpha) : (H, \Omega) \in \widetilde{\tau}\}$  intuitionistic fuzzy topology on  $U$ .

**Definition 16** Let  $(U, \widetilde{\tau}, \Delta)$  be an intuitionistic fuzzy hypersoft topological spaces over  $U$  and  $(H, \Omega)$  be a intuitionistic fuzzy hypersoft set. The intuitionistic fuzzy hypersoft interior of  $(H, \Omega)$  denoted by  $int_{IFH}(H, \Omega)$ , is defined by the intuitionistic fuzzy hypersoft union of all intuitionistic fuzzy hypersoft open subsets of  $(H, \Omega)$ .

Clearly,  $int_{IFH}(H, \Omega)$  is the largest intuitionistic fuzzy hypersoft open set that is contained in  $(H, \Omega)$ .

**Theorem 2** Let  $(U, \widetilde{\tau}, \Delta)$  be a intuitionistic fuzzy hypersoft topological space over  $U$  and  $(H_1, \Omega_1), (H_2, \Omega_2) \in IFHS(U, \Delta)$  Then,

1.  $int_{IFH}(0_{(U_{IFH}, \Delta)}) = 0_{(U_{IFH}, \Delta)}$  and  $int_{IFH}(1_{(U_{IFH}, \Delta)}) = 1_{(U_{IFH}, \Delta)}$ ,
2.  $int_{IFH}(H_1, \Omega_1) \subseteq (H_1, \Omega_1)$ ,
3.  $(H_1, \Omega_1)$  is an intuitionistic fuzzy hypersoft open set if and only if  $int_{IFH}(H_1, \Omega_1) = (H_1, \Omega_1)$ ,
4.  $int_{IFH}(int_{IFH}(H_1, \Omega_1)) = int_{IFH}(H_1, \Omega_1)$ ,
5. If  $(H_1, \Omega_1) \subseteq (H_2, \Omega_2)$ , then  $int_{IFH}(H_1, \Omega_1) \subseteq int_{IFH}(H_2, \Omega_2)$
6.  $int_{IFH}(H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2)$ , then  $int_{IFH}(H_1, \Omega_1) \widetilde{\cap} int_{IFH}(H_2, \Omega_2)$ .

**Proof.** 1 and 2 are obvious.

3. Let  $(H_1, \Omega_1)$  be a intuitionistic fuzzy hypersoft open set. Since  $int_{IFH}(H_1, \Omega_1)$  is the largest intuitionistic fuzzy hypersoft open set contained in  $(H_1, \Omega_1)$ ,  $int_{IFH}(H_1, \Omega_1) = (H_1, \Omega_1)$ . Conversely, suppose that  $int_{IFH}(H_1, \Omega_1) = (H_1, \Omega_1)$ . Since  $int_{IFH}(H_1, \Omega_1)$  is an intuitionistic fuzzy hypersoft open set,  $(H_1, \Omega_1)$  is also intuitionistic fuzzy hypersoft open set.

4. Let  $cl_{IFH}(H_1, \Omega_1) = (H_1, \Omega_1)$ . Since  $(H_2, \Omega_2)$  is an intuitionistic fuzzy hypersoft open set  $int_{IFH}(H_2, \Omega_2) = (H_2, \Omega_2)$  so  $int_{IFH}(int_{IFH}(H_1, \Omega_1)) = int_{IFH}(H_1, \Omega_1)$  is obtained.

5. Let  $(H_1, \Omega_1) \subseteq (H_2, \Omega_2)$ .  $int_{IFH}(H_1, \Omega_1) \subseteq (H_1, \Omega_1)$  and hence  $int_{IFH}(H_1, \Omega_1) \subseteq (H_2, \Omega_2)$  also  $int_{IFH}(H_2, \Omega_2)$  is the largest intuitionistic fuzzy hypersoft open set contained in  $(H_2, \Omega_2)$  and  $int_{IFH}(H_1, \Omega_1) \subseteq int_{IFH}(H_2, \Omega_2)$ .

6.  $int_{IFH}(H_1, \Omega_1) \subseteq (H_1, \Omega_1)$  and  $int_{IFH}(H_2, \Omega_2) \subseteq (H_2, \Omega_2)$ .

Hence  $int_{IFH}(H_1, \Omega_1) \widetilde{\cap} int_{IFH}(H_2, \Omega_2) \subseteq (H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2)$  Since the largest intuitionistic fuzzy hypersoft open set

contained in  $(H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2)$  is  $int_{IFH}[(H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2)]$ ,  $int_{IFH}(H_1, \Omega_1) \widetilde{\cap} int_{IFH}(H_2, \Omega_2) \subseteq int_{IFH}[(H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2)]$

Conversely  $int_{IFH}(H_1, \Omega_1) \widetilde{\cap} int_{IFH}(H_2, \Omega_2) \subseteq int_{IFH}[(H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2)]$  and  $int_{IFH}(H_1, \Omega_1) \widetilde{\cap} int_{IFH}(H_2, \Omega_2) \subseteq int_{IFH}(H_2, \Omega_2)$

Hence  $int_{IFH}[(H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2)] \subseteq int_{IFH}(H_1, \Omega_1) \widetilde{\cap} cl_{IFH}(H_2, \Omega_2)$ .

**Example 3** We consider the attributes in Example 1. Obviously

$$\tau = \{0_{(U, IFH, \Delta)} = 1_{(U, IFH, \Delta)}, (H_1, \Omega_1), (H_2, \Omega_2), (H_3, \Omega_3), (H_4, \Omega_4)\}$$

is an intuitionistic fuzzy hypersoft topology on  $U$ . Suppose that any  $(H_5, \Omega_5) \in IFHS(U, \Delta)$  be defined as follow;

$$(H_5, \Omega_5) = \left\langle \begin{array}{l} < (\alpha_2, \beta_1, \gamma_2), \left\{ \frac{u_1}{(0.3, 0.5)}, \frac{u_2}{(0.7, 0.2)}, \frac{u_3}{(0.5, 0.1)} \right\} >, \\ < (\alpha_2, \beta_3, \gamma_2), \left\{ \frac{u_1}{(0.8, 0.1)}, \frac{u_2}{(0.8, 0.1)}, \frac{u_3}{(0.6, 0.3)} \right\} >, \\ < (\alpha_3, \beta_1, \gamma_2), \left\{ \frac{u_1}{(0.7, 0.5)}, \frac{u_2}{(0.8, 0.4)} \right\} >, \\ < (\alpha_3, \beta_3, \gamma_2), \left\{ \frac{u_1}{(0.6, 0.3)}, \frac{u_2}{(0.7, 0.1)}, \frac{u_3}{(0.4, 0.1)} \right\} >, \\ < (\alpha_2, \beta_2, \gamma_1), \left\{ \frac{u_1}{(0.7, 0.1)}, \frac{u_2}{(0.9, 0.2)}, \frac{u_3}{(0.4, 0.3)} \right\} >, \\ < (\alpha_2, \beta_2, \gamma_2), \left\{ \frac{u_1}{(0.9, 0.1)}, \frac{u_2}{(0.6, 0.5)}, \frac{u_3}{(0.8, 0.1)} \right\} >, \\ < (\alpha_2, \beta_3, \gamma_1), \left\{ \frac{u_1}{(0.9, 0.1)}, \frac{u_2}{(0.7, 0.1)}, \frac{u_3}{(0.8, 0.1)} \right\} > \end{array} \right\rangle$$

Then  $0_{(U, IFH, \Sigma)}, (H_1, \Omega_1), (H_2, \Omega_2), (H_3, \Omega_3), (H_4, \Omega_4) \subseteq (H_5, \Omega_5)$ . Therefore

$$int_{IFH}(H_5, \Omega_5) = 0_{(U, IFH, \Sigma)} \widetilde{\cap} (H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2) \widetilde{\cap} (H_3, \Omega_3) \widetilde{\cap} (H_4, \Omega_4) = (H_3, \Omega_3)$$

**Definition 17** Let  $(U, \tilde{\tau}, \Delta)$  be an intuitionistic fuzzy hypersoft topological spaces over  $U$  and  $(H, \Omega)$ , be a intuitionistic fuzzy hypersoft set. The intuitionistic fuzzy hypersoft closure of  $(H, \Omega)$ , denoted by  $cl_{IFH}(H, \Omega)$ , is defined by the intuitionistic fuzzy hypersoft intersection of all intuitionistic fuzzy hypersoft closed supersets of  $(H, \Omega)$ .

Clearly,  $cl_{IFH}(H, \Omega)$  is the smallest intuitionistic fuzzy hypersoft closed set which contain  $(H, \Omega)$ .

**Proposition 3** Let  $(U, \tilde{\tau}, \Delta)$  be a intuitionistic fuzzy hypersoft topological space over  $U$ . Then the following properties are provide.

1.  $0_{(U, IFH, \Delta)}, 1_{(U, IFH, \Delta)}$  are intuitionistic fuzzy hypersoft closed sets over  $U$ .
2. The intersection of any number of intuitionistic fuzzy hypersoft closed set is a fuzzy hypersoft set over  $U$ .
3. The union of any two intuitionistic fuzzy hypersoft closed set is a fuzzy hypersoft closed set over  $U$ .

Proof. (1) For  $0_{(U, IFH, \Delta)} = \tilde{\tau}$  is IFH-open set.  $(0_{(U, IFH, \Delta)})^c = 1_{(U, IFH, \Delta)}$  then  $1_{(U, IFH, \Delta)}$  is IFH closed set. Conversely,  $1_{(U, IFH, \Delta)} \in \tilde{\tau}$  is IFH open set. Then  $(1_{(U, IFH, \Delta)})^c = 0_{(U, IFH, \Delta)}$  is IFH closed set.

(2) If  $(H_\alpha, \Omega_\alpha)^c \in \tilde{\tau}$  for  $\alpha \in I$  then  $\bigcup_{\alpha \in I} (H_\alpha, \Omega_\alpha)^c \in \tilde{\tau}$ . So  $\bigcup_{\alpha \in I} (H_\alpha, \Omega_\alpha)^c = \left( \bigcap_{\alpha \in I} (H_\alpha, \Omega_\alpha) \right)^c$ , we have  $\left( \bigcap_{\alpha \in I} (H_\alpha, \Omega_\alpha) \right)^c \in \tilde{\tau}$ .

Hence  $\bigcap_{\alpha \in I} (H_\alpha, \Omega_\alpha)$  is a IFH close set over  $U$ .

(3) Let  $(H_1, \Omega_1), (H_2, \Omega_2) \in \tilde{\tau}^c$ . Then  $(H_1, \Omega_1)^c, (H_2, \Omega_2)^c \in \tilde{\tau}$ . Also we can write  $((H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2))^c = (H_1, \Omega_1)^c \widetilde{\cap} (H_2, \Omega_2)^c$ . According to definition of IFH topology,  $((H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2))^c \in \tilde{\tau}$  and hence  $(H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2) \in \tilde{\tau}^c$ .

**Theorem 3** Let  $(U, \tilde{\tau}, \Delta)$  be a intuitionistic fuzzy hypersoft topological space over  $U$  and  $(H_1, \Omega_1), (H_2, \Omega_2) \in IFHS(U, \Delta)$  Then,

1.  $cl_{IFH}(0_{(U, IFH, \Delta)}) = 0_{(U, IFH, \Delta)}$  and  $cl_{IFH}(1_{(U, IFH, \Delta)}) = 1_{(U, IFH, \Delta)}$ ,
2.  $(H_1, \Omega_1) \subseteq cl_{IFH}(H_1, \Omega_1)$ ,
3.  $(H_1, \Omega_1)$  is a intuitionistic fuzzy hypersoft closed set if and only if  $(H_1, \Omega_1) = cl_{IFH}(H_1, \Omega_1)$ ,
4.  $cl_{IFH}(cl_{IFH}(H_1, \Omega_1)) = cl_{IFH}(H_1, \Omega_1)$ ,
5. If  $(H_1, \Omega_1) \subseteq (H_2, \Omega_2)$ , then  $cl_{IFH}(H_1, \Omega_1) \subseteq cl_{IFH}(H_2, \Omega_2)$ ,
6.  $cl_{IFH}[(H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2)] \subseteq cl_{IFH}(H_1, \Omega_1) \widetilde{\cap} cl_{IFH}(H_2, \Omega_2)$ .

Proof. 1 and 2 are obvious.

3. Let  $(H_1, \Omega_1)$  be a intuitionistic fuzzy hypersoft closed set. By (2), we have  $(H_1, \Omega_1) \subseteq cl_{IFH}(H_1, \Omega_1)$ . Since  $cl_{IFH}(H_1, \Omega_1)$  is the smallest intuitionistic fuzzy hypersoft closed set over  $U$  which contain  $(H_1, \Omega_1)$ , then  $cl_{IFH}(H_1, \Omega_1) \subseteq (H_1, \Omega_1)$ . Hence  $(H_1, \Omega_1) = cl_{IFH}(H_1, \Omega_1)$ . Conversely, suppose that  $(H_1, \Omega_1) = cl_{IFH}(H_1, \Omega_1)$ . Since  $cl_{IFH}(H_1, \Omega_1)$  is a intuitionistic fuzzy hypersoft closed set, then  $(H_1, \Omega_1)$  is closed.

4. Let  $(H_1, \Omega_1) = cl_{IFH}(H_1, \Omega_1)$ . Then,  $(H_1, \Omega_1)$  is a fuzzy hypersoft closed set. So, we have  $cl_{IFH}(cl_{IFH}(H_1, \Omega_1)) = cl_{IFH}(H_1, \Omega_1)$ .

5. If  $(H_1, \Omega_1) \subseteq (H_2, \Omega_2)$ , then  $(H_2, \Omega_2) = (H_1, \Omega_1) \widetilde{\cap} (H_1, \Omega_1) \Rightarrow cl_{IFH}(H_2, \Omega_2) = [(H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2)] = cl_{IFH}(H_1, \Omega_1) \widetilde{\cap} cl_{IFH}(H_2, \Omega_2) \Rightarrow cl_{IFH}(H_1, \Omega_1) \subseteq cl_{IFH}(H_2, \Omega_2)$ .

6. Since  $(H_1, \Omega_1) \subseteq (H_1, \Omega_1) \cup (H_2, \Omega_2)$  and  $(H_2, \Omega_2) \subseteq (H_1, \Omega_1) \cup (H_2, \Omega_2)$ , from the (5),  $(H_2, \Omega_2) \subseteq cl_{IFH}[(H_1, \Omega_1) \cup (H_2, \Omega_2)]$  and  $cl_{IFH}(H_1, \Omega_1) \subseteq cl_{IFH}[(H_1, \Omega_1) \cup (H_2, \Omega_2)]$ . Therefore

$cl_{IFH}(H_1, \Omega_1) \widetilde{\cap} cl_{IFH}(H_2, \Omega_2) \subseteq cl_{IFH}[(H_1, \Omega_1) \cup (H_2, \Omega_2)]$ . Conversely, since  $(H_1, \Omega_1) \subseteq cl_{IFH}(H_1, \Omega_1)$  and  $(H_2, \Omega_2) \subseteq cl_{IFH}(H_2, \Omega_2)$  are intuitionistic fuzzy hypersoft closed sets,  $cl_{IFH}(H_1, \Omega_1) \widetilde{\cap} cl_{IFH}(H_2, \Omega_2)$  is a intuitionistic fuzzy hypersoft closed set over  $U$  being the union of two intuitionistic fuzzy hypersoft fuzzy soft closed sets. Then,

$$cl_{IFH}[(H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2)] \subseteq cl_{IFH}(H_1, \Omega_1) \widetilde{\cap} cl_{IFH}(H_2, \Omega_2).$$

Hence  $cl_{IFH}[(H_1, \Omega_1) \widetilde{\cap} (H_2, \Omega_2)] = cl_{IFH}(H_1, \Omega_1) \widetilde{\cap} cl_{IFH}(H_2, \Omega_2)$  is obtained.

**Example 4** Let us consider the intuitionistic fuzzy hypersoft topology  $\tilde{\tau}$  given in Example-1. Suppose that any  $(H_5, \Omega_5) \in IFHS(U, \Delta)$  be defined as follow;

$$(H_5, \Omega_5) = \left\langle (\alpha_2, \beta_3, \gamma_2), \left\{ \frac{u_1}{(0.1, 0.9)}, \frac{u_2}{(0.2, 0.6)} \right\} \right\rangle$$

Now we find the complement of intuitionistic fuzzy hypersoft open sets in  $\tilde{\tau}$ ,

$$(H_1, \Omega_1)^c = \left\{ \begin{aligned} &\left\langle (\alpha_2, \beta_1, \gamma_2), \left\{ \frac{u_1}{(0.7, 0.2)}, \frac{u_2}{(0.6, 0.5)}, \frac{u_3}{(0.2, 0.4)} \right\} \right\rangle, \\ &\left\langle (\alpha_2, \beta_3, \gamma_2), \left\{ \frac{u_1}{(1, 0)}, \frac{u_2}{(0.3, 0.6)}, \frac{u_3}{(0.4, 0.5)} \right\} \right\rangle, \\ &\left\langle (\alpha_3, \beta_1, \gamma_2), \left\{ \frac{u_1}{(0.6, 0.6)}, \frac{u_2}{(0.5, 0.8)}, \frac{u_3}{(1, 0)} \right\} \right\rangle, \\ &\left\langle (\alpha_3, \beta_3, \gamma_2), \left\{ \frac{u_1}{(0.7, 0.5)}, \frac{u_2}{(0.2, 0.6)}, \frac{u_3}{(0.5, 0.3)} \right\} \right\rangle \end{aligned} \right\},$$

$$(H_2, \Omega_2)^c = \left\{ \begin{aligned} &\left\langle (\alpha_2, \beta_2, \gamma_1), \left\{ \frac{u_1}{(0.2, 0.5)}, \frac{u_2}{(0.3, 0.7)}, \frac{u_3}{(0.8, 0.3)} \right\} \right\rangle, \\ &\left\langle (\alpha_2, \beta_2, \gamma_2), \left\{ \frac{u_1}{(1, 0)}, \frac{u_2}{(0.2, 0.1)}, \frac{u_3}{(0.9, 0.3)} \right\} \right\rangle, \\ &\left\langle (\alpha_2, \beta_3, \gamma_1), \left\{ \frac{u_1}{(0.8, 0.9)}, \frac{u_2}{(0.6, 0.5)}, \frac{u_3}{(0.2, 0.8)} \right\} \right\rangle, \\ &\left\langle (\alpha_2, \beta_3, \gamma_2), \left\{ \frac{u_1}{(0.1, 0.8)}, \frac{u_2}{(0.2, 0.5)}, \frac{u_3}{(1, 0)} \right\} \right\rangle \end{aligned} \right\},$$

$$(H_3, \Omega_3)^c = \left\{ \begin{aligned} &\left\langle (\alpha_2, \beta_1, \gamma_2), \left\{ \frac{u_1}{(0.7, 0.2)}, \frac{u_2}{(0.6, 0.5)}, \frac{u_3}{(0.2, 0.4)} \right\} \right\rangle, \\ &\left\langle (\alpha_2, \beta_3, \gamma_2), \left\{ \frac{u_1}{(0.1, 0.8)}, \frac{u_2}{(0.2, 0.6)}, \frac{u_3}{(0.4, 0.5)} \right\} \right\rangle, \\ &\left\langle (\alpha_3, \beta_1, \gamma_2), \left\{ \frac{u_1}{(0.6, 0.6)}, \frac{u_2}{(0.5, 0.8)}, \frac{u_3}{(1, 0)} \right\} \right\rangle, \\ &\left\langle (\alpha_3, \beta_3, \gamma_2), \left\{ \frac{u_1}{(0.7, 0.5)}, \frac{u_2}{(0.2, 0.6)}, \frac{u_3}{(0.5, 0.3)} \right\} \right\rangle, \\ &\left\langle (\alpha_2, \beta_2, \gamma_1), \left\{ \frac{u_1}{(0.2, 0.5)}, \frac{u_2}{(0.3, 0.7)}, \frac{u_3}{(0.8, 0.3)} \right\} \right\rangle, \\ &\left\langle (\alpha_2, \beta_2, \gamma_2), \left\{ \frac{u_1}{(1, 0)}, \frac{u_2}{(0.2, 0.1)}, \frac{u_3}{(0.9, 0.3)} \right\} \right\rangle, \\ &\left\langle (\alpha_2, \beta_3, \gamma_1), \left\{ \frac{u_1}{(0.8, 0.9)}, \frac{u_2}{(0.6, 0.5)}, \frac{u_3}{(0.2, 0.8)} \right\} \right\rangle \end{aligned} \right\},$$

$$(H_4, \Omega_4)^c = \left\langle (\alpha_2, \beta_3, \gamma_2), \left\{ \frac{u_1}{(1, 0)}, \frac{u_2}{(0.3, 0.5)}, \frac{u_3}{(1, 0)} \right\} \right\rangle,$$

$$(0_{(U_{FH}, \Sigma)})^c = 1_{(U_{FH}, \Sigma)}, (1_{(U_{FH}, \Sigma)})^c = 0_{(U_{FH}, \Sigma)}$$

Obviously,  $(0_{(U_{FH}, \Sigma)})^c, (1_{(U_{FH}, \Sigma)})^c, (\Theta, \Gamma_1)^c, (\Theta, \Gamma_2)^c$  are all fuzzy hypersoft closed sets over  $(U, \tilde{\tau}, \Sigma)$ . Then  $(H_5, \Omega_5) = (0_{(U_{IFH}, \Sigma)})^c, (H_1, \Omega_1)^c, (H_2, \Omega_2)^c, (H_3, \Omega_3)^c, (H_4, \Omega_4)^c$ .

Therefore

$$cl_{IFH}(H_5, \Omega_5) = \left\langle (\alpha_2, \beta_3, \gamma_2), \left\{ \frac{u_1}{(0.1, 0.8)}, \frac{u_2}{(0.2, 0.6)}, \frac{u_3}{(0.4, 0.5)} \right\} \right\rangle$$

**Theorem 4** Let  $(U, \tilde{\tau}, \Delta)$  be an intuitionistic fuzzy hypersoft topological space over  $U$  and  $(H, \Omega) \in IFHS(U, \Delta)$ . Then,

1.  $(cl_{IFH}(H, \Omega))^c = int_{IFH}((H, \Omega)^c)$ ,
2.  $(int_{IFH}(H, \Omega))^c = cl_{IFH}((H, \Omega)^c)$ .

Proof. 1.

$$\begin{aligned} cl_{IFH}(H, \Omega) &= \tilde{\cap} \{ (H, \Omega) \in \tilde{\tau} : (H_2, \Omega_2) \tilde{\subset} (H, \Omega) \} \\ &\Rightarrow (cl_{IFH}(H, \Omega))^c = (\tilde{\cap} \{ (H, \Omega) \in \tilde{\tau} : (H_2, \Omega_2) \tilde{\subset} (H, \Omega) \})^c \\ &= \tilde{\cup} \{ (H, \Omega) \in \tilde{\tau} : (H, \Omega)^c \tilde{\subset} (H_2, \Omega_2)^c \} = int_{IFH}((H, \Omega)^c) \\ int_{IFH}(\Theta, \Gamma) &= \tilde{\cup} \{ (H, \Omega) \in \tilde{\tau} : (H, \Omega) \tilde{\subset} (H_2, \Omega_2) \} \\ &\Rightarrow (int_{IFH}(H, \Omega))^c = (\tilde{\cup} \{ (H, \Omega) \in \tilde{\tau} : (H, \Omega) \tilde{\subset} (H_2, \Omega_2) \})^c \\ &= \tilde{\cap} \{ (H, \Omega) \in \tilde{\tau} : (H_2, \Omega_2)^c \tilde{\subset} (H, \Omega)^c \} = cl_{IFH}((H, \Omega)^c). \end{aligned}$$

**Definition 18** Let  $(U, \tilde{\tau}, \Delta)$  be an intuitionistic fuzzy hypersoft topological space over  $U$  and  $\tilde{B} \subseteq \tilde{\tau}$ .  $\tilde{B}$  is called an intuitionistic fuzzy hypersoft basis for the intuitionistic fuzzy hypersoft topology  $\tilde{\tau}$  if every element of  $\tilde{\tau}$  can be written as the intuitionistic fuzzy hypersoft union of elements of  $\tilde{B}$ .

**Proposition 4** Let  $(U, \tilde{\tau}, \Delta)$  be an intuitionistic fuzzy hypersoft topological space over  $U$  and  $\tilde{B}$  be an intuitionistic fuzzy hypersoft basis for  $\tilde{\tau}$ . Then  $\tilde{\tau}$  equals the collection of intuitionistic fuzzy hypersoft union of elements of  $\tilde{B}$ .

Proof. The proof is clear from the definition of intuitionistic fuzzy hypersoft basis.

**Example 5** We consider that the example 1. Then

$$\tilde{B} = \{ 0_{(U_{IFH}, \Delta)}, 1_{(U_{IFH}, \Delta)}, (H_1, \Omega_1), (H_2, \Omega_2), (H_4, \Omega_4) \}$$

is an intuitionistic fuzzy hypersoft basis for the intuitionistic fuzzy hypersoft topology  $\tilde{\tau}$ .

**Theorem 5** Let  $(U, \tilde{\tau}, \Delta)$  be an intuitionistic fuzzy hypersoft topological space over  $U$  and  $(H, \Omega)$  be an intuitionistic fuzzy hypersoft set over  $U$ . Then the collection  $\tilde{\tau}_{(H, \Omega)} = \{ (H, \Omega) \tilde{\cap} (G_p, \Gamma_p) : (G_p, \Gamma_p) \in \tilde{\tau} \text{ for } p \in I \}$  is an intuitionistic fuzzy hypersoft topology on the intuitionistic fuzzy hypersoft subset  $(\Theta, \Gamma)$  relative parameter set  $\Gamma$ .

Proof.  $0_{(U_{IFH}, \Delta)}, 1_{(U_{IFH}, \Delta)} \in \tilde{\tau}_{(H, \Omega)}$  Besides,

$$\bigcap_{i=1}^n ((H, \Omega) \tilde{\cap} (G_i, \Gamma_i)) = \left( \bigcap_{i=1}^n (G_i, \Gamma_i) \right) \tilde{\cap} (H, \Omega)$$

and

$$\bigcup_{i=1}^n ((H, \Omega) \tilde{\cap} (G_i, \Gamma_i)) = \left( \bigcup_{i=1}^n (G_i, \Gamma_i) \right) \tilde{\cap} (H, \Omega)$$

for  $\tilde{\tau}_{(H, \Omega)} = \{(G_i, \Gamma_i) : i \in I\}$ . Therefore, the intuitionistic fuzzy hypersoft union of any number of intuitionistic fuzzy hypersoft set in  $\tilde{\tau}_{(H, \Omega)}$  belong to  $\tilde{\tau}_{(H, \Omega)}$  and the finite intuitionistic fuzzy hypersoft intersection of intuitionistic fuzzy hypersoft set in  $\tilde{\tau}_{(H, \Omega)}$  belong to  $\tilde{\tau}_{(H, \Omega)}$ . Hence is an intuitionistic fuzzy hypersoft topology on  $(H, \Omega)$ .

**Definition 19** Let  $(U, \tilde{\tau}, \Delta)$  be a intuitionistic fuzzy hypersoft topological space over  $U$  and  $(H, \Omega)$  be a intuitionistic fuzzy hypersoft set over  $U$ . Then the intuitionistic fuzzy hypersoft topology  $\tilde{\tau}_{(H, \Omega)} = \{(H, \Omega) \tilde{\cap} (G_i, \Gamma_i) : (G_i, \Gamma_i) \in \tilde{\tau} \text{ for } i \in I\}$  is called intuitionistic fuzzy hypersoft subspace topology and  $((H, \Omega) \tilde{\tau}_{(H, \Omega)}, \Omega)$  is called a intuitionistic fuzzy hypersoft subspace of  $(U, \tilde{\tau}, \Delta)$ .

**Example 6** Let  $(U, \tilde{\tau}, \Delta)$  be an intuitionistic fuzzy hypersoft topological space over  $U$  and  $(H, \Omega) \in IFHS(U, \Delta)$ . We consider the intuitionistic fuzzy hypersoft topology in Example 1 and  $(H, \Omega)$  be defined as follow;

$$(H, \Omega) = \left\{ \begin{array}{l} \langle (\alpha_2, \beta_1, \gamma_2), \left\{ \frac{u_1}{(0.2, 0.7)}, \frac{u_3}{(0.4, 0.2)} \right\} \rangle, \\ \langle (\alpha_2, \beta_2, \gamma_2), \left\{ \frac{u_1}{(0.2, 0.1)}, \frac{u_3}{(0.2, 0.6)} \right\} \rangle, \\ \langle (\alpha_2, \beta_3, \gamma_2), \left\{ \frac{u_1}{(0.2, 0.6)}, \frac{u_2}{(0.5, 0.2)}, \frac{u_3}{(0.2, 0.6)} \right\} \rangle, \\ \langle (\alpha_3, \beta_3, \gamma_2), \left\{ \frac{u_1}{(0.6, 0.8)}, \frac{u_3}{(0.3, 0.1)} \right\} \rangle \end{array} \right\}$$

Then the collection

$$\tilde{\tau}_{(H, \Omega)} = \left\{ \begin{array}{l} 0_{(U_{IFH}, \Delta)} \tilde{\cap} (H, \Omega), 1_{(U_{IFH}, \Delta)} \tilde{\cap} (H, \Omega), \\ (H_1, \Omega_1) \tilde{\cap} (H, \Omega), \\ (H_2, \Omega_2) \tilde{\cap} (H, \Omega), (H_3, \Omega_3) \tilde{\cap} (H, \Omega), \\ (H_4, \Omega_4) \tilde{\cap} (H, \Omega) \end{array} \right\}$$

is a intuitionistic fuzzy hypersoft subspace topology and  $((H, \Omega) \tilde{\tau}_{(H, \Omega)}, \Omega)$  is a intuitionistic fuzzy hypersoft topological subspace of  $(U, \tilde{\tau}, \Delta)$ .

### MCDM PROBLEM BASED ON IFH-TOPOLOGY

There are various kinds of decision-making strategies for selecting the right option. It is sometimes very difficult to choose an effective decision-making strategy in our real life issues with a similar scenario. However, the

IFH-topology based MCDM approach plays an enthusiastic role in our everyday lives and this is very helpful in selecting the best alternative. In this section, firstly, the problem is solved using IFH set structure. Later, these IFH sets were accepted as sub-base and a topology was created and the problem was solved again by using the open sets of this topology. As a result, the role of topology in MDCM was obtained by comparing the findings obtained in two ways.

### Definition of the problem

Facility location selection is one of the first and most important problems of not only hospitals but also all businesses during the establishment phase. Facility location is a situation that will affect many units, especially suppliers. Therefore, the establishment locations of hospitals are at least as important as having good internal equipment. In this problem, it will be tried to choose the most suitable place for a hospital to be opened.

Let  $U = \{u_1, u_2, u_3, u_4, u_5\}$  be a universe set where  $u_i \cdot i = 1, 5$  represent locations considered for the hospital establishment site and  $E_1, E_2, E_3$  be the set of attributes.  $E_1, E_2, E_3$  are defined as follows;

$$\begin{array}{l} E_1 = \text{Land/building cost} = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}, \\ E_2 = \text{Population density} = \{\beta_1, \beta_2, \beta_3\} \\ E_3 = \text{Distance to suppliers} = \{\gamma_1, \gamma_2, \gamma_3\} \end{array}$$

Suppose that  $A_i$  and  $B_i \cdot i = 1, 2, 3$  sets be the sets of the choices they have formed based on basic criteria by eliminating different decision makers from all attributes.

$$\begin{array}{l} A_1 = \{\alpha_1\}, A_2 = \{\beta_3\}, A_3 = \{\gamma_2, \gamma_3\}, \\ B_1 = \{\alpha_1, \alpha_2\}, B_2 = \{\beta_3\}, B_3 = \{\gamma_2\} \end{array}$$

are subset of  $E_i$  for each  $i = 1, 2, 3$ .

### Solving the problem with IFHSs

#### Algorithm 1

- Step-1 : Input the IFH sets  $(H_1, \Omega_1), (H_2, \Omega_2)$  over  $U$ .
- Step-2 : Find resultant intuitionistic fuzzy hypersoft set  $(H_1, \Omega_1) \vee (H_2, \Omega_2)$
- Step-3 : Construct comparison table of intuitionistic fuzzy hypersoft set and compute row sum ( $r_i$ ) and column sum ( $t_j$ )
- Step-4 : Calculate the resulting score  $R_i$  of  $u_p \forall i$ .
- Step-5 : Optimal choice is  $u_j$  that has  $\max\{R_i\}$ .

Figure-1 shows a brief flow-chart of Algorithm 1.

Suppose that intuitionistic fuzzy hypersoft sets  $(H_1, \Omega_1)$  and  $(H_2, \Omega_2)$  defined as follows:

$$(H_1, \Omega_1) = \left\{ \begin{array}{l} \langle (\alpha_1, \beta_3, \gamma_2), \left\{ \frac{u_1}{(0.3, 0.5)}, \frac{u_2}{(0.2, 0.8)} \right\} \rangle, \\ \langle (\alpha_1, \beta_3, \gamma_3), \left\{ \frac{u_2}{(0.6, 0.1)}, \frac{u_4}{(0.7, 0.5)}, \frac{u_5}{(0.3, 0.1)} \right\} \rangle \end{array} \right\}$$






Figure 1. Graphical representation of Algorithm-1.

Table 1: Tabular form of  $(H_1, \Omega_1)$

$(H_1, \Omega_1)$	$u_1$	$u_2$	$u_3$	$u_4$	$u_5$
$(\alpha_1, \beta_3, \gamma_2) = a_1$	(0.3,0.5)	(0.2,0.8)	(0,1)	(0,1)	(0,1)
$(\alpha_1, \beta_3, \gamma_3) = a_2$	(0,1)	(0.6,0.1)	(0,1)	(0.7,0.5)	(0.3,0.1)

Table 2: Tabular form of  $(H_2, \Omega_2)$

$(H_2, \Omega_2)$	$u_1$	$u_2$	$u_3$	$u_4$	$u_5$
$(\alpha_1, \beta_3, \gamma_2) = b_1$	(0.5,0.7)	(0.4,0.1)	(0,1)	(0,1,0.7)	(0,1)
$(\alpha_2, \beta_2, \gamma_2) = b_2$	(0,1,0.7)	(0,1,0.7)	(0,1)	(0.7,0.3)	(0.4,0.1)

Table 3: Tabular form of  $(H_1, \Omega_1) \vee (H_2, \Omega_2)$

$(H_1, \Omega_1) \vee (H_2, \Omega_2)$	$u_1$	$u_2$	$u_3$	$u_4$	$u_5$
$(a_1 \times b_1) = x_1$	(0.5,0.5)	(0.4,0.1)	(0,1)	(0,1,0.7)	(0,1)
$(a_1 \times b_2) = x_2$	(0,3,0.5)	(0.2,0.8)	(0,1)	(0.7,0.3)	(0.4,0.1)
$(a_2 \times b_1) = x_3$	(0.5,0.7)	(0.6,0.1)	(0,1)	(0.7,0.5)	(0.3,0.1)
$(a_2 \times b_2) = x_4$	(0,1,0.7)	(0.5,0.5)	(0,1)	(0.7,0.3)	(0.4,0.1)

$$(H_2, \Omega_2) = \left\langle (\alpha_1, \beta_3, \gamma_2), \left\{ \frac{u_1}{(0.5,0.7)}, \frac{u_2}{(0.4,0.1)}, \frac{u_4}{(0,1,0.7)} \right\} \right\rangle, \left\langle (\alpha_2, \beta_2, \gamma_2), \left\{ \frac{u_1}{(0,1,0.7)}, \frac{u_4}{(0.7,0.3)}, \frac{u_5}{(0.4,0.1)} \right\} \right\rangle$$

The tabular representations of  $(H_1, \Omega_1)$  and  $(H_2, \Omega_2)$  are shown in below.

Now, we find resultant intuitionistic fuzzy hypersoft set  $(H_1, \Omega_1) \vee (H_2, \Omega_2)$ .

Table 4: Comparison table of intuitionistic fuzzy hypersoft set  $(H_1, \Omega_1) \vee (H_2, \Omega_2)$

$(H_1, \Omega_1) \vee (H_2, \Omega_2)$	$u_1$	$u_2$	$u_3$	$u_4$	$u_5$
$u_1$	4	1	4	1	1
$u_2$	2	4	4	0	1
$u_3$	0	0	4	0	1
$u_4$	3	1	4	4	1
$u_5$	2	1	4	0	4

Now we find the comparison table of intuitionistic fuzzy hypersoft set  $(H_1, \Omega_1) \vee (H_2, \Omega_2)$  by using the algorithm which is given by Roy and Maji [24]. Comparison table is a square table in which the number of rows and number of columns are equal, rows and columns both are labelled by the object names  $u_1, u_2, \dots, u_n$  on of the universe, and the entries are  $x_i, i = 1, 2, \dots, n$  given by  $x_i =$  the number of parameters for which the membership value of  $o_i$  exceeds or equal to the membership value of  $u_i$ . The comparison table is given below.

Here we calculate the column sum ( $t_i$ ) and row sum ( $r_i$ ) after that we calculate the score  $R_i$  for each  $u_i, i = 1, 2, 3, 4, 5$ .

According to Table-5, it is clear that the most suitable location for the hospital is  $u_4$ . In the next section, the same problem will be solved by constructing the IFH topology and the results will be discussed.

### Solving the problem with IFHS-topology

#### Algorithm 2

Step-1 : Consider a universe of  $U$ .

Step-2 : A set E of attributes.

- Step-3 : Construct the IFH sets  $(H_1, \Omega_1), (H_2, \Omega_2)$  over  $U$ .
- Step-4 : Write IFH-topology  $\tilde{\tau}$  in which  $(H_1, \Omega_1)$  and  $(H_2, \Omega_2)$  are open IFHs in  $\tilde{\tau}$ .
- Step-5 : Find resultant intuitionistic fuzzy hypersoft set  $(H_1, \Omega_1) \vee (H_2, \Omega_2)$  and other open IFHs in  $\tilde{\tau}$  with “OR” operation.
- Step-6 : Construct comparison table of intuitionistic fuzzy hypersoft set and compute row sum ( $r_i$ ) and column sum ( $t_i$ )
- Step-7 : Calculate the resulting score  $R_i$  of  $u_i, \forall i$ .
- Step-8 : Optimal choice is  $u_j$  that has  $\max\{R_i\}$ .

Figure-2 shows a brief flow-chart of Algorithm 2. Let's build the IFH topology now. We have  $(H_1, \Omega_1), (H_2, \Omega_2)$ . Let's create a topology so that these sets are open sets.


$$\tilde{\tau} = \{0_{(U_{IFH}, \Delta)}, 1_{(U_{IFH}, \Delta)}, (H_1, \Omega_1), (H_2, \Omega_2), (H_3, \Omega_3), (H_4, \Omega_4)\}$$

where  $(H_3, \Omega_3), (H_4, \Omega_4)$  defined as follows.

$$(H_3, \Omega_3) = (H_1, \Omega_1) \tilde{\cup} (H_2, \Omega_2) = \left\langle (\alpha_1, \beta_3, \gamma_2), \left\{ \frac{u_1}{(0.5, 0.5)}, \frac{u_2}{(0.4, 0.1)}, \frac{u_4}{(0.1, 0.7)} \right\} \right\rangle, \left\langle (\alpha_1, \beta_3, \gamma_3), \left\{ \frac{u_2}{(0.6, 0.1)}, \frac{u_4}{(0.7, 0.5)}, \frac{u_5}{(0.3, 0.1)} \right\} \right\rangle, \left\langle (\alpha_1, \beta_2, \gamma_2), \left\{ \frac{u_1}{(0.1, 0.7)}, \frac{u_4}{(0.7, 0.3)}, \frac{u_5}{(0.4, 0.1)} \right\} \right\rangle$$

**Table 5:** Tabular form of score value

	Rowsum( $r_i$ )	Columnsum( $t_i$ )	Score( $R_i = r_i - t_i$ )
$u_1$	11	11	0
$u_2$	11	7	4
$u_3$	5	20	-15
$u_4$	12	5	8
$u_5$	11	8	3



**Figure 2.** Graphical representation of Algorithm-2.

**Table 6:** Tabular form of  $(H_3, \Omega_3)$

$(H_3, \Omega_3)$	$u_1$	$u_2$	$u_3$	$u_4$	$u_5$
$(\alpha_1, \beta_3, \gamma_2) = a_1$	(0.5,0.5)	(0.4,0.1)	(0,1)	(0,1,0.7)	(0,1)
$(\alpha_1, \beta_3, \gamma_3) = a_2$	(0,1)	(0.6,0.1)	(0,1)	(0.7,0.5)	(0.3,0.1)
$(\alpha_1, \beta_2, \gamma_2) = a_3$	(0,1,0.7)	(0,1)	(0,1)	(0.7,0.3)	(0.4,0.1)

**Table 7:** Tabular form of  $(H_4, \Omega_4)$

$(H_4, \Omega_4)$	$u_1$	$u_2$	$u_3$	$u_4$	$u_5$
$(\alpha_1, \beta_3, \gamma_2) = b_1$	(0.3,0.7)	(0.2,0.8)	(0,1)	(0,1)	(0,1)

**Table 8:** Tabular form of  $(H_3, \Omega_3)$

$(H_3, \Omega_3) \vee (H_4, \Omega_4)$	$u_1$	$u_2$	$u_3$	$u_4$	$u_5$
$(a_1 \times b_1) = y_1$	(0.5,0.5)	(0.4,0.1)	(0,1)	(0,1,0.7)	(0,1)
$(a_2 \times b_1) = y_2$	(0,3,0.7)	(0.6,0.1)	(0,1)	(0.7,0.5)	(0.3,0.1)
$(a_3 \times b_1) = y_3$	(0.3,0.7)	(0.2,0.8)	(0,1)	(0.7,0.3)	(0.4,0.1)

**Table 9:** Tabular form of  $(H_1, \Omega_1) \vee (H_2, \Omega_2) \vee (H_3, \Omega_3) \vee (H_4, \Omega_4)$

$(H_1, \Omega_1) \vee (H_2, \Omega_2) \vee (H_3, \Omega_3) \vee (H_4, \Omega_4)$	$u_1$	$u_2$	$u_3$	$u_4$	$u_5$
$(x_1 \times y_1)$	(0.5,0.5)	(0.4,0.1)	(0,1)	(0,1,0.7)	(0,1)
$(x_1 \times y_2)$	(0.5,0.5)	(0.6,0.1)	(0,1)	(0.7,0.5)	(0.3,0.1)
$(x_1 \times y_3)$	(0.5,0.5)	(0.4,0.1)	(0,1)	(0.7,0.3)	(0.4,0.1)
$(x_2 \times y_1)$	(0.5,0.5)	(0.4,0.1)	(0,1)	(0.7,0.3)	(0.4,0.1)
$(x_2 \times y_2)$	(0,3,0.5)	(0.6,0.1)	(0,1)	(0.7,0.3)	(0.4,0.1)
$(x_2 \times y_3)$	(0,3,0.5)	(0.2,0.8)	(0,1)	(0.7,0.3)	(0.4,0.1)
$(x_3 \times y_1)$	(0.5,0.7)	(0.6,0.1)	(0,1)	(0.7,0.5)	(0.3,0.1)
$(x_3 \times y_2)$	(0.5,0.7)	(0.6,0.1)	(0,1)	(0.7,0.5)	(0.3,0.1)
$(x_3 \times y_3)$	(0.5,0.5)	(0.6,0.1)	(0,1)	(0.7,0.3)	(0.4,0.1)
$(x_4 \times y_1)$	(0,3,0.7)	(0.6,0.1)	(0,1)	(0.7,0.3)	(0.4,0.1)
$(x_4 \times y_2)$	(0,3,0.7)	(0.6,0.1)	(0,1)	(0.7,0.3)	(0.4,0.1)
$(x_4 \times y_3)$	(0,3,0.7)	(0.6,0.1)	(0,1)	(0.7,0.3)	(0.4,0.1)

$$(H_4, \Omega_4) = (H_1, \Omega_1) \tilde{\cap} (H_2, \Omega_2) = \left\langle (\alpha_1, \beta_3, \gamma_2), \left\{ \frac{u_1}{(0.3,0.7)}, \frac{u_2}{(0.2,0.8)} \right\} \right\rangle$$

The tabular representations of  $(H_3, \Omega_3)$  and  $(H_4, \Omega_4)$  are shown in below.

**Table 10:** Comparison table of intuitionistic fuzzy hypersoft set  $(H_1, \Omega_1) \vee (H_2, \Omega_2) \vee (H_3, \Omega_3) \vee (H_4, \Omega_4)$

$(H_1, \Omega_1) \vee (H_2, \Omega_2)$	$u_1$	$u_2$	$u_3$	$u_4$	$u_5$
$u_1$	12	1	12	1	1
$u_2$	8	12	12	1	11
$u_3$	0	0	12	0	1
$u_4$	11	1	12	12	1
$u_5$	4	3	12	0	12

**Table 11:** Tabular form of score value

	Rowsum( $r_i$ )	Columnsum( $t_i$ )	Score( $R_i = r_i - t_i$ )
$u_1$	27	35	-8
$u_2$	44	17	27
$u_3$	13	60	-4
$u_4$	37	14	23
$u_5$	31	25	6

Now we calculate  $(H_1, \Omega_1) \vee (H_2, \Omega_2) \vee (H_3, \Omega_3) \vee (H_4, \Omega_4)$ . We have  $(H_3, \Omega_3) \vee (H_4, \Omega_4)$ . Then we find  $(H_1, \Omega_1) \vee (H_2, \Omega_2)$  now.

Now, we find  $(H_1, \Omega_1) \vee (H_2, \Omega_2) \vee (H_3, \Omega_3) \vee (H_4, \Omega_4)$ .

Now we find the comparison table of intuitionistic fuzzy hypersoft set  $(H_1, \Omega_1) \vee (H_2, \Omega_2) \vee (H_3, \Omega_3) \vee (H_4, \Omega_4)$ . The comparison table is given below.

Here we calculate the column sum ( $t_i$ ) and row sum ( $r_i$ ) after that we calculate the score  $R_i$  for each  $u_i, i = 1, 2, 3, 4, 5$ .

According to Table 11, it is clear that the most suitable location for the hospital is  $u_2$ .

## RESULT

In this problem, the most suitable place is sought for the selection of the hospital location. When the problem is solved according to Algorithm 1, it is seen that  $u_4$  location is the most suitable place. Then the topology was built so that the sets used in Algorithm 1 are open sets in Algorithm 2. Here, by creating a topology, the finite combination of sets and arbitrary intersection was used first. Therefore, the decision-making problem has been enriched in terms of both parameters and other examined items. While the location is selected over 4 different parameters according to Algorithm 1, the number of parameters handled with the help of the topology in Algorithm 2 has increased to 12. Thus, the problem has been studied in more depth. The findings obtained in Algorithm 1 and Algorithm 2 are compared in the graph below.




Figure 3. Comparison of Algorithm-1 and Algorithm-2.

When all these findings are examined, it may be more appropriate to choose  $u_2$  as the most suitable location for the hospital.

## CONCLUSION

The analysis of hypersoft topological spaces is of considerable significance because it offers a general structure composed of parametrized classical topological spaces. The aim of present paper is to study the concept of intuitionistic fuzzy hypersoft topological spaces. We investigated some properties of intuitionistic fuzzy hypersoft topological spaces and we introduced some notions like that interior, closure, basis, subspace topology on intuitionistic fuzzy hypersoft topological spaces. These are illustrated with appropriate examples. Additionally, the concept of intuitionistic fuzzy hypersoft topology is extended to develop multi criteria decision making problems. To better understand the importance of the study, firstly, Algorithm 1 solved the problem using the IFH set structure. Then, the findings obtained by solving the problem with IFH topology (Algorithm 2) were compared. Such a final result was achieved. Continuity, connectedness, compactness and many other topological concepts can be studied on IFH topological spaces as a continuation of this work in the future.

## DATA AVAILABILITY STATEMENT

The authors confirm that the data that supports the findings of this study are available within the article. Raw data that support the finding of this study are available from the corresponding author, upon reasonable request.

## CONFLICT OF INTEREST

The author declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## ETHICS

There are no ethical issues with the publication of this manuscript.

## REFERENCES

- [1] Abbas M, Murtaza G, Smarandache F. Basic operations on hypersoft sets and hypersoft point. *Neutrosophic Sets Sys* 2020;35:407–421.
- [2] Alcantud JCR, Khameneh AZ, Kilicman A. Aggregation of finite chains of intuitionistic fuzzy sets and their application to choices with

- temporal intuitionistic fuzzy information. *Inf Sci* 2020;514:106–117. [\[CrossRef\]](#)
- [3] Alcantud JCR. Soft open bases and a novel construction of soft topologies from bases for topologies. *Mathematics* 2020;8:672. [\[CrossRef\]](#)
- [4] Atanassov KT, Intuitionistic fuzzy sets. *Fuzzy Sets Syst* 1986;20:87–96. [\[CrossRef\]](#)
- [5] Atanassov K, Gargov G. Interval valued intuitionistic fuzzy sets. *Fuzzy Sets Syst* 1989;31:343–349. [\[CrossRef\]](#)
- [6] Aygunoglu A, Aygun H. Some notes on soft topological spaces. *Neural Comput Appl* 2012;21:113–119. [\[CrossRef\]](#)
- [7] Cagman N, Karatas S, Enginoglu S. Soft topology. *Comput Math Appl* 2011;62:351–358. [\[CrossRef\]](#)
- [8] Coker D. An introduction to intuitionistic fuzzy topological spaces. *Fuzzy Sets Syst* 1997;88:81–89. [\[CrossRef\]](#)
- [9] Deli I. Hybrid Set Structures Under Uncertainly Parameterized Hypersoft Sets: Theory and Applications In: Smarandache F, Abdel-Baset M, Saeed M, Saqlain M, editors. *Theory and application of hypersoft set*. 1st ed. Brussels: Pons Publishing House; 2021. pp. 24–49.
- [10] Enginoglu S, Arslan B. Intuitionistic fuzzy parameterized intuitionistic fuzzy soft matrices and their application in decision-making. *Comput Appl Math* 2020;39:325. [\[CrossRef\]](#)
- [11] Faizi S, Salabun W, Rashid T, Zafar S, Watrobski J. Intuitionistic fuzzy sets in multicriteria group decision making problems using the characteristic objects method. *Symmetry* 2020;12:1382. [\[CrossRef\]](#)
- [12] Feng F, Zheng Y, Alcantud JCR, Wang Q. Minkowski weighted score functions of intuitionistic fuzzy values. *Mathematics* 2020;8:1143. [\[CrossRef\]](#)
- [13] Garg H, Kaur G. Cubic intuitionistic fuzzy sets and its fundamental properties. *J. Multivalued Log S* 2019;33:507–537.
- [14] Garg H, Kumar K. Linguistic interval-valued Atanassov intuitionistic fuzzy sets and their applications to group decision-making problems. *IEEE Trans Fuzzy Syst* 2019;27:2302–2311. [\[CrossRef\]](#)
- [15] Gayen S, Smarandache F, Jha S, Singh MK, Broumi S, Kumar R. Introduction to plithogenic hypersoft subgroup. *Neutrosophic Sets Sys* 2020;33:14. [\[CrossRef\]](#)
- [16] Hussain S. On some properties of Intuitionistic fuzzy soft boundary. *Commun Fac Sci Univ Ank Ser A1 Math Stat* 2020;69:39–50. [\[CrossRef\]](#)
- [17] Li Z, Cui R. On the topological structures of intuitionistic fuzzy soft sets. *Ann Fuzzy Math Inform* 2013;5:229–239.
- [18] Maji PK, Biswas R, Roy AR. Fuzzy soft sets. *J Fuzzy Math* 2001;9:589–602.
- [19] Molodtsov DA. Soft set theory first results. *Comput Math Appl* 1999;37:19–31. [\[CrossRef\]](#)
- [20] Ozturk TY, Yolcu A. On neutrosophic hypersoft topological spaces. In: Smarandache F, Abdel-Baset M, Saeed M, Saqlain M, editors. *Theory and application of hypersoft set*. 1st ed. Brussels: Pons Publishing House; 2021. pp. 215–234.
- [21] Ozturk TY, Gunduz Aras C, Bayramov S. A new approach to operations on neutrosophic soft sets and to neutrosophic soft topological spaces. *Commun Math Appl* 2019;10:481–493. [\[CrossRef\]](#)
- [22] Ozturk TY. On bipolar soft topological spaces. *J New Theory* 2018;20:64–75.
- [23] Ozturk TY, Yolcu A. Some structures on pythagorean fuzzy topological spaces. *J New Theory* 2020;33:15–25.
- [24] Roy AR, Maji PK. A fuzzy soft set theoretic approach to decision making problems. *J Comput Appl Math* 2007;203:412–418. [\[CrossRef\]](#)
- [25] Saeed M, Ahsan M, Siddique MK, Ahmad MR. A study of the fundamentals of hypersoft set theory. *Int J Sci Eng Res* 2020;11:230.
- [26] Smarandache F. Extension of soft set to hypersoft set, and then to plithogenic hypersoft set. *Neutrosophic Sets Syst* 2018;22:168–170.
- [27] Simsekler Dizman T, Ozturk TY. Fuzzy bipolar soft topological spaces. *TWMS J App Eng Math* 2021;11:151–159.
- [28] Terepeta M. On separating axioms and similarity of soft topological spaces. *Soft Comput* 2019;23:1049–1057. [\[CrossRef\]](#)
- [29] Xu ZS, Zhao N. Information fusion for intuitionistic fuzzy decision making: an overview. *Inf Fusion* 2016;28:10–23. [\[CrossRef\]](#)
- [30] Yolcu A, Smarandache F, Ozturk TY. Intuitionistic fuzzy hypersoft set. *Commun Fac Sci Univ Ank Ser A1 Math Stat* 2021;70:443–455. [\[CrossRef\]](#)
- [31] Yolcu A, Ozturk TY, Fuzzy Hypersoft Sets and Its Application to Decision-Making. In: Smarandache F, Abdel-Baset M, Saeed M, Saqlain M, editors. *Theory and application of hypersoft set*. 1st ed. Brussels: Pons Publishing House; 2021. pp. 50–64.
- [32] Zadeh LA. Fuzzy sets. *Inf Control* 1965;8:338–353. [\[CrossRef\]](#)
- [33] Zulqarnain RM, Xin XL, Saqlain M, Smarandache F. Generalized aggregate operators on neutrosophic hypersoft set. *Neutrosophic Sets Sys* 2020;36:271–281. [\[CrossRef\]](#)
- [34] Zulqarnain RM, Xin XL, Saeed M. Extension of TOPSIS method under intuitionistic fuzzy hypersoft environment based on correlation coefficient and aggregation operators to solve decision making problem. *AIMS Mathematics* 2021;6:2732–2755. [\[CrossRef\]](#)