

STRONGLY GRADED MODULES AND POSITIVELY GRADED MODULES WHICH ARE UNIQUE FACTORIZATION MODULES

I. Ernanto, A. Ueda and I. E. Wijayanti

Received: 15 November 2022; Revised: 10 August 2023; Accepted: 18 October 2023

Communicated by Meltem Altun Özarslan

ABSTRACT. Let $M = \bigoplus_{n \in \mathbb{Z}} M_n$ be a strongly graded module over strongly graded ring $D = \bigoplus_{n \in \mathbb{Z}} D_n$. In this paper, we prove that if M_0 is a unique factorization module (UFM for short) over D_0 and D is a unique factorization domain (UFD for short), then M is a UFM over D . Furthermore, if D_0 is a Noetherian domain, we give a necessary and sufficient condition for a positively graded module $L = \bigoplus_{n \in \mathbb{Z}_0} M_n$ to be a UFM over positively graded domain $R = \bigoplus_{n \in \mathbb{Z}_0} D_n$.

Mathematics Subject Classification (2020): 13A05, 13C05, 13C70, 13E15, 13F15

Keywords: Graded ring, graded module, unique factorization module

1. Introduction

Throughout this paper $R = \bigoplus_{n \in \mathbb{Z}_0} D_n$ is a positively graded domain which is a sub-domain of the strongly graded domain $D = \bigoplus_{n \in \mathbb{Z}} D_n$ and $L = \bigoplus_{n \in \mathbb{Z}_0} M_n$ is a positively graded module over R which is a subset of the strongly graded module $M = \bigoplus_{n \in \mathbb{Z}} M_n$.

In [12], the authors defined a concept of a Unique Factorization Module (UFM for short) by submodule approach. They also proved that the concept of UFM by [12] is equivalent to Nicolas's UFM (see [8] for the definition of a UFM by Nicolas), which is defined in terms of irreducible elements of D and M . In [12], the authors proved that if M is a UFM, then the polynomial module $M[x]$ is also a UFM. There are several papers on UFM, see for example [1], [5], [8] and [9].

This work was done with the support from Universitas Gadjah Mada under Research Grant Year 2022 (Hibah Penelitian Dosen Dana Masyarakat Alokasi Fakultas MIPA Nomor 155/J01.1.28/PL.06.02/2022) as the implementation of JSPS RONPAKU (Dissertation Ph.D.) Program and the continuation of the research cooperation between the second author and the members of the Algebra Research Group at the Department of Mathematics UGM.

On the other hand, in [11], it is shown that $D = \bigoplus_{n \in \mathbb{Z}} D_n$, a strongly graded domain, is a G-Dedekind domain if and only if D_0 is a G-Dedekind domain. Moreover, in [2], it is shown that $R = \bigoplus_{n \in \mathbb{Z}_0} D_n$, a positively graded domain, is a Unique Factorization Domain (UFD for short) if and only if D_0 is a UFD and D_1 is a principal D_0 -module.

The aim of this paper is to extend the main results of [11] and [12] to a strongly graded module $M = \bigoplus_{n \in \mathbb{Z}} M_n$. The paper is organized as follows. In Section 2, we give some characterizations of a unique factorization module that will be very useful in Section 3 and Section 4. We refer the reader to [2], [6], [7], [11] and [13], and for details regarding graded rings, and to [3] regarding graded modules, that are not mentioned in this paper.

In Section 3, we prove that if M_0 is a UFM over D_0 , then $M = \bigoplus_{n \in \mathbb{Z}} M_n$ is a UFM over D as a generalization of the result in [12]. We give some examples of strongly graded modules which are UFM.

In Section 4, we give the necessary and sufficient condition for a positively graded module $L = \bigoplus_{n \in \mathbb{Z}_0} M_n$ to be a UFM. We end Section 4 with some examples of positively graded modules which are UFM as the application of the main result of this paper, that is, Theorem 4.8.

We refer the reader to [10], [14], [15] for the concept and properties of v -submodules and [4] to some properties of multiplicative ideal theory which are not mentioned in this paper.

2. Unique factorization modules

Let M be a torsion-free module over an integral domain D with the quotient field K . In [15], the authors defined the following concepts.

- Definition 2.1.** (1) A non-zero D -submodule N of KM is called a fractional D -submodule in KM if there is a non-zero $r \in D$ such that $rN \subseteq M$ and $KN = KM$.
- (2) A non-zero D -submodule \mathfrak{a} of K is called a fractional M -ideal in K if there is a non-zero $m \in M$ such that $\mathfrak{a}m \subseteq M$.

We represent $F(M)$ as the collection of all fractional D -submodules in KM , while $F_M(D)$ refers to the set containing all fractional M -ideals in K . Suppose $N \in F(M)$ and $\mathfrak{a} \in F_M(D)$. We define $N^- = \{k \in K \mid kN \subseteq M\}$ and $\mathfrak{a}^+ = \{m \in KM \mid \mathfrak{a}m \subseteq M\}$. Then, it easily follows that $N^- \in F_M(D)$ and $\mathfrak{a}^+ \in F(M)$.

For $N \in F(M)$ and $\mathfrak{a} \in F_M(D)$, we define $N_v = (N^-)^+$ and $\mathfrak{a}_{v1} = (\mathfrak{a}^+)^-$. Consequently, $N_v \in F(M)$, and it satisfies $N_v \supseteq N$. Similarly, $\mathfrak{a}_{v1} \in F_M(D)$, and

it satisfies $\mathfrak{a}_{v_1} \supseteq \mathfrak{a}$. When $N = N_v$, we classify N as a fractional v -submodule in KM. Moreover, if $\mathfrak{a} = \mathfrak{a}_{v_1}$, we refer to \mathfrak{a} as a v_1 -ideal (with respect to M). We refer the reader to [10], [14], [15] for details regarding v -submodules and v_1 -operation.

In [12], the authors defined a concept of a unique factorization module by a submodule approach. The authors gave the definition and characterization of unique factorization modules as follows.

Definition 2.2. [[12], Definition 2] A torsion-free module M over an integral domain D is called a unique factorization module (UFM for short) if

- (1) M is completely integrally closed (CIC for short), that is, $O_K(N) = \{k \in K \mid kN \subseteq N\} = D$ for every non-zero submodule N of M , where K is the quotient field of D ,
- (2) every v -submodule of M is principal,
- (3) M satisfies the ascending chain condition on v -submodules of M .

Theorem 2.3. [[12], Theorem 1] *Suppose $O_K(M) = D$. The following conditions are equivalent:*

- (1) M is a UFM.
- (2) M is a v -multiplication module and D is a UFD.
- (3) (a) D is a UFD.
 (b) For every prime element p of D , pM is a maximal v -submodule.
 (c) For every v -submodule N of M , $\mathfrak{n} = (N : M) \neq \{0\}$, where $(N : M) = \{r \in D \mid rM \subseteq N\}$.
- (4) Every v -submodule of M is principal and D is a UFD.

Note that if M is a finitely generated torsion-free D -module, then $O_K(M) = D$ by Lemma 2.1 of [3]. Throughout this paper, M is a finitely generated torsion-free D -module and satisfies the ascending chain condition on v -submodules of M .

Lemma 2.4. *Let P be a maximal v -submodule of M . Then P is a prime submodule.*

Proof. Let $rm \in P$ where $r \in D$ and $m \in M$. If $m \notin P$, then $P \subset Dm + P \subseteq (Dm + P)_v \subseteq M$ and so $(Dm + P)_v = M$. Then $P \supseteq (Drm + rP)_v = (r(Dm + P))_v = r(Dm + P)_v = rM$. Hence P is prime. \square

The following theorem will be very useful for Section 3 and Section 4.

Theorem 2.5. *Suppose that D is a UFD and M is a CIC module that satisfies the ascending chain condition on v -submodules of M . Then M is a UFM if and only if every prime v -submodule of M is principal.*

Proof. If M is a UFM, then it is clear that every prime v -submodule of M is principal, by Theorem 2.3. Conversely, we assume on the contrary that M is not a UFM. Let N be a v -submodule of M which is not principal and we may assume that N is maximal with this property because M satisfies the ascending chain condition on v -submodules of M . Let P be a maximal v -submodule of M containing N . Then $P = pM$ for some non-zero $p \in D$ by Lemma 2.4. Since $N \subset P \subset M$, we have $N \subseteq p^{-1}N \subset M$ and so $(p^{-1}N)_v = p^{-1}N_v = p^{-1}N$. Then $N = p^{-1}N$ or $p^{-1}N$ is principal by the maximality of N . If $p^{-1}N$ is principal, then $p^{-1}N = tM$ for some $t \in D$ and so $N = ptM$, a contradiction. Hence $N = p^{-1}N$, which implies $p^{-1} \in O_K(N) = D$. Then $P = pM \supseteq p(p^{-1}M) = M$, a contradiction. Hence every v -submodule N of M is principal and so M is a UFM. \square

In a UFD, the concept of a principal ideal, a v -ideal, and an invertible ideal are equivalent.

Remark 2.6. Let D be a UFD and A be a v -ideal of D . Then

- (1) D is a UFM over D .
- (2) A is a UFM over D .
- (3) If M is a finitely generated projective module over D , then M is a UFM. In particular, every finite direct sum of D is a UFM.

Proof. (1) It is clear.

- (2) Note that A is principal since A is a v -ideal of D . Then A is isomorphic to D as a D -module. Hence A is a UFM by (1).
- (3) By Theorem 3.1 of [15], M is a v -multiplication module. Then by Theorem 2.3, M is a UFM since D is a UFD. \square

3. Strongly graded modules which are UFMs

Throughout this section, $D = \bigoplus_{n \in \mathbb{Z}} D_n$ is a strongly graded domain. It is known that D is a G-Dedekind domain if and only if D_0 is a G-Dedekind domain by Theorem 2.1 of [11]. Assume that K_0 and K are the quotient fields of D_0 and D respectively. Let $M = \bigoplus_{n \in \mathbb{Z}} M_n$ be a strongly graded module over D where M_0 is a finitely generated torsion-free D_0 -module and we assume that M satisfies the ascending chain condition on v -submodules of M . In this section, we will prove that M is a UFM over D if M_0 is a UFM over D_0 .

Note that the concept of a principal ideal, a v -ideal, and an invertible ideal are equivalent in a UFD. We begin this section with the following proposition.

Proposition 3.1. *If D_0 is a UFD then $D = \bigoplus_{n \in \mathbb{Z}} D_n$ is a UFD.*

Proof. Suppose that D_0 is a UFD, that is, D_0 is a maximal order and each prime v -ideal P_0 of D_0 is principal (see Proposition 1 of [3]). Then D is a maximal order by Theorem 1 of [6]. Let P be a non-zero prime v -ideal of D . If $P_0 = P \cap D_0 \neq \{0\}$, then $P = P_0 D$ and P_0 is a v -ideal of D_0 (see the proof of Theorem 2.1 and Lemma 1.2 of [11]). So $P_0 = p_0 D_0 = D_0 p_0$ for some $p_0 \in P_0$ and $P = p_0 D = D p_0$ follows. If $P_0 = P \cap D_0 = (0)$, then $P = w A_0^{-1} B_0 D$ for some invertible ideals A_0, B_0 of D_0 , which implies P is principal since D_0 is a UFD. Thus P is principal and hence D is a UFD by Proposition 1 of [2]. \square

Recall that a module M over a CIC domain D is a UFM if and only if every prime v -submodule P of M is principal, that is, $P = pM$ for some element $p \in D$ by Theorem 2.5 (see [12] for the definition of a UFM).

Note that M is a finitely generated torsion-free D -module since M_0 is a finitely generated torsion-free D_0 -module. Furthermore, M_0 is CIC if and only if M is CIC by Theorem 3.1 of [3].

In the rest of this section, we assume that M_0 is a UFM. Then D_0 is a UFD (see Theorem 2.3).

Now, we study the structure of v -submodule P of M with $P \cap M_0 \neq \{0\}$.

Lemma 3.2. [Lemma 5.1 of [3]] *Let N_0 be a fractional D_0 -submodule of M_0 with $N_0 \subseteq M_0$ and $N = DN_0$. Then*

- (1) $N^- = D(N_0)^-$, and
- (2) $N_v = D(N_0)_v$.

Lemma 3.3. *Let P be a prime D -submodule of M with $P_0 = P \cap M_0 \neq \{0\}$. Then*

- (1) P_0 is a prime submodule of M_0 , and
- (2) $P' = DP_0$ is a prime submodule of M .
- (3) If P is a prime v -submodule, then P_0 is a prime v -submodule of M_0 , and $P = DP_0$.

Proof. (1) See Lemma 5.2 of [3].

(2) See Lemma 5.2 of [3].

(3) Let $P' = DP_0 \subseteq M$. Consider that $P = P_v \supseteq (P')_v = (DP_0)_v = D(P_0)_v$ by Lemma 3.2. Thus $P_0 = P \cap M_0 \supseteq D(P_0)_v \cap M_0 = (P_0)_v$. Hence $P_0 = (P_0)_v$ and so P_0 is a prime v -submodule by (1).

Note that $P' = DP_0 = Dp_0 M_0$ for some non-zero $p_0 \in D_0$ because M_0 is a UFM. Since Dp_0 is an invertible ideal, $(P')^- = (Dp_0)^{-1} = Dp_0^{-1} \supseteq P^-$,

which implies $D \supseteq Dp_0P^-$ and $P' = Dp_0M_0 = Dp_0M \supseteq Dp_0P^-P$. If $P \supset P'$ then $Dp_0P^-M \subseteq P' \subseteq Dp_0M$ since P' is a prime submodule by (2). Then $P^-M \subseteq M$ and so $P^- = D$ since M is CIC. Thus $P = P_v = (P^-)^+ = (D)^+ = M$, a contradiction. Hence $P = DP_0$. \square

In the rest of this section, we assume that M satisfies the ascending chain condition on v -submodules of M .

Proposition 3.4. *Let N be a v -submodule of M with $N_0 = N \cap M_0 \neq \{0\}$. Then*

- (1) N_0 is a v -submodule of M_0 and $N_0 = \mathfrak{n}_0M_0$ for some ideal \mathfrak{n}_0 of D_0 .
- (2) $N = D\mathfrak{n}_0M$ and $D\mathfrak{n}_0 = (N : M)$.

Proof. (1) Similar to the previous lemma, we get that N_0 is a v -submodule of M_0 . Furthermore, $N_0 = \mathfrak{n}_0M_0$ for some ideal \mathfrak{n}_0 of D_0 follows since M_0 is a UFM over D_0 and by Theorem 2.3.

(2) Suppose there is a v -submodule N such that $N \neq D\mathfrak{n}_0M$ where \mathfrak{n}_0 is an ideal of D_0 . We may assume that N is maximal with this property because M satisfies the ascending chain condition on v -submodules of M . Then there is a maximal v -submodule P with $P \supseteq N$ and $P = D\mathfrak{p}_0M$, where \mathfrak{p}_0 is a maximal ideal of D_0 . It follows that $M \supseteq (D\mathfrak{p}_0)^{-1}N \supseteq N$. If $(D\mathfrak{p}_0)^{-1}N = N$, then $(D\mathfrak{p}_0)^{-1} \subseteq D$, a contradiction because M is CIC. Thus $(D\mathfrak{p}_0)^{-1}N \supset N$ and it follows from Lemma 3.2 of [14] that $((D\mathfrak{p}_0)^{-1}N)_v = (D\mathfrak{p}_0)^{-1}N$. By the choice of N , $(D\mathfrak{p}_0)^{-1}N = D\mathfrak{t}_0M$ for some ideal \mathfrak{t}_0 of D_0 . Hence $N = D\mathfrak{p}_0\mathfrak{t}_0M$, a contradiction. Hence $N = D\mathfrak{n}_0M$ for some ideal \mathfrak{n}_0 of D_0 . The last statement easily follows since $D\mathfrak{n}_0$ is invertible. \square

Next we study the structure of a prime v -submodule P of M such that $P \cap M_0 = \{0\}$. Since $K^g = \bigoplus_{n \in \mathbb{Z}} K_0D_n = K_0D$ is a principal ideal domain by [6] and K_0M is a finitely generated torsion-free K^g -module, we have that a v -submodule P_1 of K_0M is prime if and only if $P_1 = \mathfrak{p}_1K_0M$, where \mathfrak{p}_1 is a maximal ideal of K^g such that $\mathfrak{p}_1 = (P_1 : K_0M)$ by Theorem 3.3 of [14].

Note that if D_0 is a UFD and \mathfrak{p} is a prime v -ideal of D , then $\mathfrak{p} = \mathfrak{p}_0D$ for some prime v -ideal \mathfrak{p}_0 of D_0 or $\mathfrak{p} = \mathfrak{p}_1 \cap D$ for some prime ideal \mathfrak{p}_1 of K_0D by Lemma 2.6 of [11], and moreover $\mathfrak{p} = pD$ for some $p \in D$ by Proposition 3.1.

The following lemma is a graded version of Lemma 4.5 of [14].

Lemma 3.5. *Let N be a D -submodule of M . Then*

- (1) $(K_0N : K_0M) = K_0\mathfrak{n}$, where $\mathfrak{n} = (N : M)$ and $K_0N^- = (K_0N)^-$.
- (2) $(K_0N)_v = K_0N_v$.

Proof. See the proof of Lemma 5.4 of [3]. □

The following lemma is a graded version of Lemma 4.6 of [14]. We write the proof because we need v_1 -operation to prove the last properties (see [10], [14], [15] for details regarding v -submodules and v_1 -operation).

Lemma 3.6. *Let M_0 be a UFM over D_0 and $P_1 = \mathfrak{p}_1K_0M$ be a prime v -submodule of K_0M , where \mathfrak{p}_1 is a maximal ideal of K_0D , $P = P_1 \cap M$ and $\mathfrak{p} = \mathfrak{p}_1 \cap D$. Then*

- (1) P is a prime submodule of M and $\mathfrak{p} = (P : M)$.
- (2) $K_0P = P_1$ and $P \cap M_0 = \{0\}$.
- (3) $P = \mathfrak{p}M$ and P is a maximal v -submodule of M

Proof. (1) See Lemma 5.5 (1) of [3].

(2) See Lemma 5.5 (2) of [3].

(3) By Lemma 3.5 and (2), we have $P_1 = (P_1)_v = (K_0P)_v = K_0P_v$, so P is a v -submodule of M . Since M is a v -Noetherian D -module, there are finite elements $m_i \in P$ such that $P = (Dm_1 + \dots + Dm_k)_v$. Note that $K_0P = K_0(Dm_1 + \dots + Dm_k)_v = (K_0Dm_1 + \dots + K_0Dm_k)_v$ by Lemma 3.5. Further since $K_0P = P_1 = K_0\mathfrak{p}_1K_0M = \mathfrak{p}_1K_0M$, for m_i there are finite $p_{ij} \in \mathfrak{p}$ and $l_{ij} \in K_0M$ such that $m_i = \sum_j p_{ij}l_{ij}$. Then there is a non-zero $c \in D_0$ with $cl_{ij} \in M$ for all l_{ij} so that $cm_i \in \mathfrak{p}M$. Put $\mathfrak{a} = \{r_0 \in D_0 \mid r_0P \subseteq \mathfrak{p}M\}$, an ideal of D_0 with $\mathfrak{a}P \subseteq \mathfrak{p}M$. If $\mathfrak{a} = D_0$, then $P = \mathfrak{p}M$ and we are done. If $\mathfrak{a} \subset D_0$, by Lemma 3.2 of [10], $\mathfrak{a}_{v_1}P \subseteq (\mathfrak{a}_{v_1}P)_v = (\mathfrak{a}P)_v \subseteq (\mathfrak{p}M)_v = \mathfrak{p}M_v = \mathfrak{p}M$ because \mathfrak{p} is an invertible ideal. By the definition of \mathfrak{a} , we have that $\mathfrak{a}_{v_1} \subseteq \mathfrak{a}$, which implies $\mathfrak{a}_{v_1} = \mathfrak{a}$, that is, \mathfrak{a} is a v_1 -ideal of D_0 . Since \mathfrak{a} is a v_1 -ideal of D_0 , \mathfrak{a}^+ is a v -submodule of M_0 by Lemma 2.3 of [15], which implies $\mathfrak{a}^+ = r_0M_0$ because M_0 is a UFM. Then $\mathfrak{a} = \mathfrak{a}_{v_1} = (\mathfrak{a}^+)^- = (r_0M_0)^- = r_0^{-1}D_0$ and so \mathfrak{a} is an invertible ideal. Note that $\mathfrak{p}^{-1}\mathfrak{a}P \subseteq M$ and $K_0\mathfrak{p}^{-1}\mathfrak{a}P = K_0\mathfrak{p}^{-1}\mathfrak{p}_1K_0M = K_0M$, since $K_0D\mathfrak{p} = \mathfrak{p}_1$. It follows that $\mathfrak{p}^{-1}\mathfrak{a}P \cap M \neq \{0\}$ and $(\mathfrak{p}^{-1}\mathfrak{a}P)_v = \mathfrak{p}^{-1}\mathfrak{a}P_v = \mathfrak{p}^{-1}\mathfrak{a}P$ by Lemma 3.2 of [14] since $\mathfrak{p}^{-1}\mathfrak{a}$ is an invertible D -ideal in K^g . Then by Proposition 3.4, $\mathfrak{p}^{-1}\mathfrak{a}P = \mathfrak{n}DM$ for some ideal \mathfrak{n} of D_0 and $P = \mathfrak{p}\mathfrak{a}^{-1}\mathfrak{n}DM$. It follows that $\mathfrak{p} = (P : M) = \mathfrak{p}\mathfrak{a}^{-1}\mathfrak{n}D$ and that $D = \mathfrak{a}^{-1}\mathfrak{n}D$. Hence $P = \mathfrak{p}M$.

To prove that P is a maximal v -submodule of M , let N be a maximal

v -submodule of M containing P . Then K_0N is a v -submodule of K_0M containing $K_0P = P_1$ by Lemma 3.5, so $K_0N = P_1$ by the assumption. Thus $P = P_1 \cap M \supseteq N$ and $N = P$ follows. Hence P is a maximal v -submodule of M . \square

Lemma 3.7. *Let M_0 be a UFM over D_0 and P be a prime v -submodule of M such that $P \cap M_0 = \{0\}$. Then there is a maximal v -submodule P_1 of K_0M such that $P = P_1 \cap M$.*

Proof. Let $\mathfrak{p} = (P : M)$. Then \mathfrak{p} is a prime v -ideal of D , so \mathfrak{p} is a non-zero minimal prime ideal. Thus \mathfrak{p} is in one of the following form: $\mathfrak{p} = \mathfrak{p}_0D$ for some prime ideal \mathfrak{p}_0 of D_0 or $\mathfrak{p} = \mathfrak{p}_1 \cap D$ for some prime ideal \mathfrak{p}_1 of K_0D by Theorem 2.1 and Lemma 2.6 of [11]. In the first case, $P \supseteq \mathfrak{p}_0DM \supseteq \mathfrak{p}_0M_0 \neq \{0\}$, a contradiction. Hence $\mathfrak{p} = \mathfrak{p}_1 \cap D$ with $K_0\mathfrak{p} = \mathfrak{p}_1$. Since $P \cap M_0 = \{0\}$, $K_0M \supset K_0P = (K_0P)_v$ by Lemma 3.5. Thus there is a maximal v -submodule P_1 of K_0M such that $P_1 \supseteq K_0P$. By Lemma 3.5, $(P_1 : K_0M) \supseteq (K_0P : K_0M) = K_0(P : M) = K_0\mathfrak{p} = \mathfrak{p}_1$. Since $(P_1 : K_0M)$ is a prime ideal of K_0D , $\mathfrak{p}_1 = (P_1 : K_0M)$. Hence $P_1 = \mathfrak{p}_1K_0M$ and $P_1 \cap M \supseteq P$. By Lemma 3.6, $P_1 \cap M = \mathfrak{p}M \subseteq P$ and hence $P = P_1 \cap M$ and $P = \mathfrak{p}M$. \square

Proposition 3.8. *Let P be a prime v -submodule of M with $P \cap M_0 = \{0\}$. Then $P = \mathfrak{p}M$ for some prime v -ideal D where $\mathfrak{p} \cap D_0 = \{0\}$.*

From Lemma 3.3 and Proposition 3.8 we get the following theorem.

Theorem 3.9. *Let $D = \bigoplus_{n \in \mathbb{Z}} D_n$ be a strongly graded domain and $M = \bigoplus_{n \in \mathbb{Z}} M_n$ be a strongly graded module over D where M satisfies the ascending chain condition on v -submodules of M . If M_0 is a UFM over D_0 then M is a UFM over D .*

Proof. Since D_0 is a UFD, D is a UFD by Proposition 3.1, and so every prime v -ideal of D is principal. It is clear that D is a maximal order by Proposition 1 of [2], which implies D_0 is a maximal order by Theorem 1 of [6] and so M is CIC by Theorem 3.1 of [3]. To prove M is a UFM, let P be a prime v -submodule of M and $P_0 = P \cap M_0$.

- (1) Case $P_0 \neq \{0\}$. Then $P = DP_0$ by Lemma 3.3. Note that P_0 is a prime v -submodule of D_0 by Lemma 3.3. Since M_0 is a UFM, $P_0 = p_0M_0$ for some $p_0 \in D_0$ and so $P = DP_0 = Dp_0M_0 = p_0DM_0 = p_0M$.

- (2) Case $P_0 = \{0\}$. Then $P = \mathfrak{p}M$ for some v -ideal \mathfrak{p} of D where $\mathfrak{p} \cap D_0 = \{0\}$ by Proposition 3.8. Since D is a UFD, $\mathfrak{p} = pD$ for some element $p \in D$ and so $P = \mathfrak{p}M = pDM = pM$, for some element $p \in D$.

Hence every prime v -submodule of M is principal and so M is a UFM by Theorem 2.5. \square

As an application of Theorem 3.9, we have the following examples.

Example 3.10. If M is a UFM over an integral domain D then the Laurent polynomial module $M[x, x^{-1}]$ is a UFM over $D[x, x^{-1}]$.

Example 3.11. Let T be an arbitrary UFD and A, B be two non-zero v -ideals of T . Let K be the quotient field of T . Then $M = \bigoplus_{n \in \mathbb{Z}} AB^n x^n = \dots + AB^{-2}x^{-2} + AB^{-1}x^{-1} + A + ABx + AB^2x^2 + \dots$ is a UFM over $D = \bigoplus_{n \in \mathbb{Z}} B^n x^n = \dots + B^{-2}x^{-2}B^{-1}x^{-1} + T + Bx + B^2x^2 + \dots \subseteq K[x, x^{-1}]$, a Laurent polynomial ring over K .

4. Positively graded modules which are UFM's

Let $R = \bigoplus_{n \in \mathbb{Z}_0} D_n$ be a positively graded domain which is a sub-domain of the strongly graded domain $D = \bigoplus_{n \in \mathbb{Z}} D_n$. It is known that R is Noetherian if and only if D_0 is Noetherian by Proposition 2.1 of [13]. In this section, we will prove that $L = \bigoplus_{n \in \mathbb{Z}_0} M_n$, a positively graded module over R , is a UFM if and only if M_0 is a UFM over D_0 when D_0 is a Noetherian domain.

In the rest of this section, $R = \bigoplus_{n \in \mathbb{Z}_0} D_n$ and $L = \bigoplus_{n \in \mathbb{Z}_0} M_n$, where D_0 is a Noetherian domain and M_0 is a finitely generated torsion-free D_0 -module.

In [2], it is shown that R is a UFR if and only if D_0 is a UFR and D_1 is a principal D_0 -module. We begin this section with the following proposition that is a commutative case of Theorem 1 of [2].

Proposition 4.1. *A positively graded domain $R = \bigoplus_{n \in \mathbb{Z}_0} D_n$ is a UFD if and only if*

- (1) D_0 is a UFD, and
- (2) D_1 is a principal D_0 -module, that is, there is $p_1 \in D_1$ such that $D_1 = D_0 p_1$.

Note that L is a finitely generated torsion-free R -module since M_0 is a finitely generated torsion-free D_0 -module (see [3], Lemma 4.4.). Furthermore, M_0 is CIC if and only if L is CIC by Theorem 4.1 of [3].

The following lemma is a module version of Lemma 2.5 (2) of [13] and can be proved in a similar way as in Lemma 5.1 of [3].

Lemma 4.2. *Let N_0 be a fractional D_0 -submodule of M_0 with $N_0 \subseteq M_0$ and $N = RN_0$. Then*

- (1) $N^- = R(N_0)^-$,
- (2) $N_v = R(N_0)_v$.

The following lemma is a graded version of Lemma 4.2 and Lemma 4.3 of [14].

Lemma 4.3. *Let M_0 be a UFM over D_0 and P be a prime R -submodule of L with $P_0 = P \cap M_0 \neq \{0\}$. Then*

- (1) P_0 is a prime submodule of M_0 , and
- (2) $P' = RP_0$ is a prime submodule of L .
- (3) If P is a prime v -submodule, then P_0 is a prime v -submodule of M_0 , and $P = RP_0$.

Proof. (1) The proof is similar to the proof of Lemma 4.2 (1) of [14].

(2) The proof is similar to the proof of Lemma 4.2 (2) of [14].

(3) Let $P' = RP_0 \subseteq L$. Consider that $P = P_v \supseteq (P')_v = (RP_0)_v = R(P_0)_v$ by Lemma 4.2. Thus $P_0 = P \cap M_0 \supseteq R(P_0)_v \cap M_0 = (P_0)_v$. Hence $P_0 = (P_0)_v$ and so P_0 is a prime v -submodule by (1).

Note that $P' = RP_0 = Rp_0M_0$ for some non-zero $p_0 \in D_0$ because M_0 is a UFM. Since Rp_0 is an invertible ideal, $(P')^- = (Rp_0)^{-1} = Rp_0^{-1} \supseteq P^-$, which implies $R \supseteq Rp_0P^-$ and $P' = Rp_0M_0 = Rp_0L \supseteq Rp_0P^-P$. If $P \supset P'$ then $Rp_0P^-L \subseteq P' = Rp_0L$ since P' is a prime submodule by Lemma 4.3 (2). Then $P^-L \subseteq L$ and so $P^- = D$. Thus $P = P_v = (P^-)^+ = (D)^+ = L$, a contradiction. Hence $P = RP_0$. \square

The following proposition is a graded version of Proposition 4.4 of [14].

Proposition 4.4. *Let M_0 be a UFM over D_0 and N be a v -submodule of L with $N_0 = N \cap M_0 \neq \{0\}$. Then*

- (1) N_0 is a v -submodule of M_0 and $N_0 = \mathfrak{n}_0M_0$ for some ideal \mathfrak{n}_0 of D_0 .
- (2) $N = R\mathfrak{n}_0L$ and $R\mathfrak{n}_0 = (N : L)$.

Proof. (1) Similar to the previous lemma, we get that N_0 is a v -submodule of M_0 . Furthermore, $N_0 = \mathfrak{n}_0M_0$ for some ideal \mathfrak{n}_0 of D_0 follows since M_0 is a UFM over D_0 and by Theorem 2.3.

(2) Suppose there is a v -submodule N such that $N \neq R\mathfrak{n}_0L$ where \mathfrak{n}_0 is an ideal of D_0 . We may assume that N is maximal with this property because

M is Noetherian. Then there is a maximal v -submodule P with $P \supseteq N$ and $P = R\mathfrak{p}_0L$, where \mathfrak{p}_0 is a maximal ideal of D_0 . It follows that $L \supseteq (R\mathfrak{p}_0)^{-1}N \supseteq N$. If $(R\mathfrak{p}_0)^{-1}N = N$, then $(R\mathfrak{p}_0)^{-1} \subseteq R$, a contradiction because L is CIC. Thus $(R\mathfrak{p}_0)^{-1}N \supset N$ and it follows from Lemma 3.2 of [14] that $((R\mathfrak{p}_0)^{-1}N)_v = (R\mathfrak{p}_0)^{-1}N$. By the choice of N , $(R\mathfrak{p}_0)^{-1}N = R\mathfrak{t}_0L$ for some ideal \mathfrak{t}_0 of D_0 . Hence $N = R\mathfrak{p}_0\mathfrak{t}_0L$, a contradiction. Hence $N = R\mathfrak{n}_0L$ for some ideal \mathfrak{n}_0 of D_0 . The last statement easily follows since $R\mathfrak{n}_0$ is invertible. \square

Next we study the structure of a prime v -submodule P of L such that $P \cap M_0 = \{0\}$. Since $Q^g = \bigoplus_{n \in \mathbb{Z}_0} K_0D_n = K_0R$ is a principal ideal domain by Lemma 2.1 of [13] and K_0L is a finitely generated torsion-free Q^g -module, we have that a v -submodule P_1 of K_0L is prime if and only if $P_1 = \mathfrak{p}_1K_0L$, where \mathfrak{p}_1 is a maximal ideal of Q^g such that $\mathfrak{p}_1 = (P_1 : K_0L)$ by Theorem 3.3 of [14].

The following lemma is a graded version of Lemma 4.5 of [14].

Lemma 4.5. *Let N be an R -submodule of L . Then*

- (1) $(K_0N : K_0L) = K_0\mathfrak{n}$, where $\mathfrak{n} = (N : L)$ and $K_0N^- = (K_0N)^-$.
- (2) $(K_0N)_v = K_0N_v$.

Proof. (1) The proof is similar to the proof of Lemma 4.5 (1) of [14].

- (2) Let $m' \in (K_0N)_v = ((K_0N)^-)^+ = (K_0N^-)^+$, that is, $K_0L \supseteq K_0N^-m' \supseteq N^-m'$. Then there is $r \in D_0$ such that $N^-rm' = rN^-m' \subseteq L$. Thus $rm' \in (N^-)^+ = N_v$ and so $m' \in r^{-1}N_v \subseteq K_0N_v$.

Conversely, let $m' \in K_0N_v$. We write $m' = \sum_{i=1}^t k_{0i}m_i$ where $k_{0i} \in K_0$ and $m_i \in N_v$ for all $i = 1, 2, \dots, t$. Then for all $i = 1, 2, \dots, t$, we have $N^-m_i \subseteq L$ and so $K_0N^-m' = K_0N^- \left(\sum_{i=1}^t k_{0i}m_i \right) \subseteq N^-(K_0m_1 + \dots + K_0m_t) \subseteq K_0L$. Then

$$m' \in (K_0N^-)^+ = ((K_0N)^-)^+ = (K_0N)_v.$$

Hence $(K_0N)_v = K_0N_v$. \square

The following lemma is a graded version of lemma 4.6 of [14]. We write the proof because we need v_1 -operation to prove the last properties (see [10], [14], [15] for details regarding v -submodules and v_1 -operation).

Lemma 4.6. *Let M_0 be a UFM over D_0 and $P_1 = \mathfrak{p}_1 K_0 L$ be a prime v -submodule of $K_0 L$, where \mathfrak{p}_1 is a maximal ideal of $K_0 R$, $P = P_1 \cap L$ and $\mathfrak{p} = \mathfrak{p}_1 \cap R$. Then*

- (1) P is a prime submodule of L and $\mathfrak{p} = (P : L)$.
- (2) $K_0 P = P_1$ and $P \cap M_0 = \{0\}$.
- (3) $P = \mathfrak{p}L$ and P is a maximal v -submodule of L .

Proof. (1) Let $r \in R$ and $m \in L$ such that $rm \in P$ and $m \notin P$. Since $m \notin P_1$ and P_1 is prime, we have $rL \subseteq rK_0 L \subseteq P_1$ and so $rL \subseteq P$. Hence P is a prime submodule of L .

Since $\mathfrak{p}L \subseteq \mathfrak{p}K_0 L = P_1$, we have $\mathfrak{p}L \subseteq P$, so $\mathfrak{p} \subseteq (P : L)$. Conversely let $r \in (P : L)$, that is, $r \in R$ and $rL \subseteq P$. Then $rK_0 L \subseteq K_0 P \subseteq P_1$, so $r \in (P_1 : K_0 L) = \mathfrak{p}_1$. Thus $r \in \mathfrak{p}_1 \cap R = \mathfrak{p}$. Hence $\mathfrak{p} = (P : L)$.

- (2) Let $m' \in P_1$ and we write $m' = \sum_{i=1}^n t_i m_i$ where $t_i \in \mathfrak{p}_1$ and $m_i \in K_0 L$. Then there are $\alpha, \beta \in D_0$ such that $\alpha t_i \in \mathfrak{p}$ and $\beta m_i \in L$ and so $\alpha \beta m_i \in \mathfrak{p}L \subseteq P$. Thus $m' \in (\alpha \beta)^{-1} P \subseteq K_0 P$. Hence $K_0 P = P_1$.

Note that $\mathfrak{p}_1 = \langle t \rangle = tK_0 R$ for some prime element $t \in K_0 R$ with $\deg(t) \geq 1$. If $P \cap M_0 \neq \{0\}$ and let $0 \neq m \in P \cap M_0$. Then $m = tm'$ for some $m' \in K_0 L$, since $K_0 P = P_1 = tK_0 L$. Write $t = t_n + t_{n-1} + \dots + t_0$ ($t_i \in K_0 D_i$, with $t_n \neq 0$) and $m' = m_l + \dots + m_0$ ($m_j \in K_0 M_j$). Then we get $t_n m_l = 0$, so $m_l = 0$ and so on. Then we have $m = 0$, a contradiction. Hence $P \cap M_0 = \{0\}$.

- (3) The proof is similar to Lemma 3.6 (3). □

Lemma 4.7. *Let M_0 be a UFM over D_0 and P be a prime v -submodule of L such that $P \cap M_0 = \{0\}$. Then $P = \bigoplus_{n \geq 1} M_n = D_1 L$ or there is a maximal v -submodule P_1 of $K_0 L$ such that $P = P_1 \cap L$.*

Proof. Let $\mathfrak{p} = (P : L)$. Then \mathfrak{p} is a prime v -ideal of R , so \mathfrak{p} is a non-zero minimal prime ideal. Thus \mathfrak{p} is in one of the following form: $\mathfrak{p} = \mathfrak{p}_0 R$ for some prime ideal \mathfrak{p}_0 of D_0 , $\mathfrak{p} = \bigoplus_{n \geq 1} D_n$ or $\mathfrak{p} = \mathfrak{p}_1 \cap R$ for some prime ideal \mathfrak{p}_1 of $K_0 R$ by Proposition 3.1 of [13]. In the first case, $P \supseteq \mathfrak{p}_0 R L \supseteq \mathfrak{p}_0 M_0 \neq \{0\}$, a contradiction. In the second case, if $P \supseteq (\bigoplus_{n \geq 1} D_n) L = R D_1 L = D_1 L = \bigoplus_{n \geq 1} M_n$. If $P \supset \bigoplus_{n \geq 1} M_n$, there is a non-zero submodule T_0 of M_0 such that $P = T_0 + \bigoplus_{n \geq 1} M_n$. Then $P \cap M_0 \supseteq T_0 \neq \{0\}$, a contradiction. Hence $P = \bigoplus_{n \geq 1} M_n$. In the last case, $\mathfrak{p} = \mathfrak{p}_1 \cap R$ with $K_0 \mathfrak{p} = \mathfrak{p}_1$. Since $P \cap M_0 = \{0\}$, $K_0 L \supset K_0 P = (K_0 P)_v$ by Lemma 4.5. Thus there is a maximal v -submodule P_1 of $K_0 L$ such that $P_1 \supseteq K_0 P$. By Lemma 4.5, $(P_1 : K_0 L) \supseteq (K_0 P : K_0 L) = K_0(P : L) = K_0 \mathfrak{p} = \mathfrak{p}_1$. Since $(P_1 : K_0 L)$

is a prime ideal of K_0R , $\mathfrak{p}_1 = (P_1 : K_0L)$. Hence $P_1 = \mathfrak{p}_1K_0L$ and $P_1 \cap L \supseteq P$. By Lemma 4.6, $P_1 \cap L = \mathfrak{p}L \subseteq P$ and hence $P = P_1 \cap L$ and $P = \mathfrak{p}L$. So by the last two cases, $P = \bigoplus_{n \geq 1} M_n$ or there is a maximal v -submodule P_1 of K_0L such that $P = P_1 \cap L$. \square

Note that if $R = \bigoplus_{n \in \mathbb{Z}_0} D_n$ is a Noetherian UFD, then $R = D_0[p_1]$ for some element $p_1 \in D_1$ by Theorem 1 of [2] and so $M = M_0[p_1]$, a polynomial module. Then the necessary condition of Theorem 4.8 is already proved in [12], but we give another proof by using v_1 -operator.

Theorem 4.8. *Let $R = \bigoplus_{n \in \mathbb{Z}_0} D_n$ be a Noetherian UFD and $L = \bigoplus_{n \in \mathbb{Z}_0} M_n$ be a positively graded module over R . Then L is a UFM if and only if M_0 is a UFM.*

Proof. (\Rightarrow) Suppose that $L = \bigoplus_{n \in \mathbb{Z}_0} M_n$ is a UFM over R . Then L is CIC and so M_0 is CIC by Theorem 4.1 of [3]. Let P_0 be a non-zero prime v -submodule of M_0 . Then $P = RP_0$ is a v -submodule of L by Lemma 4.2. Furthermore, as in Lemma 4.3 (2), RP_0 is a prime submodule of L . Hence RP_0 is a prime v -submodule of L and so it is principal since L is a UFM. Then $RP_0 = rL$ for some $r \in R$. Since $\{0\} \neq P_0 \subset RP_0 = rL$, $r \in D_0$, which implies $P_0 = rM_0$. Hence P_0 is principal and so M_0 is a UFM by Theorem 2.5.

(\Leftarrow) Suppose that M_0 is a UFM over D_0 . Since R is a UFD, it is clear that R is a maximal order by Proposition 1 of [2], which implies D_0 is a maximal order by Theorem 2.1 of [13] and so L is CIC by Theorem 4.1 of [3]. Note that D_0 is a UFD and D_1 is a principal D_0 -module since R is a UFD. To prove L is a UFM, let P be a prime v -submodule of L and $P_0 = P \cap M_0$. By Lemma 4.3 (3), P_0 is a prime v -submodule.

- (1) Case $P_0 \neq \{0\}$. Then $P = RP_0$ by Lemma 4.3 (3). Since M_0 is a UFM, $P_0 = p_0M_0$ for some $p_0 \in D_0$ and so $P = RP_0 = Rp_0M_0 = p_0RM_0 = p_0L$.
- (2) Case $P_0 = \{0\}$. Then $P = \bigoplus_{n \geq 1} M_n = D_1L$ or $P = \mathfrak{p}L$ for some v -ideal \mathfrak{p} of R by Lemma 4.7. If $P = \bigoplus_{n \geq 1} M_n = D_1L$, then $P = d_1D_0L = d_1L$ for some $d_1 \in D_1$ since D_1 is a principal D_0 -module. If $P = \mathfrak{p}L$, then $P = \mathfrak{p}L = pRL = pL$ for some $p \in R$ since R is a UFD.

Hence every prime v -submodule of L is principal and so L is a UFM by Theorem 2.5. \square

We end this section with examples of a positively graded module which is a UFM.

Example 4.9. Let $R = \bigoplus_{n \in \mathbb{Z}_0} D_n$ be a positively graded domain where D_0 is a Noetherian UFD and D_1 is a principal D_0 -module. Let $M = R \oplus R \oplus \dots \oplus R$ be a positively graded module over R and P be a graded submodule of M such that $M = P \oplus T$. Then P is a UFM.

Proof. Note that P is a projective module. Then P is a G-Dedekind module and it is a v -multiplication module by Theorem 3.1 of [15]. Furthermore, since P is a v -multiplication module and R is a UFD, P is a UFM by Theorem 2.3. \square

Lemma 4.10. Let D be a domain, B be an invertible ideal of D and A be a non-zero ideal of D . Let $R = D + Bx + B^2x^2 + \dots \subseteq D[x]$, where $D[x]$ is a polynomial ring over D and $L = A + ABx + AB^2x^2 + \dots = AR$. Then L is a positively graded module over positively graded domain R .

From Remark 2.6 and Lemma 4.10 we have the following example.

Example 4.11. Let D be an arbitrary Noetherian UFD and A, B be two non-zero v -ideals of D . Then $L = A + ABx + AB^2x^2 + \dots$ is a UFM over $R = D + Bx + B^2x^2 + \dots$.

Proof. Note that R is a UFD since D is a UFD and Bx is a principal D -module. Since A is a non-zero v -ideal of D , A is a UFM by Remark 2.6. Then by Theorem 4.8, L is a UFM over R . \square

Acknowledgement. The authors would like to thank the referee for the valuable comments and suggestions that significantly improved this manuscript.

References

- [1] D. L. Costa, *Unique factorization in modules and symmetric algebras*, Trans. Amer. Math. Soc., 224(2) (1976), 267-280.
- [2] I. Ernanto, H. Marubayashi, A. Ueda and S. Wahyuni, *Positively graded rings which are unique factorization rings*, Vietnam J. Math., 49 (2021), 1037-1041.
- [3] I. Ernanto, A. Ueda, I. E. Wijayanti and Sutopo, *Some remarks on strongly graded modules*, submitted for publication, 2022.
- [4] R. Gilmer, *Multiplicative Ideal Theory*, Marcel Dekker, New York, 1972.
- [5] C. P. Lu, *Factorial modules*, Rocky Mountain J. Math., 7 (1977), 125-139.
- [6] H. Marubayashi, S. Wahyuni, I. E. Wijayanti and I. Ernanto, *Strongly graded rings which are maximal orders*, Sci. Math. Jpn., 82 (2019), 207-210.

- [7] C. Nastasescu and F. van Oystaeyen, *Graded Ring Theory*, North-Holland Mathematical Library, 28, North-Holland Publishing Co., Amsterdam-New York, 1982.
- [8] A. M. Nicolas, *Modules factoriels*, Bull. Sci. Math. (2), 95 (1971), 33-52.
- [9] A. M. Nicolas, *Extensions factorielles et modules factorables*, Bull. Sci. Math. (2), 98 (1974), 117-143.
- [10] M. M. Nurwigantara, I. E. Wijayanti, H. Marubayashi and S. Wahyuni, *Krull modules and completely integrally closed modules*, J. Algebra Appl., 21(1) (2022), 2350038 (14 pp).
- [11] S. Wahyuni, H. Marubayashi, I. Ernanto and Sutopo, *Strongly graded rings which are generalized Dedekind rings*, J. Algebra Appl., 19(3) (2020), 2050043 (8 pp).
- [12] S. Wahyuni, H. Marubayashi, I. Ernanto and I. P. Y. Prabhadika, *On unique factorization modules: a submodule approach*, Axioms, 11(6) (2022), 288 (7 pp).
- [13] I. E. Wijayanti, H. Marubayashi and Sutopo, *Positively graded rings which are maximal orders and generalized Dedekind prime rings*, J. Algebra Appl., 19(8) (2020), 2050143 (11 pp).
- [14] I. E. Wijayanti, H. Marubayashi, I. Ernanto and Sutopo, *Finitely generated torsion-free modules over integrally closed domains*, Comm. Algebra, 48(8) (2020), 3597-3607.
- [15] I. E. Wijayanti, H. Marubayashi, I. Ernanto and Sutopo, *Arithmetic modules over generalized Dedekind domains*, J. Algebra Appl., 21(3) (2022), 2250045 (14 pp).

Iwan Ernanto (Corresponding Author) and **Indah E. Wijayanti**

Department of Mathematics

Faculty of Mathematics and Natural Sciences

Universitas Gadjah Mada

Sekip Utara Yogyakarta, Indonesia

e-mails: iwan.ernanto@ugm.ac.id (I. Ernanto)

ind_wijayanti@ugm.ac.id (I. E. Wijayanti)

Akira Ueda

Department of Mathematics

Interdisciplinary Faculty of Science and Engineering

Shimane University

1060 Nishikawatsu-cho, Matsue, Shimane 690-8504, Japan

e-mail: ueda@riko.shimane-u.ac.jp (A. Ueda)