


Uterine Tumor Resembling Ovarian Sex Cord Tumor; A Rare Case Report with Histopathology, Immunohistochemical Findings and P53 Expression

Overin Seks Kord Tümörüne Benzeyen Uterin Tümör; Histopatoloji, İmmunohistokimyasal Bulgular ve P53 İfadesi İçeren Nadir Bir Olgu Sunumu

^{1*} Ebru Altındal Susem 

¹ Niğde Omer Halisdemir University, School of Medicine, Department of Surgical Medicine, Department of Pathology

*Sorumlu Yazar/Corresponding Author
ebruasusem@gmail.com

Geliş Tarihi/ Date of Submission: 08.01.2024

Kabul Tarihi/ Date of Acceptance: 12.03.2024

Yayın Tarihi/ Date of Publication: 14.03.2024

Değerlendirme/ Peer-Review: İki Dış Hakem, Çift Taraflı Körleme / Two external, Double anonymized

Etik Beyan / Ethical Statement: Bu çalışmanın hazırlanma sürecinde bilimsel ve etik ilkelere uyulduğu ve yararlanılan tüm çalışmaların kaynakçada belirtildiği beyan olunur./It is declared that scientific and ethical principles have been followed while carrying out and writing this study and that all the sources used have been properly cited.

BenzerlikTaraması/ Plagiarism checks: Yapıldı – Turnitin/ Yes – Turnitin

Çıkar Çatışması/ Conflicts of Interest: Çıkar çatışması beyan edilmemiştir/The author(s) has no conflict of interest to declare

Finansman/ Grant Support: Bu araştırmayı desteklemek için dış fon kullanılmamıştır/The author(s) acknowledge that they received no external funding in support of this research

Telif Hakkı & Lisans/Copyright & License: Yazarlar dergide yayınlanan çalışmalarının telif hakkına sahiptirler ve çalışmalarını CC BY-NC 4.0 lisansı altında yayımlanmaktadır/Authors publishing with the journal retain the copyright to their work licensed under the CC BY-NC4.0

Abstract

Uterine tumors resembling ovarian sex cord tumors, is a rare mesenchymal tumor of the uterus that displays similar histological features of sex cord tumors. The pathogenesis for this tumor, which was first described by Morehead and Bowman in 1945 and about 100 cases of which have been published so far, is still unknown. It is frequently seen in perimenopausal or menopausal women around the age of 45. It can manifest itself with gynecological symptoms such as menorrhagia, postmenopausal bleeding, abdominal pain, or it can be detected incidentally with no symptoms. Although it behaves like benign tumors, its attitude is unpredictable due to publications reporting that it recurs and metastases and it can be considered as a low malignancy potential tumor. Therefore, bilateral salpingooforectomy and hysterectomy is the most reliable method in the treatment, except for the reproductive age where only tumor resection is performed. Here, I present a case of this rare tumor accompanied by histomorphological findings and immunohistochemical studies including P53. P53 marker study in uterine tumor resembling ovarian sex cord tumor has been reported in only four cases in the literature; three of them showed P53 marker positivity and one mentioned that it could be associated with recurrence or tumor progression. More studies need to be done in terms of prognostic or diagnostic value between uterine tumors resembling ovarian sex cord tumors and P53 marker. Our case showed overt expression with p53.

Keywords: Sex cord tumor, uterine tumor, P53 marker

Özet

Overin seks kord-stromal tümörüne benzeyen uterin tümör, seks kord tümörlerine benzer histolojik özellikler gösteren, uterusun nadir görülen mezankimal tümörleridir. İlk kez 1945 yılında Morehead ve Bowman tarafından tanımlanan ve bugüne kadar 100'e yakın vakası yayınlanmış olan bu tümörün patogenezi halen bilinmemektedir. Sıklıkla 45 yaş civarında perimenopozal veya menopozal kadınlarda görülür. Menoraji, postmenopozal kanama, karın ağrısı gibi jinekolojik semptomlarla kendini gösterebileceği gibi herhangi bir semptom olmadan tesadüfi tesbit edilebilir. İyi huylu tümörler gibi davranmasına rağmen, rekürrens ve metastaz yaptığını bildiren yayımlar nedeniyle tutumu öngörülemez ve düşük malignite potansiyelli bir tümör olarak kabul edilebilir. Bu nedenle üreme yaşı dışında sadece tümör rezeksiyonu yapılan olgularda, tedavide iki taraflı salpingooforektomi ve histerektomi en güvenilir yöntemdir. Burada nadir görülen bu tümörün bir olgusunu histomorfolojik bulgular ve P53 dahil immünohistokimyasal çalışmalar eşliğinde sunuyorum. Overin seks kord-stromal tümörüne benzeyen uterin tümöre ait P53 marker çalışması literatürde sadece dört olguda rapor edilmiş olup; üçü P53 marker pozitifliği göstermiş ve biri bunun nüks veya tümör ilerlemesi ile ilişkili olabileceğinden bahsetmiştir. Overin seks kord-stromal tümörüne benzeyen uterin tümörü ile P53 belirteci arasında prognostik veya tanısal değer açısından daha fazla çalışma yapılması gerekmektedir. Bizim olgumuzda p53 ile belirgin ekspresyon görüldü.

Anahtar Kelimeler: Seks kord tümör, uterin tümör, P53 marker

Introduction

Uterine tumors resembling ovarian sex cord tumors, is a rare mesenchymal tumor of the uterus that displays similar histological features of sex cord tumors. The pathogenesis for this tumor, which was first described by Morehead and Bowman in 1945 and about 100 cases of which have been published so far, is still unknown (1). It is frequently seen in perimenopausal or menopausal women around the age of 45 (2). It can manifest itself with gynecological symptoms such as menorrhagia, postmenopausal bleeding, abdominal pain, or it can be detected incidentally with no symptoms (3). Although it behaves like benign tumors, its attitude is unpredictable due to publications reporting that it recurs and metastases and it can be considered as a low malignancy potential tumor (4-6). Therefore, bilateral salpingo-oophorectomy and hysterectomy is the most reliable method in the treatment, except for the reproductive age where only tumor resection is performed (7).

Histopathologically, they show varying features such as glandular, macrofollicular, microfollicular, trabecular, tubular, cords, retiform, solid clusters, diffuse or mixed. Immunohistochemically, it displays a pattern of unknown origin due to its staining with epithelial, smooth muscle and sex cord markers. For this reason, it is among the miscellaneous mesenchymal tumors of the corpus uteri in the World Health Organization classification (8).

Here, it is presented a case of this rare tumor accompanied by histomorphological findings and immunohistochemical studies including P53. P53 marker study in uterine tumor resembling ovarian sex cord tumor has been reported in only four cases in the literature; three of them showed P53 marker positivity and one mentioned that it could be associated with recurrence or tumor progression (3,7,9). More studies need to be done in terms of prognostic or

diagnostic value between uterine tumors resembling ovarian sex cord tumors and P53 marker. Our case showed overt expression with p53.

Case Report

In a 47-year-old patient who did not have any complaints and came for intrauterine contraceptive device control, a 16 mm diameter polyps in the uterine cavity was detected in ultrasound. Sampling is done by curettage from the patient.

Microscopic examination of 1.5 cc endometrial sampling revealed two endocervical polyps showing squamous metaplasia between endometrial tissues in the early secretory phase. Besides, very dense cellular and vascularized polypoid tumor fragments were observed. These fragments were mostly solid layers with a focal trabecular and retiform pattern of neoplastic cells with focal pleomorphism, usually with monotonous oval-round nuclei with faint nucleoli and scant to abundant eosinophilic cytoplasm. There was no mitosis and necrosis (Figure 1).




Figure 1. Histology of the uterine tumor resembling ovarian sex-cord tumor (a) Polypoid tumor, (b) trabecular pattern, (c,g) retiform pattern, (d,e) solid areas, (f) densely cellular and vascular (Hematoxylin-eosin staining, a: $\times 40$, d, e: $\times 100$, the others $\times 200$)

In immunohistochemical studies, neoplastic cells showed patch staining with the epithelial marker pancytokeratin, diffuse staining with desmin and calponin, the myoid markers, diffuse staining with the ovarian sex-cord marker CD56, focal staining with calretinin and cell clusters with melan A. In addition, diffuse staining with vimentin and patchy staining with BCL2 was observed (Figure 2).




Figure 2. Immunohistochemistry of the uterine tumor resembling ovarian sex-cord tumor ($\times 40$)

Diffuse nuclear staining with estrogen, progesterone and P53, and point nuclear staining with cyclin D1 was remarkable. No staining was seen with EMA, CD99, inhibin, SMA, P16, CD34, CD117, S100, HMB45, β catenin. Ki 67 proliferation index was evaluated as 5-10%. This staining pattern supports the diagnosis of a uterine tumors resembling ovarian sex cord tumors.

Discussion

First described by Morehead and Bowman in 1945 (1), it was divided into two groups according to the proportion of ovarian sex cord tumors-like components in 1976 by Clement and Scully, and this tumor, which was named as ovarian sex cord tumor (UTROSCT), was 50% -100%. sex cord-like histomorphology is in question. The other group was named as endometrial stromal tumors with sex cord-like elements (ESTSCLE), with less than 50% sex cord tumor-like components (10).

UTROSCT; it often presents clinical symptoms such as menorrhagia, postmenopausal bleeding, abdominal distention, abdominal or pelvic pain (11). It may also appear asymptotically. Usually manifests as an intramural, submucosal or subserosal mass in the uterine corpus (12). It is often interpreted as intracavitary polyps or intramural myomatous lesions. The diagnosis is made by histopathological and immunohistochemical examinations.

It may show histomorphological variants as solid layers or small clusters, or as retiform or plexiform pattern, or as intersecting cords of two cell widths, or in a trabecular or tubular form, or with macrofollicular or microfollicular characteristics, or diffuse or mixed pattern within the hyalinized or fibroblastic stroma (11). Rarely, formations similar to Call-Exner bodies observed in granulosa cell tumors and cells with eosinophilic or foamy cytoplasm resembling luteinized stromal cells can be seen (13-16). Mild cellular atypia may be observed, but it is generally composed of cells with round or oval nuclei in uniform appearance, indeterminate nucleolus and eosinophilic cytoplasm. Nuclei can be hyperchromatic or a nuclear groove can be observed in between (11,17-19). Rare lymphocytic infiltration, foamy histiocytes, focal

hemosiderin may appear as other uncommon microscopic findings (20). Necrosis, vascular invasion, 2-3 (10 hpf) mitosis are extremely rare; their presence indicates aggressive progression (21)

Immunohistochemically, UTROSCT has a highly variable staining pattern just like its histology. For diagnosis, it should be stained with one or more of the markers of both ovarian sex cord, smooth muscle and epithelial. The sex cord markers it is dyed frequently include CD56, inhibin, calretinin, WT1, CD99, melan A. It is also expected to be stained with SMA, caldesmon, calponin, desmin as myoid and pancytokeratin, EMA, CK AE1 / 3 as epithelial markers. Additionally, it stains with hormone markers such as estrogen, progesterone, androgen and various markers such as vimentin, CD10, BCL2, CD117, S100. Staining with P53 has been reported in the literature in a total of three cases, one of which was associated with recurrence (3,7,22). Immunohistochemical markers and staining diffusions for UTROSCT are different from each other (16,23). Not staining with CD34, chromogranin and HMB45 is important in differential diagnosis (4-6,24).

UTROSCT can be similar to many uterine tumors. Our case was predominantly solid component histologically. Therefore, endometrial polyp, ESTSCLE, endometrial stromal tumors, PEComa, cellular leiomyoma, epithelioid hemangioendothelioma were considered in the differential diagnosis.

ESTSCLE is more similar to endometrial stromal tumors. Its molecular structure, clinical course and treatment protocol are different from UTROSCT (17). However, it is microscopically quite similar to UTROSCT. JAZF1 and JJAZ1 fusion monitored in ESTSCLE. It is not seen in UTROSCT and its molecular structure is still unresolved. In UTROSCT, it is expected to be painted with at least two sex cord markers, especially the calretinin. In

ESTSCLE, staining is observed with only one sex cord marker (inhibin, CD99, Melan-A and Wilms tumor 1), mostly calretinin (17).

In 2020, the World Health Organization; Endometrial Stromal Tumors classified in 4 groups as endometrial stromal nodüle, low grade endometrial stromal sarcoma, high grade endometrial stromal sarcoma and undifferentiated stromal sarcoma (25). Endometrial stromal nodules usually show diffuse staining with CD10, SMA, vimentin, CD56, estrogen receptors, and focal staining with pancytokeratin, desmin, and progesterone receptors. No staining expected with WT1, S100, CD117, EMA, CD34, Cyclin D1 (26). Diffuse or focal staining with CD10, SMA, desmin, estrogen receptor, progesterone receptor, cyclin D1, pancytokeratin, WT1, CD34, BCL2, vimentin may occur in low grade endometrial stromal sarcoma; however, they do not stain with S100, CD117, EMA, HMB45, CD99, melan A, synaptophysin, chromogranin, inhibin and P53 (27-29). High grade endometrial stromal sarcomas are stained with CD56, CD117, CD99, cyclin D1, staining with CD10 is variable. Usually staining is not expected with pancytokeratin, DOG1, EMA, SMA, desmin, calretinin, inhibin, estrogen receptor and progesterone receptor (30,31). While stromal cells were stained with P16, CD10, P63 in endometrial polyps; no staining with pancytokeratin, EMA (32). PEComa is almost always positive with SMA, cathepsin K, HMB45, desmin, and melan A markers (33). They are not stained with the markers pancytokeratin, PAX8, S100, CD10 (34). Epithelioid hemangioendothelioma is expected to be CD34 positive, but not staining with desmin (35).

There are a number of studies done with P53 marker in UTROSCT (9). The differential diagnosis and prognostic value of p53 marker positivity are not yet known. New studies may clarify this issue in the future.

As a last word, mastering the pathology of this rare tumor, which is considered to be tumors with uncertain attitude or low malignant potential, enables accurate clinical follow-up and treatment.

Acknowledgement

Written informed consent was obtained from the patient for this case report and publication of the microscopic images of the case.

References

1. Morehead RP, Bowman MC. Heterologous mesodermal tumors of the uterus: Report of a neoplasm resembling a granulosa cell tumor. *Am J Pathol* 1945; 21(1):53–61.
2. Nyugen CV, Phung HT, Dao LT, TaDHH, Tran MN. Uterine tumor resembling ovarian sex cord tumor: clinicopathological characteristics of a rare case. *Case Rep Oncol* 2020; 13(2):807-12
3. Fan LL, Shen Y, Chanda K, Ren ML. Uterine tumors resembling ovarian sex-cord tumor: A case report and literature review. *J Can Res Ther* 2018; 14 (Suppl S5):1209–12.
4. Mačák J, Dundr P, Dvořáčková J, KlátJ. Uterine tumors resembling ovarian sex cord tumors (UTROSCT). Report of a case with lymph node metastasis. *Cesk Patol* 2014; 50:46–49.
5. Umeda S, Tateno M, Miyagi E, Sakurai K, Tanaka R, et al. Uterine tumors resembling ovarian sex cord tumors (UTROSCT) with metastasis: Clinicopathological study of two cases. *Int J Clin Exp Pathol* 2014; 7:1051–9.
6. Biermann K, Heukamp LC, Büttner R, Zhou H. Uterine tumor resembling an ovarian sex cord tumor associated with metastasis. *Int J Gynecol Pathol* 2008; 27:58–60.
7. Çetinkaya N, Bas S, Cuyilan ZF, Erdem O, Erkaya S, et al. Uterine tumor resembling ovarian sex cord tumors: A case report and literature review. *Oncol Lett* 2016; 11(2):1496-8.
8. Kim K-R, Lax SF, Lazar AJ, Longacre TA, Malpica A, et al. Tumours of uterine corpus. Female genital tumours, WHO Classification of tumours, 5th Ed. Lyon, France: IARC Press; 2020: 294-5.
9. Zhang X, Zou S, Gao B, Qu W. Uterine tumor resembling ovarian sex cord tumor: Clinicopathological and immunochemical analysis of two cases and a literature review. *J Int Med Res* 2019; 47(3):1339–47.
10. Clement PB, Scully RE. Uterine tumors resembling ovarian sex-cord tumors: A clinicopathologic analysis of fourteen cases. *Am J Clin Pathol* 1976; 66(3):512–5.
11. Sutak J, Lazic D, Cullimore JE. A uterine tumor that looks like an ovarian sex cord tumor. *Journal of Clinical Pathology* 2005; 58 (8):888–90.
12. Pradhan D, Mohanty SK. Uterine tumors resembling ovarian sex cord tumors. *Arch Pathol Lab Med* 2013;137:1832-6.
13. Fekete PS, Vellios F, Patterson BD . Uterine tumor resembling ovarian sex-cord tumor: report of a case of an endometrial stromal tumor with foam cells and ultrastructural evidence of epithelial differentiation. *Int J Gynecol Pathol* 1985;4:378-87.
14. Baker RJ, Hildebrandt RH, Rouse RV, Hendrickson MR, Longacre TA. Inhibin and CD99 (MIC2) expression in uterine stromal neoplasms with sex-cord-like elements. *Hum Pathol* 1999;30:671-9.
15. Nogales FF, Isaac MA . Functioning uterine sex cord tumour. *Histopathol* 2002;41:277-9.
16. Hashmi AA, Faridi N, Edhi MM, Khan M. Uterine tumor resembling ovarian sex cord tumor (UTROSCT), case report with literature review. *Int Arch Med* 2014; 7:47.
17. Czernobilsky B. Uterine tumors resembling ovarian sex cord tumors: an

update. *Int J Gynecol Pathol* 2008;27(2):229–35.

18. Gupta M, de Leval L, Selig M, Oliva E, Nielsen GP. Uterine tumors resembling ovarian sex cord tumors: An ultrastructural analysis of 13 cases. *Ultrastruct Pathol* 2010;34(1):16–24.

19. Hauptmann S, Nadjari B, Kraus J, Turnwald W, Dietel M. Uterine tumor resembling ovarian sex-cord tumor: A case report and review of the literature. *Virchows Arch* 2001;439(1):97–101.

20. Symmonds RE, Dockerty MB, Pratt JH. Sarcoma and sarcoma-like proliferations of the endometrial stroma. III. Stromal hyperplasia and stromatosis (stromal endometriosis). *Am J Obstet Gynecol* 1957;73:1054-70.

21. Moore M, McCluggage WG. Uterine tumour resembling ovarian sex cord tumour: First report of a large series with follow-up. *Histopathol* 2017;71:751-59.

22. Wei X-Q, Tang M, Liu L, Zhao S-P. Uterine tumor resembling ovarian sex cord tumor: A clinicopathological and immunohistochemical analysis of a recurrent case and a literature review. 2021;assets.researchsquare.com.

23. Marucci O, Nicoletti P, Mauriello A, Facchetti S, Patrizi L, et al. Uterine tumor resembling ovarian sex cord tumors type II with vaginal vault recurrence. *Case Rep Obstet Gynecol* 2019;2019:5231219.

24. Hurrell DP, McCluggage WG. Uterine tumour resembling ovarian sex cord tumour is an immunohistochemically polyphenotypic neoplasm which exhibits coexpression of epithelial, myoid and sex cord markers. *Journal of Clinical Pathology* 2007;60(10):1148–54.

25. Akaev L, Yeoh CC, Rahimi. Update on endometrial stromal tumours of the uterus. *Diagnostics* 2021;11:429.

26. Nomura Y, Tamura D, Horie M, Sato M, Sasaki S, et al. Detection of MEAF6-PHF1 translocation in an endometrial stromal nodule. *Genes Chromosomes Cancer* 2020;59:702–8.

27. Alan S, Yilmaz E, Tecellioglu F, Akatli AN, Coskun EI, et al. Multifocal low-grade endometrial stromal sarcoma arising from pre-existing endometriosis in a hysterectomised patient: A case report. *J. Obstet. Gynaecol* 2019;39:1177–80.

28. Xu Y, Liang ZX, Guo JT, Su X, Lu YL, et al. Cystic and solitary nodular pulmonary metastases in a patient with low-grade endometrial stromal sarcoma: A case report and literature review. *Oncol Lett* 2019;18:1133–44.

29. Subbaraya S, Murthy SS, Devi GS. Immunohistochemical and molecular characterization of endometrial stromal sarcomas. *Clin Pathol* 2020;13:2632010X20916736.

30. Mutter GL, Prat J. Pathology of the Female Reproductive Tract. Churchill Livingstone Elsevier; 2014;3rd ed.

31. Chiang S, Lee CH, Stewart CJR, Oliva E, Hoang LN, et al. BCOR is a robust diagnostic immunohistochemical marker of genetically diverse high-grade endometrial stromal sarcoma, including tumors exhibiting variant morphology. *Mod Pathol* 2017;30:1251–61.

32. Stewart CJR, Bharat C, Crook M. p16 immunoreactivity in endometrial stromal cells: stromal p16 expression characterises but is not specific for endometrial polyps. *Pathol* 2015;47(2):112-7.

33. Bennett JA, Braga AC, Pinto A, Van de Vijver K, Cornejo K, et al. Uterine PEComas: A morphologic, immunohistochemical, and molecular analysis of 32 tumors. *Am J Surg Pathol* 2018;42(10):1370-83.

34. Conlon N, Soslow RA, Murali R. Perivascular epithelioid tumours (PEComas) of the gynaecological tract. *J Clin Pathol* 2015;68(6):418-26.

35. Flucke U, Vogels RJC, Somerhausen NSA, Creytens DH, Riedl GR, et al. Epithelioid Hemangioendothelioma: clinicopathologic, immunohistochemical, and molecular genetic analysis of 39 cases. *Diagn Pathol* 2014;9:131.