

Effects of humic acid and mineral Zn fertilizer on Zn uptake and maize yield in soils formed on limestone and marl parent materials

Mehmet Keçeci^{1*} , Gamze Depel¹ , Nuray Güneş¹ , Sadık Usta² , Veli Uygur³ ,
Muhittin Onur Akça² , Zeynep Demir¹ 

¹ Soil, Fertilizer and Water Resources Central Research Institute, 06172 Ankara, Türkiye

² Faculty of Agriculture, Department of Soil Science and Plant Nutrition, Ankara University, 06110 Ankara, Türkiye

³ Faculty of Agriculture, Department of Soil Science and Plant Nutrition, Isparta University of Applied Sciences, 32200 Isparta, Türkiye

How to cite

Keçeci, M., Depel, G., Güneş, N., Usta, S., Uygur, V., Akça, O. M., & Demir, Z. (2024). Effects of humic acid and mineral Zn fertilizer on Zn availability and maize yield in soils formed on limestone and marl parent materials. *Soil Studies* 13(1), 1-16. <http://doi.org/1021657/soilst.1520537>

Article History

Received 07 March 2024

Accepted 01 May 2024

First Online 23 July 2024

*Corresponding Author

Tel.: +90 312 315 65 60

E-mail:

mehmet.kececi@tarimorman.gov.tr

Keywords

Zn

Adsorption-Desorption

Marl soils

Lime soils

Maize

Abstract

In this study, the effects of zinc (Zn) and humic acid (HA) applications on soils (12 Aridisol soil samples) formed on limestone and marl parent materials, which are very common in Türkiye, were tested by incubation, adsorption, and greenhouse experiments. Adsorption experiments were carried out using the batch sorption technique, and the Langmuir and Freundlich isotherm parameters were calculated. In the incubation, soils were incubated with control, HA, Zn, and HA+Zn, and the DTPA-Zn was tested in five different periods (1, 7, 15, 30, and 90th days). Maize was grown for eight weeks in pots in the greenhouse. The maximum Zn adsorption for all soils ranged between 3333 and 6250 mg kg⁻¹ in marl soils and 1042 and 5263 mg kg⁻¹ in limestone soils, which decreased to ranges between 2631 and 5555 mg kg⁻¹ in marl soils and 1052 and 5000 mg kg⁻¹ in limestone soils with HA application, respectively. The desorption rate (%) of adsorbed Zn increased as the initial Zn concentration increased in all soil series formed on marl (smectite clay mineral) and limestone (kaolinite clay mineral) parent materials. The effects of humic acid and Zn application on the fresh and dry weights of maize were found to be statistically significant at the 5% level for marl and limestone parent material. Humic acid application increased the fresh weight of maize compared with that of the control. The highest available Zn was determined in Başkuyu series (0.79 mg kg⁻¹) formed on marl parent material, while the lowest available Zn was obtained in Saraççeşme series (0.60 mg kg⁻¹) formed on limestone parent material. In the greenhouse, HA increased the fresh weight of maize in soils formed on both parent materials by 0.4 and 19.6%, respectively, compared to the control. Zn fertilization with HA further increased the fresh and dry yields, with 12 mg kg⁻¹ performing better. S_{max} (maximum Zn adsorption of the soil) parameters of soils were negatively correlated with EC, P_{av}, K_{av}, organic matter, silt, clay, total N, and cation exchange capacity (CEC) of the samples. Although HA alone increases the availability of Zn in plants, HA + Zn applications should be applied together to obtain higher yields.

Introduction

The formation and composition of soils involve the combined effects of parent material, topography, vegetation, time, and climate (Weil and Brady, 2017; Yavitt, 2000). At a relatively large scale, climate and

parent material are the most effective soil-forming factors to gain a specific character in soils. However, hilly topography causes a significant difference in soil properties even on similar types of parent material

over a very short distance because of its control over water, heat, and vegetation. The other key soil-forming factor, parent material, regulates the speed of physical, chemical, and biological weathering processes in a specific environment ([Delgado and Gomez, 2016](#)). The parent material is also a primary source of nutrients necessary for plant growth and a habitat for soil microorganisms in natural soils ([Trettin et al., 1999](#)). The mineralogical composition of the parent material can influence the movement of fine soil particles and plant nutrients along the soil profile through water, weathering, growing vegetation types, and soil-dwelling organisms ([Jacobs, 1998](#)).

Marl is a rock containing clay-sized (<0.002 mm) clastic material and carbonate (calcite). Therefore, it is defined as a transitional rock material between clastic and chemical sedimentary rocks ([Šestanović, 2001](#)). Smectite, which predominates in the marl parent material classified as aridisol, is a phyllosilicate mineral characterized by a 2:1 (Si: Al) layer structure ([Brown, 1980](#); [Bailey et al., 2015](#)). Smectite clays have a variable net negative charge stabilized by the outward sorption of metal cations (Na, Ca, Mg, and H) on the interlamellar surfaces. The smectite structure has numerous unique properties, including chemical and exchangeable ion structures and small crystal sizes. Furthermore, the structure of smectite is responsible for its highly active surface area, high cation exchange capacity, rare hydration properties, and ability to strongly modify the movement behavior of solutions ([Mortland, 1970](#); [Sawhney and Singh, 1997](#)). Furthermore, Al³⁺-saturated clays have strong water polarity associated with Al³⁺, which leads to better adsorption by forming strong H-bonds with insecticides ([Sawhney and Singh, 1997](#)). [Johnston et al., \(2001\)](#) and [Sheng et al., \(2002\)](#) reported that some hydrated metal cations, such as Ca²⁺ and Mg²⁺, surrounded by water molecules reduce ion-dipole bonds, preventing direct contact between exchangeable cations and polar functional groups. Furthermore, nonpolar organic compounds can interact with the siloxane surfaces of smectite ([Laird and Fleming, 1999](#)). Saturated clays with weakly hydrated cations, such as NH⁴⁺ and K⁺, enhanced the adsorption of organic compounds by increasing the size of the adsorptive spheres between exchangeable cations. Marl parent materials have the most erodible surfaces in arid environments, including arid regions ([Cerdeira 2002](#); [Martinez-Mena et al., 2002](#)).

Limestone is a sedimentary rock that accounts for approximately 10% of the sedimentary rocks found on Earth's surface. It is formed by the deposition of shells and shell fragments, or by the direct crystallization of calcium carbonate from water ([Blatt and Tracy, 1996](#)). The remaining components may include other carbonate minerals such as dolomite magnesite and aragonite ([Jackson, 1997](#)). Kaolinite, which predominates in the limestone parent material, is a weathering product of feldspars. Because kaolinite is electrically stable, its ability to adsorb ions is lower than that of other clay

minerals. Kaolinite is a mineral commonly found in soil. It tends to be particularly abundant in weathered soils, such as Ultisols and Oxisols. Very few isomorphous substituents are present in the tetrahedral and/or octahedral layers of kaolinites. Most kaolinites were close to the ideal formula for Al₂Si₂O₅(OH)₄. The 1:1 layer had little or no permanent charge owing to the low amount of substitution. The cation exchange capacities and surface areas are typically low. Soils with a high kaolinite ratio are generally less fertile than soils dominated by 2:1 clay minerals ([Milliken, 2003](#)).

Illite is a closely related group of nonexpansive clay minerals. It is a secondary mineral precipitate and an example of phyllosilicates or layered aluminosilicates. Its structure is a 2:1 sandwich of silica tetrahedron (T)-alumina octahedron (O)-T layers ([USGS, 2019](#)). The formation of illite is generally favored under alkaline conditions and high Al and K concentrations. The cation exchange capacity (CEC) of illite is smaller than that of smectite, but higher than that of kaolinite, typically around 20-40 cmol kg⁻¹ ([Weil and Brady, 2017](#)).

Different parent materials influence the morphology and mineralogical and physicochemical properties of soils under the same conditions, such as the biosphere, topography, and climate, especially in arid and semi-arid regions. Therefore, differences in soil properties are primarily related to the parent material ([Washer and Collins, 1988](#)). Zinc (Zn) content in soil ranges from 10 to 300 mg kg⁻¹, depending on the chemistry of the parent material and soil texture, organic matter, and pH ([Mihaljevic, 1999](#)), with an estimated global average of 64 mg kg⁻¹ ([Kabata-Pendias, 2011](#)). Zn is the second most abundant transition metal in living organisms. Zn is critical for generative growth (seed formation) rather than vegetative growth. Zn may be important for pollen grain development, possibly by improving the phytohormone status of plants and protein synthesis ([Brown et al., 1993](#)).

Zinc deficiency in soils has been reported worldwide, especially in calcareous soils in arid and semiarid regions. Zinc deficiency in soils and crop plants occurs in almost all countries, especially in cereal-growing areas. Türkiye is among the countries with the most severely Zn-deficient soils. Since Zn deficiency is a major micronutrient deficiency in humans in Türkiye, increases in cereal Zn concentrations by Zn fertilization have significant impacts on human health. In Türkiye, Zn deficiency is a widespread problem in crops ([Çakmak et al. 1999](#)) and humans ([Çavdar et al., 1983](#)). Zinc concentrations in Turkish soils are some of the lowest ever recorded ([Sillanpää, 1990](#)). According to the results obtained from 1,511 soil samples using the Soil-DTPA test ([Lindsay and Norvell, 1978](#)), 49.8% of the cultivated soils in Türkiye were classified as Zn-deficient ([Eyupoglu et al., 1994](#)). The prevalence of Zn deficiency in Turkish soils increased with increasing pH and decreasing organic matter levels in soils collected from different regions. Soils in different regions of Türkiye

are generally deficient in Zn, which is reflected in the poor growth of plants under greenhouse conditions (Kacar, 1998). Zinc deficiency is common in plants growing in highly weathered acidic and calcareous soils. In the latter case, Zn deficiency was often associated with Fe deficiency (lime chlorosis). The low availability of Zn in calcareous soils with high pH is due to the adsorption of Zn to clay or CaCO_3 rather than the formation of poorly soluble $\text{Zn}(\text{OH})_2$ or ZnCO_3 (Trehan and Sekhon, 1977). In addition, Zn uptake and translocation to the shoot are inhibited by high bicarbonate and HCO_3^- concentrations (Dogar and van Hai, 1980).

The organic matter (OM) content controls the magnitude of Zn sorption in soil (Montalvo et al., 2016). Organic matter is composed of different fractions such as humic acid (HA) and fulvic acid. Therefore, the surface of organic colloids is dominated by negative charges at low soil pH (Stevenson, 1994). As the OM and soil pH increased, the amount of organic ligands and their charge density increased. It also increases Zn adsorption because of the increased stability of the organic complexes with HA as the pH increases (Boguta and Sokołowska, 2016). Humic acid, a pool of humic substances, tends to form complexes with zinc. However, the stability and solubility of the formed complexes depend on the Zn-HA stoichiometry, plant species, proportion of HA added to the soil, growth medium characteristics, and soil type (Olaetxea et al., 2018). Therefore, depending on the combination of these factors and soil management practices, crop Zn uptake can be improved or reduced (de Morais et al., 2021). Humic substances improve nutrient uptake and plant growth by adding HA and Zn to the soil (Justi, 2019). The effects of humic substances on plant tissues include stimulation of root proliferation, enhancement of H^+ -ATPase activity in the cell membrane, and changes in the magnitude of nutrient uptake, assimilation, and utilization efficiency (Aguiar et al., 2013; Nardi et al., 2021). HA in soil can improve Zn nutrition (Boguta and Sokołowska, 2016) and plant growth by forming a Zn complex (De Morais et al., 2021). Organometallic complexes increase Zn content in the soil solution and Zn diffusion from the solution to the cell root surface. This is related to the complexation of Zn by organic ligands and the improvement in Zn uptake by plants (Justi et al., 2019). Dissolved Zn in the soil liquid phase is readily available to plants as the Zn concentration in solution contains a small fraction of the total Zn content of the soil (Montalvo et al., 2016). Therefore, HA is expected to contain organic radicals capable of complexing Zn into soluble forms. The use of HA effectively increased the Zn content in the solution, especially at a high soil pH (7.2) (Justi et al., 2019). HA can also buffer soil acidity. It maintains the pH within the optimum range required to provide crops with abundant nutrients, especially Zn, whose availability decreases as pH increases (Montalvo et al., 2016).

The buffering effect of HA on soil acidity is more frequent. When pH is in the alkaline range, less dis-

solved Zn is present in the soil, while low insoluble forms (ZnOH^+ and ZnOH_2) are formed in the soil (Alloway, 2009; Montalvo et al., 2016). HA buffers soil acidity and prevents oscillations in soil reaction (Aguiar et al., 2013; Rosa et al., 2018), and consequently promotes the formation of soluble Zn^{2+} free hydrated forms and soluble Zn-organic complexes (Montalvo et al., 2016). HA also depend on the structure-organic functional groups-function properties of HA (Nardi et al., 2021). It is linked to the type and concentration of oxygen, nitrogen and sulphur-containing functional groups in the HA structure (Aguiar et al., 2013; Zanin et al., 2019; Nardi et al., 2021; de Morais et al., 2021).

Materials and Methods

Materials

The taxonomy of the experimental soils

The characteristics of the experimental soils formed on marl (6 soils) and limestone parent materials belonging to the Aridisol order in the U.S. Soil Taxonomy are given in Table 1. Marl soils are classified as Xeric Haplocambic, Xeric Pedrocalcic and Typic Calciorthid and limestone soils are Vertic Calciorthid, Typic Paleorthisid and Typic Calciorthisid. The name of marl soils are Atınova, Hacifakıl, Çatalca, Başkuyu, Kuyubaşı and Seyrantepe soil series and Limestone's soils are Saraççeşme, Tine, Karataş-1, Tuvem-1, Tuvem-1, Tuvem-2 and Karataş-2 soil series. The soil series is characterized by arid and semiarid climate.

Some physical and chemical properties of soils

Composite surface soil samples (0-20 cm depth) were brought to laboratory in plastic bags. Air-dried samples were then passed through 2 mm sieve to performe the following analysis: particle size distribution (sand, silt and clay contents) by a hydrometer method (Bouyoucus, 1951), pH and electrical conductivity (EC) in saturation paste (Richard, 1954) calcium carbonate equivalent (CCE) by a manametic method using the Scheibler calcimeter (Soil Survey Staff, 2014) organic matter by modified Walkley-Black method (Sparks, 1996), cation exchange capacity (CEC) by molar Na-Acetate saturation (Richard 1954), total nitrogen (N) by conctional Kjeldahl method (Bremner 1965), DTPA-extractable cationic microelements (Fe, Cu, Mn, Zn) (Lindsay and Norwel, 1978) including certain heavy metals (Ni, Cr, Pb and Cd) by ICP-OES (Perkin-Elmer, DV, 2100).

The physical and chemical properties of the experimental soils were presented in Table 2. The limestone soils had finer texture than the marl soils.

Electrical conductivity values ranged similarly in both parent materials. Soil organic matter contents were higher with limestone soils (1.61 - 2.69%) than the marl soils' (0.89 - 1.69%). The CCE contents of marl

Table 1. The soil taxonomy classification of the experimental soils

Soils	Soil Family	Soil Series	Parent material	Order
S1	Xeric Haplocambic	Atınova		
S2	Xeric Haplocambic	Hacıfakıl		
S3	Xeric Haplocalcid	Çatalca	Marl	
S4	Xeric Haplocalcid	Başkuyu		
S5	Xeric Pedocalcid	Kuyubaşı		
S6	Typit Calciorthid	Seyrantepe		Aridisol
S7	Vertic Calciorthid	Saraççeşme		
S8	Typit Paleorthid	Tine		
S9	Typit Calciorthid	Karataş-1	Limestone	
S10	Vertic Calciorthid	Tüvem-1		
S11	Vertic Calciorthid	Tüvem-2		
S12	Typit Calciorthid	Karataş-2		

soils (27.41 - 43.40%) were higher than the limestone soils' (5.28 - 24.61%). The total nutrient element concentrations of soils were given in Table 3. The element concentrations were highly affected from the parent materials depending on the overall effects of soil form-

ing factors and agricultural practices.

Properties of humic acid (HA)

Some chemical properties of HA used in the experiment are given in Table 4. Organic matter content

Table 2. Some chemical and physical properties of the experimental soils

Soil	Sand %	Silt %	Clay %	Texture Class	OM %	CEC cmol kg ⁻¹	pH	CCE %	EC dS m ⁻¹	P _{av} kg ha ⁻¹	K _{av} kg ha ⁻¹
S1	43.0	24.0	33.0	CL	1.28	24.2	7.54	27.4	0.782	7.84	134
S2	43.0	26.7	30.3	CL	0.89	23.9	7.50	35.6	0.716	4.66	96
S3	41.0	34.2	24.9	L	0.79	19.3	7.69	35.0	0.619	11.9	146
S4	39.0	33.5	27.6	CL	1.69	23.6	7.56	35.1	0.721	0.93	106
S5	42.2	31.5	26.3	L	1.50	20.9	7.49	43.4	0.656	4.27	158
S6	45.0	31.5	23.6	L	1.44	19.3	7.72	39.5	0.499	5.36	102
S7	26.2	38.2	35.6	CL	1.97	25.6	7.51	38.7	0.717	5.74	158
S8	29.0	35.5	35.6	CL	1.61	28.4	7.61	9.01	0.358	10.5	134
S9	23.0	38.0	39.0	CL	2.69	29.2	7.24	7.47	0.855	24.6	201
S10	42.2	26.2	31.6	CL	2.20	25.7	7.76	16.3	0.506	10.3	154
S11	26.6	37.8	35.6	CL	2.10	24.9	7.53	20.1	0.691	5.59	179
S12	40.2	32.2	27.6	CL	2.27	24.8	7.49	5.28	0.594	4.35	206

S1: Atınova soil series, S2: Hacıfakıl soil series, S3: Çatalca soil series, S4: Başkuyu soil series, S5: Kuyubaşı soil series, S6: Seyrantepe soil series, S7: Saraççeşme soil series, S8: Tine soil series, S9: Karataş-1 soil series, S10: Tüvem-1 soil series, S11: Tüvem-2 soil series, S12: Karataş-2 soil series, OM: Organic matter, CEC: Cation exchange capacity, pH: Soil reaction, CCE: Calcium carbonate equivalent, EC: Electrical conductivity, P_{av}: Available P₂O₅, K_{av}: Available K₂O, C: Clay, L: Loam

Table 3. Total concentration of some plant nutrients in the experimental soils

Soil Series	g kg ⁻¹					mg kg ⁻¹	
	P	Mn	N	K	Fe	Cu	Zn
S1	0.42	0.51	0.64	18.07	16.85	18.4	32.2
S2	0.43	0.95	0.45	15.04	17.94	30.7	83.1
S3	0.62	0.76	0.40	17.96	10.64	20.6	52.7
S4	0.69	0.97	0.85	11.75	12.93	35.1	96.1
S5	0.72	0.82	0.75	15.73	22.34	29.1	52.6
S6	0.52	0.9	0.72	1.211	18.06	29.8	97.0
S7	0.51	0.96	0.99	12.38	14.04	34.3	88.7
S8	0.55	0.79	0.81	10.36	12.57	22.8	66.1
S9	0.11	0.56	1.35	9.05	10.20	12.0	28.0
S10	0.17	0.91	1.10	14.72	10.20	26.4	78.7
S11	0.52	0.82	1.05	11.43	19.67	35.4	58.9
S12	0.24	0.80	1.14	8.42	10.21	29.3	68.0

S1: Atınova soil series, S2: Hacırakıl soil series, S3: Çatalca soil series, S4: Başkuyu soil series, S5: Kuyubaşı soil series, S6: Seyrantepe soil series, S7: Saraççeşme soil series, S8: Tine soil series, S9: Karataş-1 soil series, S10: Tüvem-1 soil series, S11: Tüvem-2 soil series, S12: Karataş-2 soil series

was 43.3%, total humic acid value was 20.8% and electrical conductivity was 0.27 dS m⁻¹. The HA was highly acidic with a pH of 3.3 and its heavy metal contents were at trace amounts below the detection limits of ICP-OES.

Table 4. Chemical properties of the humic acid

Parameters	Results
Organic matter (%)	43.3
Σ Humic acid (%)	20.8
Σ Nitrogen (%)	0.17
pH	3.3
Electrical conductivity (dS m ⁻¹)	0.27
Σ Phosphorus (%)	1.71
Σ Potassium (mg kg ⁻¹)	395
Σ Calcium (mg kg ⁻¹)	83.5
Σ Magnesium (mg kg ⁻¹)	2125
Σ Iron (mg kg ⁻¹)	3.12
Σ Zinc (mg kg ⁻¹)	1.87
Σ Manganese (mg kg ⁻¹)	<0.02
Σ Nickel (mg kg ⁻¹)	<0.02
Σ Chromium (mg kg ⁻¹)	<0.02
Σ Lead (mg kg ⁻¹)	<0.02
Σ Cadmium (mg kg ⁻¹)	<0.02

Method

Adsorption and desorption experiments

Triplicate scoops of 2 g of the soils were equilibrated with 25 mL of Zn solutions containing of 0, 6.5 - 32.5 - 65 - 162.5 - 325 - 487.5 - 650 mg L⁻¹ prepared in 0.01 M Ca(NO₃)₂ background solution to keep ionic strength constant. The suspensions were shaken for 24 h at 25°C on an end-over-end shaker. Then the solution phase were separated by sequential centrifugation at 3000 rpm for 10 min and filtering through Whatman 2 filter paper. Immediately after adsorption, the desorption batches were performed by equilibrating the Zn-loaded-soils with 20 mL of 0.01 Ca(NO₃)₂ solutions for 24 h at 25°C. Then, the Zn concentration of the supernatants obtained from the adsorption and desorption batches were determined by ICP-OES. The amounts of adsorbed and desorbed Zn was calculated from the difference in the equilibrium and initial Zn concentrations (Alumaa, 2001). The sorption data were then tested for the conformation of linear Langmuir and Freundlich isotherms as given below:

$$\frac{C_e}{S} = \frac{1}{k S_{max}} + \frac{1}{S_{max}}$$

Where S amount of Zn adsorbed (mg kg⁻¹), C_e Zn concentration of the equilibrium solution (mg L⁻¹), S_{max} maximum Zn adsorption of the soil (mg kg⁻¹), k bonding energy coefficient (mg mL⁻¹).

The exponential forms to the Freundlich model is:

$$S = K_F C_e^{1/n}$$

Where K_F Freundlich constant and n constant.

In order to reduce the deviation from the Langmuir model, the total amounts of the Zn was included into the S . This improved the determination coefficient ($R^2 \geq 0.948$) of the regression line without omitting any sorption data point at the lower-end.

Clay mineralogy

In order to carry out clay analyses in soils, carbonates, oxide minerals and organic matter were sequentially removed by the following methods: carbonates by 1 N acetate buffer (pH. 4.5-5), organic matter by 30% H_2O_2 on a hot plate, sesquioxides by citrate-bicarbonate-dithionite on a hot plate at 80°C.

The X-ray diffractograms of the soil samples were obtained to determine clay types (Whittig and Allardice, 1986). The semi-quantitative percentages of the minerals were calculated by the ratio of the individual peaks to total peak area. The results were presented in Table 5. The content of smectite clay minerals varied between 37.18% and 29.75%, kaolinite clay minerals between 59.08% and 43.87% and illite clay minerals between 18.95-11.17%. The dominance of the clays in the soils were smectite 42.86 %, kaolinite 35.71% and illite 21.43%. The smectite clays were dominant in the limestone soils whereas kaoline was dominant in the marl soils. Illite however were present at equal amounts in soils developed on both parent materials.

Table 5. Semi-quantitative clay contents (%) of the experimental soils

Soil	Dominance	Smectite	Illite	Kaolinite
S1	Smectite > Kaolinite > Illite	41.6	21.9	36.5
S2	Smectite > Kaolinite > Illite	45.4	11.5	43.2
S3	Smectite > Kaolinite > Illite	58.2	4.2	37.6
S4	Smectite > Kaolinite > Illite	55.2	8.2	36.6
S5	Kaolinite > Smectite > Illite	31.6	12.4	55.9
S6	Kaolinite > Illite > Smectite	16.2	32.7	51.0
S7	Kaolinite > Illite > Smectite	5.33	44.1	50.6
S8	Kaolinite > Illite > Smectite	20.4	20.9	58.6
S9	Kaolinite > Smectite > Illite	31.8	3.53	64.7
S10	Smectite > Kaolinite > Illite	50.6	16.4	32.9
S11	Kaolinite > Smectite > Illite	29.7	11.2	59.1
S12	Kaolinite > Smectite > Illite	37.2	18.9	43.9

S1: Atınova soil series, S2: Hacıfakıl soil series, S3: Çatalca soil series, S4: Başkuyu soil series, S5: Kuyubaşı soil series, S6: Seyrantepe soil series, S7: Saraççeşme soil series, S8: Tine soil series, S9: Karataş-1 soil series, S10: Tüvem-1n soil series, S11: Tüvem-2 soil series, S12: Karataş-2 soil series

Incubation experiments

According to the differences in adsorption isotherms (lime, composition, etc.), four soils (S2 and S4 soil series on marl parent material and S7 and S8 soil series on limestone parent material) were selected to represent both marl and limestone parent materials. These soils were used to test time-dependent effects of HA acid treatments on the DTPA extractability of Zn over 90 days. The experiment consisted of HA, Zn chemical fertiliser ($ZnSO_4$), humic acid+Zn chemical fertiliser, and the control (without HA and Zn) treatments in triplicates. The experiment was set up in completely-randomised design. The soil were incubated at constant moisture content in the field capacity and $25 \pm 2^\circ C$. Soil samples were taken on the 1, 7, 15, 30 and 90th days of incubation and analysed for DTPA-Zn (Lindsay and Norwel, 1978).

Greenhouse experiments

A representative soil samples were selected from each parent material by taking into account the results of adsorption/desorption and incubation experiments. The greenhouse experiment was conducted in triplicates with two soils (Baskuyu soil series (S4) on marl parent material and Tine soil series (S8) on limestone parent material), two doses of humic acid (0% and 2%) and 5 levels of $ZnSO_4$ (0 - 4 - 8 - 12 - 16 mg Zn kg^{-1}). Soil samples were passed through a 5-mesh sieve and 2 kg soil was filled into plastic pots on dry weight basis. Plant nutrient solutions were prepared and sprayed on the soil and mixed thoroughly to maintain homogeneity. Maize seeds (*Zea mays* var. *indentata*) were sown after soaking 24 hours before sowing. The plant were irrigated in field capacity. At the end of the vegetation period of 8 weeks, they were harvested just above the soil and their heights and fresh weights were determined and then washed with distilled water and dried

at 65°C to a constant weight and dry weights were determined. Plant samples were then ground and homogenized to a particle size of ≤ 0.5 mm and prepared for analysis.

Plant analysis

Scoops of 0.5 g plant samples were digested in a microwave oven in $\text{HNO}_3/\text{H}_2\text{O}_2$ mixture (8/2, v/v). Then the plant nutrient concentration of the digests were analysed for plant nutrients by means of ICP-OES. The total nitrogen was determined by the conventional Kjeldahl method.

Statistical analysis

Data were analyzed using SPSS 19.0 statistical software. The differences between the treatment

means were tested using the Duncan's multiple range test. The underlying coherence between the investigated parameters were revealed by Pearson's correlation (SPSS 19.0, SPSS Inc. 2011). The incubation data were subjected to repeated measure analysis procedures to simultaneously test the between (treatment effects) and within effects (the time-effect) and the mean separation was performed by Bonferroni test at $p \leq 0.05$.

Results

Zn adsorption isotherms

Adsorption isotherms of 12 soil samples without and with humic acid application are given in Figure 1. Zn adsorption data were characterized at acceptable




Figure 1. The adsorption isotherms of the experimental soils with and without humic acid treatment. Soils numbered 1-6 are marl soils and 7-12 are limestone soils

levels with both isotherms. The adsorption maximum values of Langmuir isotherm (b) without and with humic acid application ranged between 1042-6250 mg kg⁻¹ and 1052-5555 mg kg⁻¹, respectively. The maximum adsorption values were significantly reduced by the HA treatment. This was apparent at higher Zn loadings (Figure 1). The binding energy (k) varied between 0.061-0.137 and 0.056-0.267 in soils without and with humic acid application, respectively. In the Freundlich isotherms, the adsorption power (n) varied between 0.171-0.231 and 0.225-0.384 in soils without and with humic acid treatment, respectively. K_f values varied between 326-1564 and 550-2019 in soils without and with humic acid application, respectively. The regression coefficients for the Langmuir isotherm without and with humic acid application were R²=0.958-0.997 and R²=0.990-0.999, while the regression coefficients for the Freundlich isotherm were R²=0.472-0.968 and R²=0.653-0.968, respectively.

Zn desorption isotherms

In general, the amount of Zn desorbed increased

with increasing initial Zn concentration and HA treatment (Table 6). In the non-HA treatments in the marl parent material in the Kuyubaşı series Xeric Pedrocalcic soil (S5), 16.9% of the highest amount of Zn adsorbed was desorbed and this rate increased to 25.6% with humic acid application. In general, the percentages of desorbed Zn remained constant after 325 mg L⁻¹ Zn treatment. At the 650 mg L⁻¹ initial concentration, 22.2% of the Zn amount was desorbed in the limestone parent material in the Saraççeşme series Vertic Calcic soil (S7), and this rate increased to 28.7% with the application of humic acid.

Incubation period

During incubation, Zn contents of the soil samples were taken on 1, 7, 15, 30 and 90 days. In the Zn incubation experiment, the difference between soils (formed on marl and limestone parent materials) and treatments (humic acid, chemical fertilizer, humic acid + chemical fertilizer application and the control) was statistically significant at 1% level (Table 7). Soil and treatment interaction was found to be significant at 1% level. The

Table 6. Desorption ratio (%) of the adsorbed Zn as a function of the initial Zn concentrations, humic acid treatments and parent material

		S1		S2		S3		S4		S5		S6	
	Init. Zn (mg L ⁻¹)	Humic acid treatments											
		(-)	(+)	(-)	(+)	(-)	(+)	(-)	(+)	(-)	(+)	(-)	(+)
Marl soils	6.5	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	32.5	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	65	0.34	0.63	0.91	1.10	1.94	4.67	1.31	2.40	2.00	2.27	2.22	2.42
	162.5	2.19	3.48	3.73	4.63	2.93	5.51	4.80	6.61	5.29	7.19	6.07	6.62
	325	3.76	6.57	6.82	7.96	6.96	10.1	7.26	10.2	9.88	12.1	10.1	11.0
	487.5	3.05	10.6	7.20	9.77	7.49	12.2	9.22	14.9	14.2	18.6	16.0	17.6
	650	3.18	10.6	8.49	13.5	10.3	19.4	9.54	17.2	16.9	25.6	16.1	17.6
		S7		S8		S9		S10		S11		S12	
	Init. Zn (mg L ⁻¹)	Humic acid treatments											
		(-)	(+)	(-)	(+)	(-)	(+)	(-)	(+)	(-)	(+)	(-)	(+)
Limestone soils	6.5	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	32.5	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	65	2.02	2.22	0.90	0.98	1.51	1.65	1.59	1.75	1.24	1.74	1.59	1.37
	162.5	5.48	6.68	3.78	5.60	5.53	6.33	4.43	5.07	4.37	6.96	5.24	5.03
	325	12.8	14.6	7.69	11.2	7.81	8.13	7.39	9.59	5.69	11.0	8.48	10.1
	487.5	20.5	26.1	11.0	14.2	15.9	18.1	11.6	15.1	10.6	15.3	13.0	15.6
	650	22.2	28.7	10.2	16.5	18.0	22.6	13.3	19.2	10.6	18.9	14.7	19.4

S1: Atınova soil series, S2: Hacıfakıl soil series, S3: Çatalca soil series, S4: Başkuyu soil series, S5: Kuyubaşı soil series, S6: Seyrantepe soil series, S7: Saraççeşme soil series, S8: Tine soil series, S9: Karataş-1 soil series, S10: Tüvem-1 soil series, S11: Tüvem-2 soil series, S12: Karataş-2 soil series

Table 7. Time-dependent DTPA-Zn concentrations (mg kg^{-1}) of the experimental

Soils	Treatments	Incubation periods (Days)					Treat.mean	Soil mean	Grand mean
		1	7	15	30	90			
S4	WS	0.08	0.09	0.19	0.09	0.12	0.12 b	0.79 A	0.68
	WS + Zn	1.17	1.51	1.30	1.58	1.13	1.34 a		
	WS + HA	0.08	0.11	1.57	0.08	0.14	0.40 b		
	WS + Zn + HA	1.17	1.42	1.27	1.49	1.20	1.31 a		
	Incubation mean	0.63	0.78	1.08	0.81	0.65			
S2	WS	0.08	0.15	0.14	0.08	0.12	0.12 b	0.70 B	
	WS + Zn	1.23	1.52	1.33	1.20	0.28	1.11 a		
	WS + HA	0.07	1.13	0.13	0.07	0.12	0.30 b		
	WS + Zn + HA	1.15	1.46	1.33	1.19	1.26	1.28 a		
	Incubation mean	0.63	1.07	0.73	0.64	0.44			
S7	WS	0.06	0.08	0.15	0.07	0.11	0.10 b	0.60 C	
	WS + Zn	1.05	1.55	1.19	1.19	1.16	1.23 a		
	WS + HA	0.06	0.10	0.13	0.06	0.10	0.09 b		
	WS + Zn + HA	1.04	1.41	1.17	0.17	1.15	0.99 a		
	Incubation mean	0.55	0.79	0.66	0.37	0.63			
S8	WS	0.06	0.08	0.19	0.06	0.12	0.10 b	0.62 C	
	WS + Zn	1.06	1.31	1.13	1.11	1.10	1.14 a		
	WS + HA	0.05	0.08	0.12	0.06	0.11	0.08 b		
	WS + Zn + HA	1.08	1.30	1.13	1.10	1.08	1.14 a		
	Incubation mean	0.56	0.69	0.64	0.58	0.60			
Grand mean of incubation periods		0.59	0.83	0.78	0.60	0.58			
		b	a	a	b	b			
F_i	12.028						P_i	0.000**	
F_t	375.859						P_t	0.000**	
F_{pm}	8.128						P_{pm}	0.000**	
$F_{t \times pm}$	2.293						$P_{t \times pm}$	0.018*	

S4: Başkuyu soil series, S2: Hacifakıl soil series, S7: Saraççeşme soil series, S8: Tine soil series, WS: Whole soil, S7: Saraççeşme soil series, i: Incubation periods, t: treatments, pm: parent material

effects of Zn on soils formed on marl and limestone parent materials due to humic acid application were insignificant. Humic acid applications increased the available Zn content compared to the control in the soil formed on marl parent material. The highest available Zn content was determined in Başkuyu series (0.79 mg kg^{-1}) formed on marl parent material, while the lowest DTPA-Zn content was obtained in Saraççeşme series (0.60 mg kg^{-1}) formed on limestone parent material. The highest Zn content of the soils (1.34 mg kg^{-1}) was increased by humic acid application in Başkuyu series formed on marl parent material, while the lowest available Zn content (0.08 mg kg^{-1}) was determined in chemical fertilizer application in Tine series formed on limestone parent material. Humic acid and humic acid +

chemical fertilizer applications statistically affected at the same level the available Zn content of the soils. In addition, control and only chemical fertilizer applications had statistically increased the available Zn content of soils.

Greenhouse Experiment

Maize performance on the marl soil

The effect of humic acid and Zn applications on the fresh and dry weights of maize was found to be statistically significant at the 5% level (Table 8). Humic acid applications (HA 2%) increased the fresh weight of maize compared to the control (HA 0%). The lowest fresh and dry weights of maize for marl parent material

were 58.0 and 24.3 g for the S4, 65.5 and 26.4 g for the 2% HA obtained from the control treatment, whereas the highest fresh and dry weights of maize were 82.2 and 34.9 g for the marl parent material (S4), 86.3 and 35.7 g for the 2% HA obtained from the 2% HA + 12 mg kg⁻¹ Zn treatment, respectively.

Maize performance on the limestone soil

The effect of humic acid and Zn applications on the fresh and dry weights of maize was statistically significant ($p \leq 5\%$) (Table 8). Humic acid applications increased the fresh weight of maize compared to the control. The lowest fresh and dry weights were 65.8 and 24.2 g for the limestone parent material soil (S8),

78.7 and 26.2 g for the 2% HA obtained from the control treatment; whereas the highest fresh and dry weights were 86.0 and 34.6 g for the the limestone parent material soil (S8), 84.8 and 35.0 g for the 2% HA obtained from the 2% HA + 12 mg kg⁻¹ Zn treatment, respectively.

Discussion

In soil, Zn availability is controlled by the intensity of adsorption and precipitation reactions. Zn sorption increases as the soil pH are higher due to the rise of negative charges on soil colloid surfaces and the incre-

Table 8. The effects of humic acid (HA) and Zn treatments on the fresh and dry biomasses of maize grown on the marl (S4) and limestone soil (S8) in the greenhouse experiment

Zn treatments, mg kg ⁻¹	Marl parent material (S4)				Limestone parent material (S8)			
	Fresh biomass, g pot ⁻¹ HA (0%)	Fresh biomass, g pot ⁻¹ HA (2%)	Dry biomass, g pot ⁻¹ HA (0%)	Dry biomass, g pot ⁻¹ HA (2%)	Fresh biomass, g pot ⁻¹ HA (0%)	Fresh biomass, g pot ⁻¹ HA (2%)	Dry biomass, g pot ⁻¹ HA (0%)	Dry biomass, g pot ⁻¹ HA (2%)
0	58.0 e	65.5 d	24.3 d	26.4 e	65.8 d	78.7 b	24.2 d	26.2 e
4	71.2 d	74.3 c	29.2 c	29.0 d	82.2 b	83.5 a	28.4 c	28.6 d
8	78.3 b	77.5 b	32.2 b	32.1 c	82.5 b	84.7 a	31.7 b	31.9 c
12	82.2 a	86.3 a	34.9 a	35.7 a	86.0 a	84.8 a	34.6 a	35.0 a
16	75.3 c	79.5 b	33.0 b	34.1 b	75.2 c	75.5 c	31.8 b	33.5 b

Different letter in the same column indicate a significant difference at $p \leq 0.05$

ased presence of adsorbed and precipitated Zn forms in soil (Montalvo et al., 2016). Langmuir and Freundlich adsorption models were applied in Zn adsorption study, and adsorption parameters were determined from these models (Table 9). Langmuir and Freundlich adsorption isotherms of 12 soil samples without and with humic acid application were determined. In these soils, adsorption data for applications are described at acceptable levels with both isotherms. The adsorption data of all the soils were better described by the linear Langmuir model compared with the Freundlich model. For all soils, the Langmuir isotherm showed significant results both in the control soils without humic acid and in the soils with 2% humic acid. Freundlich isotherms showed significant results only in soils with humic acid. Regarding the physical and physicochemical properties of the soil, texture (clay content), organic matter and pH were the factors that sharply influenced soil Zn availability. Organic matter content controls the magnitude of Zn sorption in soil (Montalvo et al., 2016). Organic matter is composed of different fractions, including humic acid and fulvic acids, which are organic matter pools characterized by low isoelectric points; thus, even at low soil pH levels, negative charges prevail on the surface of organic colloids (Stevenson, 1994). Soil properties can modulate Zn interaction, adsorption and precipitation with soil components, including organic matter pools (Broadley et al., 2012; Alloway, 2009;

Montalvo et al., 2016). When associated with organic ligands, Zn availability for crops depends on the chemical stability, solubility and reactivity of the bonds formed between the humic matrix and the metal in the complexes formed (Boguta and Sokółowska, 2016; Rose et al., 2014).

In this study, soils formed on marl and limestone parent material showed differences in the desorption rates of adsorbed Zn at initial Zn concentrations (6.5-32.5-65-162.5-325-487.5-650 mg L⁻¹). These differences in soil properties (Table 2 and 3) may have caused differences in the effects of humic acid application on sorption isotherm parameters. As a function of initial concentrations, the soil scavenge the Zn by variety of sorption/precipitation reactions that each has different irreversibility and bounding strength. The desorption rate (%) of adsorbed Zn increased as the initial Zn concentration increased in the all soil series formed on marl (smectite clay mineral) and limestone (kaolinite clay mineral) parent material (Table 6). An increase in clay content is a key factor to increase Zn adsorption in soils. However, the density of negative charges in tropical clay minerals and Al and Fe oxides is lower than those in organic colloids (Alloway, 2009; Broadley et al., 2012; Montalvo et al., 2016). Moreover, Zn availability to plants is controlled by soil type, minerals associated with clay and Fe and Al oxides, soil parent material, total Zn content, soil pH, concentrations of organic

Table 9. The effects of humic acid treatment on the sorption isotherm parameters

Soil	Treatment	Langmuir isotherm			Freundlich isotherm		
		S _{max} (mg kg ⁻¹)	k	R ²	K _F (mg kg ⁻¹)	1/n	R ²
S1	HA	5555	0.069	0.991	674	0.384	0.898
	WS	5882	0.090	0.990	1482	0.231	0.886
S2	HA	5263	0.066	0.990	769	0.342	0.901
	WS	6250	0.065	0.969	1564	0.222	0.905
S3	HA	4347	0.116	0.998	804	0.304	0.873
	WS	5000	0.087	0.991	1383	0.213	0.896
S4	HA	3846	0.090	0.992	883	0.256	0.890
	WS	5000	0.064	0.973	1414	0.196	0.902
S5	HA	2631	0.060	0.995	550	0.258	0.816
	WS	3333	0.061	0.993	1077	0.173	0.845
S6	HA	2857	0.267	0.999	845	0.225	0.842
	WS	3704	0.083	0.988	1259	0.174	0.886
S7	HA	3030	0.095	0.993	790	0.233	0.853
	WS	3571	0.075	0.979	1213	0.171	0.875
S8	HA	4348	0.106	0.993	795	0.311	0.892
	WS	5263	0.072	0.959	1469	0.210	0.901
S9	HA	1052	0.059	0.979	2019	0.264	0.935
	WS	1042	0.069	0.976	326	0.183	0.947
S10	HA	5000	0.056	0.970	766	0.326	0.880
	WS	5000	0.071	0.948	1249	0.227	0.872
S11	HA	4167	0.084	0.976	832	0.282	0.871
	WS	4762	0.071	0.976	1336	0.202	0.890
S12	HA	4000	0.153	0.999	840	0.283	0.888
	WS	4545	0.137	0.997	1386	0.203	0.895

S1: Atınova soil series, S2: Hacifakıl soil series, S3: Çatalca soil series, S4: Başkuyu soil series, S5: Kuyubaşı soil series, S6: Seyrantepe soil series, S7: Saraççeşme soil series, S8: Tine soil series, S9: Karataş-1 soil series, S10: Tüvem-1 soil series, S11: Tüvem-2 soil series, S12: Karataş-2 soil series, HA: Humic acid, WS: Whole soil, S_{max}: Maximum Zn adsorption of the soil, k: Bonding energy coefficient, K_F: Freundlich K, n: Constant

matter, Ca, calcite, bicarbonate, and phosphate found in the soil, solution or in labile forms prone to be solubilized and to react with zinc with subsequent formation of precipitates and high-stability organometallic complexes (Alloway, 2009; Boguta and Sokołowska, 2016; Broadley et al., 2012; Montalvo et al., 2016). Smectite is one of the largest and most important classes of the phyllosilicate clay-mineral group. They are common in temperate soils and tend to dominate the cationic adsorption chemistry of these soils due to their cation exchange capacity and very high specific surface (Klopogge and Frost, 1999). Desorption is an important property that determines the mobility of adsorbed elements or metals in contaminated areas. The amount of desorption varies depending on the nature of the adsorption event, i.e. whether the binding is by physical, electrostatic bonds or covalent bonds. Physically adsorbed cations are usually capable of being readily replaced or desorbed by other cations present in the soil solution. On the other hand, ions or molecules bound by covalent bonds are not easily desorbed because they bind to the adsorbent with high energy and usually released back to the solution phase with

the complete breakdown of the adsorbent. In soil 9, none of the Zn absorbed at the lowest concentration was desorbed. At the 650 mg L⁻¹ initial concentration, 18.0% of the Zn amount was desorbed in the limestone parent material in the Karataş-1 series Typit Calciorthid soil family (S9) in the Aridisol and this rate increased to 22.6% with the application of humic acid. The amount of Zn that could be desorbed or replaced by Ca was calculated from the adsorbed Zn at each initial concentration applied. Accordingly, at the lowest concentrations applied, almost none of the absorbed Zn was desorbed. The amount of absorbed Zn was almost not desorbed at the lowest concentrations applied (6.5 - 32.5 - 65 mg L⁻¹). This shows that especially Zn adsorbs with a high energy and is retained with a high energy even at high concentrations. The other point is that the adsorption or precipitation mechanisms of Zn cause the release of 2 H⁺ ion on a stoichiometric base (Uygur and Rimmer, 2000). This reduces the pH of equilibrium solution below 6.5, in turn, the physisorption processes take place that increase the desorption rate. In addition, humic acid application showed that desorption was higher than the application without humic acid

because of chelation reaction between the HA and Zn and possible blockage of specific sorption site on the colloidal surfaces (Spark et al., 1997). This suggests that Zn may be highly immobilized in these soils, especially at high concentrations. The fact that Zn is not desorbed at low concentrations indicates that Zn deficiency is a key fact in soils with poor organic matter content and high pH under insufficient fertilization practices, to limit plant performance. The release of trace elements over time (desorption rate) is important for plant nutrition. In general, desorption of soils increased Zn mobility due to humic acid application.

Bivariate Pearson correlations coefficients between the soil properties and the sorption parameters were given in Table 10. Smax parameter was negatively correlated with soil EC, P_{av} , K_{av} , OM, silt, clay, total N and CEC. The properties negatively correlated with Smax indicates that the property function a reduced sorption site, co-input of Zn by previously applied agricultural practices and chelate with it. The parameters such as pH, CCE, sand, total K, P, Fe, Zn, Cu, Mn and CEC positively correlated with has a primary rol over accumulation or fixation of Zn in the soil. Regarding the soil's physical and physicochemical properties, the texture (clay content), organic matter and pH were the factors that sharply influenced soil Zn availability. Soil properties are capable of modulating Zn interaction,

adsorption, and precipitation with soil components, including organic matter pools (Alloway, 2009; Broadley et al., 2012; Montalvo et al., 2016). When associated with organic ligands, the Zn availability to crops relies on the chemical stability, solubility, and reactivity of the bonds formed between the humic matrix and the metal in the complexes formed. Generally, Fe and Al oxides have a larger and stronger sorption sites due to a chemisorption process take place on their surface, can coat relatively lesser soil component such as carbonates and sand sized fractions (Uygur and Rimmer, 2000). Smax parameter was significantly and positively correlated with K_F (0.940, $p < 0.01$) and n (0.755, $p < 0.01$) values.

The DTPA-Zn measured throughout the entire incubation period of 90-day are given in Table 7. The availability of plant nutrients during the growth period ensures a good performance along with a desirable yield and yield quality. The release of trace elements retained by sorption sites is required to sustainably meet the trace element needs of the plant (Uygur and Rimmer, 2000). In this study, HA+Zn application increased the DTPA-Zn concentration of the soils compared to only HA application in all soil series formed on marl (smectite clay mineral) and limestone (kaolinite clay mineral) parent material during the incubation period. When Zn was mixed with HA, free sulfates were

Table 10. Bivariate Pearson correlations between the soil properties and the sorption parameters

Parameters	Smax	k	K_F	n
k	0.110			
K_F	0.940**	0.206		
n	0.755**	0.154	0.514	
EC (dS m ⁻¹)	-0.301	-0.136	-0.413	-0.114
pH	0.559	0.038	0.628*	0.314
CCE (%)	0.137	-0.408	0.259	-0.323
P_{av} (kg ha ⁻¹)	-0.584*	-0.107	-0.761**	0.017
K_{av} (kg ha ⁻¹)	-0.541	0.421	-0.551	-0.171
OM (%)	-0.619*	0.138	-0.629*	-0.300
Sand (%)	0.526	0.213	0.530	0.339
Silt (%)	-0.582*	-0.101	-0.439	-0.631*
Clay (%)	-0.252	-0.252	-0.397	0.080
ΣN (%)	-0.620*	0.139	-0.629*	-0.300
ΣK (%)	0.459	-0.240	0.359	0.410
ΣP (%)	0.277	-0.338	0.449	-0.267
ΣFe (%)	0.117	-0.373	0.196	-0.255
ΣZn (mg kg ⁻¹)	0.279	-0.085	0.453	-0.202
ΣCu (mg kg ⁻¹)	0.313	-0.076	0.504	-0.248
ΣMn (mg kg ⁻¹)	0.250	-0.219	0.404	-0.223
CEC (cmol kg ⁻¹)	-0.210	-0.069	-0.352	0.149

Σ : Total, k: Bonding energy coefficient, K_F : Freundlich K, n: Constant, EC: Electrical conductivity, pH: Soil reaction in saturation paste, CCE: Calcium carbonate equivalent, P_{av} : Available P₂O₅, K_{av} : Available K₂O, OM: Organic matter, CEC: Cation exchange capacity

retained in the complex formed. In addition, bonds related to sulfone groups were formed, which are indicative of the interaction of HA with $ZnSO_4$ and the synthesis of Zn-HA complexes. Sulfur groups present in the HA+ $ZnSO_4$ mixture are indicative of Zn-HA complexation. This is probably due to the presence of functional groups containing oxygen, nitrogen and sulfur on the HA surface (Aguiar et al., 2013; Nardi et al., 2021; Zanin et al., 2019). There was no statistical difference between Zn treatment and Zn+HA, treatment in terms of DTPA-Zn concentrations of soils in both soil series (Başkuyu, Hacifakıl, Saraççeşme and Tine) regardless the parent material during the incubation period. The DTPA-Zn concentration of the soils was higher in the soil series formed in the marl parent material dominated by smectite clay mineral than in the limestone parent material, especially in Başkuyu soil series. The mechanisms and processes of Zn interaction with HA include ionic or uncoordinated forms, undefined complexes, bidentate chelates and bidentate bridging coordination bonds (Boguta and Sokołowska, 2016). The interaction of organic ligands with Zn is beneficial for providing Zn to plants, considering that free Zn ions are prone to react with soil constituents through sorption and precipitation, which are the main processes regulating Zn availability in soil (Montalvo et al., 2016; Olaetxea et al., 2018). Organic matter is also a source of soluble organic compounds that can bind micronutrients through the formation of soluble organo-metal complexes (Wang and Xing, 2005). In this study, S4, which had a higher Zn content than S2, S7 and S8, presented a higher soil Zn-DTPA content in the Başkuyu soil series formed on the marl parent material with smectite clay mineral during the incubation period.

The effects of humic acid (HA) and Zn treatments on the fresh and dry biomasses of maize grown on the marl (S4) and limestone soil (S8) in the greenhouse experiment were given in Table 8. The highest fresh

and dry weight of maize was 86.3 and 35.7 g from 2% HA+12 mg kg^{-1} Zn application in the Başkuyu soil series formed on the marl parent material with smectite clay mineral, respectively. Humic acid applications in soil formed on marl (smectite clay mineral) and limestone (kaolinite clay mineral) parent material increased the fresh weight of maize compared to the control. Humic substances are formed by the chemical and biological breakdown of plant and animal residues and provide an important source of organic carbon in soil. The most prominent biostimulant is humic acid, which has direct and indirect effects on the morpho-physiological growth of plants (Peña-Méndez et al., 2005). HA is mainly derived from humic substances and the final decomposed material contains 60% of the organic matter in the soil (Muscolo et al., 2013). On the other hand, application of HA to soil not only improves various soil properties (such as structural stability, biological activity and affecting soil pH), but also promotes better root development, water holding capacity, carbon sequestration, cation exchange capacity and nutrient availability in the rhizosphere (Gümüş and Şeker, 2015). Solution cations have an influence on the rate and amount of HA sorbed onto smectite. The adsorption of HA by mineral surfaces is mostly influenced by solution pH, ionic strength and the type of exchangeable cations. These factors also influence the structure conformation of HA (Essington, 2015). HA and clay complexes contribute to soil structure and water holding capacity and play an important role in the cycling of various nutrients and pollutants by taking part in reductive and oxidative reactions (Sparks, 2003). The presence of Ca^{2+} improves the binding between the mineral surface and HA. Ca^{2+} is more effective than K^+ in eliminating the repulsive charges between the smectite surface and weathered humates and fulvates. Furthermore, the Ca^{2+} ions reduce the HA solubility (Li et




Figure 2. Scatter diagram for the experimental max adsorption vs. predicted ones with (HA) and without (WS) humic acid (HA) treatment

al., 2004) by bridging clay and surfaces. This infact further decreases the efficiency of potentially high energy sites for Zn sorption by changing the nature of colloidal surfaces. Therefore, higher calcium carbonate containing Başkuyu soil (S4, 35.1%) had higher DTPA-Zn in overall than the Tine soil series (S8, 9.01%) due to apparent drawback the effect of excessive soil carbonates in the soil. Similarly, the heavy fraction of soil organic carbon such as HA accumulated twice as much SOC at the CaCO₃ bearing site, in Ca prevalent soils than the soils with less Ca (Rowley et al., 2021).

Conclusion

It was determined that the clay types were different in 12 different soil samples with two different parent materials belonging to the Arisol order. While the dominant clay mineral in the soils formed on marl parent material was smectite, the dominant clay mineral in the soils formed on limestone parent material was kaolinite. Because of adsorption analysis, the parameters of Langmuir and Freundlich models were determined. Langmuir model was found to be more suitable for Zn adsorption than Freundlich model. There was no significant difference between Langmuir and Freundlich isotherm parameters in soils formed on marl and limestone parent material with and without humic acid. The desorption rate (%) of adsorbed Zn increased as the initial Zn concentration increased in the all soil series formed on marl (smectite clay mineral) and limestone (kaolinite clay mineral) parent material. In the incubation study, humic acid applications increased the available Zn content in soils formed on both parent materials compared to the control. Humic acid applications increased the fresh weight of maize compared to the control. The highest available Zn was determined in Başkuyu series (0.79 mg kg⁻¹) formed on marl parent material, while the lowest available Zn was obtained in Saraççeşme series (0.60 mg kg⁻¹) formed on limestone parent material. In the greenhouse experiment, humic acid applications increased the fresh weight of maize in soils formed on both parent materials by 0.4 and 19.6% compared to to the control. Zn fertilization with HA further increased the fresh and dry yield that 12 mg kg⁻¹ performed better. The lowest wet and dry weight values of maize were obtained from the control in the soil formed on marl parent material, while the highest wet and dry weight values were obtained from 2% humic acid + 12 mg kg⁻¹ Zn treatment. Smax (maximum Zn adsorption of the soil) parameter of soils were negatively correlated with EC, P_{av}, K_{av}, organic matter, silt, clay, total N and cation exchange capacity (CEC) of the samples. Although humic acid alone increases the plant availability of Zn, HA + Zn applications should be applied together in order to obtain higher yields. It is

necessary to improve the results obtained by conducting studies for soils with different parent material and soil families.

Funding

The authors would like to thank the Republic of Turkey Ministry of Agriculture and Forestry General Directorate of Agricultural Research and Policies (TAGEM) for the financial support for the Project TAGEM-BB-0833-1.

Author Contribution

M.K.: Conceptualization, investigation, methodology, validation, software, validation, formal analysis, investigation, resources, data curation, writing-original draft preparation, writing-review and editing, visualization, supervision, statistical analysis, project administration. **G.D. and N.G.:** methodology, formal analysis. **V.U. and M.O.A.:** Methodology, validation, software, validation, formal analysis, investigation, resources, data curation, writing-original draft preparation, writing-review and editing, visualization, supervision, statistical analysis. **Z.D.:** Writing-original draft preparation, writing-review and editing, visualization, statistical analysis, supervision, resources, data curation, all authors reviewed the manuscript.

Conflicts of Interest

The authors declare no conflict of interest.

Acknowledgement

The authors would like to thank the Republic of Turkey Ministry of Agriculture and Forestry General Directorate of Agricultural Research and Policies (TAGEM) for the financial support for the Project TAGEM-BB-0833-1.

References

- Aguiar, N. O., Novotny, E. H., Oliveira, A. L., Rumjanek, V. M., Olivares, F. L., & Canellas, L. P. (2013). Prediction of humic acids bioactivity using spectroscopy and multivariate analysis. *Journal of Geochemical Exploration*, 129, 95-102. <http://dx.doi.org/10.1016/j.gexplo.2012.10.005>
- Alloway, B.J. (2009). Soil Factors Associated with Zinc Deficiency in Crops and Humans. *Environmental Geochemistry and Health*. 31:537-548. <http://dx.doi.org/10.1007/s10653-009-9255-4>
- Alumaa, P., Steinnes, E., Kirso, U., & Petersell, V. (2001). Heavy metal sorption by different Estonian soil types at low equilibrium solution concentrations. *Proc. Estonian Acad. Sci. Chem*, 50(2), 104-115. <http://dx.doi.org/10.3176/chem.2001.2.05>

- Bailey, L., Lekkerkerker, H. N., & Maitland, G. C. (2015). Smectite clay-inorganic nanoparticle mixed suspensions: phase behaviour and rheology. *Soft Matter*, 11(2), 222-236. <http://dx.doi.org/10.1039/c4sm01717j>
- Blatt, H., & Tracy, R.J. (1996). *Petrology*, 2nd. edit. W.H. Freeman & Co., 529pp.
- Boguta, P., & Sokołowska, Z. (2016). Interactions of Zn (II) ions with humic acids isolated from various type of soils. Effect of pH, Zn concentrations and humic acids chemical properties. *PLoS One*, 11(4), e0153626. <https://doi.org/10.1371/journal.pone.0153626>.
- Bouyoucos, G. J. (1951). A recalibration of the hydrometer method for making mechanical analysis of soils. <http://dx.doi.org/10.2134/agronj1951.00021962004300090005x>
- Black, C. A., Evans, D. D., White, J. L., Ensminger, L. E., & Clark, F. E. (1965). Methods of soil analysis, part (1) and part (2). *Am. Soc. Agron. Inc., Publ. Madison, Wisconsin, USA*. <https://doi.org/10.12691/ijebb-3-1-3>
- Broadley, M., Brown, P., Cakmak, I., Rengel, Z., & Zhao, F. (2012). Function of nutrients: micronutrients. In *Marschner's mineral nutrition of higher plants* (pp. 191-248). Academic Press. <https://doi.org/10.1016/B978-0-12-384905-2.00007-8>
- Brown, G., & Brindley, G. W. (1980). X-ray diffraction procedures for clay mineral identification.
- Brown, P. H., Cakmak, I., & Zhang, Q. (1993). Form and function of zinc plants. In *Zinc in Soils and Plants: Proceedings of the International Symposium on 'Zinc in Soils and Plants' held at The University of Western Australia, 27-28 September, 1993* (pp. 93-106). Dordrecht: Springer Netherlands.
- Çakmak, İ., Kalaycı, M., Ekiz, H., Braun, H. J., Kılınc, Y., & Yılmaz, A. (1999). Zinc deficiency as a practical problem in plant and human nutrition in Turkey: a NATO-science for stability project. *Field Crops Research*, 60(1-2), 175-188. [https://doi.org/10.1016/S0378-4290\(98\)00139-7](https://doi.org/10.1016/S0378-4290(98)00139-7)
- Cerda, A. (2002). The effect of season and parent material on water erosion on highly eroded soils in eastern Spain. *J. Arid. Environ.* 52:319-337. <https://doi.org/10.1006/jare.2002.1009>
- Çavdar, A. O., Arcasoy, A., Cin, S., Babacan, E., & Gözdasoğlu, S. (1983). Geophagia in Turkey: iron and zinc deficiency, iron and zinc absorption studies and response to treatment with zinc in geophagia cases. *Progress in clinical and biological research*, 129, 71-97.
- de Morais, T. M. O., Berenguer, E., Barlow, J., França, F., Lennox, G. D., Malhi, Y., ... & Ferreira, J. (2021). Leaf-litter production in human-modified Amazonian forests following the El Niño-mediated drought and fires of 2015–2016. *Forest Ecology and Management*, 496, 119441. <https://doi.org/10.1016/j.foreco.2021.119441>
- Delgado, A., & Gómez, J. A. (2016). The soil. Physical, chemical and biological properties. *Principles of agronomy for sustainable agriculture*, 15-26. https://doi.org/10.1007/978-3-319-46116-8_2
- Dogar, M. A., & Van Hai, T. (1980). Effect of P, N and HCO₃-Levels in the Nutrient Solution on Rate of Zn Absorption by Rice Roots and Zn Content in Plants. *Zeitschrift für Pflanzenphysiologie*, 98(3), 203-212. <https://doi.org/10.1080/01904167.2014.920377>
- Essington, M.E. (2015). *Soil and Water Chemistry: An Integrative Approach*. 2nd edition. CRC Press, Boca Raton, Florida, USA, 656 pp.
- Eyupoglu, F., Kurucu, N., & Sanyasag, U., (1994). Status of plant available micronutrients in Turkish soils. In: *Soil and Fertilizer Research Institute 1993 annual report*. Report No: 118, Ankara, Turkey.
- Gümüş, İ., & Şeker, C. (2015). Influence of humic acid applications on modulus of rupture, aggregate stability, electrical conductivity, carbon and nitrogen content of a crusting problem soil. *Solid Earth*, 6(4), 1231-1236. <https://doi.org/10.5194/se-6-1231-2015>
- Jackson, J.B.C. (1997). Reefs since Columbus. *Coral Reefs* 16:23-32.
- Jacobs, P. M. (1998). Influence of parent material grain size on genesis of the Sangamon Geosol in south-central Indiana. *Quaternary International*, 51, 127-132.
- Johnston, C. T., de Oliveira, M. F., Teppen, B. J., Sheng, G., & Boyd, S. A. (2001). Spectroscopic study of nitroaromatic-smectite sorption mechanisms. *Environmental science & technology*, 35(24), 4767-4772. <http://doi.org/10.1021/es010909x>
- Justi, M., Morais, E. G., & Silva, C. A. (2019). Fulvic acid in foliar spray is more effective than humic acid via soil in improving coffee seedlings growth. *Archives of Agronomy and Soil Science*. <http://doi.org/10.1080/03650340.2019.1584396>
- Kabata-Pendias, A. (2011). *Trace Elements in Soils and Plants*, 4th ed. CRC Press, Boca Raton, Florida, USA
- Kacar, B. (1998). Toprakta Çinkonun Bulunuşu, Yarıyışliliği ve Tepkimeleri. 1.Ulusal Çinko Kongresi, 12–16 Mayıs 1997, Eskişehir. 47-60. Adana.
- Kloprogge, J. T., & Frost, R. L. (1999). Fourier transform infrared and Raman spectroscopic study of the local structure of Mg-, Ni-, and Co-hydrotalcites. *Journal of Solid State Chemistry*, 146(2), 506-515. <https://doi.org/10.1006/jssc.1999.8413>
- Laird, D. A., & Fleming, P. D. (1999). Mechanisms for adsorption of organic bases on hydrated smectite surfaces. *Environmental Toxicology and Chemistry: An International Journal*, 18(8), 1668-1672. [https://doi.org/10.1897/15515028\(1999\)018<1668:MF AOOB>2.3.CO;2](https://doi.org/10.1897/15515028(1999)018<1668:MF AOOB>2.3.CO;2)
- Li, H., Teppen, B. J., Laird, D. A., Johnston, C. T., & Boyd, S. A. (2004). Geochemical modulation of pesticide sorption on smectite clay. *Environmental science & technology*, 38(20), 5393-5399. <https://doi.org/10.1021/es0494555>
- Lindsay, W.L., & Norvell, W.A. (1978). Development of DTPA soil test for zinc, iron, manganese and copper. *Soil Science Society of America Journal*, 42:421-28. <https://doi.org/10.2136/sssaj1978.03615995004200030009x>
- Martinez-Mena, M., Castillo, V., & Albaladejo, J. (2002). Relations between interrill erosion processes and sediment particle size distribution in a semiarid Mediterranean area of SE of Spain. *Geomorphology*, 45(3-4), 261-275. [https://doi.org/10.1016/S0169-555X\(01\)00158-1](https://doi.org/10.1016/S0169-555X(01)00158-1)

- Marshall, C. P., & Fairbridge, R. W. (Eds.). (1999). *Encyclopedia of geochemistry*. Springer Science & Business Media.
- Milliken, K. L. (2003). *Late diagenesis and mass transfer in sandstone shale sequences* (Vol. 7, p. 407).
- Montalvo, D., Degryse, F., Da Silva, R. C., Baird, R., & McLaughlin, M. J. (2016). Agronomic effectiveness of zinc sources as micronutrient fertilizer. *Advances in agronomy*, 139, 215-267. <https://doi.org/10.1016/bs.agron.2016.05.004>
- Mortland, M.M., (1970). Clay-organic complexes and interactions. *Adv Agron*. 22:75–117.
- Musco, A., Sidari, M., & Nardi, S. (2013). Humic substance: relationship between structure and activity. Deeper information suggests univocal findings. *Journal of Geochemical Exploration*, 129, 57-63. <https://doi.org/10.1016/j.gexplo.2012.10.012>
- Nardi, S., Schiavon, M., & Francioso, O. (2021). Chemical structure and biological activity of humic substances define their role as plant growth promoters. *Molecules*, 26(8), 2256. <https://doi.org/10.3390/molecules26082256>
- Olaetxea, M., De Hita, D., Garcia, C. A., Fuentes, M., Baigorri, R., Mora, V., & Garcia-Mina, J. M. (2018). Hypothetical framework integrating the main mechanisms involved in the promoting action of rhizospheric humic substances on plant root-and shoot-growth. *Applied Soil Ecology*, 123, 521-537. <http://dx.doi.org/10.1016/j.apsoil.2017.06.007>
- Peña-Méndez, E. M., Gajdošová, D., Novotná, K., Prošek, P., & Havel, J. (2005). Mass spectrometry of humic substances of different origin including those from Antarctica: A comparative study. *Talanta*, 67(5), 880-890. <http://dx.doi.org/10.1016/j.talanta.2005.03.032>
- Richard, L. A. (1954). Diagnosis and improvement of saline and alkaline soils. Handbook No. 60. *US Department of Agriculture*.
- Rosa, S. D., Silva, C. A., & Maluf, H. J. G. M. (2018). Wheat nutrition and growth as affected by humic acid-phosphate interaction. *Journal of Plant Nutrition and Soil Science*, 181(6), 870-877. <http://dx.doi.org/10.1002/jpln.201700532>
- Rose, T. J., Impa, S. M., Rose, M. T., Pariasca-Tanaka, J., Mori, A., Heuer, S., & Wissuwa, M. (2013). Enhancing phosphorus and zinc acquisition efficiency in rice: a critical review of root traits and their potential utility in rice breeding. *Annals of botany*, 112(2), 331-345. <http://dx.doi.org/10.1093/aob/mcs217>
- Rowley, M. C., Grand, S., Spangenberg, J. E., & Verrecchia, E. P. (2021). Evidence linking calcium to increased organo-mineral association in soils. *Biogeochemistry*, 153(3), 223-241. <http://dx.doi.org/10.1007/s10533-021-00779-7>
- Sawhney, B. L., & Singh, S. S. (1997). Sorption of atrazine by Al- and Ca-saturated smectite. *Clays and Clay Minerals*, 45(3), 333-338. <https://doi.org/10.1346/CCMN.1997.0450304>
- Šestanović, S. (2001). *Osnove geologije i petrografije*, (The basics of geology and petrography in Croatian), Građevinski fakultet Sveučilišta u Splitu, Split.
- Sheng, G., Johnston, C. T., Teppen, B. J., & Boyd, S. A. (2002). Adsorption of dinitrophenol herbicides from water by montmorillonites. *Clays and Clay Minerals*, 50(1), 25-34. <https://doi.org/10.1346/000986002761002630>
- Sillanpää, M. (1990). *Micronutrient assessment at the country level: an international study* (No. 63, pp. pp-208). <http://dx.doi.org/10.1186/s12889-016-2765-y>
- Soil Survey Staff. (2014). Kellogg soil survey laboratory methods manual. Soil Survey Investigations Report No. 42, Version 5.0, ed. R. Burt and Soil Survey Staff. Lincoln, NE: USDA Natural Resources Conservation Service.
- Spark, K. M., Wells, J. D., & Johnson, B. B. (1997). The interaction of a humic acid with heavy metals. *Soil Research*, 35(1), 89-102. <https://doi.org/10.1071/S96008>
- Sparks, D., (1996). *Methods of Soil Analysis, Part 3. Chemical Methods*, Soil Science Society of America Inc., Madison, Wisc, USA.
- Sparks, D.L., (2003). *Environmental Soil Chemistry* 2nd Ed., Academic Press, Amsterdam, The Netherlands.
- Stevenson, F. J. (1994). *Humus chemistry: genesis, composition, reactions*. John Wiley & Sons.
- Trehan, S. P., & Sekhon, G. S. (1977). Effect of clay, organic matter and CaCO₃ content on zinc adsorption by soils. *Plant and Soil*, 46, 329-336. <https://doi.org/10.1007/BF00010089>
- Trettin Jr, C. C., Johnson, D. W., & Todd, D. E. (1999). Forest Nutrient and Carbon Pools at Walker Branch Watershed Changes during a 21-Year Period. *Soil Science Society of America Journal*, 63(5), 1436-1448. <https://doi.org/10.2136/sssaj1999.6351436x>
- USGS. (2019). Illite Group Minerals. USGS Coastal and Marine Geology Program. Retrieved 3 Apr 2019.
- Uygur, V., & Rimmer, D. L. (2000). Reactions of zinc with iron-oxide coated calcite surfaces at alkaline pH. *European Journal of Soil Science*, 51(3), 511-516. <https://doi.org/10.1046/j.1365-2389.2000.00318.x>
- Wang, K., & Xing, B. (2005). Structural and sorption characteristics of adsorbed humic acid on clay minerals. *Journal of Environmental Quality*, 34(1), 342-349. <https://doi.org/10.2134/jeq2005.0342>
- Washer, N. E., & Collins, M. E. (1988). Genesis of adjacent morphologically distinct soils in northwest Florida. *Soil Science Society of America Journal*, 52(1), 191-196. <https://doi.org/10.2136/sssaj1988.03615995005200010033x>
- Weil, R.R., & Brady, N.C. (2017). *Soil organic matter. Nature and properties of soils* (15th ed). Pearson Education Limited, England, 545-601.
- Whittig, L. D., & Allardice, W. R. (1986). X-ray diffraction techniques. *Methods of Soil Analysis: Part 1 Physical and Mineralogical Methods*, 5, 331-362.
- Yavitt, J. B. (2000). Nutrient Dynamics of Soil Derived from Different Parent Material on Barro Colorado Island, Panama 1. *Biotropica*, 32(2), 198-207. <https://doi.org/10.1111/j.1744-7429.2000.tb00462.x>
- Zanin, L., Tomasi, N., Cesco, S., Varanini, Z., & Pinton, R. (2019). Humic substances contribute to plant iron nutrition acting as chelators and biostimulants. *Frontiers in Plant Science*, 10, 45287. <https://doi.org/10.3389/fpls.2019.00675>