

Original Article/Özgün Araştırma

The effect of probiotic culture on quality characteristics of sucuk

Probiyotik kültürün sucuğun kalite karakteristikleri üzerine etkisi

İrem Ece Kayar¹ , Güzin Kaban² , Barış Yalınkılıç^{1*}  

¹Department of Gastronomy and Culinary Arts, Faculty of Architecture and Design, Istanbul Gedik University, Kartal, İstanbul, Türkiye

²Atatürk University, Faculty of Agriculture, Department of Food Engineering, Erzurum, Türkiye

(Yazar sıralamasına göre)

ORCID ID: 0009-0001-5701-2294, M.Sc.

ORCID ID: 0000-0001-6720-7231, Prof. Dr.

ORCID ID: 0000-0002-6195-7821, Dr.

*Corresponding author / Sorumlu yazar: baris.yalinkilic@gedik.edu.tr

Geliş Tarihi : 17.10.2024

Kabul Tarihi : 17.12.2024

Abstract

Objective: Sucuk is a dry fermented meat product that naturally contains high numbers of lactic acid bacteria. This study investigated the feasibility of incorporating the probiotic lactic acid bacteria strain *Lactocaseibacillus casei* 431 in sucuk production.

Material and methods: For this purpose, three groups of sucuk were prepared: Group 1 with a starter culture (*Latilactobacillus sakei* S15 + *Staphylococcus xylosus* GM92), Group 2 with the probiotic culture (*L. casei* 431), and Group 3 combining both starter and probiotic cultures (*L. sakei* S15 + *S. xylosus* GM92 + *L. casei* 431). Then, the sucuk samples were subjected to physical, chemical, microbiological, and sensory analyses on the 0th, 30th, and 60th days of storage.

Results and conclusion: The results demonstrated that the microbial culture factor significantly influenced the sucuk samples' pH, L^* value, a^* value, thiobarbituric acid reactive substances (TBARS), odor, taste, and overall acceptability scores, as well as the counts of lactic acid bacteria, *Micrococcus/Staphylococcus*, yeasts-molds, and *L. casei* ($P < 0.01$). However, it had no significant impact ($P > 0.05$) on water activity, b^* value, color, or texture parameters. The storage period notably affected the TBARS, L^* value and lactic acid bacteria ($P < 0.05$) and *L. casei* counts ($P < 0.01$), whereas it did not significantly alter other examined parameters ($P > 0.05$).

Keywords: *Lactocaseibacillus casei*; probiotic sucuk; fermented sausage; starter culture

Öz

Amaç: Kuru fermente bir et ürünü olan sucuk, doğası gereği bünyesinde yüksek sayıda laktik asit bakterisi içermektedir. Bu çalışmada, probiyotik laktik asit bakteri suşu *Lactocaseibacillus casei* 431'in sucuk üretiminde kullanılabilirliği araştırılmıştır.

Materyal ve yöntem: Bu maksatla çalışma kapsamında üç grup sucuk hazırlanmıştır: Grup 1: starter kültür (*Latilactobacillus sakei* S15 + *Staphylococcus xylosus* GM92); grup 2: probiyotik kültür (*L. casei* 431); grup 3: starter kültür + probiyotik kültür (*L. sakei* S15 + *S. xylosus* GM92 + *L. casei* 431). Sucuk örneklerine üretimi müteakip depolamanın 0., 30. ve 60. günlerinde fiziksel, kimyasal, mikrobiyolojik ve duyu analizler uygulanmıştır.

Tartışma ve sonuç: Araştırma sonuçlarına göre mikrobiyal kültür faktörü sucuk örneklerinin pH, L^* , a^* , TBARS, koku, tat ve genel kabul edilebilirlik değerleri ile laktik asit bakteri, *Micrococcus/Staphylococcus*, maya-küf ve *L. casei* sayıları üzerinde önemli düzeyde etkili olmuştur ($P < 0.01$). Buna karşın mikrobiyal kültür faktörü sucuk örneklerinin su aktivitesi, b^* , renk ve tekstür parametreleri üzerinde önemli ($P > 0.05$) etkiye sahip değildir. Depolama periyodu faktörü ise örneklerin tiyobarbitürik asit reaktif maddeleri (TBARS), L^* değeri, laktik asit bakteri ($P < 0.05$) ve *L. casei* ($P < 0.01$) parametreleri üzerinde önemli etki

sergilemiştir. Buna karşın depolama periyodunun incelenen diğer parametreler üzerinde ise önemli etkisi ($P > 0.05$) olmamıştır.

Anahtar kelimeler: *Lacticaseibacillus casei*; probiyotik sucuk; fermente sosıs; starter kültür

1. Introduction

Probiotics are live microorganisms that have beneficial effects on host health when consumed in adequate amounts (Food and Agricultural Organization (FAO), 2006). These microorganisms can be directly added to food or offered as dietary supplements in the form of powder, capsules, or tablets (Coconcelli and Fontana, 2008). The rapid increase in the trend towards health-focused and healthy living foods, which started with probiotic dairy products in the 1990s, has also increased the interest in developing probiotic meat products. Probiotics require the intake of adequate amounts of live microorganisms to confer their benefits, and fermented sausages are considered an optimal delivery matrix for such probiotics. (Bağdatlı and Kundakçı, 2013; Bis-Souza et al., 2019a; Khan et al., 2011; Kröckel, 2013; Pavli et al., 2020). The most commonly used probiotic microorganism strains belong to the genera *Lactobacillus* and *Bifidobacterium*. Additionally, microorganisms from the genera *Enterococcus* and *Pediococcus* are also used as probiotic cultures, albeit less frequently (Cavalheiro et al., 2015).

Fermented sausages are generally classified into two categories as dry and semi-dry fermented sausages. Dry fermented sausages do not undergo any heat treatment, whereas the production process for semi-dry fermented sausages may include a heat treatment step (Caplice and Fitzgerald, 1999; Kumar et al., 2017). In these products, lactic acid bacteria and Gram-positive, catalase-positive cocci are technologically significant microorganisms constituting two important components of the product microbiota. Lactic acid bacteria contribute to product safety and the development of sensory attributes such as texture and color through acid production. Gram-positive, catalase-positive cocci influence color formation and stability through their nitrate reductase activities, delay oxidation with their catalase activities, and contribute to flavor formation through their proteolytic and lipolytic properties. Specific strains selected from both groups of microorganisms are used either individually or in combination as starter cultures in industrial production (Yalınkılıç, 2009; Karshoğlu et al., 2014; Kaya and Kaban, 2019; Alvarez et al., 2023; Yılmaz Topcam et al., 2024).

Sucuk is a type of dry fermented sausage commonly produced in Türkiye, in which beef and/or sheep/buffalo meat, along with sheep tail fat and/or beef fat are used as raw materials. In this product, bovine small intestine or collagen casings are used as casing. Fermentation and drying are two critical stages in the production of sucuk (Kaban and Kaya, 2009; Yalınkılıç et al., 2012). The initial fermentation temperature and ripening period vary between 12 to 26 °C and 6 to 20 days, respectively (Kaban and Kaya, 2008, 2009; Soyer et al., 2005; Yalınkılıç et al., 2012). Unlike semi-dry fermented sausages, which undergo heat treatment, the production process for dry fermented sausages does not involve any heat treatment. Therefore, sucuk provides a favorable environment for probiotic microorganisms.

In the production of probiotic meat products, strains exhibiting probiotic properties, isolated or identified from intestinal microbiota and/or fermented meat products, can be utilized. However, it is necessary to determine the resilience of these strains to processing conditions and storage environments (Kołożyn-Krajewska and Dolatowski, 2012; Neffe-Skocińska et al., 2016). To this end, studies have been conducted on the potential applications of probiotic microorganisms in both sucuk (Bağdatlı and Kundakçı, 2016; Ergönül and Kundakçı, 2011; Kaya and Aksu, 2005; Kozan and Sarıçoban, 2023; Öztürk Er, 2002; Tükel and Şengün, 2024; Ünal Turhan et al., 2019) and other non-heat-treated fermented sausages (Agüero et al., 2020; Arief et al., 2014; Bis-Souza et al., 2019b, Bis-Souza, 2020; Erkkilä et al., 2001; Holko et al., 2013; Klingberg and Budde, 2006; Klingberg et al., 2005). Moreover, each new probiotic strain, considered to have technological significance, may have varied effects on product quality characteristics, necessitating their experimental use in the product environment. As mentioned, the use of various probiotic strains in the production of sucuk has been examined. However, research is still needed to develop new strains with high potential for adaptation to the sucuk environment and to determine their effects on product quality characteristics. Thus, the aim of this study was to investigate the effects of simultaneous use of both starter culture (*L. sakei* S15 and *S. xyloso* GM92)

and probiotic culture (*L. casei* 431) on the quality characteristics of sucuk.

2. Materials and methods

2.1. Materials

Beef meat and beef fat from three different beef carcasses, obtained from a local butcher, were used as raw materials. Spices used in the formulation were purchased from a local market. Sodium nitrite and all other chemicals used in the study were obtained from a medical supply company. The study was conducted with three replicates, using one carcass for each replicate. The strains *Latilactobacillus sakei* S15 (Kaya et al., 2015) and *Staphylococcus xylosum* GM92 (Kaban and Kaya, 2008) were used as starter cultures, while *Lactocaseibacillus casei* 431 (Christian Hansen, Denmark), a commercial probiotic culture strain, was used as the probiotic culture.

The chemicals and culture media used in this study are as follows: Sodium chloride (NaCl, Merck), Trichloroacetic Acid (TCA, 1.00807, Merck), Ethylenediaminetetraacetic Acid (EDTA, 1.08418, Merck), Propyl Gallate (P3130, Sigma-Aldrich), 2-Thiobarbituric Acid (TBA, T5500, Sigma-Aldrich), and Ethanol (1.00983, Merck). The culture media include De Man Rogosa Sharpe Agar (MRS, 1.10660, Merck), Mannitol Salt Phenol-Red Agar (MSA, 1.05404, Merck), Violet Red Bile Dextrose Agar (VRBD-Agar, 1.10275, Merck), Rose Bengal Chloramphenicol Agar (RBC-Agar, 1.00467, Merck), and MRS-IM Agar.

2.2. Sucuk production

In the production of sucuk, beef meat and fat were used in an 80:20 ratio as raw materials, and spice varieties were included in the formulation as specified by Yalınkılıç et al. (2012). Sucrose (4 g/kg of sucuk dough) was added as a source of sugar and 150 mg/kg of sodium nitrite as a curing agent. In the study, three groups of sucuk were prepared: Group 1 with a starter culture (*Latilactobacillus sakei* S15 + *Staphylococcus xylosum* GM92), Group 2 with the probiotic culture (*Lactocaseibacillus casei* 431), and Group 3 combining both starter and probiotic cultures (*L. sakei* S15 + *S. xylosum* GM92 + *L. casei* 431). A total of 9 sucuk doughs were prepared by making 3 production runs (replicate) for each experimental group. The sucuk doughs were

prepared in a small-scale cutter (MADO, Germany) and stuffed into collagen casings (Naturin Darm, 38 mm) using a small-scale stuffer (MADO, Germany). After stuffing, the sucuk samples from each group were allowed to rest for 4 hours in the production area before being transferred to the climate chamber (Reich, Germany) for the ripening stage, where air speed, relative humidity, and temperature could be automatically adjusted. The initial fermentation temperature was set at 24 ± 1 °C and was gradually reduced from the second day onward. The relative humidity was set at 92% and was gradually decreased over the following days. Air velocity was set at 0.5 m/s for the first three days and then reduced to 0.1 m/s in subsequent days. Ripening was terminated when the water activity (a_w) of the sucuk samples dropped below 0.90. The ripened sucuk samples were vacuum-packaged using a laboratory-type packaging machine (Multivac, Germany) in a packaging material made of Polyamide/Polyethylene with low oxygen, gas, and water vapor permeability. After packaging, the samples were stored at a cold temperature (4 ± 1 °C) for 60 days, and samples from each group were subjected to microbiological, sensory, and physicochemical analyses on days 0, 30, and 60.

2.3. Analyses

2.3.1. Microbiological analyses

For microbiological analyses, 25 g of samples were taken and homogenized in a sterile stomacher bag with 225 mL of sterile physiological saline for 2 minutes (Lab Stomacher Blander 400-BA 7021, Seward, England). The homogenized samples were then analyzed for lactic acid bacteria counts using de Man, Rogosa, and Sharpe Agar (MRS) (30°C for 48 h under anaerobic conditions), *Micrococcus/Staphylococcus* counts using Mannitol Salt Agar (MSA) (30°C for 48 h under aerobic conditions), Enterobacteriaceae counts using Violet Red Bile Dextrose Agar (VRBD) (30°C for 48 h under anaerobic conditions) (Baumgart et al., 1993), and yeast-mold counts using Rose Bengal Chloramphenicol Agar (RBC) (25°C for 5 days under aerobic conditions) (Gökalp et al., 2010). For determining the count of probiotic bacteria, MRS-IM Agar specified by the manufacturer of the probiotic culture was used (Chr Hansen, Denmark, 2005). After inoculation,

the plates were incubated at 20 °C for 6 days, followed by enumeration (Chr Hansen, Denmark, 2005). Additionally, verification tests were conducted on typical colonies (Baumgart et al., 1993).

2.3.2. pH and water activity (a_w)

Ten grams of samples were weighed and homogenized with 100 mL of distilled water using a homogenizer. The pH value of the homogenized samples was measured using a previously calibrated pH meter (Gökalp et al., 2010). The water activity (a_w) of the samples was measured using a water activity device (Novasina, Pfäffikon, Switzerland).

2.3.3. Color analysis

The color measurements of the sucuk samples were determined using a colorimeter (Chroma Meter CR-400, Japan). The color values (L^* , a^* , and b^*) of the samples were measured on the cutting surfaces of the samples.

2.3.4. TBARS analysis

The analysis of thiobarbituric acid reactive substances (TBARS) was performed using the method described by Lemon (1975), and the results are presented as mg MDA/kg.

2.3.5. Sensory analysis

Sucuk samples were evaluated by 25 trained panelists in each panel. Prior to the sensory analysis, participants were informed about the products and given preliminary instructions on how to taste the sucuk. The analysis was carried out by presenting the samples coded with 3-digit numbers to the panelists along with the sensory panel form. The number of participants in each replication was 25, with a total of 75 participants. Sensory evaluation parameters including color, texture, odor, taste, and overall acceptability were scored on a 1-9 point scale.

2.3.6. Statistical analyses

The study was carried out according to a randomized complete block experimental plan with 3 replications, based on the factors of treatment (starter culture, probiotic culture, starter culture + probiotic culture) and storage period (0, 30, and 60 days). The raw data were examined for normality of distribution and homogeneity of

variances between groups, and then subjected to analysis of variance (ANOVA) using SPSS version 24 (SPSS Inc., Chicago, IL, USA). For the mean values that showed statistically significant differences ($P < 0.05$) according to the ANOVA, Duncan's multiple range test was also applied.

3. Results and discussion

3.1. pH

The effects of the microbial culture and storage period on the pH value of the sucuk are presented in Table 1. The pH value measured in all groups was below 5.0. This was probably due to the organic acids released during the metabolic activities of lactic acid bacteria used in sucuk production. Similar results have been obtained in studies utilizing starter cultures (Yalınkılıç et al., 2012) or probiotic cultures (Bağdatlı and Kundakcı, 2016). The average pH values in sucuk samples using either starter or probiotic cultures alone are higher compared to those where both were used together ($P < 0.05$). The combined use of starter and probiotic cultures has resulted in a greater reduction in product pH. The interaction of microbial culture and storage period did not significantly affect the average pH values of the sucuk samples over a 60-day storage period ($P > 0.05$). Contrary to our findings, a study using *L. casei* CRL-431 and *Lactobacillus acidophilus* as probiotic cultures in turkey meat-based probiotic sucuk found that the pH value decreased during the first two months of storage, with no change in the subsequent months (Ergönül, 2009). Another study found no variation in the pH of probiotic sucuk depending on the culture type, although a partial decrease was observed during storage (Ünal Turhan et al., 2017). However, in research using strains *Lactobacillus acidophilus* DSM 20079, *Lacticaseibacillus casei* 431, and *L. acidophilus* NCFM in probiotic sucuk, an increase in product pH values was observed in the later stages of storage. The same study also noted differences in product pH depending on the strain used (Kozan and Sarıçoban, 2023).

3.2. Water activity

The effects of the microbial culture and storage period on the water activity value of the sucuk are presented in Table 1. Water activity values in all samples were at least 0.90, indicating that the desired level of drying in the product was achieved. According to the results, neither

microbial culture nor the storage process had a statistically significant effect on the water activity values of probiotic sucuk samples ($P > 0.05$). Similarly, a study on probiotic sucuk found that the microbial culture used did not significantly affect the product's water activity, and no significant changes were observed in the water activity values up to the 60th day of storage (Ünal Turhan et al., 2017). In contrast, research conducted by Kozan and Sariçoban (2023) determined that lactic acid bacteria strains used in the production of probiotic sucuk influenced the product's water activity, and the storage duration also negatively affected the water activity values. On the other hand, in the present study, the interaction of microbial culture with the storage period did not have a statistically significant impact on the average water activity values of the sucuk samples ($P > 0.05$).

3.3. Color

The effects of the microbial culture and storage period on the L^* , a^* , and b^* values of the sucuk are presented in Table 1. The microbial culture variable significantly influenced the L^* and a^* values of sucuk samples ($P < 0.01$) while it had no

significant effect on the b^* value ($P > 0.05$). The storage period only significantly affected the L^* parameter of sucuk samples ($P < 0.01$), with no significant impact on other parameters ($P > 0.05$). Similarly, a research showed that the L^* values of turkey sucuk produced with different probiotic cultures varied depending on the type of culture used, and that an 8-month storage process had a significant effect on the product's L^* values. However, in contrast to our findings, the same study reported that the culture used did not significantly influence the a^* value. The researcher also observed that both the culture used and the period of storage had a significant effect on the b^* value (Ergönül, 2009). In sucuk samples produced using *L. acidophilus* DSM 20079, *L. casei* 431, and *L. acidophilus* NCFM strains, a regular decrease in L^* , a^* , and b^* values was observed throughout the storage period, and the cultures used did not have a significant impact on the product's L^* , a^* , and b^* values (Kozan and Sariçoban, 2023). Additionally, in the present study, the interaction of microbial culture with the storage period had no statistically significant effect on the instrumental color parameters of the samples ($P > 0.05$) (Table 1).

Table 1. Overall effect of microbial culture and storage period on the pH, a_w , instrumental color and TBARS parameters of dry-fermented probiotic sucuk.

	pH	a_w	L^*	a^*	b^*	TBARS (mg MDA/kg)
Microbial Culture (MC)						
SC	4,94±0,14 ^a	0,900±0,002 ^a	41,76±1,47 ^a	14,20±0,77 ^b	12,61±1,23 ^a	0,78±0,30 ^a
PC	4,88±0,12 ^a	0,902±0,004 ^a	39,71±0,80 ^b	16,29±0,93 ^a	12,41±1,47 ^a	0,47±0,17 ^c
SC+PC	4,78±0,07 ^b	0,899±0,005 ^a	40,94±1,51 ^a	15,90±0,90 ^a	13,05±0,84 ^a	0,63±0,28 ^b
Significance	**	NS	**	**	NS	**
Storage Period (SP)						
Day 0	4,83±0,12 ^a	0,902±0,005 ^a	41,72±1,48 ^a	15,23±1,53 ^a	11,93±1,11 ^a	0,36±0,08 ^c
Day 30	4,86±0,13 ^a	0,901±0,003 ^a	40,82±1,63 ^{ab}	15,42±1,36 ^a	13,35±0,75 ^a	0,62±0,18 ^b
Day 60	4,90±0,14 ^a	0,899±0,004 ^a	39,87±0,85 ^b	15,74±0,84 ^a	12,79±1,30 ^a	0,91±0,21 ^a
Significance	NS	NS	**	NS	NS	**
Interaction						
MCxSP	NS	NS	NS	NS	NS	*

a-c: any two means in the same column having the same letters in the same section are not significantly different. * $p < 0.05$, ** $p < 0.01$, NS: not significant. SC: starter culture, PC: probiotic culture, SC+PC: starter culture + probiotic culture.

3.4. TBARS

The effects of the microbial culture and storage period on the TBARS values of the sucuk samples are presented in Table 1. The microbial culture

had a significant effect on the TBARS value of sucuk ($P < 0.01$). The use of probiotic cultures resulted in lower TBARS values in the product compared to other groups ($P < 0.05$) (Table 1).

Ergönül (2009) found that turkey sucuk produced with *L. casei* CRL-431 had lower TBA values compared to those produced with *L. acidophilus* strain, with the highest average TBARS values observed in the group using starter cultures. Upon examining the storage period, it was determined that oxidation increased over time ($P < 0.01$), with the lowest TBARS values recorded on day 0 and the highest on day 60 of storage ($P < 0.05$). Additionally, the interaction of microbial culture with the storage period had a statistically significant effect on the TBARS values of the sausage samples ($P < 0.05$). The observed increase in TBARS values during storage is likely due to the accumulation of compounds released by oxidative reactions (Feiner, 2006). Parallel to our findings, a study on probiotic turkey sucuk noted an increase in TBA values over an eight-month storage period (Ergönül, 2009). However, in a study where three different probiotic lactic acid bacteria strains were used separately, Kozan and Sarıçoban (2023) observed a decrease in TBARS values during storage, and also noted variability in TBARS values depending on the bacterial strain used.

3.5. Microbiological properties

The effects of microbial culture and storage period on microbiologically significant groups in sucuk samples are presented in Table 2. The lactic acid bacteria count was affected by microbial culture ($P < 0.01$). The highest average number of lactic acid bacteria was found in the group using starter culture, with significant differences between this group and all others ($P < 0.05$). The lowest average number of lactic acid bacteria was observed in the group using only the probiotic culture ($P < 0.05$). Similar results were observed for the counts of *Micrococcus/Staphylococcus* ($P < 0.01$) (Table 2). However, the interaction of microbial culture and storage period had a statistically significant effect on the *Micrococcus/Staphylococcus* values of the sucuk samples ($P < 0.01$). A difference at the $P < 0.05$ level was found between the group using only the probiotic culture and the group using both starter and probiotic cultures in terms of *L. casei* counts. Similar to our findings, Bağdatlı and Kundakcı (2016) found lactic acid bacteria and *L. casei* CRL-431 counts to be above 8 log and 6 log, respectively, in their study on probiotic sucuk. This study shows that the count of *L. casei* 431 in

probiotic sucuk samples exceeded the 6 log cfu/g level as indicated by Kolozyn-Krajewska and Dolatowski (2009), demonstrating that the *L. casei* 431 strain has sufficient adaptability for probiotic sucuk production (Bağdatlı and Kundakcı, 2016). The count of yeasts/molds in all groups was below the 3 log level, with the lowest average value found in the group produced with the starter culture ($P < 0.05$). In contrast to our findings, studies on probiotic sucuk produced from beef (Bağdatlı and Kundakcı, 2016) and turkey meat (Ergönül, 2009) using the *L. casei* CRL-431 strain reported yeast/mold counts below detectable limits. In the present study, Enterobacteriaceae counts remained below the detectable limit (< 2 log) in all groups, likely due to the reduction in water activity below 0.90 and the observed decrease in pH.

The storage period had a significant ($P < 0.05$) effect on the count of lactic acid bacteria and a very significant ($P < 0.01$) effect on the count of *L. casei*, with both microbial parameters experiencing significant declines during storage. Neither *Micrococcus/Staphylococcus* nor yeast/mold counts showed significant changes during the storage period ($P > 0.05$) (Table 2). Data from the study indicate that the count of *L. casei* remained above 7 log cfu/g during storage, demonstrating the high survivability of this strain throughout the storage period. This indicates that the *L. casei* 431 strain used in the study is suitable for probiotic sucuk production. Similarly, Ünal Turhan et al. (2017) observed a decline in both lactic acid bacteria and *Lactobacillus rhamnosus* counts during the storage period in a study that used microencapsulated probiotic *L. rhamnosus*. Another study reported an increase in lactic acid bacteria during storage in probiotic sucuk and also noted that yeast/mold counts reached 4 log in the samples (Kozan and Sarıçoban, 2023).

3.6. Sensory parameters

The effects of the microbial culture and storage period on the sensory parameters of the sucuk samples are presented in Table 2. The storage period did not have a statistically significant effect on the sensory characteristics of the product ($P > 0.05$). However, the microbial culture factor had a highly significant impact ($P < 0.01$) on odor, taste, and overall acceptability parameters, but was not significant for color and texture parameters ($P >$

0.05). Average values for color and texture across all groups were above 7 points. However, the lowest average values in groups where the microbial culture factor was significant were found in samples using only the probiotic culture. The use of probiotic culture had a negative impact on product sensory parameters such as odor, taste, and overall acceptability. In contrast, a study using *L. casei* CRL-431 in sucuk production reported that this strain improved the sensory characteristics of the product (Bağdatlı and Kundakcı, 2016). The highest average values for odor and overall acceptability were found in the group using both starter and probiotic cultures. Additionally, the interaction of microbial culture and storage period had statistically significant effect on texture ($P < 0.01$) and taste ($P < 0.05$) parameters of the samples. Ergönül (2009) found that the microbial culture used in the production of probiotic turkey sucuk significantly affected the product's odor value and observed a decline in

odor value over the storage period, contrary to our findings. The researcher also stated, contrary to our results, that color and texture parameters were significantly affected by the microbial culture used and the duration of storage. In a study on sucuk production using *L. plantarum* and microencapsulated probiotic *L. rhamnosus*, it was determined that the microbial culture had no significant impact on the product's slice color, texture, pleasant odor, and overall acceptability scores (Ünal et al., 2017). Another study on sucuk production using microencapsulated probiotic *L. rhamnosus* also found no significant differences in slice color, texture, pleasant odor, and overall acceptability scores among the groups during the first 60 days of storage, but a significant decline in sensory scores was observed by the sixth month of storage. In the same study, it was determined that the varieties of cultures used did not have a significant impact on the sensory characteristics of the product (Ünal Turhan et al., 2017).

Table 2. Overall effect of microbial culture and storage period on the sensory and microbiological parameters of dry-fermented probiotic sucuk.

	Color	Texture	Odor	Taste	Overall acceptability	Lactic acid bacteria (log CFU/g)	<i>Micrococcus / Staphylococcus</i> (log CFU/g)	<i>Mold-Yeast</i> (log CFU/g)	<i>Lactocaseibacillus casei</i> (log CFU/g)
Microbial Culture (MC)									
SC	7,10±0,36 ^a	7,20±0,32 ^a	7,22±0,39 ^b	7,17±0,64 ^a	7,04±0,34 ^b	8,70±0,13 ^a	6,28±0,23 ^a	2,21±0,21 ^b	ND
PC	7,20±0,31 ^a	7,10±0,21 ^a	6,30±0,27 ^c	6,11±0,29 ^b	6,11±0,21 ^c	7,56±0,17 ^c	5,26±0,65 ^b	2,77±0,45 ^a	7,60±0,16 ^a
SC+PC	7,34±0,22 ^a	7,24±0,19 ^a	7,56±0,41 ^a	7,31±0,22 ^a	7,38±0,20 ^a	8,55±0,20 ^b	5,52±0,25 ^b	2,87±0,43 ^a	7,46±0,11 ^b
Significance	NS	NS	**	**	**	**	**	**	**
Storage Period (SP)									
Day 0	7,14±0,32 ^a	7,16±0,18 ^a	7,10±0,61 ^a	6,77±0,56 ^a	6,79±0,59 ^a	8,37±0,55 ^a	5,87±0,47 ^a	2,82±0,64 ^a	7,63±0,13 ^a
Day 30	7,20±0,36 ^a	7,11±0,28 ^a	6,84±0,55 ^a	6,74±0,67 ^a	6,89±0,62 ^a	8,27±0,54 ^{ab}	5,63±0,54 ^a	2,63±0,32 ^a	7,51±0,16 ^b
Day 60	7,30±0,25 ^a	7,28±0,26 ^a	7,13±0,79 ^a	7,08±0,81 ^a	6,86±0,65 ^a	8,17±0,57 ^b	5,57±0,77 ^a	2,40±0,32 ^a	7,45±0,14 ^b
Significance	NS	NS	NS	NS	NS	*	NS	NS	**
Interactions									
MCxSP	NS	**	NS	*	NS	NS	**	NS	NS

a-c: any two means in the same column having the same letters in the same section are not significantly different. * $p < 0.05$, ** $p < 0.01$, NS: not significant. SC: starter culture, PC: probiotic culture, SC+PC: starter culture + probiotic culture.

4. Conclusion

Considering the results obtained in this study, sucuk, a traditional dry fermented sausage, has proven to be a suitable matrix for the probiotic *L. casei* 431 during both production and storage. The findings demonstrated that *L. casei* 431 can successfully survive and maintain its viability

within the sucuk matrix throughout fermentation and storage, reinforcing its potential as a probiotic carrier food. However, it was observed that *L. casei* 431 negatively affected certain sensory properties of sucuk, particularly when used alone. Therefore, it is recommended to combine this strain with starter cultures to optimize both

sensory quality and product acceptance. Future research could explore additional probiotic strains to further enhance the sensory and functional characteristics of sucuk, paving the way for innovative fermented meat products with added health benefits.

Acknowledgment

This work was supported by Scientific Research Projects Coordination Unit of Istanbul Gedik

University with project number “GDK202410-04. The authors thank NAMET Food Industry and Trading Inc. (Kocaeli, Türkiye) for their support in the realization of this study. Ethical approval for this study was obtained from the Ethics Committee of Istanbul Gedik University (E-56365223-050.04-2024.137548.110)

Conflict of interest

The authors have no conflict of interest.

5. References

- Agüero, N. D. L., Frizzo, L. S., Ouwehand, A. C., Aleu, G., & Rosmini, M. R. (2020). Technological characterisation of probiotic lactic acid bacteria as starter cultures for dry fermented sausages. *Foods*, 9(5), 596.
- Álvarez, M., Andrade, M. J., Cebrián, E., Roncero, E., & Delgado, J. (2023). Perspectives on the probiotic potential of indigenous moulds and yeasts in dry-fermented sausages. *Microorganisms*, 11(7), 1746.
- Arief, I. I., Wulandari, Z., Aditia, E. L., & Baihaqi, M. (2014). Physicochemical and microbiological properties of fermented lamb sausages using probiotic *Lactobacillus plantarum* IIA-2C12 as starter culture. *Procedia Environmental Sciences*, 20, 352-356.
- Bağdatlı, A. B., and Kundakçı, A. (2013). Fermente et ürünlerinde probiyotik mikroorganizmaların kullanımı. *Celal Bayar University Journal of Science*, 9(1), 31-38.
- Bağdatlı, A., & Kundakci, A. (2016). Optimization of compositional and structural properties in probiotic sausage production. *Journal of Food Science and Technology*, 53, 1679-1689.
- Baumgart, J., Eigener, V., Firnhaber, J., Hildebrant, G., Reenen Hoekstra, E.S., Samson, R.A., Spicher, G., Timm, F., Yarrow, D. & Zschaler, R., 1993. Mikrobiologische Untersuchung von Lebensmitteln, (3., aktualisierte und erw. Aufl.), Hamburg, Germany.
- Bis-Souza, C. V., Barba, F. J., Lorenzo, J. M., Penna, A. B., & Barretto, A. C. S. (2019a). New strategies for the development of innovative fermented meat products: A review regarding the incorporation of probiotics and dietary fibers. *Food Reviews International*, 35(5), 467-484.
- Bis-Souza, C. V., Pateiro, M., Domínguez, R., Lorenzo, J. M., Penna, A. L. B., & da Silva Barretto, A. C. (2019b). Volatile profile of fermented sausages with commercial probiotic strains and fructooligosaccharides. *Journal of Food Science and Technology*, 56, 5465-5473.
- Bis-Souza, C. V., Penna, A. L. B., & da Silva Barretto, A. C. (2020). Applicability of potentially probiotic *Lactobacillus casei* in low-fat Italian type salami with added fructooligosaccharides: in vitro screening and technological evaluation. *Meat Science*, 168, 108186.
- Caplice, E., & Fitzgerald, G. F. (1999). Food fermentations: Role of microorganisms in food production and preservation. *International Journal of Food Microbiology*, 50(1-2), 131-149.
- Cavalheiro, C. P., Ruiz-Capillas, C., Herrero, A. M., Jiménez-Colmenero, F., de Menezes, C. R., & Fries, L. L. M. (2015). Application of probiotic delivery systems in meat products. *Trends in Food Science and Technology*, 46(1), 120-131.
- Chr. Hansen. (2005). *L. acidophilus, L. casei and Bifidobacteria in fermented milk products. Guidelines method for counting probiotic bacteria*. Bulletin F-6 LA LC BB March 2005/3:8. Chr. Hansen Denmark.
- Cocconcelli, P. S., & Fontana, C. (2008). Characteristics and applications of microbial starters in meat fermentations. In *Meat Biotechnology* (pp. 129-148). Springer, New York.

- Ergönül, B. (2009). Farklı probiyotik kültürler kullanılarak Hindi sucuğu üretimi ve kalite üzerine etkileri. Celal Bayar University Natural and Applied Sciences Institute, PhD thesis, Manisa
- Ergönül, B., & Kundakçı, A. (2011). Microbiological attributes and biogenic amine content of probiotic Turkish fermented sausage. *J. Verbr. Lebensm.*, 6, 49–56.
- Erkkilä, S., Suihko, M. L., Eerola, S., Petäjä, E., & Mattila-Sandholm, T. (2001). Dry sausage fermented by *Lactobacillus rhamnosus* strains. *International Journal of Food Microbiology*, 64(1-2), 205-210.
- FAO. (2006). *Probiotics in food: Health and nutritional properties and guidelines for evaluation* (FAO Food and Nutrition Paper, No. 85).
- Feiner, G. (2006). Meat composition and additives. In: *Meat Products Handbook Practical Science and Technology*, CRC Press: New York.
- Gökalp, H.Y., Kaya, M., Tülek Y. ve Zorba, Ö., 2010. Et ve ürünlerinde kalite kontrolü ve laboratuvar uygulama kılavuzu. Atatürk Üniv. Yayınları: Erzurum.
- Holko, I., Hrabě, J., Šalaková, A., & Rada, V. (2013). The substitution of a traditional starter culture in mutton fermented sausages by *Lactobacillus acidophilus* and *Bifidobacterium animalis*. *Meat Science*, 94(3), 275-279.
- Kaban, G., & Kaya, M. (2008). Identification of lactic acid bacteria and Gram-positive catalase-positive cocci isolated from naturally fermented sausage (sucuk). *Journal of Food Science*, 73(8), M385-M388.
- Kaban, G., & Kaya, M. (2009). Effects of *Lactobacillus plantarum* and *Staphylococcus xylosus* on the quality characteristics of dry fermented sausage “sucuk”. *Journal of Food Science*, 74(1), S58-S63.
- Karshoğlu, B., Çiçek, Ü. E., Kolsarıcı, N., & Candoğan, K. (2014). Lipolytic changes in fermented sausages produced with turkey meat: effects of starter culture and heat treatment. *Korean journal for food science of animal resources*, 34(1), 40.
- Kaya, M., & Aksu, M. I. (2005). Effect of modified atmosphere and vacuum packaging on some quality characteristics of sliced ‘sucuk’ produced using probiotics culture. *Journal of the Science of Food and Agriculture*, 85(13), 2281-2288.
- Kaya, M., Güllüce, M., Kaban, G., Çınar, K., Karadayı, M., Bozoğlu, C., Sayın, B., & Alaylar, B. (2015). Geleneksel sucuklardan izole edilen laktik asit bakteri ve koagülaz negatif stafilokok suşlarının starter kültür olarak kullanım imkânları. TAGEM-13/ARGE/7 (Gelişme Raporu), Gıda Tarım ve Hayvancılık Bakanlığı, Ankara.
- Kaya, M., & Kaban, M. (2019). Fermente et ürünleri. İçinde: *Gıda Biyoteknolojisi* (Necla ARAN Ed.) 7. Basım. Nobel Yayıncılık: İstanbul.
- Khan, M. I., Arshad, M. S., Anjum, F. M., Sameen, A., & Gill, W. T. (2011). Meat as a functional food with special reference to probiotic sausages. *Food Research International*, 44(10), 3125-3133.
- Klingberg, T. D., & Budde, B. B. (2006). The survival and persistence in the human gastrointestinal tract of five potential probiotic lactobacilli consumed as freeze-dried cultures or as probiotic sausage. *International Journal of Food Microbiology*, 109, 157-159.
- Klingberg, T. D., Axelsson, L., Naterstad, K., Elsser, D., & Budde, B. B. (2005). Identification of potential probiotic starter cultures for Scandinavian-type fermented sausages. *International Journal of Food Microbiology*, 105(3), 419-431.
- Kołożyn-Krajewska, D., & Dolatowski, Z. J. (2009). Probiotics in fermented meat products. *Acta Scientiarum Polonorum Technologia Alimentaria*, 8(2), 61-74.
- Kołożyn-Krajewska, D., & Dolatowski, Z. J. (2012). Probiotic meat products and human nutrition. *Process Biochemistry*, 47(12), 1761-1772.
- Kozan, H. İ., & Sarıçoban, C. (2023). Effect of oat bran addition on the survival of selected probiotic strains in Turkish fermented sausage during cold storage. *Food Bioscience*, 54, 102820.
- Kröckel, L. (2013). The role of lactic acid bacteria in safety and flavour development of meat and

meat products. In *Lactic Acid Bacteria - R and D for Food, Health and Livestock Purposes* (pp. 129-152).

Kumar, P., Chatli, M. K., Verma, A. K., Mehta, N., Malav, O. P., Kumar, D., & Sharma, N. (2017). Quality, functionality, and shelf life of fermented meat and meat products: A review. *Critical reviews in food science and nutrition*, 57(13), 2844-2856.

Lemon, D.W. 1975. "An Improved TBA Test for Rancidity New Series Circular" No: 21. Halifax-Laboratory, Halifax, Nova, Scotia.

Neffe-Skocińska, K., Wójciak, K., & Zielińska, D. (2016). Probiotic microorganisms in dry fermented meat products. *Probiotics and prebiotics in human nutrition and health*, 279-300.

Öztürk Er, H. (2002). The possibilities of the use of the probiotic bacteria in sucuk, Turkish dry fermented sausages. (Master's thesis, Atatürk University, Graduate Institute of Science). Erzurum, Turkey.

Pavli, F. G., Argyri, A. A., Chorianopoulos, N. G., Nychas, G. J. E., & Tassou, C. C. (2020). Effect of *Lactobacillus plantarum* L125 strain with probiotic potential on physicochemical, microbiological and sensorial characteristics of dry-fermented sausages. *LWT*, 118, 108810.

Soyer, A., Ertaş, A. H., & Üzümcüoğlu, Ü. (2005). Effect of processing conditions on the quality of naturally fermented Turkish sausages (sucuks). *Meat science*, 69(1), 135-141.

Tukel, O., & Sengun, I. (2024). Production of probiotic fermented salami using *Lactobacillus rhamnosus*, *Lactiplantibacillus plantarum*, and *Bifidobacterium lactis*. *Journal of Food Science*, 89(5), 2956-2973.

Ünal Turhan, E., Erginkaya, Z., & Selli, S. (2017). The effect of microencapsulated *Lactobacillus rhamnosus* and storage period on aroma properties of Turkish dry-fermented sausage (sucuk). *Journal of Food Measurement and Characterization*, 11, 2131-2141.

Ünal Turhan, E., Erginkaya, Z., Benli, H., Akın, M. B., & Ağçam, E. (2019). The effects of microencapsulated *L. rhamnosus* and storage on

biogenic amine amount of sucuk. *Gıda*, 44(5), 819-825.

Ünal, E., Erginkaya, Z., Polat, S., & Özer, E. A. (2017). Design of probiotic dry fermented sausage (sucuk) production with microencapsulated and free cells of *Lactobacillus rhamnosus*. *Turkish Journal of Veterinary and Animal Sciences*, 41(5), 598-603.

Yalınkılıç, B., Kaban, G., & Kaya, M. (2012). The effects of different levels of orange fiber and fat on microbiological, physical, chemical, and sensorial properties of sucuk. *Food Microbiology*, 29(2), 255-259.

Yalınkılıç, B. (2009). Sucuk üretiminde portakal lifi kullanımı. Atatürk University, Natural and Applied Sciences Institute, Master of Science Thesis, Erzurum.

Yılmaz Topcam, M. M., Arslan, B., & Soyer, A. (2024). Sucuk, Turkish-Style Fermented Sausage: Evaluation of the Effect of Bioprotective Starter Cultures on Its Microbiological, Physicochemical, and Chemical Properties. *Applied Microbiology*, 4(3), 1215-1231.