

## Two new records for the Turkish Diptera fauna from Izmir Province, Western Turkey

Hasan Sungur CIVELEK\*

### Summary

This study was carried out during 1997-1998 in Izmir province, western Turkey. The specimens of Diptera were collected from culture and non-culture plants and identified. *Oscinella nartshukiana* Beschovski, 1978 (Chloropidae) and *Scaptomyza graminum* (Fallén, 1823) were first records for the Turkish Diptera fauna.

**Key words:** *Oscinella nartshukiana*, *Scaptomyza graminum*, new records, Turkey

**Anahtar sözcükler:** *Oscinella nartshukiana*, *Scaptomyza graminum*, yeni kayıtlar, Türkiye

### Introduction

Diptera is one of the most abundant orders, consisting of approximately 120.000 species belonging to 4 suborders and 188 families (Thompson & Fairman, 1999). These insects are a major component of virtually all non-marine ecosystems; only the arctic and antarctic ice caps are without flies. In addition to being a large order, flies are also of great economic importance. Some studies about Turkish Diptera fauna has been carried out by Anonymous (1987), Campobasso et al. (1999), Civelek et al. (2000), Hayat & Alaoğlu (1990), Hayat & Özbek (1994), Kara (1999a,b), Kılıç (1992, 1996), Öncüer (1991). At the present time 328 species belonging to 18 families have been recorded from Turkey. In this paper, it was aimed to contribute to the Turkish Diptera fauna.

### Material and Methods

This study was carried out during 1997-1998, primarily from costal areas in the Izmir region (western Turkey). The specimens were collected between April

---

\* Mugla University, Ortaca Vocational School, 48600 Ortaca Mugla, Turkey

e-mail: chasan@mu.edu.tr

Alınış (Received): 18.03.2002

and July in both years by using sweeping nets on cultured and wild plants. These species were identified by Dr. John Deeming (National Museum & Galleries of Wales, Cardiff, UK). The representative specimens were deposited at the Laboratory in the Ortaca Vocational School, Mugla, Turkey.

## Results and Discussion

As a result of this study, *Oscinella nartshukiana* Beschovski, 1978 (Chloropidae) and *Scaptomyza graminum* (Fallén, 1823) (Drosophilidae) were recorded new in Turkey.

### Family: Chloropidae Meigen, 1803

World wide, 3 subfamilies, 1.200 species in the world, 300 species in Central Europe. Body 1-3 mm. The imagines are usually small flight slow-acting to tiny and quite. The body is yellow or black touched or inconspicuously black, the eyes however magnificently colored. Larvae, typical fly larvae, are hard and quite mobile. They mines predominantly in the impulses and stems of most diverse Gramineae. The larvae of some kinds live however predatory (Haar, 2000).

### Genus: *Oscinella* Becker, 1909

### Species: *Oscinella nartshukiana* Beschovski, 1978

**Description:** A small black species, wing length 1 mm (Figure 1); Ocellar triangle extending to or almost to anterior margin of frons, shining but dusted on ocellar prominence. Gena about as deep as width of fore femur. Third antennal segment with a slight dorsoapical angle and with arista dark. Mesonotum dusted with humerus shining. Male cerci semicircular in outline (rather than triangular, which is the case in *O. sensu stricto* species). Halteres pale yellow and legs dirty yellow on at least mid tibia and basal tarsal segments of mid and hind leg.




Figure 1. *Oscinella nartshukiana*, wing (Scale line: 1 mm).

**Hosts:** Unknown. This species was collected first time on grasses by Beschovski (1978).

**Distribution:** This species was described from a locality close to the Bulgarian capital city of Sofia (Beschovski, 1978).

**Material examined:** *O. nartshukiana* was collected in Bornova (Izmir) on wild grass in 3.10.1997 (1♂) by sweeping.

## Family: Drosophilidae Loew, 1862

Drosophilidae is one of the largest families of acalyptrate Diptera and are found worldwide. Members of the family are attracted to and feed upon a great variety of fermenting substances, hence the common name, vinegar, pomace, or fruit flies (the latter not to be confused with the true fruit flies of the family Tephritidae). Oviposition and larval breeding are, however, more highly specialized in fruits or flowers, slime fluxes, decaying leaves, or fungi. Some species are leaf mining, commensal, or even predatory. By the end of 1984, 62 genera and 2.822 described species of this family were known worldwide, and at least 4.000 species were predicted to exist. Evenhuis & Okada (2000) record over 1.000 species in 65 genera and subgenera from the Australasian/Oceanian Regions.

**Genus: *Scaptomyza*** Hardy, 1849

**Species: *Scaptomyza graminum*** (Fallén, 1823)

**Description:** Blackish species, wing length from 2.0 to 2.2 mm. Frons yellow, orbital bristles with 1 ors-2 ori; jowl narrow 1/8 of eyes; antennae entirely yellow, 3rd segment small and round, arista with 5 long fringe. Acrostichal bristles in 4 irregular rows in wings (Figure 2) costa extending  $R_{4+5}$ , last section of  $M_{3+4}$  5 times shorter than penultimate, squamae white margin and fringe dark, halteres white. All legs yellow.




Figure 2. *Scaptomyza graminum*, wing (Scale line: 1 mm).

**Hosts: *Aquilegia vulgaris*, *Brassica campestris*, *B. oleracea*, *B. rapa*, *Petasites officinalis* and *Triticum pratense*** (Stalker, 1945).

**Distribution:** Egypt, Formosa, U.S.A. (Stalker, 1945), Russia (Kas'yanov, 1976), Uzbekistan (Maca, 1992), England (Pennington, 1997), Faroe Islands, Iceland and Denmark (Meier et al., 2000), Hawaii (Tamura et al., 2000) and India (Thompson, 2000).

**Material examined: *S. graminum*** was collected in Menderes (İzmir) on *Triticum vulgare* in 23.5.1998 (1♀) by sweeping.

## Özet

### İzmir İlinden Türkiye Diptera faunası için iki yeni kayıt

Bu çalışma 1997-1998 yılları arasında İzmir ili ve çevresinde yürütülmüştür. Diptera takımına bağlı türler kültürü yapılan ve yapılmayan bitkiler üzerinden toplanmış ve teşhis

edilmiştir. *Oscinella nartshukiana* Beschovski, 1978 (Chloropidae) ve *Scaptomyza graminum* (Fallén, 1823) (Drosophilidae) Türkiye Diptera faunası için yeni kayıt türlerdir.

## Acknowledgements

Many thanks to Dr. John C. Deeming for the identifications of the species.

## References

- Anonymous, 1987. Türkiye'de Tarım, Orman ve Evcil Hayvanlarda Hayvansal Kökenli Zararlı ve Yararlı Türlerin Bilimsel ve Türkçe İsimleri. Tarım ve Köyşleri Bakanlığı, Mesleki Yayınlar, No: 8, Ankara, 120 s.
- Beschovski, V., 1978. Subdivision of the genus *Oscinella* Becker, 1909 (Diptera, Chloropidae) with description of a new species. *Acta Zoologica Bulgarica*, **10**: 21-29.
- Campobasso, G., E. Colonelli, L. Knutson, G. Terragitti & M. Cristofaro, 1999. Wild Plants and Their Associated Insects in the Port Royal Road, Springfield, Palaearctic Region, Primarily Europe and the Middle East. U. S. Department of Agriculture, Agricultural Research Service, ARS-147, 249 pp.
- Civelek, H.S., J. Deeming, & F. Önder, 2000. Some new records for the Turkish leafminers (Diptera: Agromyzidae) fauna from Izmir province. *Türk. entomol. derg.*, **24**(1) : 17-26.
- Evenhuis, N.L. & T. Okada, 2000. Family Drosophilidae.  
<http://www.bishopmuseum.org/bishop/ento/aocat/drosophilidae.html>
- Haar, J., 2000. Insects of Cedar Creek Natural History Area. Order Diptera, Family Chloropidae, (Grass Flies). <http://cedarcreek.umn.edu/insects/029094n.html>
- Hayat, R. & Ö. Alaoğlu, 1990. Fauna of Syrphidae (Diptera) in the vicinity of Erzurum (I), Syrphinae. *Türk. entomol. derg.*, **14**(3) : 173-182.
- Hayat, R. & H. Özbek, 1994. New records of robber flies (Diptera:Asilidae) for the Turkish fauna. *Türk. entomol. derg.*, **18**(4): 241-244.
- Kara, K., 1999a. Tokat çevresinde saptanan Tachininae (Diptera: Tachinidae) türleri. *Türk. entomol. derg.*, **23**(2): 121-134.
- Kara, K. 1999b. Tokat çevresinde saptanan Dexiinae (Diptera:Tachinidae) türleri. *Türk. entomol. derg.*, **23**(3): 203-210.
- Kas'yanov, A.I., 1976. Pests of rice crops. *Zaashchita Rasteneii*, **8**: 28-31.
- Kılıç, A.Y., 1992. Eskişehir ve çevresi Tabanidae (Diptera) faunası. *Türk. entomol. derg.*, **16**(3): 161-180.
- Kılıç, A.Y., 1996. Çamlıyayla (İçel) Tabanidae (Diptera) faunası üzerinde araştırmalar. *Türk. entomol. derg.*, **20**(2): 123-135.
- Maca, J., 1992. Addition to the fauna of Drosophilidae, Camillidae, Curtonotidae and Campichoetidae (Diptera) of Soviet Middle Asia. *Annotationes Zoologicae et Botanicae*. No. 210: 8.
- Meier, R., W. Buck, K.D. Klas, V. Michelsen & J.F.T. Petersen, 2000. Collection inventory of the Diptera collection of the Zoological Museum, Entomology Department, University of Copenhagen, Denmark. <http://www.zmuc.dk/EntoWeb/collections-databaser/Diptera/dipterasites.htm>
- Öncüer, C., 1991. Türkiye'nin Bitki Zararlısı Böceklerinin Parazit ve Predatör Kataloğu. Ege Üniversitesi, Ziraat Fakültesi Yayınları, No: 505, 354 s.

- Pennington, M., 1997. Diptera. True flies. Insects in Shetland.  
<http://www.wildlife.shetland.co.uk/insects/flies.htm>
- Stalker, H.D., 1945. On the biology and genetics of *Scaptomyza graminum* Fallén (Diptera, Drosophilidae). **Genetics**, **30**: 266-280.
- Tamura, K., K. Toba, J. Park & K. Aotsuka, 2000. Origin of Hawaiian Drosophilids inferred from alcohol dehydrogenase gene sequences. Dept. Biol. Tokyo Metropolitan Univ. 1-1 Minami-Ohsawa, Hachioji-shi, Tokyo 192-03, Japan.  
<http://evolgen.biol.metro-u.ac.jp/WWW/Adh/>
- Thompson, C. & J.E. Fairman, 1999. The Diptera Site. Systematic Entomology Laboratory, ARS, USDA Dept. Of Systematic Biology, NMNH, SI.  
<http://www.sel.barc.usda.gov/Diptera/diptera.htm>
- Thompson, J., 2000. Cumulative Drosophilid Species Index. DIS Volumes 1 to 71.  
<http://fly.bio.indiana.edu:7081/allied-data/lk/DIS/index-species.txt>