

Determination of Aphididae (Homoptera) fauna of Diyarbakır Province of Turkey*

Selime ÖLMEZ BAYHAN**

Mehmet Rifat ULUSOY***

Seval TOROS****

Summary

Aphids were surveyed in Diyarbakır province of Turkey between 1998-2000. As a result of the survey studies performed, 61 species of aphids were determined on different host plants. Their distribution in Turkey, host plants, collecting dates and occurrence were given for each species. Of them, 30 species were the first record for Diyarbakır province.

Key words: Aphididae, Diyarbakır, fauna, Turkey

Anahtar sözcükler: Aphididae, Diyarbakır, fauna, Türkiye

Introduction

Aphids (Homoptera: Aphididae), particularly from the subfamily Aphidinae, are usually the most important pests of agricultural and horticultural crops in the world, damaging plants by sucking phloem sap and by transmitting viral diseases (Kennedy et al., 1962; Blackman & Eastop, 1984; Conti, 1985).

The first records of Turkish aphids were given by Trotter (1903), and Fahringer (1922). Then there were conducted many studies on aphids in Turkey (İyriboz & İleri, 1941; Alkan, 1953; Düzgüneş & Tuatay, 1956; Börner & Heinze, 1957; Bodenheimer & Swirski, 1957; Tuatay & Remaudiére, 1964; Düzgüneş et al., 1982; Elmalı ve Toros, 1996; Toros et al., 2002). These studies have primarily concentrated on faunistic. However, no investigation on aphids which have wide

* Bu çalışma Çukurova Üniversitesi Araştırma Fonu tarafından desteklenen FBE.99.YL.6 nolu yüksek lisans tezininin bir bölümüdür.

** Dicle Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Diyarbakır
e-mail: solmez@mail.cu.edu.tr

*** Çukurova Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, 01330-Balcalı, Adana
e-mail: mrulusoy@mail.cu.edu.tr

**** Ankara Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Dışkapı, Ankara
Alınmış (Received): 03.11.2003

range has been performed in South East Anatolia region before. The aim of this study was to determine Aphididae species in cultivated and noncultivated areas in Diyarbakır province of Turkey.

Material and Methods

Aphid samples on various hosts were collected from different locations (Central province, Bismil, Çınar, Çermik, Çüngüş, Dicle, Eğil, Ergani, Hazro, Kocaköy, Lice and Silvan) in cultivated and noncultivated areas of Diyarbakır province, which is located Southeast Anatolia region of Turkey between 1998-2000. Plant parts having aphids were cut and placed separately in polyethylene bags which were brought to the laboratory. The sample of aphids transferred into ephendorph tubes containing % 70 ethyl alcohol with a fine brush. The method of Hille Ris Lambers (1950) was used for mounting technique. Identifications of the species were made by Prof. Dr. S. Toros using Bodenheimer and Swirski (1957), Hille Ris Lambers (1945, 1947a, 1947b, 1973), Börner and Heinze (1957), Tuatay and Remaudiére (1964), Shaposhnikov (1964), Stroyan (1957, 1961, 1963, 1977, 1984), Eastop (1971), Bissel (1978) and Blackman and Eastop (1984, 1994). The determined species were classified according to Remaudiére & Remaudiére (1997). Asterisk indicates new record for Diyarbakır province.

Results and Discussion

In this study, a total of 63 species were determined belonging to 6 subfamilies (Anoeciinae, Aphidinae, Chaitophorinae, Lachninae, Myzocallidinae and Pemphiginae), 11 tribe (Anoecini, Aphidini-Aphidina, Aphidini-Rhopalosiphini, Macrosiphini, Chaitophorini, Cinarini, Lachnini, Myzocallidini, Eriosomatini, Fordini and Pemphigini), 31 genus, and 7 subgenus of Aphididae.

Anoeciinae: Anoecini

Anoecia Koch, 1857

Anoecia corni (Fabricius, 1775)

Distribution in Turkey: Ankara, Bitlis, Diyarbakır, İçel, İstanbul, Tekirdağ (Tuatay & Remaudiére, 1964; Düzgüneş et al., 1982; Toros et al., 2002).

Material examined: *Lycopersicum esculentum* L., Çermik, 1.7.1999.

Occurrence: Rarely.

Aphidinae: Aphidini: Aphidina

Aphis Linnaeus, 1758

* *Aphis affinis* del Guercio, 1911

Distribution in Turkey: Adana, Ankara, Bitlis, İçel, Kahramanmaraş (Tuatay & Remaudiére, 1964; Düzgüneş et al., 1982; Zeren, 1989; Aslan, 2002; Toros et al., 2002).

Material examined: *Mentha* sp.; Central province, 4.5.1999, 12.5.1999; Çınar, 8.6.1999; Hani, 18.5.1999.

Occurrence: Occasionally.

* **Aphis avicularis** (Hille Ris Lambers, 1931)

Distribution in Turkey: Ankara, Van (Tuatay & Remaudiére, 1964).

Material examined: *Polygonum aviculare*, Çüngüş, 11.6.1999.

Occurrence: Rarely.

* **Aphis callunae** Theobald, 1915

Distribution in Turkey: Ankara (Tuatay & Remaudiére, 1964).

Material examined: *Erica* sp., Çınar, 16.7.1999.

Occurrence: Rarely.

* **Aphis chloris** Koch, 1854

Distribution in Turkey: Van (Tuatay & Remaudiére, 1964).

Material examined: *Hypericum empetrifolium* Willd. roots, Bismil, 14.6.1999.

Occurrence: Rarely.

Aphis craccivora Koch, 1854

Distribution in Turkey: Adana, Ankara, Antalya, Aydın, Balıkesir, Bolu, Burdur, Bursa, Diyarbakır, East Mediterranean region, Giresun, İstanbul, Kahramanmaraş, Manisa, Mardin, Şanlıurfa, Van (Bodenheimer & Swirski, 1957; Çanakçıoğlu, 1967; Giray, 1974; Düzgüneş et al., 1982; Akkaya & Uygun, 1996; Toros et al., 1996; Aslan, 2002; Toros et al., 2002).

Material examined: *Senecio vulgaris* L., Central province, 6.12.1998, 4.5.1999; *Centaurea* sp., Central province, 5.5.1999; *Robinia pseudacacia* L., Central province, 5.5.1999, 12.5.1999; Bismil, 17.7.1998, 21.7.1998, 07.7.1999; Çermik, 12.6.1999; Çınar, 27.8.1998; Çüngüş, 11.6.1999; Ergani, 21.7.1998; Hani, 18.5.1999; Kocaköy, 11.8.1999; *Capsella bursa-pastoris* (L.), Central province, 5.5.1999; *Amaranthus* sp., Central province, 5.6.1999; *Phaseolus vulgaris* L., Central province, 5.6.1999; Çermik, 12.6.1999; *Citrullus vulgaris*, Bismil, 12.6.1999; 7.7.1999; *Glycyrrhiza glabra* L., Bismil, 12.6.1999, 7.7.1999; Hani, 18.5.1999; *Solanum nigrum* L., Çermik, 12.6.1999; *Mentha* sp., Hani, 18.5.1999; *Lycopersicum esculentum* L., Hani, 18.5.1999; *Tribulus terrestris* L., Hani, 21.7.1999; undetermined weed, Lice, 3.5.1998; *Portulaca oleracea* L., Lice, 3.8.1999.

Occurrence: Commonly.

* **Aphis davletshinae** Hille Ris Lambers, 1966

Distribution in Turkey: Malatya, Ankara (Tuatay & Remaudiére, 1964; Düzgüneş et al., 1982).

Material examined: Undetermined weed, Hazro, 18.6.1999.

Occurrence: Rarely.

Aphis fabae Scopoli, 1763

Distribution in Turkey: Aegean region, Adana, Ankara, Antalya, Diyarbakır, Edirne, East Mediterranean region, Hatay, İçel, İstanbul, Kırklareli, Konya, Niğde,

Şanlıurfa, Tekirdağ, Van (İyriboz, 1937; Schimitschek, 1944; Bodenheimer & Swirski, 1957; Tuatay & Remaudiére, 1964; Çanakçıoğlu, 1967; Giray, 1974; Çanakçıoğlu, 1975; Düzgüneş et al., 1982; Zeren, 1989; Toros et al., 1996; Özdemir & Toros, 1997; Bayhan & Özder, 1999; Toros et al., 2002).

Material examined: *Cirsium arvense* (L.), Central province, 6.12.1998, 3.5.1999, 4.5.1999, 6.5.1999, 12.5.1999, 13.5.1999, 18.5.1999; Hani, 18.5.1999; *Urtica* sp., Central province, 6.12.1998, 3.5.1999; *Centaurea* sp., Central province, 4.5.1999; *Gallium* sp., Central province, 4.5.1999; *Osyris alba* L., Central province, 13.5.1999; *Amaranthus* sp., Central province, 30.11.1998; *Rumex crispus* L., Central province, 6.12.1998; Çüngüş, 11.6.1999; *Nicotiana tabacum* L., Bismil, 7.7.1999; *Medicago sativa* L., Çüngüş, 11.6.1999; *G. glabra*, Ergani, 21.7.1999; *P. vulgaris*, Ergani, 21.7.1999; *P. oleracea* L., Hazro, 1.8.1999; *L. esculentum*, Hazro, 1.8.1999.

Occurrence: Commonly.

***Aphis fabae solanella* Theobald, 1914**

Distribution in Turkey: Ankara, Diyarbakır, East Mediterranean region, İstanbul, İzmir, Van (Giray, 1974; Çanakçıoğlu, 1975; Düzgüneş et al., 1982; Zeren, 1989; Toros et al., 1996; Toros et al., 2002).

Material examined: *Solanum* sp., Central province, 5.6.1999; *S. nigrum*, Bismil, 25.8.1998, 7.7.1999, 25.8.1998; Çınar, 27.8.1998; Dicle, 28.7.1999, Ergani, 21.7.1999; Hani, 18.5.1999; Kocaköy, 4.6.1998, 11.8.1999.

Occurrence: Commonly.

*** *Aphis frangulae* Kaltenbach, 1845**

Distribution in Turkey: Adana, Kahramanmaraş (Aslan, 2002; Toros et al., 2002; Tezcan, 2003).

Material examined: *Lycopersicum esculentum* L., Çermik, 1.7.1999.

Occurrence: Rarely.

***Aphis gossypii* Glover, 1877**

Distribution in Turkey: Cosmopolite (Bodenheimer & Swirski, 1957; Toros et al., 2002; Tezcan, 2003).

Material examined: *Gossypium* sp., Central province, 20.10.1998, Bismil, 7.7.1999, 25.8.1999; Çermik, 29.7.1998; *T. terrestris*, Central province, 20.10.1998; *P. vulgaris*, Central province, 3.5.1999, 4.5.1999; Çınar, 1.7.1999; *Cucurbita pepo* L., Central province, 4.5.1999; Bismil, 14.8.1999; Eğil, 30.7.1999; Ergani, 21.7.1998, 16.8.1999; Hazro, 18.6.1999; Lice, 17.7.1998; Kocaköy, 2.8.1998, 11.8.1999; *Ficus carica* L., Central province, 12.5.1999, 5.6.1999; Çermik, 12.6.1999; *Solanum melongena* L., Bismil, 17.7.1998, 25.8.1998; Çermik, 1.7.1999; Eğil, 30.7.1999; Kocaköy, 11.8.1999; *Capsicum annuum* L., Bismil, 25.8.1998; Çermik, 1.7.1999, 21.7.1999; Ergani, 27.6.1998; *Cucumis sativus* L., Bismil, 7.7.1999; Eğil, 30.7.1999; Ergani, 21.7.1999; Kocaköy, 11.8.1999; *Citrullus vulgaris* L., Bismil, 7.7.1999;

Çermik, 29.7.1998, 1.7.1999; Ergani, 21.7.1998; Kocaköy, 11.8.1999;
Mentha piperita L., Çınar, 8.6.1999; Çermik, 12.6.1999, ***Cucumis melo*** L.,
Eğil, 30.7.1999; ***Polygonum*** sp., Ergani, 21.7.1999; ***P. oleracea***, Ergani,
21.7.1999; ***A. hybridus*** L., Ergani, 21.7.1999; ***A. blibides*** L., Ergani,
21.7.1999; ***Hibiscus esculentus*** L., Lice, 3.8.1999; ***L. esculentum***, Silvan,
1.8.1999; ***C. bursa-pastoris***, Ergani, 3.5.1999.

Occurrence: Commonly.

* ***Aphis nerii*** Boyer de Fonscolombe, 1841

Distribution in Turkey: Ankara, Antalya, East Mediterranean region, İzmir,
Kahramanmaraş (Bodenheimer & Swirski, 1957; Çanakçıoğlu, 1967; Giray,
1974; Aslan 2002; Toros et al., 2002).

Material examined: ***Nerium olander*** L., Central province, 18.11.1998;
Hazro, 18.6.1999.

Occurrence: Occasionally.

Aphis pomi de Geer, 1773

Distribution in Turkey: Cosmopolite (Lodos, 1982; Toros et al., 2002).

Material examined: ***Cydonia vulgaris*** Pers., Central province, 12.5.1999;
Malus communis (L.), Central province, 14.5.1999; Hazro, 18.6.1999.

Occurrence: Frequently.

Aphis punicae Passerini, 1863

Distribution in Turkey: Cosmopolite (Tuatay & Remaudiére, 1964;
Çanakçıoğlu, 1975; Toros et al., 2002).

Material examined: ***Punica granatum*** L., Central province, 3.5.1999,
6.5.1999; Bismil, 7.7.1999; Çermik, 29.7.1998, 1.7.1999; Çınar, 4.10.1998,
8.6.1999; Ergani, 27.11.1998.

Occurrence: Commonly.

* ***Aphis spiraecola*** Patch, 1914

Distribution in Turkey: Ankara, East Mediterranean region (Tuatay &
Remaudiére, 1964; Düzgüneş et al., 1982; Yumruktepe & Uygun, 1994; Toros
et al., 2002).

Material examined: ***Prunus domestica*** L., Bismil, 7.7.1999.

Occurrence: Rarely.

* ***Aphis tirucallis*** Hille Ris Lambers, 1954

Distribution in Turkey: Ankara, East Mediterranean region (Düzgüneş et
al., 1982; Toros et al., 2002).

Material examined: ***Euphorbia*** sp., Hazro, 18.6.1999.

Occurrence: Rarely.

* ***Aphis urticata*** J.F. Gmelin, 1790

Distribution in Turkey: Ankara, East Mediterranean region (Bodenheimer &
Swirski, 1957; Toros et al., 2002).

Material examined: *Urtica* sp., Central province, 3.5.1999.

Occurrence: Rarely.

Aphidinae: Aphidini: Aphidina

Hyalopterus Koch, 1854

Hyalopterus amygdali (Blanchard, 1840)

Distribution in Turkey: Ankara, Diyarbakır, East Mediterranean region, Kahramanmaraş, Siirt, Şanlıurfa (Düzungün et al., 1982; Uygun et al., 1995; Aslan, 2002; Toros et al., 2002).

Material examined: *Prunus amygdalus* (L.), Central province, 3.5.1999; Çüngüş, 11.6.1999; Hazro, 18.6.1999.

Occurrence: Frequently.

Hyalopterus pruni (Geoffroy, 1762)

Distribution in Turkey: Ankara, Antalya, East Mediterranean region, Gaziantep, İzmir, Kahramanmaraş (Düzungün & Tuatay, 1956; Bodenheimer & Swirski, 1957; Çanakçıoğlu, 1975; Düzungün et al., 1982; Aslan, 2002, Toros et al., 2002).

Material examined: *Phragmites* sp., Central province, 20.6.1998; Bismil, 7.7.1999; Kocaköy, 2.8.1998; undetermined weed, Central province, 3.5.1999; *P. domestica*, Central province, 5.5.1999, 14.5.1999, 18.5.1999; Çermik, 1.7.1999; Çınar, 4.10.1998; Lice, 3.5.1998; Silvan, 4.5.1999; *Prunus persica* (L.), Central province, 5.5.1999, 12.5.1999, 14.5.1999, 18.6.1999; Çermik, 29.7.1998; Çüngüş, 11.6.1999; Hazro, 18.5.1999; *Prunus armeniaca* L., Central province, 5.5.1999, 12.5.1999; *Prunus cerasus* L., Central province, 18.6.1999; Hani, 18.5.1999.

Occurrence: Commonly.

Rhopalosiphum Koch, 1854

* ***Rhopalosiphum nymphaeae*** (Linnaeus, 1761)

Distribution in Turkey: Cosmopolite (Çanakçıoğlu, 1975).

Material examined: *Prunus domestica* L., Central province, 6.7.1998; *P. cerasus*, Çınar, 4.10.1998.

Occurrence: Occasionally.

Rhopalosiphum maidis (Fitch, 1856)

Distribution in Turkey: Cosmopolite (Çanakçıoğlu, 1975; Toros et al., 2002).

Material examined: *Zea mays* L., Central province, 28.7.1999; Bismil, 7.7.1999; Çermik, 1.7.1999; Dicle, 12.7.1999; Ergani, 21.7.1999; Hazro, 18.6.1999; Lice, 3.8.1999; Kocaköy, 4.7.1998; *Sorghum* sp., Bismil, 7.7.1999; Lice, 3.8.1999.

Occurrence: Commonly.

Rhopalosiphum padi (Linnaeus, 1758)

Distribution in Turkey: Ankara, Diyarbakır, East Mediterranean region, Edirne, Kahramanmaraş, Konya, Kırklareli, Mardin, Ordu, Tatvan, Tekirdağ,

Sanliurfa, Şırnak (Tuatay & Remaudiére, 1964; Düzgüneş et al., 1982; Elmalı & Toros, 1994; Kırın, 1994; Tozlu & Alaoğlu, 1994; Bayhan & Ozder, 1999; Aslan 2002; Toros et al., 2002).

Material examined: *Triticum durum* L., Central province, 5.5.1999; *Hordeum sativum* L., Central province, 5.5.1999; *Sorghum* sp., Çınar, 8.6.1999.

Occurrence: Occasionally.

Macrosiphini

Acyrtosiphon Mordvilko, 1914

Acyrtosiphon pisum (Harris, 1776)

Distribution in Turkey: Cosmopolite (Lodos, 1982; Toros et al., 2002).

Material examined: *M. sativa*, Çüngüş, 11.6.1999.

Occurrence: Rarely.

Aulacorthum Mordvilko, 1914

* *Aulacorthum solani* (Kaltenbach, 1843)

Distribution in Turkey: East Mediterranean region, İstanbul (Tuatay, 1988; Toros et al., 2002)

Material examined: *L. esculentum*, Central province, 6.1.1998.

Occurrence: Rarely.

Brachycaudus van der Goot, 1913

Brachycaudus helichrysi (Kaltenbach, 1843)

Distribution in Turkey: Adana, Adiyaman, Ankara, Diyarbakır, East Mediterranean region, Gaziantep, İstanbul, Kahramanmaraş, Mardin, Siirt (Tuatay & Remaudiére, 1964; Tuatay et al., 1972; Giray, 1974; Düzgüneş et al., 1982; Uygun et al., 1995; Aslan, 2002; Toros et al., 2002).

Material examined: *P. domestica*, Central province, 7.12.1998, 4.5.1999, 5.5.1999, 6.5.1999; Hani, 18.5.1999; Hazro, 6.5.1999; Silvan, 18.5.1999;

Calendula officinalis L., Lice, 3.8.1999.

Occurrence: Commonly.

* *Brachycaudus (Acaudus) cardui* (Linnaeus, 1758)

Distribution in Turkey: Ankara, Bitlis, East Mediterranean region, Trabzon (Bodenheimer & Swirski, 1957; Tuatay & Remaudiére, 1964; Çanakçıoğlu, 1975; Toros et al., 2002).

Material examined: *Prunus domestica* L., Silvan, 18.5.1999; *Cirsium* sp., Hani, 5.5.1999.

Occurrence: Occasionally.

* *Brachycaudus (Acaudus) lycnidis* (Linnaeus, 1758)

Distribution in Turkey: Çankırı (Tuatay & Remaudiére, 1964).

Material examined: *Silene* sp., Çüngüş, 12.6.1999.

Occurrence: Rarely.

* ***Brachycaudus (Appelia) tragopogonis*** (Kaltenbach, 1843)

Distribution in Turkey: Ankara, East Mediterranean region, Eskişehir, Malatya, Kahramanmaraş, Kırşehir (Düzungüneş & Tuatay, 1956; Tuatay & Remaudiére, 1964; Düzungüneş et al., 1982; Aslan 2002; Toros et al., 2002).

Material examined: *Tragopogon longirostris* var. *longirostris* L., Çınar, 8.6.1999.

Occurrence: Rarely.

Brachycaudus (Thuleaphis) amygdalinus (Schouteden, 1905)

Distribution in Turkey: Ankara, Elazığ, Isparta (Bodenheimer & Swirski, 1957; Tuatay & Remaudiére, 1964).

Material examined: *Amygdalus communis* L., Central province, 3.5.1999; *C. officinalis*, Hani, 5.5.1999.

Occurrence: Occasionally.

Brevicoryne van der Goot., 1915

Brevicoryne brassicae (Linnaeus, 1758)

Distribution in Turkey: Ankara, East Mediterranean region, Hatay, İzmir, Kahramanmaraş, Van, Şanlıurfa (Düzungüneş & Tuatay, 1956; Giray, 1974; Düzungüneş et al., 1982; Zeren, 1989; Avcı & Özbek, 1991; Uygun et al., 1995; Toros et al., 1996; Aslan, 2002; Toros et al., 2002).

Material examined: *Brassica oleracea* var. *capitata*, Central province, 2.12.1998; Bismil, 7.7.1999; Hazro, 18.6.1999; Lice, 3.8.1999.

Occurrence: Frequently.

Dysaphis Börner, 1931

Dysaphis* (s.str.) *devecta (Walker 1849)

Distribution in Turkey: Adana, Ankara, İçel, Malatya, Van (Tuatay & Remaudiére, 1964; Düzungüneş et al., 1982; Toros et al., 1996; Toros et al., 2002).

Material examined: *M. communis*, Central province, 4.5.1999.

Occurrence: Rarely.

Dysaphis (Pomaphis) plantaginea (Passerini, 1860)

Distribution in Turkey: Ankara, Antalya, Çanakkale, Diyarbakır, East Mediterranean region, Gaziantep, Giresun, Gümüşhane, Elazığ, Niğde, Isparta, İzmir, Kahramanmaraş, Kayseri, Sakarya, Samsun, Şanlıurfa, Tekirdağ (Bodenheimer & Swirski, 1957; Tuatay, 1990; Aslan 2002; Toros et al., 2002).

Material examined: *Cydonia vulgaris* Pers., Central province, 5.5.1999; *M. communis*, Central province, 6.12.1998, 12.5.1999, 14.5.1999; Ergani, 27.11.1998; Lice, 3.5.1998.

Occurrence: Commonly.

Dysaphis (Pomaphis) pyri (Boyer de Fonscolombe, 1841)

Distribution in Turkey: Adana, Ankara, Isparta, İçel, İzmir, Konya, Niğde (Bodenheimer & Swirski, 1957; Tuatay & Remaudiére, 1964; Giray, 1974; Düzungüneş et al., 1982; Toros et al., 1996; Toros et al., 2002).

Material examined: *Cydonia vulgaris* Pers., Central province, 5.5.1999;
Prunus communis L., Central province, 14.5.1999; Çınar, 8.6.1999.

Occurrence: Occasioanly.

Hayhurstia del Guercio, 1917

* ***Hayhurstia atriplicis*** (Linnaeus, 1761)

Distribution in Turkey: Ankara, Antalya, East Mediterranean region, Edirne, Elazığ, Eskişehir, Van (Bodenheimer & Swirski, 1957; Tuatay & Remaudière, 1964; Çanakçıoğlu, 1967; Tuatay, 1990; Toros et al., 2002).

Material examined: *Chenopodium album* L., Bismil, 7.7.1999.

Occurrence: Rarely.

Hyadaphis Kirkaldy, 1904

* ***Hyadaphis tataricae*** (Aizenberg, 1935)

Distribution in Turkey: Ankara, Kahramanmaraş (Toros, 1986; Aslan, 2002).

Material examined: *Lonicera tatarica* L., Central province, 3.5.1999.

Occurrence: Rarely.

Hyperomyzus Börner, 1933

* ***Hyperomyzus lactcea*** (Linnaeus, 1758)

Distribution in Turkey: East Mediterranean region, Eskişehir, İstanbul (Tuatay & Remaudière, 1964; Toros et al., 2002).

Material examined: *Sonchus* sp., Hazro, 18.6.1999.

Occurrence: Rarely.

Lipaphis Mordilko, 1928

* ***Lipaphis erysimi*** (Kaltenbach, 1843)

Distribution in Turkey: Adana, Gaziantep (Tuatay & Remaudière, 1964; Toros et al., 2002).

Material examined: *Raphanus sativus* L., Central province, 2.12.1998.

Occurrence: Rarely.

Macrosiphoniella del Guercio, 1911

Macrosiphoniella artemisiae (Boyer de Fonscolombe, 1841)

Distribution in Turkey: Ankara, Kahramanmaraş (Tuatay, 1990; Aslan 2002).

Material examined: *Artemisia* sp., Central province, 12.9.1999; Çüngüş, 18.6.1999.

Occurrence: Occasionally.

Macrosiphum Passerini, 1860

* ***Macrosiphum funestum*** (Macchiati, 1855)

Distribution in Turkey: Artvin, Bolu, Giresun, Kastamonu, Rize (Çanakçıoğlu, 1975; Tuatay, 1990).

Material examined: *Spiraea vanhouttei* L., Central province, 12.5.1999; *Centaurea* sp., Hani, 18.5.1999.

Occurrence: Occasionally.

* *Macrosiphum rosae* (Linnaeus, 1758)

Distribution in Turkey: Ankara, East Mediterranean region, İstanbul, Isparta, İzmir, Van (Bodenheimer & Swirski, 1957; Giray, 1974; Düzgüneş et al., 1982; Toros et al., 1996; Toros et al., 2002).

Material examined: *Rosa* sp., Central province, 3.5.1999, 4.5.1999, 5.5.1999, 5.6.1999, 12.5.1999; Bismil, 7.7.1999; Çermik, 12.6.1999; Hani, 18.5.1999; Hazro, 18.6.1999; Kocaköy, 8.4.1999.

Occurrence: Commonly.

Myzus Passerini, 1860

Myzus cerasi (Fabricius, 1775)

Distribution in Turkey: Ankara, East Mediterranean region, West and Middle Anatolia, Gümüşhane, İstanbul, İznik, Kars, Kahramanmaraş, Rize (Bodenheimer & Swirski, 1957; Lodos, 1982; Aslan, 2002; Toros et al., 2002).

Material examined: *Prunus avium* L., Çüngüş, 11.6.1999.

Occurrence: Rarely.

Myzus (Nectarosiphon) persicae (Sulzer, 1776)

Distribution in Turkey: Cosmopolite (Lodos, 1982; Toros et al., 2002).

Material examined: *M. communis*, Central province, 30.11.1998; Çınar, 8.6.1999; *C. annuum*, Central province, 7.12.1998; *Duranta repens* L., Central province, 3.5.1999; *P. persica*, Central province, 5.5.1999; *Nicotiana tabacum* L., Bismil, 7.7.1999; Çermik, 1.7.1999; Kocaköy, 11.8.1999; Lice, 3.8.1999; *L. esculentum*, Çınar, 8.6.1999; Hani, 18.5.1999; *P. domestica*, Çınar, 8.6.1999; *C. sativus*, Çınar, 8.6.1999; *Mentha* sp., Hani, 18.5.1999; *C. annuum*, Hani, 18.5.1999; Kocaköy, 11.8.1999.

Occurrence: Commonly.

Ovatus van der Goot, 1913

* *Ovatus crataegarius* (Walker, 1850)

Distribution in Turkey: Ankara (Tuatay & Remaudiére, 1964).

Material examined: *Crataegus orientalis* L., Hani, 18.5.1999.

Occurrence: Rarely.

Ovatus insitus (Walker, 1849)

Distribution in Turkey: Ankara, Diyarbakır, Elazığ, İzmir, Niğde, Sakarya, Trabzon (Tuatay & Remaudiére, 1964).

Material examined: *C. vulgaris*, Central province, 6.12.1998; 5.5.1999.

Occurrence: Occasionally.

Sitobion Mordvilko, 1914

Sitobion avenae (Fabricius, 1775)

Distribution in Turkey: Adana, Ağrı, Ankara, Aydın, Bolu, Çanakkale, Denizli, Diyarbakır, Erzurum, Isparta, İzmir, Kırklareli, Konya, Malatya, Mardin, Niğde, Samsun, Sinop, Şanlıurfa, Sırmak, Yozgat, Tekirdağ, Van (İyriboz & İleri, 1941; Tuatay et al., 1972; Düzgüneş et al., 1982; Tuatay, 1991; Elmalı & Toros, 1994; Kiran, 1994; Toros et al., 1996; Özder & Toros, 1999; Toros et al., 2002).

Material examined: *Triticum durum* L., Central province, 5.5.1999; Çüngüş, 11.6.1999; Hazro, 18.6.1999.

Occurrence: Frequently.

Uroleucon Mordvilko, 1914

* **Uroleucon** (s.str.) *sonchi* Linnaeus, 1767

Distribution in Turkey: Adana, Ankara, Antalya, Aydın, Çanakkale, Hatay, İstanbul, İzmir, Kırklareli, Sakarya (Bodenheimer & Swirski, 1957; Tuatay & Remaudiére, 1964; Giray, 1974; Çanakçıoğlu, 1975; Düzgüneş et al., 1982; Tuatay, 1991; Toros et al., 2002).

Material examined: *Sonchus arvensis* L., Central province, 4.5.1999, 5.6.1999; Hani, 18.5.1999; Hazro, 18.6.1999; *Spiraea vonhautti* L., Hani, 18.5.1999.

Occurrence: Frequently.

Uroleucon (Uromelan) carthami (Hille Ris Lambers, 1848)

Distribution in Turkey: Adana, Ankara, Mersin, Muğla (Tuatay & Remaudiére, 1964; Toros et al., 2002).

Material examined: From undetermined weed, Lice, 3.5.1998.

Occurrence: Rarely.

Chaitophorinae: Chaitophorini

Chaitophorus Koch, 1854

Chaitophorus leucomelas Koch, 1854

Distribution in Turkey: Adana, Ankara, Adapazarı, Bozova (Şanlıurfa), Diyarbakır, İçel, İstanbul, Kaş (Antalya), Kırşehir, Milas (Muğla), Niğde, Van (Bodenheimer & Swirski, 1957; Tuatay & Remaudiére, 1964; Çanakçıoğlu, 1975; Toros et al., 1996; Toros et al., 2002).

Material examined: *Populus* sp., Central province, 18.11.1998, 14.8.1999, 28.10.1998; Ergani, 27.11.1998; Hani, 18.5.1999; Hazro, 18.6.1999; Lice, 3.8.1999; Kocaköy, 11.8.1999.

Occurrence: Commonly.

* **Chaitophorus salijapanicus niger** Mordvilko, 1929

Distribution in Turkey: Ankara, Antalya, Burdur, Tatvan (Tuatay & Remaudiére, 1964; Çanakçıoğlu, 1967; Tuatay et al., 1972; Düzgüneş et al., 1982; Toros et al., 2002).

Material examined: *Salix* sp., Central province, 9.11.1998, 12.5.1999; Çınar, 8.6.1999; Ergani, 27.11.1998.

Occurrence: Frequently.

Lachninae: Cinarini

Eulachnus del Guercio, 1909

* **Eulachnus tuberculostematus** (Theobald, 1915)

Distribution in Turkey: Van, Adana (Toros et al., 1996; Toros et al., 2002).

Material examined: *Pinus* sp., Central province, 4.5.1999.

Occurrence: Rarely.

Lachnini

Lachnus Burmeister, 1835

Lachnus roboris (Linnaeus, 1758)

Distribution in Turkey: Ankara, Artvin, Diyarbakır, İstanbul, Kahramanmaraş, Mardin (Düzungüneş & Tuatay, 1956; Bodenheimer & Swirski, 1957; Çanakköy, 1967; Aslan, 2002).

Material examined: *Pinus* sp., Çermik, 1.7.1999; Dicle, 28.7.1999; Ergani, 21.7.1999; Lice, 3.8.1999.

Occurrence: Commonly.

Tuberolachnus Mordvilko, 1909

Tuberolachnus salignus (J. F. Gmelin, 1790)

Distribution in Turkey: Adana, Ankara, Diyarbakır, Hazar Lake (Elazığ), İstanbul, Kahramanmaraş, Kars, Van (Bodenheimer & Swirski, 1957; Tuatay & Remaudiére, 1964; Çanakköy, 1967; Düzungüneş et al., 1982; Aslan, 2002; Toros et al., 2002).

Material examined: *Salix* sp., Central province, 20.10.1998; Eğil, 27.11.1998.

Occurrence: Occasionally.

Myzocallidinae: Myzocallidini

Chromaphis Walker, 1870

* **Chromaphis juglandicola** Kaltenbach, 1843

Distribution in Turkey: Antalya, Middle and West Anatolia, East Mediterranean region, Kahramanmaraş (Lodos, 1982; Aslan, 2002; Toros et al., 2002).

Material examined: *Juglans* sp., Central province, 5.5.1999; Hani, 18.5.1999; Hazro, 18.6.1999.

Occurrence: Occasionally.

Panaphis Kirkaldy, 1904

* **Panaphis juglandis** (Goeze, 1778)

Syn.: **Callaphis juglandis** Walker, 1870

Distribution in Turkey: Ankara, Artvin, East Mediterranean region, İstanbul, İzmir, Niğde, Antalya (Kaş), Trabzon (Düzungüneş & Tuatay, 1956; Bodenheimer & Swirski, 1957; Çanakköy, 1967; Düzungüneş et al., 1982; Toros et al., 2002).

Material examined: *Juglans* sp., Çermik, 1.7.1999; Çüngüş, 11.6.1999;
Hazro 18.6.1999.

Occurrence: Occasionally.

Myzocallis Passerini, 1860

* ***Myzocallis coryli*** (Goetze, 1778)

Distribution in Turkey: Ankara, İstanbul, Trabzon (Fahringer, 1922; Düzgüneş & Tuatay, 1956; Bodenheimer & Swirski, 1957; Çanakçıoğlu, 1967).

Material examined: *Corylus avellana* L., Çüngüş, 11.6.1999.

Occurrence: Rarely.

Myzocallidinae: Myzocallidini

Eriosoma Leach, 1818

Eriosoma lanigerum (Hausmann, 1802)

Distribution in Turkey: Cosmopolite wherever apples are found (Alkan, 1953).

Material examined: *Malus communis*, Central province, 14.5.1999; Çermik, 1.7.1999.

Occurrence: Occasionally.

Fordini

Forda van Heyden, 1837

Forda formicaria van Heyden, 1837

Distribution in Turkey: Adana, Burdur, Bursa, Diyarbakır, İçel, İstanbul, İznik, Konya (Trotter, 1903; Schimitschek, 1944; Çanakçıoğlu, 1975; Elmali ve Toros, 1996; Toros et al., 2002).

Material examined: Half-moon-shaped galls on *Pistacia vera* L., Silvan, 1.8.1999.

Occurrence: Rarely.

Kaltenbachiella Schouteden, 1906

* ***Kaltenbachiella pallida*** (Haliday, 1838)

Distribution in Turkey: Ankara, Bandırma, Bursa, Eskişehir, İstanbul, İzmir, Söke (Aydın), Trabzon (Alkan, 1953; Çanakçıoğlu, 1967; Tuatay, 1967).

Material examined: *Ulmus campestris* L., Central province, 9.7.1999.

Occurrence: Scarcely.

Patchiella Tullgren, 1927

***Patchiella* sp.**

Material examined: From undetermined weed, Central province, 30.11.1998.

Occurrence: Rarely.

Pemphigini

Pemphigus Hartig, 1839

Pemphigus bursarius (Linnaeus, 1758)

Distribution in Turkey: Ankara, Bursa, İstanbul, Eskişehir, Kahramanmaraş, Konya, Trabzon (Çanakçıoğlu, 1975).

Material examined: *Populus* sp., Çınar, 8.6.1999,

Occurrence: Rarely.

* *Pemphigus immunis* Buckton, 1896

Distribution in Turkey: Adana, Afyon, Ankara, Antalya, Artvin, Beyşehir, Burdur, Denizli, Eskişehir, İçel, İstanbul, İzmir, Kahramanmaraş, Niğde, Trabzon (Alkan, 1953; Çanakçıoğlu, 1967; Aslan, 2002; Toros et al., 2002).

Material examined: *Ulmus* sp., Central province, 13.6.1998; Dicle, 28.7.1999; Hani, 18.5.1999; Hazro, 18.6.1999.

Occurrence: Commonly.

Özet

Diyarbakır İli Aphididae (Homoptera) faunasının saptanması

Diyarbakır İlinde 1998-2000 yıllarında Aphididae familyasına bağlı türlerin surveyı yapılmıştır. Survey sonucunda farklı konukçu bitkiler üzerinde 63 tür saptanmış olup, bunların Türkiye'deki dağılımları, toplandıkları yer ve tarihler ile görünüm sıklıkları açıklanmıştır. Bunlardan 30 tür Diyarbakır İli lokal faunası için yeni kayıt niteliğindedir.

References

- Akkaya, A. & N. Uygun, 1996. Diyarbakır ve Şanlıurfa İlleri yazlık sebze ekosistemindeki insecta faunası. Türkiye III. Entomoloji Kongresi Bildirileri, 24-28 Eylül 1996, Ankara, 423-430.
- Alkan, B., 1953. Türkiye'nin Zoosesid (Zooecid)'leri (kökeni hayvansal bitki urları) üzerinde çalışmalar. Ankara Zir. Fak. Yayın., 185-291.
- Aslan, M.M., 2002. Kahramanmaraş İlinde Aphidoidea (Homoptera) Türleri ile Bunların Parazitot ve Predatörlerinin Saptanması. Çukurova Üniversitesi Fen Bilimleri Enstitüsü Bitki Koruma Bölümü Basılmamış Doktora Tezi, 136 s.
- Avcı, Ü. & H. Özbek, 1991. Erzurum İlinde Lahana unlu yaprakbitinin (*Brevicoryne brassicae* L.) (Homoptera: Aphididae) doğal düşmanları üzerine araştırmalar. *Türk. entomol. derg.*, 15 (1): 37-41.
- Bayhan, E. & N. Özder, 1999. Investigations on the corn pests and their natural enemies in the Thrace Region Province. XIV th International Plant Protection Congress (IPPC), July 25-30 1999, Jerusalem, Israel, 66.
- Bissel, T.L., 1978. Aphids on Juglandaceae in North America. Martland Agricultural Experiment Station 78 pp.
- Blackman, R.L. & V.F. Eastop, 1984. Aphids on the Word's Crops. An Identification quide. A Wiley, intencience Publication, 466 pp.
- Blackman, R. L. & V. F. Eastop, 1994. Aphids on the World's Trees. An Identification and information quide CAB International, 986+16 pp.
- Bodenheimer, F.S. & E. Swirski, 1957. The Aphidoidea of the Middle East. The Weigmann Science Press of Isreal, Jerusalem, 378 pp.
- Börner, C. & K. Heinze, 1957. Handbuch der Pflanzenkrankheiten. Band V, 4. Lieferung: Homoptera II. Teil, 577 pp.
- Conti, M., 1985. Transmission of Plant Viruses by Leafhoppers and Planthoppers. A Wiley Interscience Publication, New York, p. 289-307.

- Çanakçıoğlu, H., 1967. Türkiye'de Orman Ağaçlarına Arız Olan Yaprakbitleri (Aphidoidea) Üzerine Araştırmalar. T.C. Tarım Bakanlığı, Orman Gn. Md. Yayınlarından Sıra No: 466, Seri No: 22, VIII, 151 s.
- Çanakçıoğlu, H., 1975. The Aphidoidea of Turkey. İstanbul Üni. Orman Fak. Yayınları, Yayın No: 1751, O. F. Yayın No: 189, 309 s.
- Düzgüneş, Z. & N. Tuatay, 1956. Türkiye Aphidleri. Ziraat Vekaleti, Ank. Zir. Enst. Md. Sayı: 4, 63 s.
- Düzgüneş, Z., S. Toros, N. Kılınçer & B. Kovancı, 1982. Ankara İlinde Bulunan Aphidoidea Türlerinin Parazitoit ve Predatörlerinin Tespiti. Tarım ve Orm. Bak. Zir. Múc ve Zir. Kar. Gn. Md. Yayın Şb., 251 s.
- Eastop, V. F., 1971. Keys for the identification of *Acyrtosiphon* (Hemiptera: Aphididae). **Bull. of British Museum (Natural History), Entomology**, **26** (1): 1-115.
- Elmalı, M. & S. Toros, 1994. "Konya İlinde buğday tarlalarında yaprakbiti doğal düşmanlarının tespiti üzerinde araştırmalar. Türkiye III. Biyolojik Mücadele Kongresi Bildirileri, 25-28 Ocak 1994, İzmir, 13-28.
- Elmalı, M. & S. Toros, 1996. Konya İlinde Buğdaylarda Aphidoidea Türleri ve Bulunuş Oranları. Ank. Üniv. Zir. Fak. Yayınları No 1454. Bilimsel Araştırma ve İncelemeler 802, 40 s.
- Fahringer, J., 1922. Eine Rhynchotenausbeute aus der Türkei, Kleinasien und den Benachbarten Gebieten. **Konowia**, **1**: 137-44, 296-307.
- Giray, H., 1974. İzmir İli ve çevresinde Aphididae (Homoptera) familyası türlerine ait ilk liste ile burlann konukçu ve zarar şekilleri hakkında notlar. **Ege Üni. Zir. Fak. Derg.**, **11** (1): 39-69.
- Hille Ris Lambers, D., 1945. de Bloedvlekkenluis van Appel, *Sappaphis devecta* (Wlk.). **Tijdschr. Ov. Plantenziekt.**, **51**: 57-66.
- Hille Ris Lambers, D., 1947 a. Contributions to a monograph of the Aphididae of Europe. III. **Temminckia**, **7**: 179-319.
- Hille Ris Lambers, D., 1947 b. On some mainly Western European Aphids. **Zoologische Mededeelingen**, **28**: 291-333.
- Hille Ris Lambers, D., 1950. On mounting aphids and other soft skinned insects. **Entomologische Berichten**, **13**: 54-55.
- Hille Ris Lambers, D., 1973. Notes on some oriental Aphids. **Orient. Insects**, **7**: 239-258.
- İyriboz, N., 1937. Pamuk Hastalıkları.- Ankara Ziraat Vekaleti Neş. U. S. 237, Pamuk Bürosu, 85 s.
- İyriboz, N. & M. İleri, 1941. "Aphidoidea, s.117-119". Hububat Hastalıkları. İzmir, 182 s.
- Kennedy, J. S., M. F. Day & V. F. Eastop, 1962. A Conspectus of Aphids as Vector of Plant Viruses. Commonwealth Inst. Ent. London. 114 pp.
- Kıran, E., 1994. Güneydoğu Anadolu Bölgesi hububat ekiliş alanlarında görülen yaprakbiti türleri ve doğal düşmanları üzerinde çalışmalar. Türkiye III. Biyolojik Mücadele Kongresi Bildirileri, 25-28 Ocak 1994, İzmir, 29-37.
- Lodos, N., 1982. Türkiye Entomolojisi II (Genel, Uygulamalı ve Faunistik). Ege Üniv. Zir. Fak. Yay. No: 429, 591 s.
- Özdemir, I. & S. Toros, 1997. Ankara parklarında mevsimlik süs bitkilerinde zararlı Aphidoidea (Hom.) türleri. **Türk. Ent. Derg.**, **21** (4): 283-298.
- Özder, N. & S. Toros, 1999. Tekirdağ İlinde buğdaylarda zarar yapan yaprakbiti türlerinin doğal düşmanları üzerinde araştırmalar. Türkiye IV. Biyolojik Mücadele Kongresi Bildirileri, 26-29 Ocak 1999, Adana, 501-512.
- Remaudiére, G. & M. Remaudiére, 1997. Catalogue des Aphididae du Monde (of the World's Aphididae) Homoptera, Aphidoidea, Preface par V. F. Eastop, INRA Editions, 473 pp.
- Schimitschek, E., 1944. Forstinsekten der Türkei und ihre Umwelt. Volk. Reich, Prag, Berlin, 371 pp.

- Shaposhnikov, G. K., 1964. Suborder Aphidinae-Plant Lice. (In Keys to The Insects of the European Part on the USSR. Editör: G. Bei-Bienko. Moskow and Leningrad, 616-799.
- Stroyan, H.L.G., 1957. The British species of **Sappaphis** Matsumura, Part 1, Introduction and subgenus **Sappaphis** Sensu Stricto. Her majesty's Stationery Office, London, pp: 59.
- Stroyan, H.L.G., 1961. Identification of Aphids living on citrus. **FAO Plant Protection Bull.** **9** (4): 45-65.
- Stroyan, H.L.G., 1963. The British Species of **Dysaphis** Börner (**Sappaphis auctti** nec Mats.) Part II. Her majesty's Stationery Office, London, 119 pp.
- Stroyan, H.L.G., 1977. Homoptera, Aphidoidea (Part), Chatophoridae and Callaphidae. Handbooks for the Identification of British Insects. Vol. II, Part 6. Royal Entom. Soc. Of London, 232 pp.
- Stroyan, H.L.G., 1984. Aphids-Pterocommatinae and Aphidinae (Aphidini) Homoptera: Aphididae. Handbooks for the Identification of British Insects. Vol. II, Part 6. Royal Entom. Soc. of London, 232 pp.
- Tezcan, F., 2003. Pamuk yaprakbiti [**Aphis gossypii**] Glover (Homoptera: Aphididae)]'nın İzmir ve Manisa İlleri'nde Pamuktaki Biyolojisi ve Populasyon Dalgalanmalarını Etkileyen Faktörler Üzerinde Araştırmalar. İzmir, 54 pp.
- Toros, S., 1986. Türkiye faunası için yeni bir tür; **Hyadaphis tataricae** (Aizenberg) (Homoptera: Aphididae). **Türk. Bit. Kor. Derg.**, **10**: 3, 141-148.
- Toros, S., B. Yaşar, M.S. Özgökçe & İ. Kasap, 1996. Van ilinde Aphidoidea (Homoptera) üstfAMILYASINA bağlı türlerin saptanması üzerinde çalışmalar. Türkiye III. Entomoloji Kongresi Bildirileri, 24-28 Eylül 1996, Ankara, 549-556.
- Toros, S., N. Uygur, R. Ulusoy, S. Satar, I. Özdemir, 2002. Doğu Akdeniz Bölgesi Aphidoidea TÜRLERİ. T.C. Tarım ve Köyişleri Bakanlığı, Tarımsal Araştırmalar Genel Müdürlüğü Yayınları, 108 s.
- Tozlu, G. & Ö. Alaoğlu, 1994. Ordu İl misir (**Zea mays** L.) ekim alanlarında bulunan fitofag ve predatör böcek türleri. **Türk. Ent. Derg.**, **18** (1): 51-64.
- Trotter, A., 1903. Galle della Paninsola Balsanica e Asia Minore. **Nuovo G. Bot. Ital.**, **10**: 6-54, 202-232.
- Tuatay, N. & G. Remaudiére, 1964. Première contribution au catalogue des Aphididae (Hom.) de la Turquie. **Rev. de Path. Veg. Et Ent. Agr. de Pr.**, **43** (4): 243-278.
- Tuatay, N., 1967. Nebat Koruma Müzesi Böcek Kataloğu (1961-1966). T.C. Tar. Bak. Zir. Muc. ve Zir. Kar. Gen. Md. Yayınları Meslek Kitaplar Serisi, 66 s.
- Tuatay, N., 1988. Aphids (Homoptera: Aphididae) of Turkey I. Aphidinae: Macrosiphini (Part I). **Plant Protection Bulletin**, **28**: (1-2): 1-28.
- Tuatay, N., 1990. Aphids (Homoptera: Aphididae) of Turkey II. Aphidinae: Macrosiphini (Part II). **Plant Protection Bulletin**, **30**: (1-4): 29-44.
- Tuatay, N., 1991. Aphids (Homoptera: Aphididae) of Turkey III. Aphidinae: Macrosiphini (Part III). **Plant Protection Bulletin**, **31**: (1-4): 3-18.
- Tuatay, N., A. Kalkandelen & N. Çağatay, 1972. Insect catalogue of plant protection museum (1961-1971). Turkish Ministry of Agriculture, Research Serial, 119 pp.
- Uygur, N., H. Başpinar, E. Sekeroğlu, S. Kornoşor, A.F. Özgür, İ. Karaca, M.R. Ulusoy & C. Kazak, 1995. GAP alanında zirai mücadele politikasına esas teşkil edecek zararlı ve yararlıların saptanması. GAP Bölgesi Bitki Koruma Sorunları ve Çözüm Önerileri (Şanlıurfa) Sempozyumu Bildirileri, 27-29 Nisan 1995, Şanlıurfa, 99-119.
- Yumruktepe, R. & N. Uygur, 1994. Doğu Akdeniz Bölgesi turuncgil bahçelerinde saptanmış yaprakbiti (Hom., Aphididae) türleri ve doğal düşmanları. Türkiye III. Biyolojik Mücadele Kongresi Bildirileri, 25-28 Ocak 1994, İzmir, 1-12.
- Zeren, O., 1989. Çukurova Bölgesinde sebzelerde zararlı olan yaprakbitileri (Aphidoidea) türleri, konukçuları, zararları ve doğal düşmanları üzerinde araştırmalar. Çukurova Üniversitesi Fen Filmleri Enstitüsü Bitki Koruma Bölümü Doktora tezi, 205 s.